



Short communication

Detection of *bla*_{CTX-M}-type genes in complex class 1 integrons carried by *Enterobacteriaceae* isolated from retail chicken meat in Brazil



Tiago Casella^a, María Margarita Rodríguez^b, Juliana Tiemi Takahashi^c, Barbara Ghiglione^b, Milena Dropa^d, Ednei Assunção^e, Maurício Lacerda Nogueira^c, Nilton Lincopan^{e,f}, Gabriel Gutkind^b, Mara Corrêa Lelles Nogueira^{c,*}

^a Instituto de Biociências, Letras e Ciências Exatas, Universidade Estadual Paulista "Júlio de Mesquita Filho", São José do Rio Preto, SP, Brazil

^b Universidad de Buenos Aires, Facultad de Farmacia y Bioquímica, Ciudad Autónoma de Buenos Aires, BA, Argentina

^c Faculdade de Medicina de São José do Rio Preto, São José do Rio Preto, SP, Brazil

^d Departamento de Prática de Saúde Pública, Faculdade de Saúde Pública, Universidade de São Paulo, São Paulo, SP, Brazil

^e Departamento de Microbiologia, Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, SP, Brazil

^f Departamento de Análises Clínicas, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, São Paulo, SP, Brazil

ARTICLE INFO

Article history:

Received 5 August 2014

Received in revised form 11 November 2014

Accepted 6 December 2014

Available online 11 December 2014

Keywords:

Poultry

Chicken meat

CTX-M

ISCR1

ABSTRACT

CTX-M-type extended-spectrum β -lactamase (ESBL)-producing *Enterobacteriaceae* have been increasingly identified in humans and animals, and their potential transmission by contaminated food has been highlighted. In this study, we report for the first time the isolation of multidrug-resistant (MDR) *Escherichia coli*, *Klebsiella pneumoniae* and *Proteus mirabilis* strains harboring *bla*_{CTX-M-2} or *bla*_{CTX-M-8} gene variants in chicken meat sold in markets in southeast Brazil. In this regard, the genetic environment of the *bla*_{CTX-M-2} gene is composed of a complex class 1 integron and an ISCR1-associated sequence with *dfr* and/or *aadA* gene cassettes located within the variable region. In summary, chicken meat may be a reservoir of MDR *Enterobacteriaceae* harboring *bla*_{CTX-M}-type genes, which is a public health concern.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Healthcare-associated and community-acquired infections caused by extended-spectrum β -lactamases (ESBLs)-producing bacteria have been increasingly reported worldwide (Pitout and Laupland, 2008). Although the predominant ESBL families have been TEM, SHV and CTX-M (Bush and Jacoby, 2010), in the last years, the incidence of the CTX-M-type has risen dramatically (Wellington et al., 2013). In fact, currently the CTX-M-2, CTX-M-8 and CTX-M-15 variants are widely disseminated all over the world within *Enterobacteriaceae* from humans and animals, and their potential transmission by contaminated food, mainly chicken meat, has been highlighted (Egervärn et al., 2014; Kawamura et al., 2014; Kiiru et al., 2012). These enzymes are encoded by *bla*_{CTX-M}-type gene associated with mobile genetic elements, including insertion sequences ISEcp1 type and ISCR1, most carried on transferable plasmids (Zhao and Hu, 2013). In Brazil, although the production of

the CTX-M enzymes has become the most prevalent mechanism of acquired resistance to broad-spectrum cephalosporins in gram-negative bacteria from clinical samples, poultry, food-producing animals, hospital wastewater and urban rivers (Aizawa et al., 2014; Carvalho-Assef et al., 2014; Ferreira et al., 2014; Peirano et al., 2011; Silva et al., 2013; Tollentino et al., 2011), there are no data available regarding the detection of CTX-M-producing bacteria in chicken meat intended for sale on the Brazilian market, which is curious, since the presence of CTX-M-2- and CTX-M-8-producing *E. coli* in raw chicken and broiler meat imported into European countries from South America countries, including Brazil, has been previously reported (Dhanji et al., 2010; Egervärn et al., 2014; Warren et al., 2008). So, the aim of this study was to investigate the presence of ESBL-producing bacteria and the genetic background of genes encoding ESBLs in retail chicken meat from markets, in southeastern Brazil.

2. Materials and methods

2.1. Sampling

We investigated the presence of ESBL-producing bacteria in four pork, three beef, and eight chicken meat samples, collected between March 2011 and July 2013 from seven traditional marketplaces in the

* Corresponding author at: Departamento de Doenças Dermatológicas, Infeciosas e Parasitárias, Faculdade de Medicina de São José do Rio Preto, Av. Brigadeiro Faria Lima, 5416, São José do Rio Preto, SP, CEP 15090-000, Brazil. Tel.: +55 17 3201 5900; fax: +55 17 3229 1777.

E-mail address: ml.nogueira@famerp.br (M.C.L. Nogueira).

state of São Paulo, south-eastern Brazil. The samples were immediately transported to the laboratory for microbiological analysis. For bacterial isolation, sub samples of 200 g were dispensed on sterile plastic bags (Whirl-Pak, Nasco, WI, USA) containing 200 mL of Tetrathionate broth plus 5% of Brilliant-Green broth, and incubated at 37 °C to 24 h. After, 1 mL of this diluent was submitted to serial dilution on Lactose broth and inoculated on MacConkey agar (Oxoid, Hampshire, England) and incubated at 37 °C for 24 h.

2.2. Bacterial identification and antibiotic susceptibility testing

Bacterial identification and initial susceptibility testing and detection of ESBL producers were performed using the Vitek 2 system (bioMérieux, France). Identification of ESBL producers was based on the comparison of MIC values for ceftazidime, ceftriaxone and cefepime tested alone and in association with clavulanic acid, according to CLSI recommendation (CLSI, 2011). Additionally, antimicrobial susceptibility profiles were determined by the disc diffusion method using commercially available discs (Oxoid, UK). Interpretation was performed according to criteria of the Clinical and Laboratory Standard Institute (CLSI, 2008, 2011) or following EUCAST guidelines (EUCAST, 2011). *Escherichia coli* ATCC 25922 was used as reference strain.

2.3. Identification of ESBL genes and PCR mapping of class 1 integron

ESBL genes were screened by PCR as previously described (Tollentino et al., 2011). Sequencing was performed using the BigDye Terminator v3.1 (Applied Biosystems, Foster City, CA) with the ABI PRISM 3130 automated sequencer (Applied Biosystems, Foster City, CA). The sequences were annotated in DS Gene 2.0 software and compared with GenBank.

The architecture of the class 1 integron was assessed using the following primers: *int1* gene, I5 (5'-ACCGCCAACTTTCAGCACAT-3') and I3 (5'-GCGTTCGGTCAAGGTTCTGG-3'); gene cassettes region, 5'-CS (5'-GGCATCCAAGCAGCAAGC-3') and 3'-CS (5'-AAGCAGACTTGACCTGAT-3'); *orf513*, ORFend (5'-CCGTTAAGCTCTTATGTGGG-3'); *bla*_{CTX-M-2}, blaI (5'-TTAATGATGACTCAGACAGATT-3') and blaII (5'-GATACCTCGCTCATTATATGC-3'); *sul1* + *ISCR1*, F12F (5'-GTATTGCGCCGCTCTTAGAC-3') and F12R (5'-AAACCAGCATGGTTGGCTAC-3'); and 3'-CS (*qac*Δ1), *qac*Δ1B (5'-CAAGCTTTTGCCATGAAGC-3') (Power et al., 2005).

3. Results

During the study, six broad-spectrum cephalosporin-resistant *Enterobacteriaceae* strains were recovered from four different retail chicken meat samples, purchased from distinct markets (Table 1). These antibiotic-resistant isolates identified as *Proteus mirabilis* (*n* = 1), *Citrobacter diversus* (*n* = 1), *Klebsiella pneumoniae* (*n* = 1) and

Escherichia coli (*n* = 3) exhibited additional resistance to human and veterinary quinolones, aminoglycosides and/or trimethoprim-sulfamethoxazole and were positive for the production of ESBL. In fact, PCR screening for *bla*_{ESBL} genes showed that the six strains carried CTX-M-encoding genes. According to sequencing results, while most strains harbored the *bla*_{CTX-M-2} gene, the presence of the *bla*_{CTX-M-8} was only confirmed in one *E. coli* strain from market D (Table 1). On the other hand, the analysis of the genetic environment revealed that *bla*_{CTX-M-2} genes in *P. mirabilis* S11, *K. pneumoniae* S105, and *E. coli* S137 and Lac60 strains comprised a complex class 1 integron and an *ISCR1*-associated sequence, with *dfr* and/or *aadA* gene cassettes located within the variable region of the class 1 integron (Fig. 1). Curiously, for *C. freundii* Ec32 and *E. coli* CF1 strains the genetic environment of the *bla*_{CTX-M-2} and *bla*_{CTX-M-8} gene variants could not be determined by the methodology used herein, since the presence of class 1 integron was not observed in either of them.

4. Discussion

In this study, we report the occurrence of MDR *Enterobacteriaceae* in chicken meat intended for sale in the Brazilian market. Resistance to broad-spectrum cephalosporins was associated with CTX-M-2 and CTX-M-8 enzymes, which is a cause of clinical concern. In order to determine the genetic background of *bla*_{CTX-M} genes, different PCRs were conducted, using several primer combinations (Power et al., 2005). The results showed that *bla*_{CTX-M-2} was part of a structure similar to the complex class 1 integrons InS21 and In116, except for the variable region (Di Conza et al., 2002; Power et al., 2005). This kind of structure has been very common in South America (Andrade et al., 2010; Power et al., 2005; Quiroga et al., 2013), whereas, it has been also found sporadically in Spain (Valverde et al., 2006). Additionally, in a recent study in Belgium, similar class 1 integron structures were identified in *Salmonella enterica* isolated from poultry (Doublet et al., 2014). Thus, this study report for the first time the presence of *bla*_{CTX-M-2} genes harbored by complex class 1 integrons in *Enterobacteriaceae* strains recovered from retail chicken meat in Brazil. With regard to *P. mirabilis* strain S11, the genetic environment of *bla*_{CTX-M-2} is composed of a complex class 1 integron and an *ISCR1*-associated sequence, with *dfrA1* and *aadA1* gene cassettes located within the variable region, which confer resistance to trimethoprim and streptomycin, respectively. This arrangement of gene cassettes was found in *E. coli* and *Salmonella* isolated from poultry in Korea and in Belgium (Dessie et al., 2013; Doublet et al., 2014) and from clinical isolates from animals in Ireland (Karczmarczyk et al., 2011). *K. pneumoniae* S105 presented only the *dfrA1* gene in the variable region, which has not been previously reported; and while *E. coli* S137 presented the *dfrA7* gene, a variant of *dfrA1* previously found in a *E. coli* from an unhealthy animal in Ireland (Karczmarczyk et al., 2011), *E. coli* Lac60 carried only the *aadA1* gene in the variable

Table 1
Characteristics of multidrug-resistant CTX-M-producing *Enterobacteriaceae* isolated from chicken meat in Brazilian markets.

			Resistance profile (Kirby-Bauer) ^a																	
Strain ^b	Date	Market	AMC	CTX	CRO	EFT	CAZ	FEP	FOX	ATM	NA	ENO	CIP	LEV	MXF	SXT	GEN	AK	EST	β-lactamases
<i>Pm</i> S11	03/14/2011	A	R	R	R	R	S	R	S	S	R	R	S	S	R	R	S	S	R	CTX-M-2
<i>Cd</i> Ec32	04/25/2011	B	R	R	R	R	S	S	R	S	S	S	S	S	S	S	S	S	S	CTX-M-2
<i>Kp</i> S105	08/23/2011	C	R	R	R	R	R	R	S	R	R	R	S	S	R	R	R	S	S	CTX-M-2, SHV-11, TEM-1
<i>Ec</i> S137	08/23/2011	C	R	R	R	R	S	S	S	S	S	S	S	S	S	R	S	S	R	CTX-M-2, TEM-1
<i>Ec</i> Lac60	08/23/2011	C	R	R	R	R	S	R	S	R	R	R	R	S	R	S	S	S	S	CTX-M-2
<i>Ec</i> CF1	07/10/2013	D	R	R	R	R	S	R	S	S	R	R	R	I	R	S	R	S	S	CTX-M-8, TEM-1

^a AMC, amoxicillin-clavulanic acid; CTX, cefotaxime; CRO, ceftriaxone; EFT, ceftiofur; CAZ, ceftazidime; FEP, cefepime; FOX, cefoxitin; ATM, aztreonam; NA, nalidixic acid; ENO, enrofloxacin; CIP, ciprofloxacin; LEV, levofloxacin; MXF, moxifloxacin; SXT, trimethoprim-sulfamethoxazole; GEN, gentamicin; AK, amikacin; EST, streptomycin. Resistance is indicated in bold. The CLSI breakpoints were used to interpret the susceptibility results (CLSI, 2008, 2011), except for moxifloxacin where the EUCAST criteria (resistant <17 mm) was applied (http://www.eucast.org/clinical_breakpoints/).

^b *Pm*, *Proteus mirabilis*; *Cd*, *Citrobacter diversus*; *Kp*, *Klebsiella pneumoniae*; *Ec*, *Escherichia coli*.



Proteus mirabilis S11*Klebsiella pneumoniae* S105*Escherichia coli* S137*Escherichia coli* Lac60

Fig. 1. Schematic representation of class 1 integrons carrying *bla*_{CTX-M-2} genes in *Enterobacteriaceae* strains isolated from retail chicken meat. These integrons had approximate sizes of 7.7 kb (*P. mirabilis* S11), 7.2 kb (*K. pneumoniae* S105), 7 kb (*E. coli* S137) and 7.2 kb (*E. coli* Lac60). Fat arrows indicate the direction of transcription. Thin arrows indicate primers used to identify gene cassette arrays in integrons (Power et al., 2005).

region, such as *In2::ISCR1::bla*_{CTX-M-2} integron previously reported in Argentina (Quiroga et al., 2013).

In summary, to the best of our knowledge, this is the first Brazilian study showing retail chicken meat as an important vehicle for the dissemination of ESBL-producing enterobacteria, which is a major commercial and public health issue, since Brazil is the third largest producer of chicken meat and the largest exporter of this product, with a high domestic consumption (Alves et al., 2012). So, surveillance of antimicrobial resistance in bacteria from food-producing animals and derived food products needs to be established as a priority. Moreover, strategies for the rational use of antimicrobial agents in food animals need to be taken urgently, in order to inhibit the release of MDR bacteria harboring *bla*_{CTX-M}-type genes in food.

Funding

This study was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), grant number 2011/07778-1.

Acknowledgement

T.C. was supported by CAPES and FAPESP (2010/04861-2) fellowships.

J.T.T. was supported by CAPES fellowship.

M.L.N. was supported by CNPq fellowship.

N.L. is a research fellow of CNPq.

M.M.R. and G.G. are researchers of CONICET, Argentina.

M.C.L.N. was supported by FAPESP fellowship (2011/07778-1).

References

- Aizawa, J., Neuwirt, N., Barbato, L., Neves, P.R., Leigue, L., Padilha, J., Pestana de Castro, A.F., Gregory, L., Lincopan, N., 2014. Identification of fluoroquinolone-resistant extended-spectrum β -lactamase (CTX-M-8)-producing *Escherichia coli* ST224, ST2179 and ST2308 in buffalo (*Bubalus bubalis*). J. Antimicrob. Chemother. 69 (10), 2866–2869.
- Alves, J., Marques, V.V., Pereira, L.F.P., Hirooka, E.Y., De Oliveira, A.C.R.M., 2012. Multiplex PCR for the detection of *Campylobacter* spp. and *Salmonella* spp. in chicken meat. J. Food Saf. 32, 345–350.
- Andrade, L.N., Minarini, L.A., Pitondo-Silva, A., Clímico, E.C., Palazzo, I.C., Medeiros, M.I., Darini, A.L., 2010. Determinants of beta-lactam resistance in meningitis-causing *Enterobacteriaceae* in Brazil. Can. J. Microbiol. 56, 399–407.
- Bush, K., Jacoby, G.A., 2010. Updated functional classification of beta-lactamases. Antimicrob. Agents Chemother. 54, 969–976.
- Carvalho-Assef, A.P., Pereira, P.S., Albano, R.M., Berião, G.C., Tavares, C.P., Chagas, T.P., Marques, E.A., Timm, L.N., Da Silva, R.C., Falci, D.R., Asensi, M.D., 2014. Detection of NDM-1-, CTX-M-15-, and qnrB4-producing *Enterobacter hormaechei* isolates in Brazil. Antimicrob. Agents Chemother. 58, 2475–2476.
- CLSI, 2008. Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals; Approved Standard. third edition. Clinical and Laboratory Standards Institute.
- CLSI, 2011. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-First Informal Supplement. Clinical and Laboratory Standards Institute.
- Dessie, H.K., Bae, D.H., Lee, Y.J., 2013. Characterization of integrons and their cassettes in *Escherichia coli* and *Salmonella* isolates from poultry in Korea. Poult. Sci. 92, 3036–3043.
- Dhanji, H., Murphy, N.M., Doumith, M., Durmus, S., Lee, S.S., Hope, R., Woodford, N., Livermore, D.M., 2010. Cephalosporin resistance mechanisms in *Escherichia coli* isolated from raw chicken imported into the UK. J. Antimicrob. Chemother. 65, 2534–2537.
- Di Conza, J., Ayala, J.A., Power, P., Mollerach, M., Gutkind, G., 2002. Novel class 1 integron (*InS21*) carrying *bla*_{CTX-M-2} in *Salmonella enterica* serovar infantis. Antimicrob. Agents Chemother. 46, 2257–2261.
- Doublet, B., Praud, K., Nguyen-Ho-Bao, T., Argudín, M.A., Bertrand, S., Butaye, P., Cloeckaert, A., 2014. Extended-spectrum β -lactamase- and AmpC β -lactamase-producing D-tartrate-positive *Salmonella enterica* serovar Paratyphi B from broilers and human patients in Belgium, 2008–10. J. Antimicrob. Chemother. 69, 1257–1264.
- Egervärn, M., Börjesson, S., Byfors, S., Finn, M., Kaipe, C., Englund, S., Lindblad, M., 2014. *Escherichia coli* with extended-spectrum beta-lactamases or transferable AmpC beta-lactamases and *Salmonella* on meat imported into Sweden. Int. J. Food Microbiol. 171, 8–14.

- EUCAST, 2011. Breakpoint tables for interpretation of MICs and zone diameters. Version 1.3, 2011. http://www.eucast.org/clinical_breakpoints/.
- Ferreira, J.C., Penha Filho, R.A., Andrade, L.N., Berchieri, A.J., Darini, A.L., 2014. Detection of chromosomal *bla*_{CTX-M-2} in diverse *Escherichia coli* isolates from healthy broiler chickens. *Clin. Microbiol. Infect.* 20 (10), O623–O626.
- Karczmarczyk, M., Abbott, Y., Walsh, C., Leonard, N., Fanning, S., 2011. Characterization of multidrug-resistant *Escherichia coli* isolates from animals presenting at a university veterinary hospital. *Appl. Environ. Microbiol.* 77, 7104–7112.
- Kawamura, K., Goto, K., Nakane, K., Arakawa, Y., 2014. Molecular epidemiology of extended-spectrum β -lactamases and *Escherichia coli* isolated from retail foods including chicken meat in Japan. *Foodborne Pathog. Dis.* 11, 104–110.
- Kiiru, J., Kariuki, S., Goddeeris, B.M., Butaye, P., 2012. Analysis of β -lactamase phenotypes and carriage of selected β -lactamase genes among *Escherichia coli* strains obtained from Kenyan patients during an 18-year period. *BMC Microbiol.* 28 (12), 155.
- Peirano, G., Asensi, M.D., Pitondo-Silva, A., Pitout, J.D., 2011. Molecular characteristics of extended-spectrum beta-lactamase-producing *Escherichia coli* from Rio de Janeiro, Brazil. *Clin. Microbiol. Infect.* 17, 1039–1043.
- Pitout, J.D., Laupland, K.B., 2008. Extended-spectrum beta-lactamase-producing *Enterobacteriaceae*: an emerging public-health concern. *Lancet Infect. Dis.* 8, 159–166.
- Power, P., Galleni, M., Di Conza, J., Ayala, J.A., Gutkind, G., 2005. Description of In116, the first *bla*_{CTX-M-2}-containing complex class 1 integron found in *Morganella morganii* isolates from Buenos Aires. *J. Antimicrob. Chemother.* 55, 461–465.
- Quiroga, M.P., Arduino, S.M., Merkier, A.K., Quiroga, C., Petroni, A., Argentinian Integron Study Group, Roy, P.H., Centrón, D., 2013. Distribution and functional identification of complex class 1 integrons. *Infect. Genet. Evol.* 19, 88–96.
- Silva, K.C., Fontes, L.C., Moreno, A.M., Astolfi-Ferreira, C.S., Ferreira, A.J., Lincopan, N., 2013. Emergence of extended-spectrum- β -lactamase CTX-M-2-producing *Salmonella enterica* serovars Schwarzengrund and Agona in poultry farms. *Antimicrob. Agents Chemother.* 57, 3458–3459.
- Tollentino, F.M., Polotto, M., Nogueira, M.L., Lincopan, N., Neves, P., Mamizuka, E.M., Remeli, G.A., De Almeida, M.T., Rúbio, F.G., Nogueira, M.C., 2011. High prevalence of *bla*_{CTX-M} extended spectrum beta-lactamase genes in *Klebsiella pneumoniae* isolates from a tertiary care hospital: first report of *bla*_{SHV-12}, *bla*_{SHV-31}, *bla*_{SHV-38}, and *bla*_{CTX-M-15} in Brazil. *Microb. Drug Resist.* 17, 7–16.
- Valverde, A., Cantón, R., Galán, J.C., Nordmann, P., Baquero, F., Coque, T.M., 2006. In117, an unusual InO-like class 1 integron containing CR1 and *bla*_{CTX-M-2} and associated with a Tn21-like element. *Antimicrob. Agents Chemother.* 50, 799–802.
- Warren, R.E., Ensor, V.M., O'Neill, P., Butler, V., Taylor, J., Nye, K., Harvey, M., Livermore, D.M., Woodford, N., Hawkey, P.M., 2008. Imported chicken meat as a potential source of quinolone-resistant *Escherichia coli* producing extended-spectrum β -lactamases in the UK. *J. Antimicrob. Chemother.* 61, 504–508.
- Wellington, E.M., Boxall, A.B., Cross, P., Feil, E.J., Gaze, W.H., Hawkey, P.M., Johnson-Rollings, A.S., Jones, D.L., Lee, N.M., Otten, W., Thomas, C.M., Williams, A.P., 2013. The role of the natural environment in the emergence of antibiotic resistance in gram-negative bacteria. *Lancet Infect. Dis.* 13, 155–165.
- Zhao, W.H., Hu, Z.Q., 2013. Epidemiology and genetics of CTX-M extended-spectrum β -lactamases in Gram-negative bacteria. *Crit. Rev. Microbiol.* 39, 79–101.