



# Prevalence and antimicrobial susceptibility of *Escherichia coli* O157:H7 on beef cattle slaughtered in Amman abattoir

Tareq M. Osaili <sup>a,\*</sup>, Akram R. Alaboudi <sup>b</sup>, Majdi Rahahlah <sup>b</sup>

<sup>a</sup> Department of Nutrition and Food Technology, Faculty of Agriculture, Jordan University of Science and Technology, Irbid, 22110, Jordan

<sup>b</sup> Department of Pathology and Animal Health, Faculty of Veterinary Medicine, Jordan University of Science and Technology, Irbid, 22110, Jordan

## ARTICLE INFO

### Article history:

Received 1 March 2012

Received in revised form 23 July 2012

Accepted 10 November 2012

### Keywords:

*Escherichia coli* O157:H7

PCR

Multiplex PCR

Virulence factors

Antimicrobial susceptibility

## ABSTRACT

Cattle are the main asymptomatic reservoir of *Escherichia coli* O157:H7 which can cause illness to human. The objectives of the study were to measure the prevalence of *E. coli* O157:H7 on cattle slaughtered in Amman abattoir, detect virulence factors in the isolates, determine antibacterial resistance of the isolates, and know how the isolates are different or similar when compared to characterized isolates from developed countries. A total of 540 samples (feces, hide, and carcass) were tested for *E. coli* O157:H7 using the method of ISO 16654:(E). Conventional and multiplex PCR assays were used for serotype confirmation and virulence factor detection, respectively. Fifty *E. coli* O157:H7 isolates were identified and virulence factors *eaeA* and *hlyA* were present in all of the isolates. 60%, 12%, and 22% of the isolates harbored *stx*<sub>1</sub>, *stx*<sub>2</sub>, and *stx*<sub>1</sub> and *stx*<sub>2</sub>, respectively. The prevalence rates of enterotoxigenic *E. coli* O157:H7 (*n*=47) were 8.3%, 10%, and 7.8% in feces, hides and carcasses, respectively. The antimicrobial profiles of the isolates showed an extensive resistance to erythromycin, neomycin and vancomycin and high sensitivity to ampicillin, ciprofloxacin, gentamicin, kanamycin and tetracycline.

© 2012 Elsevier Ltd. All rights reserved.

## 1. Introduction

*Escherichia coli* O157:H7 is foodborne pathogen that can cause hemorrhagic colitis (HC) and hemolytic uremic syndrome (HUS) to human (Gooding & Choudary, 1997). The virulence factor associated with these illnesses is the Shiga-like toxin which is encoded by *stx*<sub>1</sub> and *stx*<sub>2</sub> genes of the pathogen (Nataro & Kaper, 1998; O'Brien & Holmes, 1987). *E. coli* O157:H7 strains also harbor a virulence factor called intimin for attachment to the host intestinal epithelial cells encoded by the *eaeA* gene (Boerling, McEwen, Wilson, Johnson, & Gyles, 1999) and may have additional virulence factors such as the enterohemolysin encoded by *ehxA* gene (Schmidt, Beutin, & Karch, 1995).

Cattle are considered the main asymptomatic reservoir of *E. coli* O157:H7 and play a vital role in human infection (Chapman, Siddons, Cerdan Malo, & Harkin, 1997; Nataro & Kaper, 1998). The organism is carried in the gastrointestinal tract and is shed in the feces (Bach, McCallister, Mears, & Schwartzkopf-Genswein, 2004), thus, this pathogen can be found on cattle hides (Arthur et al., 2004; Elder et al., 2000). Hide contamination may occur from direct or indirect fecal contamination during production and in lairage environments (Arthur et al., 2010). *E. coli* O157:H7 can be transmitted to carcass surfaces when

gut contents or fecal materials get in contact with meat surfaces during the processes of skinning and evisceration (Dickson & Anderson, 1992). Contaminated undercooked ground beef has been the most frequently identified vehicle of *E. coli* O157:H7 in the foodborne outbreaks (Barret et al., 1994).

Earlier studies have reported the prevalence rate of *E. coli* O157:H7 in cattle fecal samples, and on hide and carcasses in the range of 2.4 to 24% (Alam & Zurek, 2006; Alonso et al., 2007; Jacob, Almes, Shi, Sargeant, & Nagaraja, 2011; McEvoy et al., 2003), 11 to 76 % (Arthur et al., 2004; Elder et al., 2000), and 0.2 to 26.7 % (Alonso et al., 2007; Barkocy-Gallagher et al., 2003; FSIS, 1994; McEvoy et al., 2003; Vanderlinde, Shay, & Murray, 1998), respectively. In a local study in Jordan; the prevalence of *E. coli* O157:H7 in calves' fecal samples was 7.9% (Abu-Ashour, 2003).

Antimicrobials are widely used in cattle for disease prevention and as growth promotion (Mora et al., 2005). There are some signals indicating that antimicrobial resistance of *E. coli* O157:H7 is rising (Threlfall, Ward, Frost, & Willshaw, 2000). In Jordan, two studies were reported on the antimicrobial susceptibility of *E. coli* O157:H7 isolated from sheep and goat. The first study showed that the isolated pathogen was highly sensitive to gentamicin and co-trimoxazole and highly resistant to tetracycline and ampicillin (Tarawneh, Al-Tawarah, Abdel-Ghani, Al-Majali, & Khleifat, 2008). The second study revealed that all lamb isolates were resistant against one or more of the tested antibiotics. One isolate was resistant to chloramphenicol, and sulphonamide, another isolate was resistant to ampicillin, chloramphenicol, sulphonamide, and trimethoprim-sulfamethoxazole (co-trimoxazole) and five isolates

\* Corresponding author. Tel.: +962 2 7201000; fax: +962 2 7201078.  
E-mail address: [tosaili@just.edu.jo](mailto:tosaili@just.edu.jo) (T.M. Osaili).

were resistant to ampicillin, chloramphenicol, streptomycin, and sulphonamide. One isolate showed a multi resistance pattern against ampicillin, chloramphenicol, ceftazidime, aztreonam, gentamicin, cephalothin, streptomycin, sulphonamide, tetracycline, trimethoprim and sulfamethoxazole (Novotna et al., 2005). Harakeh et al. (2005) reported that 100% and 63.5% of tested *E. coli* O157:H7 from meat-based fast food in Lebanon were resistant to erythromycin and vancomycin, respectively.

Because of lack of information on the prevalence of *E. coli* O157:H7 in dressed beef cattle in Jordan, this work was conducted to determine: i) the prevalence of *E. coli* O157:H7 on hides and carcasses and in feces of cattle slaughtered in Amman abattoir to determine the relation between the presence of *E. coli* O157:H7 on beef meat and specific handling practices, ii) the presence of different virulence factors in isolated *E. coli* O157:H7, and iii) and the antibacterial resistance profiles of the isolates. The study also aimed in keeping up with isolate characteristics for any global changes in antimicrobial susceptibility patterns, isolate distribution and similarities and prevalence of common virulence factors.

## 2. Materials and methods

### 2.1. Reference bacterial strains

The reference bacterial strains used in this study were *E. coli* O157:H7 (ATCC 43895) as positive control and non-toxicogenic *E. coli* O157:H7 (NCTC 12900) as a negative control.

### 2.2. Sample collection

A total of 540 samples representing 180 feces, 180 hide, and 180 carcass samples were collected from calves slaughtered at Amman slaughter house during the period from January to May 2010 to investigate the presence of *E. coli* O157:H7. Amman slaughterhouse, a manually operated gravity rail system, is a public abattoir that is located in Amman City with a design capacity of 25–30 head per h and with slaughtering average of 80–100 calves per day. Most slaughtered animals were imported calves (Australia, Romania, Uruguay and Brazil) that were reared for 3–6 months in farms located 80–100 km eastern to Amman before being delivered for slaughtering. Each of the above 180 samples represents a composite sample from three different animals. The fecal sample (100 g) was collected from colon contents of slaughtered animals; longitudinal incisions were made aseptically in the colon to obtain a representative sample of the fecal content. Hide was sampled by cutting a piece representing an area of 10×10 cm (100 cm<sup>2</sup>) from the hide from each of three different locations (back, belly, and thigh). The meat was sampled by slicing a thin meat slices (thickness 1–1.5 cm) from external surfaces from each of three different locations (leg, flank and neck) of the post-washed carcass. The samples were then transferred to the laboratory in sterile containers under aseptic conditions using an ice box. Samples were processed within 2–3 h after collection.

### 2.3. Isolation of *E. coli* O157:H7 using conventional bacteriological methods

#### 2.3.1. Pre-enrichment, selective enrichment and immunomagnetic separation

The method described by ISO 16654: (E) (2001) was followed for the isolation of *E. coli* O157:H7. Twenty five grams of each sample (hide, meat and feces) was placed in Stomacher bag (Seward, UK) containing 225 ml of modified tryptone soy broth (m-TSB) (Oxoid, Ltd., Basingstoke, UK) supplemented with 20 mg/l novobiocine (Oxoid). After stomacher homogenization for 2 min at 320 rpm, the samples were incubated for 6 h at 41.5 °C. After that, 1 ml from the pre-enriched homogenate was added to 20 µl magnetic beads coated with specific antibody against O157 (Dynabeads anti *E. coli* O157, Prod. Dynal, Norway) in a 1.5 ml

Eppendorf tube. Afterward, immunomagnetic separation (IMS) was performed according to the manufacturer's instructions.

#### 2.3.2. Isolation of *E. coli* O157:H7

Fifty microliter quantities of the suspension obtained after performing the IMS were plated on Cefixime Tellurite-Sorbitol MacConkey agar (CT-SMAC) (Oxoid) and CHROMagar (Dynal Prod.). The plates were incubated at 37 °C for 22 h for CT-SMAC agar and for 24 h for CHROMagar. Five typical colorless colonies (sorbitol negative) from CT-SMAC or mauve colonies from CHROMagar were plated on tryptone soy agar + yeast extract (TSA-YE) (Oxoid), and incubated at 37 °C for 24 h, for purification and running confirmatory tests as described below.

#### 2.3.3. Confirmatory tests for *E. coli* O157 and *E. coli* O157:H7

The pinpoint indole assay was conducted on the suspected isolates. One typical colony was picked up and placed in one drop of indole reagent (Oxoid) using filter paper. Mauve color reaction indicated positive results. Indole-positive colonies were streaked onto Sorbitol MacConkey agar with BCIG (SMAC- BCIG), (Oxoid) and incubated at 37 °C for overnight. Colonies that did not show fluorescence on the SMAC- BCIG agar when illuminated with UV light (366 nm) were recognized as the distinctive characteristics of *E. coli* O157 species.

All sorbitol-negative, indole-positive, BCIG-negative isolates were sub-cultured on TSA-YE and incubated for 24 h at 37 °C. The obtained isolates were inspected by latex agglutination using the *E. coli* O157 antiserum (Oxoid) followed by H7 antisera (Remel Inc, KS, USA). Colonies that showed a positive agglutination reaction with O157 antiserum were considered as *E. coli* O157, whereas colonies that showed positive agglutination reaction with the H7 antiserum were considered as *E. coli* O157:H7.

### 2.4. Molecular confirmation and virulence factor detection of *E. coli* O157:H7

*E. coli* O157:H7 reference strain and *E. coli* O157:H7 isolates were cultured on TSA-YE at 37 °C for 24 h, and then sub-cultured into soya broth (SB) (Oxoid) for additional 24 h. Thereafter, 1 ml of the suspension was transferred into 1.5 ml microcentrifuge tubes to extract the DNA. The genomic DNA was isolated from the reference strain and bacterial isolates using Promega Wizard® Genomic DNA Purification Kit (Promega, USA) as per manufacturer's instructions.

The extracted DNAs were used for serotype (O157 and H7) confirmation (presence of *rfbE*<sub>O157</sub> and *fliCh7* genes) and virulence factor (*stx*<sub>1</sub>, *stx*<sub>2</sub>, *eaeA* and *hlyA*) detection in *E. coli* O157:H7 isolates. The primers (Thermo Electron, MA, USA) used in this analysis are presented in Table 1. The methods described by Desmarchelier et al. (1998) and Fratamico, Bagi, and Pepe (2000) were followed in conventional PCR assay for serotype confirmation and the methods described by Fratamico et al. (2000) and Sarimehmetoglu et al. (2009) were followed in multiplex PCR assay for virulence factor detection in the *E. coli* O157:

**Table 1**

Primers used for serotype confirmation and virulence factor determination of *Escherichia coli* O157:H7 isolates.

Primer	Sequence (5'–3')	Target gene	PCR product (bp)
SLT1-F	TGTAAGTGGAAAGGTGGAGTATACA	<i>stx</i> <sub>1</sub>	210
SLT1-R	GCTATTCTGAGTCAACGAAAAATAAC		
SLTII-F	GTTTTCTTCGGTATCCTATTC	<i>stx</i> <sub>2</sub>	484
SLTII-R	GATGCATCTCTGGTCATTGTATTAC		
AE22	ATTACCATCCACACAGACGGT	<i>eaeA</i>	397
AE20-2	ACAGCGTGGTGGATCAACCT	<i>hlyA</i>	166
MFS1-F	ACGATGTGGTTTATTCTGGA		
MFS1-R	CTTCACGTCACCATACATAT	<i>fliCh7</i>	625
FLICH7-F	GCGCTGTCGAGTTCTATCGAGC		
FLICH7-R	CAACGGTGACTTTATCGCCATTC	<i>rfbE</i>	497
O157-AF <sup>a</sup>	AAGATTGCGCTGAAGCCTTTG		
O157-AR <sup>a</sup>	CATTGGCATCGTGGACAG		

<sup>a</sup> Sarimehmetoglu et al. (2009) and Desmarchelier et al. (1998).

H7 isolates. Thermal cycling was conducted in Veriti Thermal cycler (Applied Biosystems, USA) and PCR products were visualized with UV transilluminator and photographed with the gel documentation system (Gel Doc 2000, BIO-RAD, USA).

### 2.5. Susceptibility to antimicrobial agents-microbroth dilution method

*E. coli* O157:H7 isolates in addition to the reference bacterial strain *E. coli* O157:H7 (ATCC 43895) were maintained on TSA-YE. Active cultures were prepared by growing the isolates and the reference strain in 10 ml quantities of Mueller–Hinton broth (MHB) (Oxoid). The densities of bacterial growth in MHB were standardized at ca.  $1.5 \times 10^8$  CFU/ml and were adjusted according to McFarland standard solution. To determine the minimal inhibitory concentration (MIC) of antimicrobials against tested strains in MHB at 37 °C, the microbroth dilution method described by Lalitha (2008) was followed. The MIC values of the isolates and the reference strain were compared to breakpoint values from international standards or published researches (BSAC, 2008; Fitzgerald et al., 2003; NCCLS, 1999; Soussy, Cluzel, Courvalin, & the Comité de Antibiotogramme de la Société Française de Microbiologie, 1994).

#### 2.5.1. Statistical analysis

The differences among numbers of contaminated samples (carcasses, hides and feces) with enterotoxigenic *E. coli* O157:H7 were tested by Chi-square test using SAS program, version 8.1 (SAS Institute Inc., Cary, NC). Significance level is considered at  $\leq 0.05$ .

## 3. Results

### 3.1. Isolation of *E. coli* O157:H7 by conventional bacteriological method

The number of samples with typical *E. coli* O157:H7 characteristics on CT-SMAC or CHROMagar was 63 samples (22 feces samples, 17 hide samples, and 24 carcass samples). All of the presumptively identified *E. coli* O157:H7 isolates from the tested samples were positive for spot indole test and presence of O157 antigen, whereas, 55 of them (87%) were positive for presence of H7 antigen using latex agglutination antisera (Table 2).

### 3.2. Serotype confirmation of *E. coli* O157:H7 using conventional PCR

All of the presumptive isolates (63 isolates) were subjected to PCR analyses for serotype confirmation. The results showed that 51 isolates (80.9%) harbored *rfbE*<sub>O157</sub> gene (O157) and only 50 isolates (79%) contained both *rfbE*<sub>O157</sub> and *fliC*<sub>H7</sub> genes (O157:H7). Accordingly, only one isolate out of the 51 of the confirmed serotype O157 did not harbor the H7 gene. To compare latex agglutination data with PCR results, 51 out of the 63, O157 latex positive isolates were confirmed as *E. coli* O157 (83%), whereas 50 out of the 55 latex H7 positive were confirmed as *E. coli* O157:H7 (91%). The overall prevalence of PCR confirmed *E. coli* O157:H7 in feces and on hide and carcass samples was 17 (9.4%), 19 (10.6%), and 14 (7.8%), respectively (Table 2).

### 3.3. Detection of virulence factors in the confirmed *E. coli* O157:H7 using multiplex PCR

Multiplex PCR results revealed that all of the 50 confirmed *E. coli* O157:H7 isolates harbored both intimin and enterohemolysin (*eaeA* and *hlyA*) virulent genes while 47 (94%) of the isolates revealed the presence of *stx*<sub>1</sub> and/or *stx*<sub>2</sub> virulence genes; 30 (60%), 6 (12%), and 11 (22%) isolates harbored *stx*<sub>1</sub>, *stx*<sub>2</sub>, and both genes, respectively. The 3 isolates (6%) that did not have *stx*s genes were isolated from hides (1 isolate) and feces (2 isolates). The 47 molecularly confirmed enterotoxigenic *E. coli* O157:H7 were distributed between 15 isolates from feces (8.3%), 18 isolates for hides (10%) and 14 isolates from carcasses (7.8%) (Table 3). There were no significant differences ( $P > 0.05$ ) among the contamination level of enterotoxigenic *E. coli* O157:H7 on carcasses and hides and in feces. The *stx*<sub>1</sub> gene was of high prevalence among hides isolates (68%,  $n = 13$ ) followed by carcass isolates (64%,  $n = 9$ ) and finally feces isolates (47%,  $n = 8$ ). Again and in the same meaning the *stx*<sub>2</sub> gene was the highest one (15.8%,  $n = 3$ ) among hide isolates followed by feces (11.8%,  $n = 2$ ) and the carcass (7%,  $n = 1$ ) isolates. On the other hand, both *stx*<sub>1</sub> and *stx*<sub>2</sub> genes were of high prevalence in feces isolates (29.4%,  $n = 5$ ) followed by carcass (28.5%,  $n = 4$ ) and hide isolates (10.5%,  $n = 2$ ).

### 3.4. Distribution of toxigenic *E. coli* O157:H7 in feces and on hide and carcass samples

Out of the 14 positive carcasses-animal for *E. coli* O157:H7, 2 composite samples were only carcass positive; 5 composite samples were positive for both carcasses and hides; 2 composite samples were positive for carcasses and feces and finally 5 composite samples were positive for carcass, hide and feces simultaneously. In the same meaning out of 18 positive hide-animal, 6 composite samples were only hide positive, 5 composite samples were positive for hides and carcasses; 2 composite samples were positive for hides and feces and the other 5 composite samples were positive for hides, feces and carcasses simultaneously. On the other hand out of the 15 positive feces-animal, 6 composite samples were only positive for feces samples; 2 composite samples were positive for both feces and hides; 2 composite samples were positive for feces and carcasses and finally 5 composite samples were positive for feces, hides and carcasses simultaneously (Table 4).

### 3.5. Susceptibility to antimicrobial agents

None of the 40 *E. coli* O157:H7 isolates were resistant to ampicillin, ciprofloxacin or tetracycline, whereas all were resistant to erythromycin, doxycycline, vancomycin and, neomycin. 50% of the isolates were resistant to three antibiotics namely erythromycin, vancomycin and neomycin. 10 out of 40 (25%) of the isolates were multi-resistant to four antibiotics namely erythromycin, vancomycin, doxycycline and neomycin, and another 10 (25%) were multi-resistant to other four different patterns (erythromycin, vancomycin, neomycin and streptomycin). On the other hand, 2 out of 40 (5%) isolates were multi-resistant to five antibiotics namely erythromycin, vancomycin, doxycycline, neomycin and streptomycin (Table 5).

**Table 2**  
Presumptive and confirmed *Escherichia coli* O157 and *Escherichia coli* O157:H7 isolates (%).

Sample	No. of samples	Presumptive <i>E. coli</i> O157:H7 on selective media	Latex agglutination		Serotype confirmation by PCR	
			<i>E. coli</i> O157	<i>E. coli</i> O157:H7	<i>E. coli</i> O157	<i>E. coli</i> O157:H7
Carcasses	180	17 (9.4)	17 (9.4)	14 (7.2)	14 (7.8)	14 (7.8)
Hides	180	24 (13.3)	24 (13.3)	21 (11.7)	19 (10.6)	19 (10.6)
Feces	180	22 (12.2)	22 (12.2)	20 (11.1)	18 (10.0)	17 (9.4)
Total	540	63	63	55	51	50



**Table 3**  
Virulence genes detected in *Escherichia coli* O157:H7 isolates by multiplex PCR.

Source of isolates	Number of isolates	Virulence gene profiles (%)					
		<i>eaeA</i>	<i>hlyA</i>	<i>stx</i>			
				<i>stx</i> <sub>1</sub>	<i>stx</i> <sub>2</sub>	<i>stx</i> <sub>1</sub> + <i>stx</i> <sub>2</sub>	No <i>stx</i>
Carcasses	14	14 (100)	14 (100)	9 (64.3)	1 (7.1)	4 (28.6)	0 (0)
Hides	19	19 (100)	19 (100)	13 (68.4)	3 (15.8)	2 (10.5)	1 (5.3)
Feces	17	17 (100)	17 (100)	8 (47)	2 (11.8)	5 (29.4)	2 (11.8)
Total	50	50 (100)	50 (100)	30 (60)	6 (12)	11 (22)	3 (6)

#### 4. Discussion

The prevalence rates of *E. coli* O157:H7 (9.4%, 10.6% and 7.8%) or toxigenic *E. coli* O157:H7 (8.3%, 10% and 7.8%) in feces and on hide and carcass samples are within the rates reported in literature. The prevalence rate of *E. coli* O157:H7 in feces samples has been reported in the range of 2.4 to 24% (Alam & Zurek, 2006; Alonso et al., 2007; Jacob et al., 2011; McEvoy et al., 2003). The prevalence rate of *E. coli* O157:H7 on hide and carcass samples has been reported in the range of 11 to 76 % (Arthur et al., 2004; Elder et al., 2000) and 0.2 to 26.7 % (Alonso et al., 2007; Barkocy-Gallagher et al., 2003; FSIS, 1994; McEvoy et al., 2003; Vanderlinde et al., 1998), respectively. The low prevalence rates of *E. coli* O157:H7 reported in this study compared with those reported in the previously mentioned studies may be related to differences in methodology of sampling. However, the prevalence of *E. coli* O157:H7 on beef carcasses was higher in the current study than that in a number of published studies. In the United States, the prevalence of *E. coli* O157:H7 on beef carcasses after dressing was determined to be 0.2% (FSIS, 1994). Two studies that were conducted in the United Kingdom have reported that 0.47% and 1.4% of carcasses were contaminated with *E. coli* O157:H7 (Chapman, Cerdán Malo, Ellin, Ashton, & Harkin, 2001; Richards et al., 1998). But, a study in northern Italy found that 12 out of 100 (12%) carcasses sampled were contaminated with *E. coli* O157:H7 (Bonardi, Maggi, Pizzin, Morabito, & Caprioli, 2001). The higher isolation rate (7.8%) observed for *E. coli* O157:H7 on carcasses in this study might be related to the absence of antimicrobial interventions through dressing processes at abattoirs level or due to cross contamination during the slaughter process which in overall reflect the general unhygienic conditions in employees, utensils and environmental sanitation of the slaughter house under study.

The contamination levels of feces, hides and carcasses are closely related. This seems to be quite logical as the main source of contamination is the fecal material which found its way to animal external surface due to poor hygienic conditions during slaughtering process of the animals. Contamination of carcasses with *E. coli* O157:H7 can occur when gut contents, fecal matter or contaminated hides come in contact with meat surfaces. These observations come in close agreement with other findings in that a significant correlation does exist between the prevalence of *E. coli* O157 in feces and on hides and carcasses (Bonardi et al., 2001; Elder et al., 2000). The distribution of *E. coli* O157:H7 among hides and carcasses depends mainly on level of hygienic precautions taken during slaughtering to avoid cross contamination during hide

removal, evisceration and carcass dressing. This could be shown when six animals revealed the presence of *E. coli* O157:H7 from intestinal contents but not from either hide or carcasses (Table 4). Therefore, the main sources of possible carcass contamination in abattoir are both of hide and fecal contents. However, possible cross contamination from utensils and employers is not to be neglected. This might be apparent from following observation recorded in this study, where in two cases the pathogen was recovered from carcasses only, but not from hides or feces. Pathogenicity of *E. coli* O157:H7 is related to many virulence factors like Shiga toxins, intimin and enterohemolysin (Nataro & Kaper, 1998). In the present study, a total of 50 PCR identified isolates from 540 animals were tested for virulence factor by multiplex PCR procedure. Out of these, 47 toxigenic isolates (harbored *stx*<sub>1</sub> and/or *stx*<sub>2</sub> genes) were confirmed. 30 (64%) isolates harbored *stx*<sub>1</sub>; 6 (13%) isolates harbored *stx*<sub>2</sub>; and 11(23%) isolates harbored both toxins. The *stx*<sub>2</sub> producing *E. coli* O157:H7 strains are more virulent than *stx*<sub>1</sub> or *stx*<sub>1</sub> and *stx*<sub>2</sub> producing strains (Donnenberg, 2002), and being more often linked with human disease problems (Boerling et al., 1999).

Similar to our study, non- pathogenic *E. coli* O157:H7 strains (without toxins factors) have also been identified from feces or beef carcass (Guyon et al., 2001; Rogerie et al., 2001).

Cagney et al. (2004) studied the prevalence of *E. coli* O157:H7 in ground beef and beef burger (n = 1533) from supermarkets and butcher shops. In all of the 43 isolates, *eaeA*, *hlyA* and *fliCh7* genes were detected and in 41 isolates *stx*<sub>1</sub> and *stx*<sub>2</sub> genes were determined. The presence of the two (*eaeA* and *hlyA*) virulence factors comes in agreement with the isolates identified in our study, though *fliCh7* genes were only present in 91% of our isolates. The toxigenicity percentage among all isolates (95%) was in close agreement with the 94% reported in our study. In another study on ground beef in Turkey, Sarimehmetoglu et al. (2009) identified 2 *E. coli* O157:H7 isolates. Both isolates have *eaeA*, *hlyA* and *fliCh7* genes, one of the isolates has *stx*<sub>1</sub> and the other one has *stx*<sub>2</sub> only. Whereas, Vernozy-Rozand et al. (2002) recovered 4 *E. coli* O157:H7 from the 3450 ground beef samples in France and all the strains had both *stx*<sub>1</sub> and *stx*<sub>2</sub>. Therefore, it appears that the distribution of *stx*s in isolates might be different from food to other or from different geographical locations.

Antimicrobial resistance may arise either spontaneously by selective pressure or due to antimicrobial misuse by humans or overuse in feeding or treatment of beef cattle by farmers (Schroeder et al., 2002). Resistance development also might be related to exchange of resistance factors between related bacteria (Tenover, 2006). The resistance of *E. coli* O157:H7 to both erythromycin and vancomycin also comes in agreement with the results of Harakeh et al. (2005). The results of the present study indicated that *E. coli* O157:H7 were susceptible to gentamicin and ciprofloxacin, which comes in agreement with a local Jordanian study on isolates from sheep and goat (Tarawneh et al., 2008) and other studies in the United States on isolates from human, cattle, swine and food (Harakeh et al., 2005; Schroeder et al., 2002). In the present study all *E. coli* O157:H7 tested were also susceptible to tetracycline which again comes in parallel with the results of Dontorou et al. (2003) on three isolates obtained from ewes' milk, fresh sausages and swine intestines in Greece. Whereas this observation contradicts Tarawneh et al.'s (2008) report on local Jordanian isolates from sheep and goat. Ali, Farooqui, Khan, Khan, and Kazmi (2010) showed that 15 out of 51

**Table 4**  
Distribution of toxigenic *Escherichia coli* O157:H7 in feces and on hides and carcasses of the tested animals.

Sample type	Carcass (C) (14)	Hides (H) (18)	Feces (F) (15)
No of isolates for each location	C = 2	H = 6	F = 6
	C + H = 5	H + C = 5	F + H = 2
	C + F = 2	H + F = 2	F + C = 2
	C + H + F = 5	H + F + C = 5	F + H + C = 5

**Table 5**Antimicrobial minimum inhibitory concentration (MIC) values for *Escherichia coli* O157:H7 (ATCC 43895) and 40 selected toxigenic *Escherichia coli* O157:H7 isolates.

Total no. of isolates	Ery <sup>a</sup> (≤0.5–≥8) <sup>b</sup>	Gen (≤2–>4)	Amp (≤8–≥16)	Cip (≤1–≥4)	Kan (≤8–≥16)	Tet (≤4–≥16)	Dox (<1–≥1)	Str (≤16–≥64)	Van (≤4–≥16)	Neo (≥16)
<i>E. coli</i> O157:H7 (ATCC 43895)	32	0.25	4	0.03	0.5	1	1	0.5	64	128
2	32	0.25	2	0.25	2	2	1	8	>512	>512
1	32	0.25	4	0.25	1	0.5	1	2	>512	>512
1	32	0.5	4	0.015	4	0.5	1	64	>512	>512
4	32	1	4	0.015	8	0.5	1	128	>512	>512
1	32	0.5	8	0.25	8	0.5	2	2	>512	>512
1	32	1	8	0.25	8	0.5	1	2	>512	>512
3	64	2	4	0.015	8	1	2	64	>512	>512
1	32	4	2	0.015	8	1	1	64	>512	>512
2	32	2	2	0.25	8	0.5	2	32	>512	>512
1	32	4	8	0.25	8	1	2	8	>512	>512
2	64	8	4	0.015	16	1	2	16	>512	>512
3	64	1	8	0.25	16	1	2	8	>512	>512
1	32	1	4	0.5	4	1	1	4	>512	>512
4	32	0.25	4	0.25	1	0.5	1	2	>512	>512
2	32	0.125	4	0.125	8	1	1	1	>512	>512
1	32	0.25	8	0.5	1	1	1	2	>512	>512
3	32	1	4	0.015	8	1	1	64	>512	>512
2	32	0.5	4	0.25	2	0.5	1	2	>512	>512
1	32	0.5	4	0.5	2	1	1	2	>512	>512
4	32	2	4	0.5	8	1	2	8	>512	>512
Resistance %	100%	5%	0%	0%	5%	0%	100%	12.5%	100%	100%

<sup>a</sup> Ery: erythromycin, Gen: gentamicin, Amp: ampicillin, Cip: ciprofloxacin, Kan: kanamycin, Tet: tetracycline, Dox: doxycycline, Str: streptomycin, Van: vancomycin, Neo: neomycin.<sup>b</sup> Breakpoints: (BSAC, 2008, 2012; Fitzgerald et al., 2003; NCCLS, 1999; Soussy et al., 1994).

(29%) *E. coli* O157:H7 meat isolates in Pakistan were resistant to doxycycline, which is quite highly lower than the 100% resistance reported in the present study. The resistance patterns for our tested isolates for kanamycin, streptomycin and ampicillin come on the same line with a study in Japan where 15 isolates (16.9%) of enterohemorrhagic *E. coli* O157:H7 isolates obtained from diarrhea patients, were resistant to ampicillin, kanamycin or streptomycin (Miwa et al., 2002). Our observation for ampicillin susceptibility disagreed with other local studies on *E. coli* O157:H7 isolates from sheep and goat (Tarawneh et al., 2008). Once more, all tested *E. coli* O157:H7 were resistant to neomycin, which also comes in accordance with Nizza et al. (2010), as 100% of tested isolates from buffalo calves with diarrhea in Italy were resistant to neomycin.

Multiple antimicrobial resistances in *E. coli* may arise from the spread of genetic materials including integrons, transposons, and plasmids (Zhao et al., 2001). The increasing developing multi-drug resistant bacteria is signaling a serious alarm from treatment point of view or the possible transforming of resistance genes to other related pathogens (Okeke, Lamikanra, & Edelman, 1999). Multiple antimicrobial resistance is a common phenomenon among *E. coli* O157:H7 isolates drawn from different sources. Schroeder et al. (2002) in the USA, showed that 17%, 7.5%, 8%, 5%, 2% and 0.1% of the *E. coli* O157 isolates were resistant to one, two, three, four, five and six drugs, respectively. Also, Zhao et al. (2001) studied multiple resistance of isolates recovered from human, animals and food in the USA. They found that out of the twenty nine tested *E. coli* O157:H7, four isolates showed multiple resistance to five antimicrobials: tetracycline ampicillin, streptomycin, kanamycin, and sulfamethoxazole. Two isolates were from cattle, and two isolates were from human and ground beef.

It is essential to keep up with isolate characteristics for any global changes in isolate distribution and similarities and prevalence of common virulence factors. Also, it is essential to track the resistance pattern recorded globally to follow changes in antimicrobial sensitivity patterns that may require a reassessment of zoonotic control strategy. Monitoring of antimicrobial resistance in *E. coli* O157:H7 isolates is valuable for epidemiological uses and for monitoring the increase of antimicrobial resistance among different microbial species. Continuous surveillance of emerging antimicrobial resistance among zoonotic foodborne pathogens, including *E. coli* O157:H7 is required to ensure global one inspiration public health.

## Acknowledgment

The authors thank the Deanship of Scientific Research at Jordan University of Science and Technology for funding the project.

## References

- Abu-Ashour, H. H. (2003). Studies on enterohaemorrhagic *Escherichia coli* isolated from human, calves and fast food samples in Northern and middle Jordan's thesis, Faculty of Veterinary Medicine, Jordan University of Science and Technology.
- Alam, M. J., & Zurek, L. (2006). Seasonal prevalence of *Escherichia coli* O157:H7 in beef cattle feces. *Journal of Food Protection*, 69, 3018–3020.
- Ali, N. H., Farooqui, A., Khan, A., Khan, A. Y., & Kazmi, S. U. (2010). Microbial contamination of raw meat and its environment in retail shops in Karachi, Pakistan. *Journal of Infection in Developing Countries*, 4, 382–388.
- Alonso, S., Mora, A., Blanco, M., Blanco, J. E., Dahbi, G., Ferreira, M. T., et al. (2007). Fecal carriage of *Escherichia coli* O157:H7 and carcass contamination in cattle at slaughter in northern Italy. *International Microbiology*, 10, 109–116.
- Arthur, T. M., Bosilevac, J. M., Nou, X., Shackelford, S. D., Wheeler, T. L., Kent, M. P., et al. (2004). *Escherichia coli* O157:H7 prevalence and enumeration of aerobic bacteria, *Enterobacteriaceae*, and *Escherichia coli* O157:H7 at various steps in commercial beef processing plants. *Journal of Food Protection*, 67, 658–665.
- Arthur, T. M., Brichta-Harhay, D. M., Bosilevac, J. M., Kalchayanand, N., Shackelford, S. D., & Wheeler, T. L. (2010). Super shedding of *Escherichia coli* O157:H7 by cattle and the impact on beef carcass contamination: Review. *Meat Science*, 86, 32–37.
- Bach, S. J., McCallister, T. A., Mears, G. J., & Schwartzkopf-Genswein, K. S. (2004). Long-haul transport and lack of preconditioning increases fecal shedding of *Escherichia coli* and *Escherichia coli* O157:H7. *Journal of Food Protection*, 67, 672–678.
- Barkocy-Gallagher, G. A., Arthur, T. M., Rivera-Betancourt, M., Nou, X., Shackelford, S. D., Wheeler, T. L., et al. (2003). Seasonal prevalence of Shiga toxin-producing *Escherichia coli*, including O157:H7 and non-O157:H7 serotypes, and *Salmonella* in commercial beef processing plants. *Journal of Food Protection*, 66, 1978–1986.
- Barret, T. J., Lior, H., Green, J. H., Khakhria, R., Wells, J. G., Bell, B. P., et al. (1994). Laboratory investigation of a multistate food-borne outbreak of *Escherichia coli* O157:H7. *Journal of Clinical Microbiology*, 32, 3013–3017.
- Boerling, P., McEwen, S. A., Wilson, J. B., Johnson, R. P., & Gyles, C. L. (1999). Association between virulence factors of Shiga toxin-producing *Escherichia coli* and disease in humans. *Journal of Clinical Microbiology*, 37, 497–503.
- Bonardi, S., Maggi, E., Pizzini, G., Morabito, S., & Caprioli, A. (2001). Fecal carriage of verocytotoxin producing *Escherichia coli* O157 and carcass contamination in cattle at slaughter in northern Italy. *International Journal of Food Microbiology*, 66, 47–53.
- British Society for Antimicrobial Chemotherapy (BSAC) (2008). Method for antimicrobial susceptibility testing. Available at: <http://www.bsac.org.uk>.
- British Society for Antimicrobial Chemotherapy (BSAC) (2012). Method for antimicrobial susceptibility testing. Available at: <http://www.bsac.org.uk>.
- Cagney, C., Crowley, H., Duffy, G., Sheridan, J. J., O'Brien, S., Carney, E., et al. (2004). Prevalence and numbers of *Escherichia coli* O157:H7 in minced beef and beef burgers from butcher shops and supermarkets in the Republic of Ireland. *Food Microbiology*, 21, 203–212.

- Chapman, P. A., Cerdán Malo, A. T., Ellin, M., Ashton, R., & Harkin, M. A. (2001). *Escherichia coli* O157 in cattle and sheep at slaughter, on beef and lamb carcasses and in raw beef and lamb products in South Yorkshire, UK. *International Journal of Food Microbiology*, 64, 139–150.
- Chapman, P. A., Siddons, C. A., Cerdan Malo, A. T., & Harkin, M. A. (1997). A 1-year study of *Escherichia coli* O157 in cattle, sheep, pigs and poultry. *Epidemiology and Infection*, 119, 245–250.
- Desmarchelier, P. M., Bilge, S. S., Fegan, N., Mills, L., Vary, J. C., & Tarr, P. I. (1998). A PCR specific for *Escherichia coli* O157 based on the rfb locus encoding O157 lipopolysaccharide. *Journal of Clinical Microbiology*, 36, 1801–1804.
- Dickson, J. S., & Anderson, M. E. (1992). Microbiological decontamination of food animal carcasses by washing and sanitizing systems: A review. *Journal of Food Protection*, 55, 133–140.
- Donnenberg, M. S. (Ed.). (2002). *Escherichia coli: Virulence mechanisms of a versatile pathogen*. San Diego, CA: Academic Press.
- Dontorou, C., Papadopoulou, C., Filioussis, G., Economou, V., Apostolou, I., Zakkas, G., et al. (2003). Isolation of *Escherichia coli* O157:H7 from foods in Greece. *International Journal of Food Microbiology*, 82, 273–279.
- Elder, R. O., Keen, U. E., Siragusa, G. R., Barkocy-Gallagher, G. A., Koohmaraie, M., & Laegreid, W. W. (2000). Correlation of enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proceedings of the National Academy of Sciences*, 97, 2999–3003.
- Fitzgerald, A. C., Edrington, T. S., Looper, M. L., Callaway, T. R., Genovese, K. J., Bischoff, K. M., et al. (2003). Antimicrobial susceptibility and factor affecting the shedding of *E. coli* O157 H7 and *salmonella* in dairy cattle. *Journal of Applied Microbiology*, 37, 392–398.
- Food Safety and Inspection Service (FSIS) (1994). *Nationwide Beef Microbiological Baseline Data Collection Program: Steers and Heifers, October 1992–September 1993*. U.S. Dept. of Agriculture Available from: <http://www.aphis.usda.gov/vs/ceah/cahm>.
- Fratamico, P. M., Bagi, L. K., & Pepe, T. (2000). A multiplex polymerase chain reaction assay for rapid detection and identification of *Escherichia coli* O157:H7 in foods and bovine feces. *Journal of Food Protection*, 63, 1032–1037.
- Gooding, C. M., & Choudary, P. V. (1997). Rapid and sensitive immunomagnetic separation polymerase chain reaction method for the detection of *Escherichia coli* O157:H7 in raw milk and ice-cream. *The Journal of Dairy Research*, 64, 87–93.
- Guyon, R., Dorey, F., Malas, J. P., Grimont, F., Foret, J., Rouvière, B., et al. (2001). Superficial contamination of bovine carcasses by *Escherichia coli* O157:H7 in a slaughterhouse in Normandy (France). *Meat Science*, 58, 329–331.
- Harakeh, S., Yassine, H., Gharios, M., Barbour, E., Hajjar, S., El-Fadel, M., et al. (2005). Isolation, molecular characterization and antimicrobial resistance patterns of *Salmonella* and *Escherichia coli* isolates from meat-based fast food in Lebanon. *Science of the Total Environment*, 341, 33–44.
- ISO (International Organization for Standardization) (2001). (1st ed.). *Microbiology of food and animal feeding stuff-Horizontal method for the detection of Escherichia coli O157*, 16654, (E).
- Jacob, M. E., Almes, K. M., Shi, X., Sargeant, J. M., & Nagaraja, T. G. (2011). *Escherichia coli* O157:H7 genetic diversity in bovine fecal samples. *Journal of Food Protection*, 74, 1186–1188.
- Lalitha, M. K. (2008). Manual on antimicrobial susceptibility testing. *Indian Journal of Medical Research*, 88, 455–460.
- McEvoy, J. M., Dehorty, A. M., Sheridan, J. J., Thomson-Carter, F. M., Gravey, P., McGuire, L., et al. (2003). The prevalence and spread of *Escherichia coli* O157:H7 at commercial beef abattoir. *Journal of Applied Microbiology*, 95, 256–266.
- Miwa, Y., Matsumoto, M., Hiramatsu, R., Yamazaki, M., Saito, H., Saito, M., et al. (2002). Drug resistance of enterohemorrhagic *Escherichia coli* O157 and a possible relation of plasmids to the drug-resistance (abstract). *Kansenshōgaku Zasshi*, 76, 285–290.
- Mora, A., Blanco, J. E., Blanco, M., Alonso, M. P., Dhabhi, G., Echeita, A., et al. (2005). Antimicrobial resistance of Shiga toxin (verotoxin)-producing *Escherichia coli* O157:H7 and non-O157 strains isolated from humans, cattle, sheep and food in Spain. *Research in Microbiology*, 156, 793–806.
- Nataro, J. P., & Kaper, J. B. (1998). Diarrheagenic *Escherichia coli*. *Clinical Microbiology Reviews*, 11, 142–201.
- National Committee for Clinical Laboratory Standards (NCCLS) (1999). Performance Standards for Antimicrobial Disk Susceptibility Tests, Approved standard. NCCLS document M7-A3. National Committee for Clinical Laboratory Standards, Wayne, Pa.USA.
- Nizza, S., Mallardo, K., Marullo, A., Iovane, V., De Martino, L., & Pagnini, U. (2010). Antibiotic susceptibility of haemolytic *E. coli* strains isolated from diarrhoeic feces of buffalo calves. *Italian Journal of Animal Science*, 9, e26.
- Novotna, R., Alexa, P., Hamrik, J., Madanat, A., Smola, J., & Cizek, A. (2005). Isolation and characterization shiga toxin producing *Escherichia coli* from sheep and goats in Jordan with evidence of multi resistant serotype O157:H7. *Veterinarni Medicina Czech Republic*, 50, 111–118.
- O'Brien, A. D., & Holmes, R. K. (1987). Shiga and Shiga-like toxins. *Microbiology Reviews*, 51, 206–220.
- Okeke, I. N., Lamikanra, A., & Edelman, R. (1999). Socioeconomic and behavioral factors leading to acquired bacterial resistance to antibiotics in developing countries. *Emerging Infectious Diseases*, 5, 18–27.
- Richards, M. S., Corkish, J. D., Sayers, A. R., McLaren, I. M., Evans, S. J., & Wray, C. (1998). Studies of the presence of verocytotoxic *Escherichia coli* O157 in bovine feces submitted for diagnostic purposes in England and Wales and on beef carcasses in abattoirs in the United Kingdom. *Epidemiology and Infections*, 120, 187–192.
- Rogerie, F., Marecat, A., Ganbade, S., Dupond, F., Beaubois, P., & Lange, M. (2001). Characterization of Shiga toxin producing *E. coli* O157 H7 serotype isolated in France from healthy domestic cattle. *International Journal of Food Microbiology*, 63, 217–223.
- Sarimehmetoglu, B., Aksoy, M. H., Ayaz, N. D., Ayaz, Y., Kuplulu, O., & Kaplan, Y. Z. (2009). Detection of *Escherichia coli* O157:H7 in ground beef using immunomagnetic separation and multiplex PCR. *Food Control*, 20, 357–361.
- Schmidt, H., Beutin, L., & Karch, H. (1995). Molecular analysis of the plasmid-encoded hemolysin of *Escherichia coli* O157:H7 strain EDL 933. *Infection and Immunology*, 63, 1055–1061.
- Schroeder, C. M., Zhao, C., Debroy, C., Torcolini, J., Zhao, S., White, D. G., et al. (2002). Antimicrobial resistance of *Escherichia coli* O157 isolated from humans, cattle, swine, and food. *Applied and Environmental Microbiology*, 68, 576–581.
- Soussy, C. J., Cluzel, R., Courvalin, P., & The Comité de AntibioGramme de la Société Française de Microbiologie (1994). Definition and determination of in vitro antibiotic susceptibility breakpoints for bacteria in France. *European Journal of Clinical Microbiology and Infectious Diseases*, 13, 238–246.
- Tarawneh, K. A., Al-Tawarah, N. M., Abdel-Ghani, A. H., Al-Majali, A. M., & Khleifat, K. M. (2008). Characterization of verotoxigenic *Escherichia coli* (VTEC) isolates from feces of small ruminants and environmental samples in Southern Jordan. *Journal of Basic Microbiology*, 48, 1–8.
- Tenover, F. C. (2006). Mechanisms of antimicrobial resistance in bacteria. *American Journal of Medicine*, 119(6A), S3–S10.
- Threlfall, E. J., Ward, L. R., Frost, J. A., & Willshaw, G. A. (2000). The emergence and spread of antibiotic resistance in food-borne bacteria. *International Journal of Food Microbiology*, 62, 1–5.
- Vanderlinde, P. B., Shay, B., & Murray, J. (1998). Microbiological quality of Australian beef carcass meat and frozen bulk packed beef. *Journal of Food Protection*, 61, 437–443.
- Vernozy-Rozand, C., Ray-Gueniot, S., Ragot, C., Bavai, C., Mazuy, C., Montet, M. P., et al. (2002). Prevalence of *Escherichia coli* O157:H7 in industrial minced beef. *Letters in Applied Microbiology*, 35, 7–11.
- Zhao, C., Ge, B., De Villena, J., Sudler, R., Yeh, E., Zhao, S., et al. (2001). Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* serovars in retail chicken, turkey, pork, and beef from the Greater Washington, D.C. area. *Applied and Environmental Microbiology*, 67, 5431–5436.