

Accepted Manuscript

Occurrence of extended spectrum β -lactamase and AmpC genes among multidrug-resistant *Escherichia coli* and emergence of ST131 from poultry meat in Thailand

Uttapoln Tansawai, Donruedee Sanguansermisri, Anamai Na-udom, Timothy R. Walsh, Pannika R. Niumsup



PII: S0956-7135(17)30376-6

DOI: [10.1016/j.foodcont.2017.07.028](https://doi.org/10.1016/j.foodcont.2017.07.028)

Reference: JFCO 5721

To appear in: *Food Control*

Please cite this article as: Uttapoln Tansawai, Donruedee Sanguansermisri, Anamai Na-udom, Timothy R. Walsh, Pannika R. Niumsup, Occurrence of extended spectrum β -lactamase and AmpC genes among multidrug-resistant *Escherichia coli* and emergence of ST131 from poultry meat in Thailand, (2017), doi: 10.1016/j.foodcont.2017.07.028

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Occurrence of extended spectrum β -lactamase and AmpC genes among multidrug-resistant *Escherichia coli* and emergence of ST131 from poultry meat in Thailand

Uttapoln Tansawai ^a, Donruedee Sanguansermisri ^{a,b}, Anamai Na-udom ^c, Timothy R. Walsh ^d and Pannika R. Niumsup ^{a,b,*}

^a Department of Microbiology and Parasitology, Faculty of Medical Science, Naresuan University, Phitsanulok, 65000, Thailand.

^b Center of Excellence in Medical Biotechnology, Faculty of Medical Science, Naresuan University, Phitsanulok, 65000, Thailand.

^c Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok, 65000, Thailand.

^d Department of Medical Microbiology and Infectious Disease, Institute of Infection and Immunity, UHW Main Building, Heath Park Hospital, Cardiff, CF14 4XN, UK.

*** Corresponding author.** Department of Microbiology and Parasitology, Faculty of Medical Science, Naresuan University, Phitsanulok, 65000, Thailand. Tel: +66 89 8565063; Fax: 66 55 964770.

E-mail address: pannikan@nu.ac.th (P.R. Niumsup).

ABSTRACT

This study investigated the prevalence of antibiotic-resistant *Escherichia coli* using extended-spectrum β -lactamase (ESBL) and AmpC β -lactamase as exemplars of multidrug-resistant phenotypes in poultry meat samples taken from open-air and supermarkets in Phitsanulok province, Northern Thailand. Two hundred and fifty poultry meat samples from open-air ($n = 147$) and supermarkets ($n = 103$) were analyzed. In total, 143 cefotaxime-resistant *E. coli* isolates comprising 78 isolates (53.1%) from open-air markets and 65 isolates (63.1%) from supermarkets were obtained. No significant difference could be observed in the prevalence of ESBL-positive *E. coli* between samples taken from open-air (70.5%) and supermarkets (69.2%). ESBL genotypes comprised of *bla*_{CTX-M-group 1} (69%), *bla*_{CTX-M-group 9} (13%), *bla*_{TEM-116} (1%), *bla*_{SHV-2a} (1%) and *bla*_{SHV-12} (1%) were detected. 39.5% of the ESBL-negative *E. coli* possessed *bla*_{CMY-2}. The *bla*_{CTX-M-group 1}, *bla*_{CTX-M-group 9} and *bla*_{CMY-2} were successfully transferred into *E. coli* by conjugation at high frequencies. Repetitive palindromic-PCR of some *bla*_{CTX-M} and *bla*_{CMY-2}-positive *E. coli* isolates revealed identical DNA patterns suggesting clonal spread. Phylogenetic grouping and MLST analysis revealed that 3 isolates were *E. coli* ST131. Of these, 2 isolates were ESBL-negative and carried *bla*_{CMY-2}. The other isolate was ESBL-positive and carried *bla*_{TEM-116}. This is the first study to demonstrate ESBL and AmpC genotypes in *E. coli* and the first discovery of human pathogen ST131 from Thai poultry meat. Our data raises serious concerns for food safety and biosecurity in the Thai food industry.

Keywords: poultry; meat; ESBL; CTX-M; AmpC; CMY-2; *E. coli*; ST131

1. Introduction

Resistance to broad spectrum β -lactams such as third-generation cephalosporins, monobactam and carbapenems in Enterobacteriaceae, especially *Escherichia coli*, is rapidly increasing (Pitout, 2013). Reports on extended-spectrum β -lactamase (ESBL) and/or AmpC β -lactamase-producing Enterobacteriaceae from clinical and environmental samples are continuously published from the majority of countries indicating a worldwide dissemination. Several types of ESBLs have been reported such as SHV, TEM and CTX-M. CTX-M is the most prevalent ESBL while CMY-2 is frequently encountered AmpC in human infections. ESBL- and AmpC-encoding genes are usually associated with mobile genetic elements which strongly facilitate their spread within a bacterial population (Jacoby 2009; Poirel, Bonnin, & Nordmann, 2012).

Contamination of meat with antibiotic-resistant bacteria has the potential to transfer to humans and is a clear public health concern. Several studies have shown that meat, especially poultry meat, is an important reservoir of antibiotic-resistant *E. coli* (Nguyen et al., 2016; Schwaiger et al., 2012). The prevalence of ESBL- and AmpC-positive *E. coli* as well as their respective resistant genes in different types of meat even in organic meat has been reported from several countries (Cohen Stuart et al., 2012; Egea et al., 2012; Ghodousi, Bonura, Di Noto, & Mammina, 2015; Kawamura, Goto, Nakane, & Arakawa, 2014). Furthermore, transmission of ESBL and AmpC-positive *E. coli* from meat to human has been previously reported outside Thailand (Overdevest et al., 2011; Vincent et al., 2010).

In Thailand, consumption of poultry meat is popular. Meat is often sold in open-air markets, traditional Thai markets which are seen extensively throughout the country, and western-style supermarkets which are becoming increasingly popular throughout Thailand.

Previous studies in Thailand have revealed the presence of antibiotic-resistant Enterobacteriaceae isolates in chicken meat and chicken rectal swab including those producing ESBL (Boonyasiri et al., 2014; Chaisatit, Tribuddharat, Pulsrikarn, & Dejsirilert, 2012; Trongjit, Angkittitrakul, & Chuanchuen, 2016). However, the prevalence of ESBL- and/or AmpC-producing *E. coli* from meat samples in Thailand remains poorly understood. This study investigated the prevalence of ESBL- and AmpC-encoding genes from poultry meat samples obtained from both open-air and supermarkets as well as determined *E. coli* pathogenicity groups.

2. Materials and Methods

2.1 Samplings

Samplings of fresh poultry meat were performed in 29 open-air markets and 22 supermarkets in Phitsanulok province, Northern Thailand. A total of 250 poultry meat samples (chicken = 218, duck = 14, bird = 18) from open-air markets ($n = 147$) and supermarkets ($n = 103$) were sampled. Frozen poultry meat was excluded from the study. All samples were originated from Thailand. Samples were maintained at 4 °C and processed immediately.

2.2 Isolation and identification of third generation cephalosporin-resistant *E. coli*

Twenty-five grams of each sample were homogenized with a Stomacher in 225 mL buffered peptone water (Oxoid, Basingstroke, UK). Then, 10 mL of this homogenate were enriched in 90 ml EE broth (Becton, Dickinson and Company, MD, USA) for 24 h at 37 °C. The enrichment was plated onto EMB agar (Oxoid) supplemented with 2 µg/mL cefotaxime (as an exemplar of broad-spectrum cephalosporins) (Sigma Aldrich, MO, USA) and incubated under aerobic condition for 24 h at 37 °C. Presumptive *E. coli*

colonies isolated from each sample were subcultured on Tryptic Soy Agar and incubated as described above for further characterizations. Species identification was performed by using RapID™ ONE System (REMEL Inc., KS, USA) according to the manufacturers' instructions and confirmed by sequencing of 16S rRNA gene (Lane, 1991).

2.3 Antimicrobial susceptibility and ESBL detection

All isolates were tested for susceptibility to 18 antimicrobial agents by disk diffusion method according to Clinical Laboratory Standards Institute (CLSI) protocols and the results were evaluated according to CLSI criteria (CLSI, 2013). The antibiotics tested were ampicillin, cefoxitin, ceftazidime, cefotaxime, cefpodoxime, cefepime, aztreonam, imipenem, amoxicillin/clavulanic acid, ampicillin/sulbactam, amikacin, gentamicin, doxycycline, tetracycline, ciprofloxacin, levofloxacin, chloramphenicol, and trimethoprim-sulfamethoxazole. Isolates showing intermediate results were considered as resistant. Minimum Inhibitory Concentration (MICs) were determined by broth microdilution method according to CLSI guidelines (CLSI, 2013). The MIC of each antimicrobial agent was defined as the lowest concentration, which inhibited visible growth of the organism.

Isolates were tested for ESBL production by combination disk method with ceftazidime and cefotaxime in the presence or absence of clavulanic acid, according to CLSI guidelines (CLSI, 2013).

2.4 Screening for ESBL- and AmpC-encoding genes by PCR and sequencing

All ESBL-producing isolates were investigated for the presence of ESBL-encoding genes; *bla*_{TEM}, *bla*_{SHV} and *bla*_{CTX-M}, by PCR as previously described (Dallenne, Da Costa, Decré, Favier, & Arlet, 2010; Woodford, Fagan, & Ellington, 2006). Isolates showing

resistance to cefoxitin were examined for the presence of AmpC-encoding genes by multiplex PCR (Pérez-Pérez, & Hanson, 2002). PCR products were analyzed by agarose gel electrophoresis.

Selected PCR products were analyzed by DNA sequencing. Amplicons were purified using a DNA purification kit (RBC Bioscience, New Taipei City, Taiwan) and sequenced by First BASE Laboratories (Selangor, Malaysia). The obtained sequences were compared with those available in the GenBank database using the BLAST algorithm available on the National Center for Biotechnology Information (NCBI) website (<http://www.ncbi.nlm.nih.gov>).

2.5 Conjugation experiments

To investigate the transfer of antibiotic resistance, conjugation experiments were carried out by broth mating method using rifampin-resistant *E. coli* DH5 α as the recipient. Cultures of donor and recipient cells were mixed and incubated overnight at 37 °C without shaking. Transconjugants were selected on Tryptic Soy Agar supplemented with rifampin (16 μ g/mL) and cefotaxime (1 μ g/mL). Conjugation frequency was expressed as the number of transconjugants divided by the number of recipient cells. Transferrable of antibiotic-resistant gene was confirmed by PCR. MICs of transconjugants were determined by broth microdilution method.

2.6 Repetitive-palindromic polymerase chain reaction (Rep-PCR)

E. coli isolates carrying either ESBL- or AmpC genes were typed by rep-PCR as described previously by Versalovic, Koeuth, and Lupski (1991).

2.7 Phylogenetic grouping and multilocus sequence typing (MLST) analysis

Phylogenetic group (A, B1, B2 and D) of ESBL- and pAmpC-producing *E. coli* was performed by a multiplex PCR assay for *chuA*, *yjaA* and DNA fragment TspE4C2 as previously described (Clermont, Bonacorsi, & Bingen, 2000). Isolate belonging to group B2 was investigated for the presence of *pabB* gene by PCR (Clermont et al., 2009). MLST was performed by amplification and sequencing of 7 housekeeping genes (*adhA*, *fumC*, *gyrB*, *icd*, *mdh*, *purA* and *recA*) according to the protocols from *E. coli* MLST website (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>).

2.8 Statistical analysis

Fisher's exact test was used to compare proportions using Minitab software version 15. The differences were considered statistically significant at $p < 0.05$.

3. Results

3.1 Isolation of cefotaxime-resistant (Ctx-R) *E. coli* from poultry meat samples and ESBL production

In this study, 250 poultry meat samples from open-air markets and supermarkets in Phitsanulok province, Northern Thailand were analyzed (Table 1). In total, 143 Ctx-R *E. coli* isolates comprising 78 isolates (53.1%, 95% CI = 45.0–61.1%) from open-air markets and 65 isolates (63.1%, 95% CI = 53.8–72.4%) from supermarkets were obtained. The percentages of Ctx-R *E. coli* isolates were not statistically different between samples obtained from open-air and supermarkets ($p = 0.121$). Of the 143 isolates, 70.5% (55/78, 95% CI = 60.4–80.6%) and 69.2% (45/65, 95% CI = 58.0–80.5%) from open-air and supermarkets, respectively, were ESBL-positive.

3.2 Antimicrobial susceptibility testing

Antimicrobial susceptibility of *E. coli* isolates recovered from poultry meat samples was determined (Table 2). β -Lactam susceptibility test revealed that all 143 *E. coli* isolates were resistant to ampicillin, cefotaxime and cefpodoxime and >90% were resistant to aztreonam and ceftazidime. Resistance to amoxicillin/clavulanate and ampicillin/sulbactam was 60.1% and 82.5%, respectively. A small number of isolates were resistant to imipenem (16.1%). High resistant rates (>60%) were also observed for cefepime, amikacin, doxycycline, tetracycline and ciprofloxacin. Furthermore, we observed that ESBL-positive isolates showed significantly higher resistant rates to cefepime, gentamicin, doxycycline, tetracycline, levofloxacin and trimethoprim/sulfamethoxazole ($p < 0.05$) than the ESBL-negative isolates (Table 2). In contrast, ESBL-negative isolates showed significantly higher resistant rates to ceftazidime, amoxicillin/clavulanate and ampicillin/sulbactam ($p < 0.05$).

3.3 Detection of antibiotic-resistant genes

Of the 143 Ctx-R isolates, 100 were shown to be ESBL-positive (Table 2). Detection of ESBL- and AmpC-encoding genes in Ctx-R *E. coli* isolates was performed and results were shown in Table 3. Of the 100 ESBL-positive isolates, 82 were found to carry genes encoding for *bla*_{CTX-M}. *bla*_{CTX-M-group 1} was the most prevalent ($n = 69$, 69%), followed by *bla*_{CTX-M-group 9} ($n = 13$, 13%). *bla*_{TEM} and *bla*_{SHV}-related ESBL-genes were also found i.e. *bla*_{TEM-116} ($n = 1$, 1%), *bla*_{SHV-2a} ($n = 1$, 1%) and *bla*_{SHV-12} ($n = 1$, 1%). Additionally, from the 43 ESBL-negative isolates, 17 (39.5%) were shown to be *bla*_{CMY-2}-positive.

3.4 Conjugation experiments

Conjugation experiments were performed with randomly selected *E. coli* containing *bla*_{CTX-M-group 1} (7 isolates), *bla*_{CTX-M-group 9} (2 isolates) and *bla*_{CMY-2} (3 isolates) as donors. The *bla*_{CTX-M-group 1}, *bla*_{CTX-M-group 9} and *bla*_{CMY-2} genes from donor isolates could be transferred to the recipient strain *E. coli* DH5 α with conjugation frequencies of 10^{-5} - 10^{-2} , 10^{-4} and 10^{-7} - 10^{-5} , respectively (Table 4). The presence of *bla*_{CTX-M-group 1}, *bla*_{CTX-M-group 9} and *bla*_{CMY-2} in transconjugants was confirmed by PCR. The cefotaxime MICs were markedly increased in transconjugants carrying *bla*_{CTX-M} (> 128 fold) and *bla*_{CMY-2} (> 16-32 fold) compared with the recipient strain *E. coli* DH5 α (Table 4).

3.5 Typing of ESBL- and AmpC-positive *E. coli* by rep-PCR

The DNA profiles of *bla*_{CTX-M} and *bla*_{CMY-2}-positive *E. coli* generated with rep-PCR primers showed diverse banding patterns. However, the identical rep-PCR profiles in some *E. coli* isolates were detected. The representative rep-PCR patterns of the isolates were shown in Fig. 1. Rep-PCR analysis differentiated 69 and 13 isolates of *bla*_{CTX-M group 1}- and *bla*_{CTX-M group 9}-positive *E. coli* into 34 and 9 distinct patterns, respectively. However, the identical patterns among isolates carrying *bla*_{CTX-M group 1} (Fig. 1A, Lanes 1 & 5, 2 & 3 and 4 & 6) and *bla*_{CTX-M group 9} (Fig. 1B, Lanes 1 & 2 and 3 & 4) were found. In addition, a total of 11 distinct DNA profiles were found among 17 *bla*_{CMY-2}-positive *E. coli* and identical DNA patterns were detected (Fig. 1C, Lanes 1-3). Importantly, *bla*_{CTX-M group 1}-positive *E. coli* isolates recovered from different open-air and supermarkets shared the same rep-PCR pattern (Fig. 1A). In addition, we found that 22 isolates carried an identical rep-PCR type but different CTX-M genes i.e. *bla*_{CTX-M group 1} ($n = 19$) and *bla*_{CTX-M group 9} ($n = 3$).

3.6 Phylogenetic grouping and analysis of sequence type (ST)

Phylogenetic grouping of 85 ESBL- and 17 AmpC-positive *E. coli* isolates were determined. The results showed that 51% belonged to group A, 27.5% to group B1 and 17.6% to group D. Only 4 isolates (3.9%) belonged to group B2 (Table 5). The prevalence of *bla*_{CTX-M} was significantly higher in the commensal phylogroups A and B1 than pathogenic groups B2 and D ($p < 0.001$). In contrast, the prevalence of *bla*_{CMY-2} was significantly higher in the pathogenic phylogroups D and B2 than commensal groups A and B1 ($p = 0.002$). Of the 4 isolates, belonging to group B2, 3 isolates contained the *pabB* gene, which is specific for subgroup I/ O25b. These 3 isolates were identified as ST131, by MLST analysis. One carried *bla*_{TEM-116} and 2 carried *bla*_{CMY-2}.

4. Discussion

In this study, poultry meat samples from open-air and supermarkets in Phitsanulok province, Northern Thailand were analyzed and a total of 143 Ctx-R *E. coli* isolates were recovered from all types of poultry meat (chicken, duck and bird, Table 1). A previous study in Thailand revealed the presence of ESBL-positive *E. coli* from chicken meat however a limited number of meat samples were investigated (Boonyasiri et al., 2014). In contrast, we found a comparatively high prevalence of ESBL-positive *E. coli* isolates from open-air (70.5%) and supermarkets (69.2%).

Antimicrobial susceptibility test of 143 Ctx-R *E. coli* isolates revealed the high resistant rates for broad-spectrum cephalosporins, aztreonam and ciprofloxacin (Table 2). These results were different from those reported previously from Thailand where *E. coli* obtained from chicken meat was mainly susceptible to third generation cephalosporins and ciprofloxacin (Chaisatit, Tribuddharat, Pulsrikarn, & Dejsirilert, 2012). Interestingly, ESBL-positive *E. coli* isolates showed significantly higher resistance rates to many

antibiotics than ESBL-negative isolates, consistent with the common feature of ESBL-positive bacteria (Woerther, Burdet, Chachaty, & Andremon, 2013). In addition, we found that all *E. coli* isolates, regardless of the ESBL production, exhibited multidrug-resistant (MDR) phenotype as defined by Magiorakos et al. (2012).

*bla*_{CTX-M} was commonly found among ESBL-positive isolates, although genes encoding for TEM-, and SHV-type ESBLs such as *bla*_{TEM-116}, *bla*_{SHV-2a} and *bla*_{SHV-12} were also detected (Table 3). Similar results were found in other studies which reported the predominance of *bla*_{CTX-M} in poultry meat samples (Cohen Stuart et al., 2012; Kawamura, Goto, Nakane, & Arakawa, 2014; Nguyen et al., 2016). The high prevalence of *bla*_{CTX-M} seen in this study coincided with the report on community-acquired CTX-M-positive infections in Thailand (Apisarnthanarak et al., 2007). For almost all ESBL-negative *E. coli* isolates, resistance to cefoxitin was noted (Table 2) suggesting that AmpC may be responsible for cefotaxime-resistant phenotypes. Accordingly, *bla*_{CMY-2} was found in 39.5% of ESBL-negative isolates.

For the rest of *E. coli* isolates, neither ESBL- nor AmpC-encoding genes were detected and were not further investigated. It is possible that overexpression of chromosomally-encoded AmpC β -lactamases, efflux pump overexpression or outer membrane alteration may contribute to reduced susceptibility to antibiotics (Pitout, 2013). It is also interesting to note that distribution of ESBL-genes was similar between open-air and supermarkets. In contrast, *bla*_{CMY-2} was found more frequently in poultry meat taken from open-air markets (Table 3).

Conjugation experiments revealed that *bla*_{CTX-M} and *bla*_{CMY-2} were successfully transferred to *E. coli* DH5 α at high frequencies. The increased cefotaxime MICs in transconjugants suggested that *bla*_{CTX-M} and *bla*_{CMY-2} are responsible for cefotaxime resistance. These results are consistent with the fact that horizontal gene transfer plays a

major role in the spread of *bla*_{CTX-M} and *bla*_{CMY-2} in *E. coli* (Woerther, Burdet, Chachaty, & Andremont, 2013; Sidjabat et al., 2014). Genotypic analysis of *bla*_{CTX-M} and *bla*_{CMY-2}-positive *E. coli* by rep-PCR revealed the genetic diversity among these isolates. However, identical genotypes were found among isolates carrying the same genes suggesting the clonal spread of MDR *E. coli* in poultry meat samples. This observation is further supported by the same genotypes being found in isolates obtained from open-air and supermarkets. Moreover, we found 22 identical isolates carrying different *bla*_{CTX-M} inferring the recent transfer of *bla*_{CTX-M} among dominant Thai *E. coli* clades.

Typing divided the *E. coli* isolates into 4 phylogenetic groups. Groups A and B1 constitute mainly the commensal strains while extra-intestinal pathogenic strains belong to groups B2 and D (Clermont, Bonacorsi, & Bingen, 2000). In this study, the majority of *E. coli* isolates carrying ESBL-genes (*bla*_{CTX-M}, *bla*_{SHV} or *bla*_{TEM}) significantly belonged to the low virulent groups A and B1. Similar results had been reported from *E. coli* isolates recovered from poultry meat in Spain and the Netherlands (Egea et al., 2012; Kluytmans et al., 2013). Our results suggest that *E. coli* isolates of the commensal origin may contribute to the dissemination of MDR *E. coli* within the community. It is also interesting to note that most *bla*_{CMY-2}-positive *E. coli* belonged to phylogroup D, consistent with previous reports (Oteo et al., 2010; Tamang et al., 2012).

Among the extraintestinal *E. coli*, O25b-ST 131, belonging to the phylogenetic group B2, has emerged as a highly virulent human pathogen worldwide. *E. coli* O25b-ST131 usually exhibits multidrug resistance and produces various types of ESBLs and AmpC. ST131 has been reported from different origins such as animals, environments and humans. Although the widespread occurrence of *E. coli* ST131 has been documented, its presence in raw meat is rare (Nicolas-Chanoine, Bertrand, & Madec, 2014). Previous studies in the Netherlands and Spain which many ESBL-producing *E. coli* isolates from

poultry meat were investigated but no ST131 was identified (Cohen Stuart et al., 2012; Egea et al., 2012; Overdevest et al., 2011). However other studies in Italy, Japan and Canada have reported the contamination of *E. coli* ST131 in poultry meat samples (Ghodousi, Bonura, Di Noto, & Mammina, 2015; Kawamura, Goto, Nakane, & Arakawa, 2014; Vincent et al., 2010). In our study, 3 MDR *E. coli* isolates were identified as *E. coli* B2-O25b ST131. Of these, 2 isolates were ESBL-negative and carried *bla*_{CMY-2}. The other ST131 isolate was an ESBL-positive and carried *bla*_{TEM-116}, the infrequent ESBL-gene found among ST131 isolates. In Thailand, all *E. coli* ST131 reported, to date, were recovered from clinical specimen (Netikul et al., 2014). Given the fact that *E. coli* ST131 is associated predominantly with community-onset urinary tract infections, our finding of ST131 isolates suggest that poultry meat could contribute to the dissemination of MDR and virulent *E. coli* within Thai community.

Our data demonstrates that poultry meat in Thailand acts as a reservoir for MDR *E. coli* especially those producing ESBL and AmpC. This is the first analysis of ESBL and AmpC genes and the first discovery of human pathogen ST131 from Thai poultry meat. Our results indicate that poultry meat may be a source of transmission of the MDR ST131 in Thailand.

325

326 **Acknowledgments**

327 This work was supported by Naresuan University (R2558B072). U. Tansawai was
328 supported by the Royal Golden Jubilee-PhD Program from Thailand Research Fund and
329 Rajamangala University of Technology Lanna (PHD/0054/2555). We would like to thank
330 staffs at the Department of Microbiology and Parasitology, Faculty of Medical Science,
331 Naresuan University for their technical assistance.

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

References

- Apisarnthanarak, A., Kiratisin, P., Saifon, P., Kitphati, R., Dejsirilert, S., & Mundy, L.M. (2007). Clinical and molecular epidemiology of community-onset, extended-spectrum β -lactamase-producing *Escherichia coli* infections in Thailand: a case-control study. *American Journal of Infection Control*, 35, 606-612.
- Boonyasiri, A., Tangkoskul, T., Seenama, C., Saiyarin, J., Tiengrim, S., & Thamlikitkul, V. (2014). Prevalence of antibiotic resistant bacteria in healthy adults, foods, food animals, and the environment in selected areas in Thailand. *Pathogens and Global Health*, 108, 235-245.
- Chaisatit, C., Tribuddharat, C., Pulsrikarn, C., & Dejsirilert, S. (2012). Molecular characterization of antibiotic-resistant bacteria in contaminated chicken meat sold at supermarkets in Bangkok, Thailand. *Japanese Journal of Infectious Disease*, 65, 527-534.
- Clermont, O., Bonacorsi, S., & Bingen E. (2000). Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Applied and Environmental Microbiology*, 66, 4555-4558.
- Clermont, O., Dhanji, H., Upton, M., Gibreel, T., Fox, A., Boyd, D., et al. (2009). Rapid detection of the O25b-ST131 clone of *Escherichia coli* encompassing the CTX-M-15-producing strains. *Journal of Antimicrobial Chemotherapy*, 64, 274-277.
- Clinical and Laboratory Standards Institute (CLSI) (2013). *Performance standards for antimicrobial susceptibility testing; twenty-third informational supplement. Document M100-S23*. Wayne, PA: CLSI.
- Cohen Stuart, J., van den Munckhof, T., Voets, G., Scharringa, J., Fluit, A., & Hall, M.L. (2012). Comparison of ESBL contamination in organic and conventional retail chicken meat. *International Journal of Food Microbiology*, 154, 212-214.

- 375 Dallenne, C., Da Costa, A., Decré, D., Favier, C., & Arlet, G. (2010). Development of a
 376 set of multiplex PCR assays for the detection of genes encoding important β -
 377 lactamases in Enterobacteriaceae. *Journal of Antimicrobial Chemotherapy*, 65,
 378 490-495.
- 379 Egea, P., López-Cerero, L., Torres, E., Gómez-Sánchez Mdel, C., Serrano, L., Navarro
 380 Sánchez-Ortiz, M.D., et al. (2012). Increased raw poultry meat colonization by
 381 extended spectrum β -lactamase-producing *Escherichia coli* in the south of Spain.
 382 *International Journal of Food Microbiology*, 159, 69-73.
- 383 Ghodousi, A., Bonura, C., Di Noto, A.M., & Mammina, C. (2015). Extended-spectrum β -
 384 lactamase, AmpC-producing, and fluoroquinolone-resistant *Escherichia coli* in
 385 retail broiler chicken meat, Italy. *Foodborne Pathogens and Disease*, 12, 619-625.
- 386 Jacoby, G.A. (2009). AmpC β -lactamases. *Clinical Microbiology Reviews*, 22: 161-182.
- 387 Kawamura, K., Goto, K., Nakane, K., & Arakawa, Y. (2014). Molecular epidemiology of
 388 extended-spectrum β -lactamases and isolated from retail foods including chicken
 389 meat in Japan. *Foodborne Pathogens and Disease*, 11, 104-110.
- 390 Kluytmans, J.A., Overdevest, I.T., Willemsen, I., Kluytmans-van den Bergh, M.F., van der
 391 Zwaluw, K., Heck, M., et al. (2013). Extended-spectrum β -lactamase-producing
 392 *Escherichia coli* from retail chicken meat and humans: comparison of strains,
 393 plasmids, resistance genes, and virulence factors. *Clinical Infectious Diseases*, 56,
 394 478-487.
- 395 Lane, D. J. (1991). 16S/23S rRNA sequencing. In E. Stackebrandt, & M. Goodfellow
 396 (Eds.), *Nucleic acid techniques in bacterial systematics* (pp. 115-175). Chichester,
 397 United Kingdom: John Wiley and Sons.
- 398 Magiorakos, A.P., Srinivasan, A., Carey, R.B., Carmeli, Y., Falagas, M.E., Giske, C.G., et
 399 al. (2012). Multidrug-resistant, extensively drug-resistant and pandrug-resistant

- 400 bacteria: an international expert proposal for interim standard definitions for
 401 acquired resistance. *Clinical Microbiology Infection*, 18, 268-281.
- 402 Netikul, T., Sidjabat, H.E., Paterson, D.L., Kamolvit, W., Tantisiriwat, W., Steen, J.A., et
 403 al. (2014). Characterization of an IncN2-type *bla*_{NDM-1}-carrying plasmid in
 404 *Escherichia coli* ST131 and *Klebsiella pneumoniae* ST11 and ST15 isolates in
 405 Thailand. *Journal Antimicrobial Chemotherapy*, 69, 3161-3163.
- 406 Nguyen, D. P., Nguyen, T.A., Le, T.H., Tran, N.M., Ngo, T.P., Dang, V.C., et al. (2016).
 407 Dissemination of extended-spectrum β -lactamase- and AmpC β -lactamase-
 408 producing *Escherichia coli* within the food distribution system of Ho Chi Minh
 409 City, Vietnam. *BioMed Research International* 2016, 8182096.
- 410 Nicolas-Chanoine, M.H., Bertrand, X., & Madec, J.Y. (2014). *Escherichia coli* ST131, an
 411 intriguing clonal group. *Clinical Microbiology Reviews*, 27, 543-574.
- 412 Oteo, J., Cercenado, E., Cuevas, O., Bautista, V., Delgado-Iribarren, A., Orden, B., et al.
 413 (2010). AmpC β -lactamases in *Escherichia coli*: emergence of CMY-2-producing
 414 virulent phylogroup D isolates belonging mainly to STs 57, 115, 354, 393, and
 415 420, and phylogroup B2 isolates belonging to the international clone O25b-ST131.
 416 *Diagnostic Microbiology and Infectious Disease*, 67, 270-276.
- 417 Overdevest, I., Willemsen, I., Rijnsburger, M., Eustace, A., Xu, L., Hawkey, P., et al.
 418 (2011). Extended-spectrum β -lactamase genes of *Escherichia coli* in chicken meat
 419 and humans, The Netherlands. *Emerging Infectious Diseases*, 17, 1216-1222.
- 420 Pérez-Pérez, F.J., & Hanson, N.D. (2002). Detection of plasmid-mediated AmpC β -
 421 lactamase genes in clinical isolates by using multiplex PCR. *Journal of Clinical*
 422 *Microbiology*, 4, 2153-2162.

- 423 Pitout, J.D. (2013). Enterobacteriaceae that produce extended-spectrum β -lactamases and
 424 AmpC β -lactamases in the community: the tip of the iceberg?. *Current*
 425 *Pharmaceutical Design*, 19, 257-263.
- 426 Poirel, L., Bonnin, R.A., & Nordmann, P. (2012). Genetic support and diversity of
 427 acquired extended-spectrum β -lactamases in Gram-negative rods. *Infection*,
 428 *Genetics and Evolution*, 12, 883-893.
- 429 Schwaiger, K., Huther, S., Hölzel, C., Kämpf, P., & Bauer, J. (2012). Prevalence of
 430 antibiotic-resistant enterobacteriaceae isolated from chicken and pork meat
 431 purchased at the slaughterhouse and at retail in Bavaria, Germany. *International*
 432 *Journal of Food Microbiology*, 154, 206-211.
- 433 Sidjabat, H.E., Seah, K.Y., Coleman, L., Sartor, A., Derrington, P., Heney, C., et al.
 434 (2014). Expansive spread of IncII plasmids carrying *bla*_{CMY-2} amongst *Escherichia*
 435 *coli*. *International Journal of Antimicrobial Agents*, 44, 203-208.
- 436 Tamang, M.D., Nam, H.M., Jang, G.C., Kim, S.R., Chae, M.H., Jung, S.C., et al. (2012).
 437 Molecular characterization of extended-spectrum β -lactamase-producing and
 438 plasmid-mediated AmpC β -lactamase-producing *Escherichia coli* isolated from
 439 stray dogs in South Korea. *Antimicrobial Agents and Chemotherapy*, 56, 2705-
 440 2712.
- 441 Trongjit, S., Angkittitrakul, S., & Chuanchuen, R. (2016). Occurrence and molecular
 442 characteristics of antimicrobial resistance of *Escherichia coli* from broilers, pigs
 443 and meat products in Thailand and Cambodia provinces. *Microbiology and*
 444 *Immunology*, 60, 575-585.
- 445 Versalovic, J., Koeuth, T., & Lupski, J.R. (1991). Distribution of repetitive DNA
 446 sequences in eubacteria and application to fingerprinting of bacterial genomes.
 447 *Nucleic Acids Research*, 19, 6823-6831.

Vincent, C., Boerlin, P., Daignault, D., Dozois, C.M., Dutil, L., Galanakis, C., et al.

(2010). Food reservoir for *Escherichia coli* causing urinary tract infections.

Emerging infectious diseases, 16, 88-95.

Woerther, P.L., Burdet, C., Chachaty, E., & Andremont, A. (2013). Trends in human fecal

carriage of extended-spectrum β -lactamases in the community: toward the

globalization of CTX-M. *Clinical Microbiology Reviews*, 26, 744-758.

Woodford, N., Fagan, E.J., & Ellington, M.J. (2006). Multiplex PCR for rapid detection of

genes encoding CTX-M extended-spectrum β -lactamases. *Journal of Antimicrobial*

Chemotherapy, 57, 154-155.

473

474

475 **Figure Caption**

476

477 **Fig. 1.** Representative rep-PCR profiles of the cefotaxime-resistant *E. coli* carrying *bla*_{CTX}-478 M-group 1 (A), *bla*_{CTX}-M-group 9(B) and *bla*_{CMY-2} (C), respectively.

479 M1, 1 kb Sharp Ladder (RBC Bioscience, New Taipei City, Taiwan)

480 M2, 100 bp Ladder RTU (GeneDireX, Taipei City, Taiwan)

Table 1 Occurrence of cefotaxime-resistant *E. coli* from poultry meat samples

Types of Samples	No. of <i>E. coli</i>	No. of ESBL- positive <i>E.coli</i>
open air markets ($n = 147$)		
Chicken ($n = 126$)	69	46
Duck ($n = 3$)	0	0
Bird ($n = 18$)	9	9
Total	78 (53.1%)	55 (70.5%)
supermarkets ($n = 103$)		
Chicken ($n = 92$)	61	41
Duck ($n = 11$)	4	4
Total	65 (63.1%)	45 (69.2%)

Table 2 Antimicrobial resistance (%) among cefotaxime-resistant *E. coli* isolates

Antimicrobial agents	Total (<i>n</i> = 143)	ESBL production		
		ESBL-positive (<i>n</i> = 100)	ESBL-negative (<i>n</i> = 43)	p values
ampicillin	100	100	100	NS
cefoxitin	41.3	17.0	97.7	< 0.001
ceftazidime	93	90	100	0.033
cefotaxime	100	100	100	NS
cefodoxime	100	100	100	NS
cefepime	60.1	83.0	7.0	< 0.001
aztreonam	95.8	96.0	95.3	NS
imipenem	16.1	16.0	16.3	NS
amoxicillin/clavulanate	60.1	43.0	100	< 0.001
ampicillin/sulbactam	82.5	75.0	100	< 0.001
amikacin	72	72.0	72.1	1
gentamicin	57.3	67.0	34.9	< 0.001
doxycycline	62.9	78.0	27.9	< 0.001
tetracycline	71.3	86.0	37.2	< 0.001
ciprofloxacin	62.9	64.0	60.5	NS
levofloxacin	26.6	34.0	9.3	0.002
chloramphenicol	34.3	39.0	23.3	NS
trimethoprim/sulfamethoxazole	45.5	59.0	14.0	< 0.001

NS, Not significance

Table 3 Prevalence of genes encoding for ESBL and AmpC among cefotaxime-resistant *E. coli* isolates.

β -lactamase genes	ESBL-positive <i>E. coli</i> (<i>n</i> = 100)			ESBL-negative <i>E. coli</i> (<i>n</i> = 43)		
	open-air markets (<i>n</i> = 55)	supermarkets (<i>n</i> = 45)	Total	open-air markets (<i>n</i> = 23)	supermarkets (<i>n</i> = 20)	Total
<i>bla</i> _{CTX-M group 1}	25	18	43	0	0	0
<i>bla</i> _{CTX-M group 1} + <i>bla</i> _{TEM-1}	12	14	26	0	0	0
<i>bla</i> _{CTX-M group 9}	5	1	6	0	0	0
<i>bla</i> _{CTX-M group 9} + <i>bla</i> _{TEM-1}	2	5	7	0	0	0
<i>bla</i> _{SHV-2a}	1	0	1	0	0	0
<i>bla</i> _{SHV-12}	0	1	1	0	0	0
<i>bla</i> _{TEM-116}	1	0	1	0	0	0
<i>bla</i> _{CMY-2}	0	0	0	14	2	16
<i>bla</i> _{CMY-2} + <i>bla</i> _{TEM-1}	0	0	0	1	0	1
Total	46	39	85	15	2	17

Table 4 Conjugation frequency, transferred gene and cefotaxime MICs for *E. coli* DH5 α and the respective transconjugants

Strain	Frequency of transfer	Transferred gene	Cefotaxime MIC ($\mu\text{g/mL}$)
<i>E. coli</i> DH5 α^a	-	-	< 0.0625
EC2_Tc	4.2×10^{-3}	<i>bla</i> _{CTX-group 1}	> 8
EC103_Tc	2.6×10^{-3}	<i>bla</i> _{CTX-group 1}	> 8
EC107_Tc	4.4×10^{-3}	<i>bla</i> _{CTX-group 1}	> 8
EC109_Tc	6.8×10^{-5}	<i>bla</i> _{CTX-group 1}	> 8
EC121_Tc	2.5×10^{-5}	<i>bla</i> _{CTX-group 1}	> 8
EC129_Tc	8.1×10^{-2}	<i>bla</i> _{CTX-group 1}	> 8
EC146_Tc	7.5×10^{-2}	<i>bla</i> _{CTX-group 1}	> 8
EC70_Tc	2.4×10^{-4}	<i>bla</i> _{CTX-group 9}	> 8
EC119_Tc	4.7×10^{-4}	<i>bla</i> _{CTX-group 9}	> 8
EC125_Tc	1.5×10^{-5}	<i>bla</i> _{CMY-2}	2
EC128_Tc	1.6×10^{-6}	<i>bla</i> _{CMY-2}	1


EC132_Tc	1.9×10^{-7}	<i>bla</i> _{CMY-2}	1
----------	----------------------	-----------------------------	---

^a rifampin-resistant *E. coli* DH5 α was used a recipient strain.

Tc, transconjugant

Table 5 Distribution of ESBL- and AmpC-encoding genes among *E. coli* phylogroups ($n = 102$)

Phylogenetic groups	No. (%) of isolates producing:				Total (%)
	<i>bla</i> _{TEM-116}	<i>bla</i> _{SHV}	<i>bla</i> _{CTX-M}	<i>bla</i> _{CMY-2}	
A	0 (0)	1 (1.0)	50 (49.0)	1 (1.0)	52 (51.0)
B1	0 (0)	1 (1.0)	26 (25.5)	1 (1.0)	28 (27.5)
B2	1 (1.0)	0 (0)	0 (0)	3 (2.9)	4 (3.9)
D	0 (0)	0 (0)	6 (5.9)	12 (11.7)	18 (17.6)
Total	1 (1.0)	2 (2.0)	82 (80.4)	17 (16.6)	102 (100)

Fig 1

Highlights

High prevalence of ESBL-producing *E. coli* in poultry meat in Thailand was observed.

E. coli carrying *bla*_{CTX-M} and *bla*_{CMY-2} isolates were found in poultry meat samples.

Clonal spread of *bla*_{CTX-M} and *bla*_{CMY-2}-positive *E. coli* from poultry meat was detected.

E. coli ST131 isolates carrying *bla*_{TEM-116} or *bla*_{CMY-2} have emerged from poultry meat.