

## Research Note

# Resistance to Antimicrobial Agents among *Salmonella* Isolates Recovered from Layer Farms and Eggs in the Caribbean Region

ABIODUN ADESIYUN,<sup>1\*</sup> LLOYD WEBB,<sup>2</sup> LISA MUSAI,<sup>3</sup> BOWEN LOUISON,<sup>4</sup> GEORGE JOSEPH,<sup>5</sup> ALVA STEWART-JOHNSON,<sup>1</sup> SANNANDAN SAMLAL,<sup>1</sup> AND SHELLY RODRIGO<sup>6</sup>

<sup>1</sup>School of Veterinary Medicine, Faculty of Medical Sciences, University of the West Indies, St. Augustine, Trinidad and Tobago, West Indies; <sup>2</sup>Institute of Public Health Studies, College of Veterinary Medicine, Nursing and Allied Health, Tuskegee University, Tuskegee, Alabama 36088, USA; <sup>3</sup>Poultry Surveillance Unit, Ministry of Food Production, Port of Spain, Trinidad and Tobago, West Indies; <sup>4</sup>Ministry of Agriculture, Forestry and Fisheries, Ministerial Complex, Tanteen, St. George's, Grenada, West Indies; <sup>5</sup>Veterinary and Livestock Services, Ministry of Agriculture, Food Production, Rural Development and Fisheries, Castries, St. Lucia, West Indies; and <sup>6</sup>Department of Public Health and Preventive Medicine, School of Medicine, St. George's University, University Centre, Grenada, West Indies

MS 14-162: Received 7 April 2014/Accepted 16 June 2014

## ABSTRACT

This investigation determined the frequency of resistance of 84 isolates of *Salmonella* comprising 14 serotypes recovered from layer farms in three Caribbean countries (Trinidad and Tobago, Grenada, and St. Lucia) to eight antimicrobial agents, using the disc diffusion method. Resistance among isolates of *Salmonella* was related to the country of recovery, type of sample, size of layer farms, and isolate serotype. Overall, all (100.0%) of the isolates exhibited resistance to one or more of seven antimicrobial agents tested, and all were susceptible to chloramphenicol. The resistance detected ranged from 11.9% to sulphamethoxazole-trimethoprim (SXT) to 100.0% to erythromycin. The difference was, however, not statistically significant ( $P = 0.23$ ). Across countries, for types of samples that yielded *Salmonella*, significant differences in frequency of resistance were detected only to SXT ( $P = 0.002$ ) in Trinidad and Tobago and to gentamycin ( $P = 0.027$ ) in St. Lucia. For the three countries, the frequency of resistance to antimicrobial agents was significantly different for ampicillin ( $P = 0.001$ ) and SXT ( $P = 0.032$ ). A total of 83 (98.8%) of the 84 isolates exhibited 39 multidrug resistance patterns. Farm size significantly ( $P = 0.032$ ) affected the frequency of resistance to kanamycin across the countries. Overall, among the 14 serotypes of *Salmonella* tested, significant ( $P < 0.05$ ) differences in frequency of resistance were detected to kanamycin, ampicillin, and SXT. Results suggest that the relatively high frequency of resistance to six of the antimicrobial agents (erythromycin, streptomycin, gentamycin, kanamycin, ampicillin, and tetracycline) tested and the multidrug resistance detected may pose prophylactic and therapeutic concerns for chicken layer farms in the three countries studied.

Salmonellosis is a well-established disease of poultry (layers and broilers) worldwide, causing morbidity and mortality and, thus, economic losses (5, 21). Complicating control of salmonellosis in the poultry industry is the fact that, in addition to horizontal transmission on farms, vertical transmission from parent stock to offspring is also common; this makes control a challenge (9, 17) and poses a food safety hazard to consumers of contaminated eggs and egg products (3, 26).

Although vaccination to prevent salmonellosis has been used successfully on layer farms in several countries (9, 21, 36), the existence of several pathogenic non-host-adapted *Salmonella* serovars makes successful vaccination difficult. This is particularly a problem in most developing countries, where vaccines are not routinely available or are not used by poultry farmers and where prevention and treatment of salmonellosis have relied predominantly on good sanitary practices and the prophylactic and therapeutic

administration of antimicrobial agents in feeds and water (5, 29, 36). The improper use of antimicrobial agents as a result of failed preventive measures has led to the development of resistance to antimicrobial agents, with therapeutic consequences (8, 29).

In addition to the recovery of *Salmonella* isolates from several livestock species, including laying birds and their environments (1, 35), there are reports of increased resistance to antimicrobial agents among isolates (7, 12, 20). *Salmonella* isolates from the shells and contents of table eggs have been reported to exhibit resistance to antimicrobial agents (10, 16). Uncontrolled therapeutic use of antimicrobial agents in the livestock industry (15, 20) and the use of antimicrobial agents as additives in animal feeds to promote growth (11) may pose food safety, public health, and therapeutic problems in animal and human diseases (12, 31).

There are reports from various countries of variable antimicrobial resistance of *Salmonella* isolates from layers and farm environments (19, 24, 30, 33). It has been documented that antimicrobial resistance among *Salmonella*

\* Author for correspondence. Tel: (868)645-4481; Fax: (868)645-7428; E-mail: Abiodun.Adesiyun@sta.uwi.edu.

serovars is affected by various factors, such as the type of sample (cloacae, fecal, or feed) taken from poultry farms (11, 31) and the serotypes (35).

In the Caribbean region, the only report to document the antimicrobial resistance of *Salmonella* isolates recovered from poultry or table eggs was a 2005 study conducted on table eggs in Trinidad and Tobago (3). To date, there is a dearth of information on the antibiograms of *Salmonella* recovered from layer farms in the region. One objective of this study, therefore, was to determine the antibiograms of *Salmonella* recovered from layer farms in three Caribbean countries (Trinidad and Tobago, Grenada, and St. Lucia), and another was to relate the prevalence of resistance to country of origin, size of farms, type of sample, and serotypes of the isolates.

## MATERIALS AND METHODS

**Countries where study was conducted.** The study was conducted in three Caribbean countries, namely, Trinidad and Tobago, Grenada, and St. Lucia, during the periods 7 September to 3 October 2011, 17 to 21 October 2011, and 14 to 18 November 2011, respectively.

**Sample type and collection.** Layer farms (small, <5,000 birds; medium, 5,000 to 10,000 birds; and large, >10,000 birds) were visited across the three countries, and the following types of samples were collected: cloacal swabs, environmental swabs of deep litters, freshly laid eggs, and poultry feeds. The procedures used for sample collection and transportation to the laboratory were described previously (4).

**Processing of samples.** Standard methods were used to process the samples collected and to isolate *Salmonella* (4, 6, 23). Briefly, samples were preenriched in buffered peptone water (Oxoid, Ltd., Hampshire, England) for 18 to 24 h at 37°C and then were enriched in tetrathionate broth (Oxoid) and in Rappaport-Vassiliadis broth (Oxoid) incubated for 18 to 24 h at 37 and 42°C, respectively.

**Isolation and identification of *Salmonella* serovars.** The procedures used to isolate *Salmonella* serovars from the samples were described earlier (4). Briefly, the selective broths (tetrathionate and Rappaport-Vassiliadis) were subcultured onto xylose-lysine deoxycholate agar and brilliant green agar (Oxoid) and were incubated aerobically at 37°C for 24 h. All colonies suggestive of *Salmonella* serovars on the selective agar were purified on blood agar plates (Oxoid) at 37°C for 24 h and then were subjected to biochemical tests, including triple sugar iron agar, urea, citrate, methyl red, sulfide-indole-motility medium, *o*-nitrophenyl-*p*-galactopyranoside, and lysine iron agar (Oxoid) (6, 23). Initially, serological typing of all isolates biochemically identified as *Salmonella* serovars was done using a commercial slide agglutination test kit *Salmonella* polyvalent antiserum (A-I & Vi, Difco, Detroit, MI). Complete confirmation and serotyping of *Salmonella* isolates were done at the Caribbean Epidemiology Center, Port of Spain, Trinidad and Tobago, which serves as the regional laboratory for the serotyping of *Salmonella*.

**Determination of resistance to antimicrobial agents.** The antimicrobial resistance of 84 isolates of *Salmonella* recovered from 35 layer farms in the three countries (Table 1) was determined using the disc diffusion method according to the National Committee for Clinical Laboratory Standards (NCCLS)

TABLE 1. Farm sources of *Salmonella* isolates

Country	Total no. of poultry farms	No. (%) of farms sampled <sup>a</sup>	No. of <i>Salmonella</i> isolates tested
Trinidad and Tobago	23	10 (43.5) <sup>b</sup>	43
Grenada	21	10 (47.6) <sup>c</sup>	9
St. Lucia	37	15 (40.5) <sup>d</sup>	32
Total	81	35 (43.2)	84

<sup>a</sup> Types of samples collected: cloacal swabs, environmental samples, eggs, and feeds.

<sup>b</sup> Five large, two medium, and three small farms.

<sup>c</sup> One medium and nine small farms.

<sup>d</sup> Four large and 11 small farms.

(27) guidelines. For the study, the following antimicrobial agents and concentrations were used (all from Difco): erythromycin (E; 15 µg), streptomycin (S; 10 µg), gentamycin (CN; 10 µg), kanamycin (K; 30 µg), ampicillin (AMP; 10 µg), tetracycline (TE; 15 µg), sulphamethoxazole-trimethoprim (SXT; 23.75/1.25 µg), and chloramphenicol (C; 30 µg). The tests were performed on Mueller Hinton agar (Difco) followed by aerobic incubation at 35°C for 24 h. The zones of inhibition were interpreted as recommended by the disc manufacturer and NCCLS (27).

**Statistical analyses.** Data were analyzed using the Statistical Package for Social Sciences (version 22, IBM Corp., Somers, NY). Chi-square analysis was conducted to determine whether there were statistically significant differences in the frequency of resistance to antimicrobial agents among *Salmonella* isolates recovered among and within the three countries and among sources, irrespective of the country of origin. The differences in the frequency of resistance among the various serotypes of *Salmonella* and the predominant resistance patterns were also subjected to analysis. The level of significance was determined at an alpha level of 0.05.

## RESULTS AND DISCUSSION

**Frequency of resistance of *Salmonella* isolates to antimicrobial agents.** It was of therapeutic significance that all 84 *Salmonella* isolates recovered from three Caribbean countries (Trinidad and Tobago, Grenada, and St. Lucia) exhibited resistance to one or more of seven antimicrobial agents used and that all were sensitive to chloramphenicol, an antimicrobial agent not commonly used in the poultry industry in the region (Table 2). A similarly high prevalence of resistance (98.4%) was reported for isolates of *Salmonella* recovered from poultry houses in Chad (12). The use, or overuse, of antimicrobial agents in the poultry industry and on livestock farms has been shown to result in increased resistance to antimicrobial agents (8, 29), with the resultant therapeutic failures causing economic losses. Equally of therapeutic significance is that, of the six antimicrobial agents used in the poultry industry in the Caribbean region, SXT was the most effective against the *Salmonella* isolates tested from the three countries, with only 11.9% exhibiting resistance. Makaya et al. (24) in a study in Zimbabwe reported a 0.0% prevalence of resistance to SXT. In the only published report on the antibiograms of *Salmonella* isolates from eggs (shells and contents) in the

TABLE 2. Frequency of resistance of *Salmonella* isolate by type of samples from three countries<sup>a</sup>

Country	Source of sample	No. of isolates tested		No. (%) of resistant isolates <sup>b</sup>		No. (%) resistant to <sup>c</sup> :							
		E	S	P value	CN	P value	K	P value	AMP	P value	TE	P value	SXT
Trinidad and Tobago <sup>d</sup>	Feed	6	6 (100.0)	6 (100.0)	4 (66.7)	0.131	5 (83.3)	0.162	1 (16.7)	0.114	2 (33.3)	0.569	1 (16.7)
	Egg shell surface	9	9 (100.0)	9 (100.0)	9 (100.0)		7 (77.8)		7 (77.8)		6 (66.7)		2 (22.2)
	Environment	25	25 (100.0)	25 (100.0)	23 (92.0)		11 (44.0)		11 (44.0)		12 (48.0)		4 (16.0)
Grenada	Layers	3	3 (100.0)	3 (100.0)	2 (66.7)		2 (66.7)		1 (33.3)		1 (33.3)		0 (0.0)
	Total	43	43 (100.0)	43 (100.0)	38 (88.4)		25 (58.1)		20 (46.5)		21 (48.8)		7 (16.3)
	Environment	3	3 (100.0)	3 (100.0)	2 (66.7)	0.134	2 (66.7)	0.571	2 (66.7)	0.571	2 (66.7)	1.000	1 (33.3)
St. Lucia	Layers	6	6 (100.0)	6 (100.0)	6 (100.0)		5 (83.3)		5 (83.3)		4 (66.7)		0 (0.0)
	Total	9	9 (100.0)	9 (100.0)	8 (88.9)		7 (77.8)		7 (77.8)		6 (66.7)		1 (11.1)
	Environment	11	11 (100.0)	11 (100.0)	11 (100.0)	NA	3 (27.3)	0.017	2 (18.2)	0.035	1 (9.1)	0.673	9 (81.8)
All countries	Layers	21	21 (100.0)	21 (100.0)	21 (100.0)		15 (71.4)		12 (57.1)		3 (14.3)		11 (52.4)
	Total	32	32 (100.0)	32 (100.0)	32 (100.0)		18 (56.3)		14 (43.8)		4 (12.5)		20 (62.5)
	Feed	6	6 (100.0)	6 (100.0)	4 (66.7)	0.056	5 (83.3)	0.014	1 (16.7)	0.034	2 (33.3)	0.183	1 (16.7)
	Egg shell surface	9	9 (100.0)	9 (100.0)	9 (100.0)		7 (77.8)		7 (77.8)		6 (66.7)		2 (22.2)
	Environment	39	39 (100.0)	39 (100.0)	36 (92.3)		16 (41.0)		15 (38.5)		15 (38.5)		14 (35.9)
	Layers	30	30 (100.0)	30 (100.0)	29 (96.7)		22 (73.3)		18 (60.0)		8 (26.7)		11 (36.7)
All sources		84	84 (100.0)	84 (100.0)	78 (92.9)	0.137	50 (59.5)	0.491	41 (48.8)	0.179	31 (36.9)	0.001	28 (33.3)

<sup>a</sup> All isolates were sensitive to chloramphenicol (30 µg).<sup>b</sup> Resistant to one or more antimicrobial agents.<sup>c</sup> E, erythromycin (5 µg); S, streptomycin (10 µg); CN, kanamycin (10 µg); K, kanamycin (30 µg); AMP, ampicillin (10 µg); TE, tetracycline (15 µg); SXT, sulphonmethoxazole-trimethoprim (23.75/1.25 µg).<sup>d</sup> Of all the feed and freshly laid eggs sampled on farms from the three countries, *Salmonella* isolates were recovered only from Trinidad and Tobago samples.

TABLE 3. Antimicrobial resistance patterns among *Salmonella* isolates from three Caribbean countries

Country	Resistance patterns detected <sup>a</sup>	No. (%) of <i>Salmonella</i> isolates <sup>b</sup>
Trinidad and Tobago	S-E	4 (9.3)
	S-E-CN	4 (9.3)
	S-AMP-E	3 (7.0)
	S-E-K-CN	3 (7.0)
	S-AMP-E-CN	3 (7.0)
	S-AMP-E-K-CN	3 (7.0)
	SXT-S-AMP-E-K-CN	3 (7.0)
	SXT-S-E	2 (4.7)
	TE-S-E-CN	2 (4.7)
	TE-S-AMP-E	2 (4.7)
	S-AMP-E-K	2 (4.7)
Grenada	Others <sup>c</sup>	12 (27.9)
	S-AMP-E-K-CN	4 (44.4)
	S-E-K-CN	2 (22.2)
St. Lucia	Others <sup>d</sup>	3 (33.3)
	TE-S-E	9 (28.1)
	S-E-K-CN	8 (25.0)
	TE-S-E-K	3 (9.4)
	TE-S-E-CN	3 (9.4)
	S-E-CN	2 (6.3)
	TE-S-E-K-CN	2 (6.3)
Others <sup>e</sup>	Others <sup>e</sup>	5 (15.6)

<sup>a</sup> S, streptomycin; E, erythromycin; CN, gentamycin; AMP, ampicillin; K, kanamycin; SXT, sulphamethoxazole-trimethoprim; TE, tetracycline.

<sup>b</sup> Of a total of 43 isolates tested in Trinidad and Tobago, 9 isolates tested in Grenada, and 32 isolates tested in St. Lucia.

<sup>c</sup> Other patterns (one isolate each) include the following: TE-S-AMP-E-K, AMP-E, AMP-E-K-CN, SXT-S-E-K-CN, SXT-TE-S-AMP-E-K-CN, SXT-S-E-CN, TE-S-AMP-E-K-CN, E-CN, E-K, S-E-K, and SXT-S-E-K.

<sup>d</sup> Other patterns (one isolate each) include: S-AMP-E, TE-S-AMP-E-K-CN, and E.

<sup>e</sup> Other patterns (one isolate each) include: SXT-S-E-CN, S-AMP-E, TE-S-AMP-E-CN, TE-S-AMP-E-K-CN, and TE-S-E-K.

region, Adesiyun et al. (2) documented that of the 74 isolates of *Salmonella* tested, 17 (22.9%) exhibited resistance to one or more antimicrobial agents (S, TE, K, nalidixic acid, enrofloxacin, CN, and SXT) and, more importantly, that all were sensitive to SXT (compared with 79.1% of the isolates sensitive to the antimicrobial agent in the current study). There is, therefore, a possibility that the increased use of SXT is contributing to the higher frequency of resistance detected, although it is considerably lower than that detected for other antimicrobial agents tested.

Not unexpectedly, across and within countries a statistically significant frequency of resistance was detected to some of the antimicrobial agents, depending on the type of samples from which the *Salmonella* isolates were recovered. Statistically significant differences were detected in St. Lucia for CN ( $P = 0.027$ , from farm environment and cloacal swabs of layers). Across the countries and within Trinidad and Tobago, resistance was significantly higher to AMP and SXT ( $P < 0.05$ ) primarily due to *Salmonella* isolates from egg shells.

Other significant findings were that the frequency of resistance to a number of antimicrobial agents was relatively high, with a range of 48.8% (K) to 100.0% (E), and that, across the three countries studied, the frequency of antimicrobial resistance differed significantly for the individual antimicrobial agents (the lowest frequency of 11.9% to SXT; highest resistance to E). These findings suggest differences in the use of antimicrobial agents in the three countries.

The finding that 100.0% of the isolates tested in the current study were resistant to erythromycin agrees with the report of Akter et al. (5), who also reported that all isolates of *Salmonella* tested were resistant to the antibiotic. It is, however, pertinent to mention that erythromycin is not routinely used in clinical settings and animal husbandry to prevent or treat salmonellosis but was used in the current study to characterize the isolates.

The frequency of resistance to streptomycin detected in the current study (92.9%) is considerably higher than the 21.9 to 52.9% reported for poultry farms in other countries (13, 34). Of note also is the finding that the prevalence of resistance to CN found in the present study (59.5%) is much higher than that reported by others for isolates recovered from poultry farms (0.0 to 17%) (12, 25, 33).

For kanamycin, 48.8% of the isolates were resistant to the antimicrobial agent, which is comparable to the 50.0% prevalence of resistance reported for *Salmonella* isolates from layer farms in Bangladesh (18).

Of the *Salmonella* isolates from all sources in the three countries studied, 36.9% were resistant to AMP, which is very high compared with the prevalence of <3 and 3% reported by others (28, 34).

In the current study, 33.3% of the isolates of *Salmonella* were resistant to TE, a finding considerably lower than the prevalence reported for isolates from similar sources in other countries (93 to 100.0%) (14, 18, 22).

#### Frequency of resistance patterns of *Salmonella* isolates to antimicrobial agents by country of origin.

Table 3 shows the frequency of resistance patterns displayed, by country of origin. Overall, for the 84 *Salmonella* isolates tested from the three countries, 83 (98.8%) exhibited multidrug resistance. For the 43 *Salmonella* isolates tested from Trinidad and Tobago, 23 multiresistance patterns were determined, compared with 5 patterns for 9 isolates from Grenada and 11 patterns detected among 32 isolates from St. Lucia. The predominant patterns detected were S-E and S-E-CN (9.3% each), S-AMP-E-K-CN (44.4%), and TE-S-E (28.1%) for isolates from Trinidad and Tobago, Grenada, and St. Lucia, respectively. The differences were, however, not statistically significant ( $P = 0.23$ ).

The occurrence of a high frequency of multidrug resistance (98.8%) detected, in addition to the detection of a total of 39 multiresistance patterns in the current study, is further evidence of a potential for therapeutic failure that may be associated with the use of the seven antimicrobial agents across the three countries. Our findings are in agreement with the report on *Salmonella* isolates from poultry houses in Pakistan (25); from country to country,

TABLE 4. Occurrence of resistance among *Salmonella* isolates by size of farm<sup>a</sup>

Country	Size of layer farm <sup>b</sup>	No. of <i>Salmonella</i> isolates tested	No. (%) resistant to <sup>c</sup> :										
			E	S	P value	CN	P value	K	P value	AMP	P value	TE	P value
Trinidad and Tobago	Large	36	36 (100.0)	32 (88.9)	0.811	23 (63.9)	0.083	14 (38.9)	0.023	18 (50.0)	0.729	7 (19.4)	0.202
	Small	7	7 (100.0)	6 (85.7)	2 (28.6)	6 (85.7)	3 (42.9)	0 (0.0)	3 (42.9)	0 (0.0)	0 (0.0)	8 (22.2)	0.637
Grenada	Medium	3	3 (100.0)	3 (100.0)	0.453	3 (100.0)	0.257	2 (66.7)	1.000	0 (0.0)	0 (0.0)	0 (0.0)	NA
	Small	6	6 (100.0)	5 (83.3)	4 (66.7)	4 (66.7)	4 (66.7)	4 (66.7)	4 (66.7)	1 (16.7)	0 (0.0)	0 (0.0)	NA
St. Lucia	Large	27	27 (100.0)	27 (100.0)	NA	17 (63.0)	0.075	12 (44.4)	0.854	3 (11.1)	0.581	16 (59.3)	0.379
	Small	5	5 (100.0)	5 (100.0)	1 (20.0)	2 (40.0)	1 (20.0)	1 (20.0)	1 (20.0)	4 (80.0)	1 (3.7)	0 (0.0)	0.662
Total	Large	63	63 (100.0)	59 (93.7)	0.698	40 (63.5)	0.06	26 (41.3)	0.032	21 (33.3)	0.382	22 (34.9)	0.361
	Medium	3	3 (100.0)	3 (100.0)	3 (100.0)	3 (100.0)	3 (100.0)	3 (100.0)	2 (66.7)	0 (0.0)	0 (0.0)	9 (14.3)	0.487
	Small	18	18 (100.0)	16 (88.9)	7 (38.9)	12 (66.7)	8 (44.4)	8 (44.4)	5 (27.8)	1 (5.6)	0 (0.0)	0 (0.0)	1 (5.6)

<sup>a</sup> All isolates were sensitive to chloramphenicol (30 µg).<sup>b</sup> Isolates of *Salmonella* were recovered from large and small farms in Trinidad and Tobago, medium and small farms in Grenada, and large and small farms in St. Lucia.<sup>c</sup> E, erythromycin (5 µg); S, streptomycin (10 µg); CN, gentamycin (10 µg); AMP, ampicillin (10 µg); TE, tetracycline (15 µg); SXT, sulphamethoxazole-trimethoprim (27.75/1.25 µg); NA, not available.

multiresistance patterns are known to vary and to reflect the use of antimicrobial agents in the respective poultry industries (15, 19, 24, 25).

**Occurrence of resistance among *Salmonella* isolates by size of farm.** For the three countries combined, differences in the frequency of resistance for the three sizes of farm were detected to be statistically significantly different ( $P = 0.032$ ) for only kanamycin (41.3, 100.0, and 66.7% for large, medium, and small farms, respectively; Table 4).

Within countries, only in Trinidad and Tobago were statistically significant ( $P = 0.023$ ) differences detected; for kanamycin, resistance was higher for small farms (85.7%) compared with large farms (38.9%). Among countries, regardless of the source of sample, significant differences in the frequency of resistance were detected only for AMP, TE, and SXT. For AMP, the frequency of resistance differed significantly ( $P = 0.001$ ) among countries, 21 (48.8%) of 43, 6 (66.7%) of 9, and 4 (12.5%) of 32 in Trinidad and Tobago, Grenada, and St. Lucia, respectively. For TE, the frequency of resistance was statistically significantly different ( $P = 0.000$ ) across countries, 7 (16.3%) of 43, 1 (11.1%) of 9, and 20 (62.5%) of 32 in Trinidad and Tobago, Grenada, and St. Lucia, respectively. Finally, for SXT, the frequency of resistance was 9 (20.9%) of 43 in Trinidad and Tobago, 0 (0.0%) of 9 in Grenada, and 1 (3.1%) of 32 in St. Lucia; the differences were statistically significant ( $P = 0.032$ ).

In the current study, size of farm was found to have a significant effect only on the frequency of resistance to kanamycin across the three countries and within Trinidad and Tobago; this may reflect differences in management systems and the use of antimicrobial agents, as reported elsewhere (4, 32).

**Frequency of resistance of *Salmonella* isolates to antimicrobial agents by serotype.** The frequency of resistance to antimicrobial agents among the serotypes of *Salmonella* is shown in Table 5. Of the 14 *Salmonella* serovars tested against the eight antimicrobial agents, the frequency of resistance was statistically significant ( $P < 0.05$ ) only to kanamycin, AMP, and SXT. Among the predominant serovars recovered from the three countries, *Salmonella Anatum*, *Salmonella group C*, and *Salmonella Kentucky*, the frequency of resistance to antimicrobial agents was statistically significantly different ( $P < 0.05$ ) within each country. In Trinidad and Tobago, of the 11 isolates of *Salmonella Anatum* tested, the range of frequency of resistance was from 9.1% (to SXT) to 100.0% (to E); whereas, in Grenada, the lowest frequency of resistance found amongst 8 isolates of *Salmonella group C* was 0.0% (to SXT) and the highest was 100.0% (to E and S). For St. Lucia, among a total of 27 isolates of *Salmonella Kentucky* tested, the frequency of resistance ranged from 3.7% (to SXT) to 100.0% (to both E and S). The significant effect of the serotypes of *Salmonella* isolates on the frequency of resistance to kanamycin, AMP, and SXT is in agreement with published reports (22, 35).

TABLE 5. Frequency of resistance among *Salmonella* isolates by serovar<sup>a</sup>

Serotype	No. of <i>Salmonella</i> isolates tested	No. (%) of resistant isolates <sup>b</sup>	No. (%) resistant to <sup>c,d</sup> :						
			E	S	CN	K	AMP	TE	SXT
<i>Salmonella</i> group D <sup>e</sup>	1	1 (100.0)	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
<i>Salmonella</i> group C <sup>e</sup>	8	8 (100.0)	8 (100.0)	8 (100.0)	7 (87.5)	7 (87.5)	6 (75.0)	1 (12.5)	0 (0.0)
<i>Salmonella</i> Kentucky (8)20:i:z <sub>6</sub>	27	27 (100.0)	27 (100.0)	27 (100.0)	15 (55.6)	12 (44.4)	4 (14.8)	17 (63.0)	1 (3.7)
<i>Salmonella</i> Uganda 3,10:l,z <sub>13</sub> :1,5	10	10 (100.0)	10 (100.0)	8 (80.0)	7 (70.0)	2 (20.0)	4 (40.0)	1 (10.0)	0 (0.0)
<i>Salmonella</i> Anatum 3,10:e,h:1,6	11	11 (100.0)	11 (100.0)	10 (90.9)	7 (63.6)	4 (36.4)	5 (45.5)	2 (18.2)	1 (9.1)
<i>Salmonella</i> Montevideo 6,7:g,m,s:	2	2 (100.0)	2 (100.0)	0 (0.0)	1 (50.0)	1 (50.0)	0 (0.0)	0 (0.0)	1 (50.0)
<i>Salmonella</i> Mbandaka 6,7:z <sub>10</sub> e,n,Z <sub>15</sub>	4	4 (100.0)	4 (100.0)	4 (100.0)	4 (100.0)	3 (75.0)	3 (75.0)	1 (25.0)	3 (75.0)
<i>Salmonella</i> Caracas 6,14:(25):g,m,s	6	6 (100.0)	6 (100.0)	5 (83.3)	2 (33.3)	5 (83.3)	3 (50.0)	0 (0.0)	0 (0.0)
<i>Salmonella</i> Muenster 3,10:e,h:1,5	1	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
<i>Salmonella</i> Enteritidis 1,9,12:g,m:	1	1 (100.0)	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (100.0)
Polyvalent A-negative	4	4 (100.0)	4 (100.0)	4 (100.0)	3 (75.0)	4 (100.0)	2 (50.0)	1 (25.0)	3 (75.0)
<i>Salmonella</i> groups I-O <sup>e</sup>	2	2 (100.0)	2 (100.0)	2 (100.0)	1 (50.0)	0 (0.0)	1 (50.0)	1 (50.0)	0 (0.0)
<i>Salmonella</i> group E <sup>e</sup>	2	2 (100.0)	2 (100.0)	1 (50.0)	0 (0.0)	0 (0.0)	2 (100.0)	1 (50.0)	0 (0.0)
<i>Salmonella</i> group B <sup>e</sup>	5	5 (100.0)	5 (100.0)	3 (60.0)	2 (40.0)	0 (0.0)	3 (60.0)	0 (0.0)	0 (0.0)
Total	84	84 (100.0)	84 (100.0)	78 (92.9)	50 (59.5)	41 (48.8)	31 (36.9)	28 (33.3)	10 (11.9)

<sup>a</sup> All isolates were sensitive to chloramphenicol (30 µg).<sup>b</sup> Resistant to one or more antimicrobial agents.<sup>c</sup> E, erythromycin (5 µg); S, streptomycin (10 µg); CN, kanamycin (30 µg); K, kanamycin (10 µg); AMP, ampicillin (10 µg); SXT, sulphamethoxazole-trimethoprim (27.75/1.25 µg).<sup>d</sup> P values: S, 0.058; CN, 0.129; K, 0.009; AMP, 0.021; TE, 0.246; SXT, 0.000.<sup>e</sup> Classified at the level of serogroups because the regional typing center did not have the antisera for all the serotypes in the serogroups.

In conclusion, based on the relatively high frequency of resistance to most of the antimicrobial agents tested in the current study, coupled with the equally high occurrence of multidrug resistance among isolates of *Salmonella* recovered from layer birds, farm environments, and feeds, there is a risk of therapeutic failures, which could result in economic losses. There is, therefore, a need for a more judicious use of antimicrobial agents in the poultry industry in the three countries studied.

## ACKNOWLEDGMENTS

The authors are grateful to the Pan American Health Organization (PAHO) (grant 65503 52163) for funding the project. We appreciate the assistance rendered by the technical staff in the offices of the Chief Veterinary Officers in Trinidad and Tobago, Grenada, and St. Lucia and their staff in facilitating sample collection.

## REFERENCES

- Abraham, S., M. D. Groves, D. J. Trott, T. A. Chapman, B. Turner, M. Hornitzky, and D. Jordan. 2014. *Salmonella enterica* isolated from infections in Australian livestock remain susceptible to critical antimicrobials. *Int. J. Antimicrob. Agents* 43:126–130.
- Adesiyun, A., N. Offiah, N. Seepersadsingh, S. Rodrigo, V. Lashley, and L. Musai. 2007. Antimicrobial resistance of *Salmonella* spp. and *Escherichia coli* isolated from table eggs. *Food Control* 18:306–311.
- Adesiyun, A., N. Offiah, N. Seepersadsingh, S. Rodrigo, V. Lashley, L. Musai, and K. Georges. 2005. Microbial health risk posed by table eggs in Trinidad. *Epidemiol. Infect.* 133:1049–1056.
- Adesiyun, A., L. Webb, L. Musai, B. Louison, G. Joseph, A. Stewart-Johnson, S. Samlal, and S. Rodrigo. 2014. Survey of *Salmonella* contamination in chicken layer farms in three Caribbean countries. *J. Food Prot.* 77:1471–1480.
- Akter, M. R., K. A. Choudhury, M. M. Rahman, and M. S. Islam. 2007. Seroprevalence of salmonellosis in layer chickens with isolation, identification and antibiogram study of their causal agents. *Bangladesh J. Vet. Med.* 5:32–42.
- Andrews, W. 1992. *Salmonella*, chap. 4, p. 27–48. In *Manuals of food quality control: 4. FAO food and nutrition paper 14/4, rev. 1*. Food and Agriculture Organization of the United Nations, Rome.
- Bischoff, K. M., T. S. Edrington, T. R. Callaway, K. J. Genovese, and D. J. Nisbet. 2004. Characterization of antimicrobial resistant *Salmonella* Kinshasa from dairy calves in Texas. *Lett. Appl. Microbiol.* 38:140–145.
- Cantas, L., S. Q. Shah, L. M. Cavaco, C. M. Manaia, F. Walsh, M. Popowska, H. Garellick, H. Bürgmann, and H. Sørum. 2013. A brief multi-disciplinary review on antimicrobial resistance in medicine and its linkage to the global environmental microbiota. *Front. Microbiol.* 4:96.
- Dawoud, T. M., P. Herrera, I. Hanning, Y. M. Kwon, and S. C. Ricke. 2011. In vitro invasion of laying hen ovarian follicles by *Salmonella* Enteritidis strains. *Poult. Sci.* 90:1134–1137.
- Dias de Oliveira, S., F. Siqueira-Flores, L. R. Dos Santos, and A. Brandelli. 2005. Antimicrobial resistance in *Salmonella* Enteritidis strains isolated from broiler carcasses, food, human and poultry-related samples. *Int. J. Food Microbiol.* 97:297–305.
- Durand, A. M., M. L. Barnard, M. L. Swanepoeland, and M. M. Engelbrecht. 1987. Resistance to various antibiotics of *Salmonella* and *Escherichia coli* isolated from registrable farm feeds. *Onderstepoort J. Vet. Res.* 54:21–26.
- Elmadiena, M. M., A. A. El Hussein, C. A. Muckle, L. Cole, E. Wilkie, K. Mistry, and A. Perets. 2013. Antimicrobial susceptibility and multi-drug resistance of *Salmonella enterica* subspecies enterica serovars in Sudan. *Trop. Anim. Health Prod.* 45:1113–1118.
- Enabulele, S. A., P. O. Amune, and W. T. Aborisade. 2010. Antibioograms of *Salmonella* isolates from poultry farms in Ovia North East local government area, Edo State, Nigeria. *Agric. Biol. J. N. Am.* 1:1287–1290.
- Fashae, K., F. Ogunsola, F. M. Aarestrup, and R. S. Hendriksen. 2010. Antimicrobial susceptibility and serovars of *Salmonella* from chickens and humans in Ibadan, Nigeria. *J. Infect. Dev. Ctries.* 4:484–494.
- Hamer, D. H., and C. J. Gill. 2002. From the farm to the kitchen table: the negative impact of antimicrobial use in animals on humans. *Nutr. Rev.* 60:261–264.
- Hang'ombe, B. M., R. N. Sharma, E. Skjerve, and L. M. Tuchili. 1999. Occurrence of *Salmonella* Enteritidis in pooled table eggs and market-ready chicken carcasses in Zambia. *Avian Dis.* 43:597–599.
- Hannah, J. F., J. L. Wilson, N. A. Cox, L. J. Richardson, J. A. Cason, D. V. Bourassa, and R. J. Buhr. 2011. Horizontal transmission of *Salmonella* and *Campylobacter* among caged and cage-free laying hens. *Avian Dis.* 55:580–587.
- Hassan, M. M., K. B. Amin, M. Ahaduzzaman, M. Alam, M. S. Al Faruk, and I. Uddin. 2014. Antimicrobial resistance pattern against *E. coli* and *Salmonella* in layer poultry. *Res. J. Vet. Pract.* 2:30–35.
- Iwabuchi, E., N. Maruyama, A. Hara, M. Nishimura, M. Muramatsu, T. Ochiai, and K. Hirai. 2010. Nationwide survey of *Salmonella* prevalence in environmental dust from layer farms in Japan. *J. Food Prot.* 73:1993–2000.
- Khachatourians, G. G. 1998. Agricultural use of antibiotics and the evolution and transfer of antibiotic-resistant bacteria. *Can. Med. Assoc. J.* 159:1129–1136.
- Kwon, Y. K., A. Kim, M. S. Kang, M. Her, B. Y. Jung, K. M. Lee, B. K. An, and J. H. Kwon. 2010. Prevalence and characterization of *Salmonella* Gallinarum in the chicken in Korea during 2000 to 2008. *Poult. Sci.* 89:236–242.
- Lu, Y., C. M. Wu, G. J. Wu, H. Y. Zhao, T. He, X. Y. Cao, L. Dai, L. N. Xia, S. S. Qin, and J. Z. Shen. 2011. Prevalence of antimicrobial resistance among *Salmonella* isolates from chicken in China. *Foodborne Pathog. Dis.* 8:45–53.
- MacFaddin, J. F. 2002. *Biochemical tests for identification of medical bacteria*. Williams and Wilkins, New York.
- Makaya, P. V., G. Matope, and D. M. Pfukenyi. 2012. Distribution of *Salmonella* serovars and antimicrobial susceptibility of *Salmonella* Enteritidis from poultry in Zimbabwe. *Avian Pathol.* 41:221–226.
- Mir, I. A., S. A. Wani, I. Hussain, S. D. Qureshi, M. A. Bhat, and Y. Nishikawa. 2010. Molecular epidemiology and in vitro antimicrobial susceptibility of *Salmonella* isolated from poultry in Kashmir. *Rev. Sci. Tech.* 29:677–686.
- Mughini-Gras, L., R. Enserink, I. Friesema, M. Heck, Y. van Duynhoven, and W. van Pelt. 2014. Risk factors for human salmonellosis originating from pigs, cattle, broiler chickens and egg laying hens: a combined case-control and source attribution analysis. *PLoS One* 9:e87933.
- National Committee for Clinical Laboratory Standards (NCCLS). 2002. Performance standards for antimicrobial disc and dilution susceptibility for bacteria isolated from animals; approved standards, 2nd ed., vol. 22. NCCLS, Wayne, PA.
- Poppe, C., K. A. McFadden, and W. H. Demczuk. 1996. Drug resistance, plasmids, biotypes and susceptibility to bacteriophages of *Salmonella* isolated from poultry in Canada. *Int. J. Food Microbiol.* 30:325–344.
- Sasanya, J. J., J. W. Okeng, F. Ejobi, and M. Muganwa. 2005. Use of sulfonamides in layers in Kampala district, Uganda and sulfonamide residues in commercial eggs. *Afr. Health Sci.* 5:33–39.
- Schwaiger, K., E. M. Schmied, and J. Bauer. 2008. Comparative analysis of antibiotic resistance characteristics of gram-negative bacteria isolated from laying hens and eggs in conventional and organic keeping systems in Bavaria, Germany. *Zoonoses Public Health* 55:331–341.
- Singh, R., A. S. Yadav, V. Tripathi, and R. P. Singh. 2013. Antimicrobial resistance profile of *Salmonella* present in poultry and poultry environment in north India. *Food Control* 33:545–548.
- Snow, L. C., R. H. Davies, K. H. Christiansen, J. J. Carrique-Mas, A. J. C. Cook, and S. J. Evans. 2012. Investigation of risk factors of *Salmonella* on commercial egg-laying farms in Great Britain, 2004–2006. *Vet. Rec.* 166:579–586.

33. Snow, L. C., R. H. Davies, K. H. Christiansen, J. J. Carrique-Mas, A. D. Wales, J. L. O'Connor, A. J. Cook, and S. J. Evans. 2007. Survey of the prevalence of *Salmonella* species on commercial laying farms in the United Kingdom. *Vet. Rec.* 161:471–476.
34. Someya, A., K. Otsuki, and T. Murase. 2005. Antimicrobial susceptibilities of *Salmonella* isolates obtained from layer chicken houses on a commercial egg-producing farm in Japan, 1997 to 2002. *J. Food Prot.* 68:2030–2034.
35. Tabo, D., C. D. Diguimbaye, S. A. Granier, F. Moury, A. Brisabois, R. Elgroud, and Y. Millemann. 2013. Prevalence and antimicrobial resistance of non-typhoidal *Salmonella* serotypes isolated from laying hens and broiler chicken farms in N'Djamena, Chad. *Vet. Microbiol.* 166:293–298.
36. Wierup, M., B. Engström, A. Engvall, and H. Wahlström. 1995. Control of *Salmonella* Enteritidis in Sweden. *Int. J. Food Microbiol.* 25:219–226.

Copyright of Journal of Food Protection is the property of Allen Press Publishing Services Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.