



Prevalence and characterization of *Salmonella enterica* serovar in retail meats in market place in Uighur, Xinjiang, China



Mingyuan Yin ^{a, 2}, Baowei Yang ^{b, 1, 2}, Yun Wu ^{a, c, *}, Lu Wang ^c, Haotian Wu ^a, Tao Zhang ^c, Gulnazi Tuohetaribayi ^a

^a College of Food Science and Pharmaceutical Science, Xinjiang Agricultural University, Urumqi, Xinjiang, 830052, China

^b College of Food Science and Engineering, Northwest A&F University, Yangling, 712100, China

^c Xinjiang Agricultural University Institute of Science and Technology, Urumqi, Xinjiang, 830052, China

ARTICLE INFO

Article history:

Received 10 July 2015

Received in revised form

21 December 2015

Accepted 21 December 2015

Available online 24 December 2015

Keywords:

Serotyping

Antimicrobial susceptibility

Salmonella

PFGE

Substitution

ABSTRACT

Ninety nine *Salmonella* isolates recovered from 1414 retail meats were characterized by serotyping, antimicrobial susceptibility, pulsed-field gel electrophoresis (PFGE), presence of amino acid mutation in quinolone resistance determination region (QRDR) of DNA gyrase subunit A (GyrA) and topoisomerase IV subunit C (ParC), and of *qnrA*, *qnrB*, *qnrS*, *aac(6')*-Ib, *qepA*, and *oqxAB*. Among 1414 retail meat samples, 96 (6.8%) including 49 (9.0%) chickens, 22 (6.8%) lambs, 10 (4.8%) beefs, 13 (6.8%) porks and 2 (1.4%) horse meats were positive to *Salmonella*. The commonly detected *Salmonella* serotypes were *S. Hadar* (n = 21, 21.2%), *S. Enteritidis* (n = 17, 17.2%), *S. London* (n = 17, 17.2%), and *S. Havana* (n = 11, 11.1%). Eighty four (84.8%) isolates were simultaneously resistant to more than three antimicrobial agents. Antibiotic resistance was most commonly found to trimethoprim (100%), and a less extent to chloramphenicol (88.9%), tetracycline (63.6%), nalidixic acid (58.6%), sulfisoxazole (57.6%), streptomycin (43.4%), trimethoprim/tulfisoxazole (41.4%), ampicillin (25.6%), amoxicillin/clavulanate (25.6%), kanamycin (6.1%), ceftriaxone (5.1%), gentamicin (3.0%), cefoxitin (2.0%), and amikacin (1.0%). *qnrA* (11.1%), *qnrB* (34.3%), *qnrS* (8.1%), *aac(6')*-Ib (7.1%), *qepA* (7.1%), *oqxA* (10.1%) and *oqxB* (9.1%) were detected from the 99 isolates. Amino acid substitutions of Asp87Asn (4.8%), Asp87Tyr (28.6%), Asp87Val (4.8%), Ser83Phe (52.4%), Ser83Tyr (7.1%) and Gly75Phe (2.4%) in GyrA were detected, as well as Thr57Ser (98.6%) and Gly53Val (1.4%) in ParC. Mutations of Ser83Phe (GyrA)/Thr57Ser (ParC) that simultaneously detected in GyrA and ParC were found in 22 isolates. Totally 82 different DNA patterns generated after the 99 isolates were subtyped using PFGE. The results demonstrated that the prevalence of *Salmonella* in retail meats in Uighur of Xinjiang province were not common, however, the isolates exhibited multidrug resistance, phenotypical and genotypical diverse.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Foodborne *Salmonella* has been recognized as significant pathogen associated with public health worldwide. In the United States, approximately 1 million cases, 19,336 hospitalizations, 378 deaths, and \$600 million to \$3.5 billion medical expenditures were caused by non-typhoidal *Salmonella* annually (Scallan et al., 2011; Yang et al., 2010). In European, in 2009 and 2010, 108,614 and 99,020

salmonellosis were confirmed, respectively (Bonardi et al., 2013; De, Bravo, & Medina, 2012). Meanwhile, in Southeast Asia, an unofficial *Salmonella* surveillance indicated that approximate 22.8 million cases were occurred yearly with 37,600 deaths (Van, Nguyen, Smooker, & Coloe, 2012). As those uncovered, *Salmonella* infections commonly associated with consumption of contaminated foods that were mainly processed from food animals including poultry, pig, beef and lamb, which all are original sources to *Salmonella* (Aslam et al., 2012; Boonmar et al., 2013; Liu, Chen, Huang, Liu, & Shi, 2010; Mąka, Maćkiw, Ścieżyńska, Pawłowska, & Popowska, 2014; Wouafo et al., 2010).

Up to present, more than 2500 serovars have been identified among *Salmonella* (Son et al., 2013), while majorities of human infections were caused by limited number of serovars, among

* Corresponding author. College of Food Science and Pharmaceutical Science, Xinjiang Agricultural University, 311# Road, Urumqi, Xinjiang, 830052, China.

E-mail address: wuyunster@sina.com (Y. Wu).

¹ Co-first author.

² The first two authors contributed equally to this paper.

which, *S. Enteritidis* has been recognized as the most common serovar to human salmonellosis, other serovars including *S. London*, *S. Hadar* and *S. Derby* that involved in salmonellosis were reported to be prevalent in certain foods as well (Dogru, Ayaz, & Gencay, 2010; Hendriksen et al., 2011; Thong & Modarressi, 2011). With the antibiotics were widely used in food animal production, more and more *Salmonella* strains were identified as multiple resistant (MDR) ones, and the increasing MDR *Salmonella* were of great concern for food safety and public health worldwide (Boonmar et al., 2012; Kim, Park, Kwak, & Woo, 2011). Presence and dissemination of plasmid-mediated quinolone resistance (PMQR) genes (*qnrA*, *qnrB*, *qnrS*, *aac(6′)-Ib*, *qepA*, *oqxAB*) were regarded as one of the important mechanisms that contributed to quinolone resistance. Meanwhile, Amino acid alterations in GyrA and ParC of the quinolone resistance determining region (QRDR) have been demonstrated to result in decreased susceptibility to fluoroquinolones (Shariat et al., 2013; Zhang et al., 2014).

In this study, 99 *Salmonella* isolates recovered from retail meats in 2013 and 2014 in Xinjiang Uighur were characterized for better understand food safety situation in China to ensure public health.

2. Materials and methods

2.1. Isolates

A total of 1414 retail raw meat samples including 542 chickens, 325 lambs, 210 beefs, 192 porks and 145 horse meats were collected from 23 wet markets in seven districts of Uighur city of Xinjiang province, China, during 2013–2014. Detailed information for sample collection and *Salmonella* isolation were as previously described (Yin et al., 2014). Isolates with typical *Salmonella* phenotypes on XLT4 (Beijing Land Bridge Technology Co Ltd., Beijing, China) and XLD (Beijing Land Bridge Technology Co Ltd.) plates were finally identified and confirmed by O hypersera A–F (S&A Reagent Lab, Bangkok, Thailand). All isolates were stored at -80°C in Luria–Bertani broth (LB; Difco, Maryland, USA)/glycerol (50%/50%, V/V) until use.

Salmonella isolates were serotyped by slide agglutination method (GB 4789.4–2010; Yang et al. 2013) using specific O and H antisera (S&A Reagent Lab, Bangkok, Thailand) in Henan Center for Disease Control and Prevention, Zhengzhou, Henan, China.

2.2. Antimicrobial susceptibility test

All isolates were examined for their susceptibility to 15 antibiotics including ampicillin (AMP), amoxicillin/clavulanic (AMC), amikacin (AMK), cefoxitin (CFX), ceftriaxone (CRO), chloramphenicol (CHL), nalidixic acid (NAL), ciprofloxacin (CIP), gentamicin (GEN), kanamycin (KAN), streptomycin (STR), tetracycline (TET), trimethoprim (TIO), sulfisoxazole (FIS) and trimethoprim/sulfamethoxazole (SXT). The minimum inhibitory concentrations (MICs) of the antibiotics were determined by agar dilution method that described by the Clinical and Laboratory Standards Institute (CLSI) (Clinical and Laboratory Standards Institute, 2013). *Escherichia coli* ATCC25922 and *Enterococcus faecalis* ATCC29212 were used as quality control organisms in MICs determinations. The breakpoints for antimicrobial susceptible and/or resistant were interpreted and determined by CLSI guidelines except streptomycin, the breakpoint of which was interpreted according to that of the National Antimicrobial Resistance Monitoring System (NARMS) managed by the Food and Drug Administration (FDA), the U.S. Centers for Disease Control and Prevention (CDC) and the United States Department of Agriculture (USDA) (U.S. Food and Drug Administration, 2013).

2.3. Pulse field gel electrophoresis (PFGE)

PFGE was carried out for *Salmonella* genetic subtyping according to the protocol developed by the CDC (Ribot et al., 2006). Briefly, *Salmonella* isolates were cultured on Luria–Bertani agar (Beijing Land Bridge Technology Co Ltd.) at 37°C overnight, the genomic DNA was prepared by embedding the isolates in agarose plugs, after cells were lysed, the embedded DNA was digested with 50 U of XbaI (TaKaRa, Dalian, China) for 1.5–2 h in a water bath at 37°C . The DNA fragments were subsequently separated by electrophoresis in $0.5 \times \text{TBE}$ buffer at 14°C for 18 h using a Chef Mapper electrophoresis system (Bio-Rad, Hercules, CA, USA) with pulse times of 2.16–63.8 s. *Salmonella* Braenderup H9812 was used as the standard control strain. The gels were stained with ethidium bromide and the DNA bands were visualized using UV transillumination (Bio-Rad). Fingerprinting profiles were analyzed using the BioNumerics software (Version 3.0; Applied-Maths, Kortrijk, Belgium) manually, the genotype was determined by a cutoff value of 90% similarity based on the unweighted pair group method with arithmetic mean (UPGMA).

2.4. Detection of PMQR genes (*qnrA*, *qnrB*, *qnrS*, *aac(6′)-Ib-cr*, *qepA* and *oqxAB*) and QRDR (*GyrA* and *ParC*) mutations

All nalidixic acid resistant isolates were screened for presence of PMQR genes and amino acid substitutions in GyrA and ParC by polymerase chain reaction (PCR) using the primers and annealing temperatures listed in Table 1. PCRs were carried out in a 25 μL PCR mixture that contained 0.5 μM of each primer, 250 μM of each dNTP, 2.5 μL of $10 \times \text{PCR}$ buffer, 0.5 U of Taq DNA polymerase (TaKaRa), 1.5 mM MgCl_2 and 5 μL of sample template DNA, with predenaturation at 94°C for 10 min; 35 cycles of denaturation at 94°C for 30 s, at annealing temperatures for 30 s and a final extension at 72°C for 7 min. Primers were synthesized by TaKaRa Biotechnology Co., Ltd. PCR products were stained with ethidium bromide and visualized under UV light after gel electrophoresis in 1% agarose. For *gyrA* and *parC* analysis, PCR products were sequenced in Shanghai Sunny Biotechnology Co, Ltd (Shanghai, China) and the sequence were aligned using BLAST program (<http://www.ncbi.nlm.nih.gov/BLAST/>).

3. Results

3.1. Isolation and identification of *Salmonella*

Among 1414 retail meat samples, 96 (6.8%) including 49 (9.0%) of 542 chickens, 22 (6.8%) of 325 lambs, 10 (4.8%) of 210 beefs, 13 (6.8%) of 192 porks and 2 (1.4%) of 145 horse meats were detected *Salmonella* positive (Table 2). Totally 99 *Salmonella* isolates (1–2 isolates per sample) including 52 from chicken, 22 from lamb, 13 from pork, 10 from beef and 2 from horsemeat were recovered (see Fig. 1).

3.2. Serotype distribution

A total of 18 different serotypes were identified among 99 *Salmonella* isolates. Nine, 10, 8, 8, and 1 serotypes were identified among isolates recovered from retail chicken, lamb, beef, pork and horsemeat, respectively. The commonly prevalent serotypes were *Salmonella* Hadar ($n = 21$, 21.2%), *S. Enteritidis* ($n = 17$, 17.2%), *S. London* ($n = 17$, 17.2%) and *S. Havana* ($n = 11$, 11.1%). *Salmonella* Enteritidis, *S. London*, and *S. Havana* were simultaneously identified from retail chickens, lambs, beefs and porks. *Salmonella* Enteritidis, *S. Hadar*, *S. London*, *S. Paratyphi B* and *S. Havana* were commonly identified among the isolates recovered from retail chickens.

Table 1
Primers used for target genes detection and sequencing.

Target gene	Primer	Sequence (5'-3')	Annealing temperature (°C)	Product size (bp)	Reference
Amplification and sequencing primers					
<i>gyrA</i>	<i>gyrA</i> -F	ACGTACTAGGCAATGACTGG	56	190	Eaves, Liebana, Woodward, & Piddock, 2002
	<i>gyrA</i> -R	AGAAGTCGCGCTCGATAGAA			
<i>parC</i>	<i>parC</i> -F	CTATGCCATGTCAGAGCTGG	54	270	Eaves et al., 2004
	<i>parC</i> -R	TAACAGCAGCTCGGCGTATT			
Detection primers					
<i>qnrA</i>	<i>qnrA</i> -F	AGAGGATTTCTCACGCCAGG	60	580	Cattoir, Poirel, Rotimi, Soussy, & Nordmann, 2007
	<i>qnrA</i> -R	TGCCAGGCACAGATCTTGAC			
<i>qnrB</i>	<i>qnrB</i> -F	GGMATHGAAATTCGCCACTG	56	264	Cattoir et al., 2007
	<i>qnrB</i> -R	TTTGCYGYCCGCGAGTCGAA			
<i>qnrS</i>	<i>qnrS</i> -F	GCAAGTTCATTGAACAGGGT	57	428	Cattoir et al., 2007
	<i>qnrS</i> -R	TCTAAACCGTCGAGTTCGGCG			
<i>aac(6')-Ib</i>	<i>aac(6')-Ib</i> -F	TTGCGATGCTCTATGAGTGGCTA	55	482	Park, Robicsek, Jacoby, Sahm, & Hooper, 2006
	<i>aac(6')-Ib</i> -R	CTCGAATGCCTGGCGTGTTC			
<i>qepA</i>	<i>qepA</i> -F	CTGCAGGTACTGCGTCATG	60	403	Chen, Zhang, et al., 2012; Chen, Shao, et al., 2012
	<i>qepA</i> -R	CGTGTGTGTCGAGTTCTTC			
<i>oqxA</i>	<i>oqxA</i> -F	GACAGCGTCGCACAGAATG	62	339	Chen, Zhang, et al., 2012; Chen, Shao, et al., 2012
	<i>oqxA</i> -R	GGAGACGAGGTTGGTATGGA			
<i>oqxB</i>	<i>oqxB</i> -F	CGAAGAAAGACCTCCCTACCC	62	240	Chen, Zhang, et al., 2012; Chen, Shao, et al., 2012
	<i>oqxB</i> -R	CGCCGCAATGAGATACA			

Table 2
Prevalence of *Salmonella* in different source of retail meats during 2013–2014.

Retail meat	No. sample	No.(%) samples positive to <i>Salmonella</i>	No. <i>Salmonella</i> isolates recovered ^a
Chicken	542	49 (9.0)	52
Lamb	325	22 (6.8)	22
Beef	210	10 (4.8)	10
Pork	192	13 (6.8)	13
Horsemeat	145	2 (1.4)	2
Total	1414	96 (6.8)	99

^a 1–2 *Salmonella* isolates were recovered from each positive sample.

Salmonella Schwarzengrund II, *S. Agona*, *S. Assinie*, *S. Braenderup*, *S. Goldcoast*, *S. Infantis*, *S. Thompson*, *S. Typhimurium*, *S. Tennessee*, and *S. Bloomsbury* were detected from retail lambs, beefs, and porks. Two *Salmonella* isolates recovered from horsemeat were all *S. London* (Table 3).

3.3. Antimicrobial susceptibility testing

Ninety nine *Salmonella* isolates were all susceptible to ciprofloxacin (see Table 4). Antibiotic resistance was commonly observed to trimethoprim (100%), then to chloramphenicol (88.9%), tetracycline (63.6%), nalidixic acid (58.6%), sulfisoxazole (57.6%), streptomycin (43.4%) and trimethoprim/sulfisoxazole (41.4%), and a less extent to ampicillin (25.6%), amoxicillin/clavulanate (25.6%), kanamycin (6.1%), ceftriaxone (5.1%), gentamicin (3.0%), cefoxitin (2.0%) and amikacin (1.0%).

All isolates were resistant to at least one antibiotic, no *Salmonella* can resist to more than 10 antimicrobial agents (Table 5). Thirty (30.3%) *Salmonella* isolates were resistant to 1–3 antibiotics, 34 (34.3%) to 4–6 antibiotics, 31 (31.3%) to 7–9 antibiotics and 4 (4.0%) to 10 antibiotics. When analyzed by retail meat types, resistance was most frequently detected in chicken-derived *Salmonella* isolates, 23 (44.3%) of which were resistant to at least 7 antimicrobials, and 3 (5.8%) to 10 antimicrobials. *Salmonella* isolates derived from beefs ($n = 7$, 70.0%) and lambs ($n = 10$, 45.5%) were most commonly resistant to 4–6 antimicrobials, those from porks ($n = 10$, 7.7%) were to 10 antimicrobials. *Salmonella* isolates ($n = 2$) derived from horsemeats was relatively susceptible.

When analyzed by serovar, 33.3% of *Salmonella* Havana and 29.6% of *S. London* isolates frequently exhibited resistance to 1–3

antibiotics. 50.0% of *S. Hadar* and 25.0% of *S. London* isolates were simultaneously resistant to 4–6 antibiotics. Some of *S. Enteritidis* ($n = 12$, 46.1%), *S. Hadar* ($n = 7$, 26.9%) and *S. Derby* ($n = 4$, 15.4%) isolates could resist more than 7 antibiotics. Two *S. Enteritidis* and two *S. London* isolates could resist 10 antibiotics (Table 6). Generally, antibiotic resistance level of *Salmonella* Enteritidis, *S. Hadar* and *S. Derby* isolates was much higher than that of isolates of other serotypes.

3.4. PFGE

Genetic relatedness of 99 *Salmonella* isolates was analyzed using PFGE with XbaI Fig. 1. A total of 82 different PFGE patterns were observed. Using a cutoff value of 90% similarity, five typical clusters (named as A, B, C, D, and E, respectively) and several individual profiles were observed. Among each cluster, DNA profiles of the isolates exhibited high homology, although difference still could be found. Eleven isolates were grouped in cluster A, and 10 of them were *S. Havana*. Seventeen *S. Enteritidis* isolates were grouped in cluster B. Eight isolates were grouped in cluster C, and six of them were *S. Paratyphi B*. Twenty one isolates were grouped in cluster D, 20 of them were *S. Hadar*. Seventeen isolates were grouped in cluster E, and isolates in this cluster were all *S. London*. Three pairs of *S. Havana* isolates in cluster A, four *S. Enteritidis* isolates in Cluster B, four *S. Paratyphi B* isolates in Cluster C, two *S. Hadar* isolates in Cluster D and three *S. London* isolates in Cluster E, were detected 100% similarity. DNA profiles of some isolates with same serotype in same PFGE cluster were the same or similar, although these isolates were recovered from different times, different sampling places and different retail meats.




Table 3
Distribution of *Salmonella* serovars in retail meats during 2013–2014.

Serotype	No. (%) isolates (n = 99)	No. (%) type of meat				
		Chicken (n = 52)	Lamb (n = 22)	Beef (n = 10)	Pork (n = 13)	Horsemeat (n = 2)
Hadar	21 (21.2)	10 (19.2)	9 (40.9)	2 (15.4)		
Enteritidis	17 (17.2)	12 (23.1)	2 (9.1)	2 (15.4)	1 (10.0)	
London	17 (17.2)	8 (15.4)	3 (13.6)	3 (23.1)	1 (10.0)	2 (100)
Havana	11 (11.1)	7 (13.5)	1 (4.5)	1 (9.1)	2 (20.0)	
Paratyphi B	9 (9.1)	8 (15.4)	1 (4.5)			
Derby	5 (5.1)	3 (5.8)		2 (15.4)		
Mbandaka	5 (5.1)	2 (3.8)	2 (9.1)		1 (10.0)	
Oran II	3 (3.0)	1 (1.9)	1 (4.5)		1 (10.0)	
Schwarzengrund II	2 (2.0)				2 (20.0)	
Agona	1 (1.0)	1 (1.9)				
Assinie	1 (1.0)				1 (10.0)	
Braenderup	1 (1.0)		1 (4.5)			
Goldcoast	1 (1.0)			1 (9.1)		
Infantis	1 (1.0)			1 (9.1)		
Thompson	1 (1.0)		1 (4.5)			
Typhimurium	1 (1.0)				1 (10.0)	
Tennessee	1 (1.0)		1 (4.5)			
Bloomsbury	1 (1.0)			1 (9.1)		
Total	99 (100)	52 (52.5)	22 (22.2)	13 (13.1)	10 (10.1)	2 (2.0)

Table 4
Antimicrobial resistance of *Salmonella* isolated from retail meats during 2013–2014.

Antimicrobial agent	No. of isolates (%)						
		Resistant breakpoint ^a (μg/ml)	Chicken (n = 52)	Lamb (n = 22)	Pork (n = 13)	Beef (n = 10)	Horsemeat (n = 2)
Trimethoprim	R ≥ 16		52 (100)	22 (100)	13 (100)	10 (100)	2 (100)
Chloramphenicol	R ≥ 32		47 (90.4)	18 (81.8)	11 (84.6)	10 (100)	2 (100)
Tetracycline	R ≥ 16		32 (61.5)	15 (68.2)	9 (69.2)	5 (50.0)	2 (100)
Nalidixic acid	R ≥ 32		31 (59.6)	16 (72.7)	5 (38.5)	6 (60.0)	
Sulfisoxazole	R ≥ 512		33 (63.5)	10 (45.5)	8 (61.5)	6 (60.0)	
Streptomycin ^b	R ≥ 64		26 (50.0)	10 (45.5)	7 (53.8)		
Trimethoprim/Sulfisoxazole	R ≥ 4/76		22 (42.3)	7 (31.8)	8 (61.5)	4 (40.0)	
Ampicillin	R ≥ 32		19 (36.5)	1 (4.5)	4 (30.8)	1 (10.0)	
Amoxycillin/Clavulanate	R ≥ 32/16		19 (36.5)	1 (4.5)	4 (30.8)	1 (10.0)	
Kanamycin	R ≥ 64		2 (3.8)	2 (9.1)	1 (7.7)	1 (10.0)	
Ceftriaxone	R ≥ 64		4 (7.7)		1 (7.7)		
Gentamicin	R ≥ 16		1 (1.9)		2 (15.4)		
Cefoxitin	R ≥ 32			2 (9.1)			
Amikacin	R ≥ 64				1 (7.7)		
Ciprofloxacin	R ≥ 8						

^a MICs (μg/ml) were determined via agar dilution in accordance with CLSI.

^b The breakpoint was interpreted according to that of the NARMS.

3.5. Detection of *GyrA* and *ParC* mutation and *qnrA*, *qnrB*, *qnrS*, *aac(6')-Ib-cr*, *qepA*, *oqxAB*

Mutations of *GyrA* and *ParC* were detected in 89 of the 99 *Salmonella* isolates, and a total of 113 amino acid substitutions including 42 in *GyrA* and 71 in *ParC* were detected. The frequently detected mutations in *GyrA* were Ser83Phe (n = 22, 52.4%), Asp87Tyr (n = 12, 28.6%), Ser83Tyr (n = 3, 7.1%), Asp87Val (n = 2, 4.8%) and Gly75Phe (n = 1, 2.4%). Thr57Ser (n = 69, 97.2%) was most commonly detected in *ParC* in addition to Gly53Val/Thr57Ser (n = 1, 1.4%). Simultaneously mutations of Thr57Ser(*ParC*)-Ser83Phe(*GyrA*) (n = 22, 95.7%) and Thr57Ser(*ParC*)-Ser83Tyr(*GyrA*) (n = 1, 4.3%) detected in *GyrA* and *ParC* in 23 isolates. Sixty one isolates were detected harboring PMQR genes, the

most prevalent one of which was *qnrB* (34.3%), followed ones were *oqxAB* (19.2%), *qnrA* (11.1%), *qnrS* (8.1%), *aac(6')-Ib* (7.1%) and *qepA* (7.1%). Forty three (43.4%), 12 (12.1%), 3 (3.0%) and 2 (2.0%) isolates were detected simultaneously carrying 1, 2, 3 and 4 of the seven PMQR genes, respectively.

4. Discussion

In present study, retail meats collected from seven districts of Uighur, Xinjiang in China were identified for the presence of *Salmonella*, and *Salmonella* isolates were further characterized via serotyping, antimicrobial susceptibility, and PFGE.

Our results indicated that *Salmonella* was prevalent in retail meats in Xinjiang, and was more prevalent in retail chickens (9.0%)

Fig. 1. Dendrogram of PFGE patterns of 99 *Salmonella* isolates from retail meats in Xinjiang Province, China. Antimicrobial agents including Trimethoprim(TIO), Chloramphenicol(CHL), Tetracycline(TET), Nalidixic acid(NAL), Sulfisoxazole(FIS), Streptomycin(STR), Trimethoprim/Sulfisoxazole(SXT), Ampicillin(AMP), Amoxycillin/Clavulanate(AMC), Kanamycin(KAN), Ceftriaxone(CRO), Gentamicin(GEN), Cefoxitin(FOX), Amikacin(AMK), Ciprofloxacin(CIP). Districts including Shayibake(SY), Shuimogou(SM), Xinshiqu(XS), Tountunhe(TT), Changji(CJ), Midong (MD), Tianshan(TS).

Table 5Multidrug resistance (MDR) among *Salmonella* isolates recovered from retail meats during 2013–2014.

Retail meats (no. isolates)	No. (%) isolates resisted to indicated number of antimicrobials				Total resistance (≥ 1)
	1–3	4–6	7–9	10	
Chicken (n = 52)	16 (30.8)	13 (25.0)	20 (38.5)	3 (5.8)	52 (100)
Lamb (n = 22)	6 (27.3)	10 (45.5)	6 (27.3)		22 (100)
Pork (n = 13)	4 (30.8)	4 (30.8)	4 (30.8)	1 (7.7)	13 (100)
Beef (n = 10)	2 (20.0)	7 (70.0)	1 (10.0)		10 (100)
Horsemeat (n = 2)	2 (100)				2 (100)
Total (n = 99)	30 (30.3)	34 (34.3)	31 (31.3)	4 (4.0)	99 (100)

than in lambs (6.8%), porks (6.8%), beefs (4.8%) and horsemeats (1.4%). Previous studies indicated that *Salmonella* in retail meats in Shaanxi (9.15%), Shandong (5.63%), and Xinjiang (9.14%) was similar to the results acquired in this study as well as a survey carried by the Chinese National Centers for Disease Control and Prevention (9.35%) (Chen, Shao, Guan, Hu, & Dong, 2010; Wang et al., 2004; Yin et al., 2014; Zhang, Zhu, & Li, 2008). Among various retail meats, chicken was regarded as one of the dominant *Salmonella* vectors all over the world. Compared to previous studies that overall 52.2% of retail chickens were observed to be positive to *Salmonella* in six provinces and two national cities of China including Guangxi (65.3%), Guangdong (64.6%), Beijing (63.9%), Shaanxi (50.7%), Henan (47.9%), Shanghai (44.4%), Fujian (42.4%), and Sichuan Province (38.9%) in 2010 (Yang et al., 2011), and those were 31% in pork, 17% in beef, and 20% in lamb samples in supermarkets and free markets in Shaanxi Province during 2007–2008 (Yang et al., 2010), while 28.3% in retail chicken and 10% in pork meat in Sichuan during 2010–2011 (Li et al., 2013), we thought the prevalence of *Salmonella* in retail meat in Xinjiang province was much lower, and the reason to which may associate with its special geographical position.

In this study, 18 serotypes were identified among 99 *Salmonella* isolates. *Salmonella* Hadar was commonly found in the previous investigations (Maka et al., 2014; Thai and Yamaguchi, 2012; Thong & Modarressi, 2011; Wouafo, et al, 2014) and was detected to be the predominant serovar, particularly in lamb meats, the results of our study were agreement with those of the previous ones. Other than *Salmonella* Hadar, *S. Enteritidis* was verified as the most common serotype that identified in epidemiological investigations associated with consumption of contaminated livestock and poultry products (Laconcha, Baggesen, Rementeria, & Garaizar, 2000; Wang et al., 2015), while it was only identified to be the predominant serovar in chicken meats, and was the second serovar together with *Salmonella* London in this study. Meanwhile, other serovars recovered in the present study included *S. Thompson*, *S. Agona*, *S. Infantis* and *S. Derby* were often detected in other surveys (Bonardi et al., 2013; Maka et al., 2014; Wouafo et al., 2010). The prevalence of meat-derived *Salmonella* may vary from the differences in

country, types of meat sample, sampling seasons and isolation methods.

Antimicrobial resistance in foodborne pathogens has been a global problem. Surveillance data demonstrated an obvious increase in overall antimicrobial resistance among foodborne *Salmonella* isolates from 20% to 30% in the 1990s to 70% at the beginning of the century (Su, Chiu, Chu, & Jonathan, 2004). Our results indicated that none of the isolates were susceptible to the antimicrobial agents tested in addition to ciprofloxacin, however, all the isolates were resistant to at least one antimicrobial agent, 35.3% of the *Salmonella* isolates were resistant to more than 7 antibiotics, but none was resistant to more than 10 antibiotics, and only 4.0% of the isolates were resistant to 10 antibiotics, which indicated that the antibiotic resistance situation in retail meats in Xinjiang province was better than that in other districts (Shao, 2011; Wang et al., 2007; Yang et al., 2010).

Antimicrobial resistant *Salmonella* isolates were also detected in previous investigations and were found commonly resistant to the traditional antibiotics that early applied in disease prevention and health treatment, such as amoxicillin, ampicillin, streptomycin, penicillin, tetracycline, chloramphenicol (Shao, 2011; Wang et al., 2007; Yang et al., 2010). Our study was consistent with previous ones that *Salmonella* isolates could resist the most commonly used antibiotics. The reasons may be due to the selection pressure caused by the usage of traditional antimicrobials in food animal production for therapy, prophylaxis and growth promotion for a long time.

Cephalosporins and fluoroquinolones were effectively used to treat invasive infections, particularly in *Salmonella* infections (White et al., 2001). Our study indicated that just a few *Salmonella* isolates were resistant to cefoxitin (3.0%) and ceftriaxone (5.1%), and no isolate was resistant to ciprofloxacin. A previous study on meat-derived *Salmonella* in 10 provinces of China during 1999–2000 indicated that resistance to cephalosporin was similar to that in other provinces (Chen et al., 2004; Yang, 2010). Nalidixic acid was the first generation of quinolones applied in the treatment of *Salmonella* disease after chloramphenicol. In present study, resistance of *Salmonella* to nalidixic acid (58.6%) was very common, particularly of isolates recovered from lamb (72.7%), beef (60.0%) and chicken (59.6%), which indicated an increased nalidixic acid-resistant *Salmonella* and this result was consistent with other reports from China (Chen et al., 2010; Pan et al., 2009; Zhang et al., 2009; Zhu et al., 2014). Ciprofloxacin-resistant *Salmonella* usually associated with infections and was resistant to multiple drugs (Cui et al., 2008), fortunately, all isolates recovered from retail meats in Xinjiang province were susceptible to ciprofloxacin.

Resistance to fluoroquinolones was partly due to amino acid substitutions of DNA gyrase and topoisomerase IV. In this study, majorities (89 of 99) of isolates were detected harbored amino acid substitutions in their QRDR (GyrA and/or ParC), and 53.9% of these isolates were resistant to nalidixic acid, which was in agreement with the findings that point mutations in QRDR of GyrA and/or ParC

Table 6Multidrug resistance (MDR) observed among commonly detected *Salmonella* serovars during 2013–2014.

Serotype (no. isolates)	No. isolates (%) resisted to indicated number of antimicrobials			
	1–3	4–6	7–9	10
Hadar (n = 21)		14 (50.0)	7 (26.9)	
Enteritidis (n = 17)	2 (7.4)	1 (3.6)	12 (46.1)	2 (50.0)
London (n = 17)	8 (29.6)	7 (25.0)		2 (50.0)
Havana (n = 11)	9 (33.3)	1 (3.6)	1 (3.8)	
Paratyphi B (n = 9)	5 (18.5)	2 (7.1)	2 (7.7)	
Derby (n = 5)		1 (3.6)	4 (15.4)	
Mbandaka (n = 5)	3 (11.1)	2 (7.1)		
Total (n = 85)	27 (31.8)	28 (32.9)	26 (30.6)	4 (4.0)

could mediate resistance to the non fluorinated quinolone (Lee et al., 2009).

Quinolone resistance was thought to be acquired only by chromosomal mutations, until plasmid-mediated quinolone resistance (PMQR) was described in 1998 (Chen, Zhang, et al., 2012; Chen, Shao, et al., 2012). A penta-peptide repeated protein that encoded by plasmid-mediated *qnr* gene accompanied with two additional PMQR determinants, *qepA* and *oxqAB*, have also been described to involved in quinolone resistance (Chen, Zhang, et al., 2012; Chen, Shao, et al., 2012; Wang et al., 2003). Meanwhile, *aac(6')-Ib* was also described to encode aminoglycoside acetyltransferase to reduce susceptibility to ciprofloxacin (Cui et al., 2008). Although aminoglycoside acetyltransferase encoded by *aac(6')-Ib* itself was not enough to cause complete ciprofloxacin resistance, when combined with gyrase mutation, over expressed efflux pump, or *qnr*, the expression of *aac(6')-Ib* could result in strong resistance to ciprofloxacin (Robicsek, Jacoby, & Hooper, 2006). Three major groups of *qnr* determinants (*qnrA*, *qnrB* and *qnrS*) had been identified in several bacterial species such as *Escherichia coli* and *Klebsiella pneumoniae* in Europe, the United States, Africa, Australia, and Asia (Robicsek et al., 2006; Wang et al., 2003; Yang, 2010). In our study, *qnr* was commonly detected in *Salmonella* isolates recovered in different districts and meats in Xinjiang province, which were highly consistent with previous studies (Robicsek et al., 2006; Wang et al., 2003). According to the previous results that *qnr* only conferred low-level quinolone resistance and could facilitate the development of QRDR mutations, interactions between QRDR mutations and *qnr* can result in higher-level fluoroquinolone resistance (Jacoby, Chow, & Waites, 2003; Wu et al., 2010). Our results partly agreed with and strengthened previous studies. However, no *qnr* genes, amino acid substitutions of GyrA and ParC were detected in some isolates that exhibited nalidixic acid resistance, the reason of which should be further studied.

PFGE was considered a gold-standard method for *Salmonella* molecular subtyping, and was commonly used for relatedness determination to the isolates in outbreaks and for epidemiology understanding of the sporadic cases acquired from various sources (Pang et al., 2007; Yang et al., 2010). In this study, several typical PFGE clusters were found among 99 isolates, and some isolates with identical PFGE patterns were grouped in a typical PFGE cluster, which suggested that *Salmonella* isolates in each certain serovar were genetically diverse.

In a word, results of this study demonstrated that the prevalence of *Salmonella* in retail meats in Uighur of Xinjiang province were not common, however, the isolates exhibited multidrug resistance, phenotypical and genotypical diverse.

Acknowledgment

The research work was supported by High Technology Research and Development Program of the Xinjiang Uygur Autonomous Region (201317106).

References

- Aslam, M., Checkley, S., Avery, B., Chalmers, G., Bohaychuk, V., Gensler, G., et al. (2012). Phenotypic and genetic characterization of antimicrobial resistance in *Salmonella* serovars isolated from retail meats in Alberta, Canada. *Food Microbiology*, 32(1), 110–117.
- Bonardi, S., Bassi, L., Brindani, F., D'Incau, M., Barco, L., Carra, E., et al. (2013). Prevalence, characterization and antimicrobial susceptibility of *Salmonella enterica* and *Yersinia enterocolitica* in pigs at slaughter in Italy. *International Journal of Food Microbiology*, 163(2–3), 248–257.
- Boonmar, S., Morita, Y., Pulsrikarn, C., Chaichana, P., Pornruangwong, S., Chauchom, S., et al. (2013). *Salmonella* prevalence in meat at retail markets in Pakse, Champasak Province, Laos, and antimicrobial susceptibility of isolates. *Journal of Global Antimicrobial Resistance*, 1(3), 157–161.
- Cattoir, V., Poirel, L., Rotimi, V., Soussy, C. J., & Nordmann, P. (2007). Multiplex PCR for detection of plasmid-mediated quinolone resistance *qnr* genes in ESBL-producing enterobacterial isolates. *Journal of Antimicrobial Chemotherapy*, 60(2), 394–397.
- Chen, Y., Shao, K., Guan, B., Hu, B., & Dong, H. (2010). Survey on *Salmonella* contamination of food in field of distribution in Shandong in 2003 to 2008. *Chinese Journal of Public Health Management*, 2, 163–165.
- Chen, Y., Shao, K., Guan, B., Hou, P., Zhang, H., & Wang, B. (2012). An analysis on the serotypes and antibiotic sensitivities of foodborne *Salmonella* in Shandong province from 2003 to 2010. *Chinese Journal of Food Hygiene*, 24(1), 9–13.
- Chen, X., Zhang, W., Pan, W., Yin, J., Pan, Z., Gao, S., et al. (2012). Prevalence of *qnr*, *aac(6')-Ib-cr*, *qepA*, and *oxqAB* in *Escherichia coli* isolates from humans, animals, and the environment. *Antimicrobial Agents and Chemotherapy*, 56(6), 3423–3427.
- Chen, S., Zhao, S., White, D. G., Schroeder, C. M., Ran, L., Yang, H., et al. (2004). Characterization of multiple-antimicrobial-resistant *Salmonella* serovars isolated from retail meats. *Applied and Environmental Microbiology*, 70(1), 1–7.
- Clinical and Laboratory Standards Institute. (2013). *Performance standards for antimicrobial susceptibility testing 2013*. PA: CLSI.
- Cui, S., Li, J., Sun, Z., Hu, C., Jin, S., Guo, Y., et al. (2008). Ciprofloxacin-resistant *Salmonella enterica* serotype typhimurium, China. *Emerging Infectious Diseases*, 14(3), 493–495.
- De, A. M., Bravo, D., & Medina, M. (2012). High pressure treatments on the inactivation of *Salmonella enteritidis* and the characteristics of beef Carpaccio. *Meat Science*, 92(4), 823–828.
- Dogru, A. K., Ayaz, N. D., & Gencay, Y. E. (2010). Serotype identification and antimicrobial resistance profiles of *Salmonella* spp. isolated from chicken carcasses. *Tropical Animal Health and Production*, 42(5), 893–897.
- Eaves, D. J., Liebana, E., Woodward, M. J., & Piddock, L. J. V. (2002). Detection of *gyrA* mutations in quinolone-resistant *Salmonella enterica* by denaturing high-performance liquid chromatography. *Journal of Clinical Microbiology*, 40(11), 4121–4125.
- Eaves, D. J., Luke, R., Gray, D. T., Antony, B., Woodward, M. J., White, A. P., et al. (2004). Prevalence of mutations within the quinolone resistance-determining region of *gyrA*, *gyrB*, *parC* and *parE* and association with antibiotic resistance in quinolone-resistant *Salmonella enterica*. *Antimicrobial Agents and Chemotherapy*, 48, 4012–4015.
- GB 4789.4-2010 Food safety detection methods of National Standards of the People's Republic of China, Food microbiology identification, *Salmonella* identification. (<http://down.foodmate.net/standard/sort/3/21703.html>).
- Hendriksen, R. S., Vieira, A. R., Karlsmose, S., Dm, L. F. W., Jensen, A. B., & Wegener, H. C. (2011). Global monitoring of *Salmonella* serovar distribution from the world health organization global foodborne infections network country data bank: results of quality assured laboratories from 2001 to 2007. *Foodborne Pathogens and Disease*, 8(8), 887–900.
- Jacoby, G. A., Chow, N., & Waites, K. B. (2003). Prevalence of plasmid-mediated quinolone resistance. *Antimicrobial Agents and Chemotherapy*, 47(2), 559–562.
- Kim, K. Y., Park, J. H., Kwak, H. S., & Woo, G. J. (2011). Characterization of the quinolone resistance mechanism in foodborne *Salmonella* isolates with high nalidixic acid resistance. *International Journal of Food Microbiology*, 146(1), 52–56.
- Laconcha, I., Baggesen, D. L., Rementeria, A., & Garaizar, J. (2000). Genotypic characterisation by PFGE of *Salmonella enterica* serotype enteritidis phage types 1, 4, 6, and 8 isolated from animal and human sources in three European countries. *Veterinary Microbiology*, 75(2), 155–165.
- Lee, H., Su, L., Tsai, M., Kim, S., Chang, H., Jung, S., et al. (2009). High rate of reduced susceptibility to ciprofloxacin and ceftriaxone among nontyphoid *Salmonella* clinical isolates in Asia. *Antimicrobial Agents and Chemotherapy*, 53(6), 2696–2699.
- Li, R., Jing, L., Yang, W., Liu, S., Yun, L., & Liu, K. (2013). Prevalence and characterization of *Salmonella* species isolated from pigs, ducks and chickens in Sichuan province, China. *International Journal of Food Microbiology*, 163(1), 14–18.
- Liu, W., Chen, J., Huang, Y., Liu, B., & Shi, X. (2010). Serotype, genotype, and antimicrobial susceptibility profiles of *Salmonella* from chicken farms in Shanghai. *Journal of Food Protection*, 73(3), 562–567 (6).
- Mąka, Ł., Maćkiw, E., Ścieżyńska, H., Pawłowska, K., & Popowska, M. (2014). Antimicrobial susceptibility of *Salmonella* strains isolated from retail meat products in Poland between 2008 and 2012. *Food Control*, 36, 199–204.
- Pan, W., Chen, X., Wang, X., Cong, Q., Pan, Z., Gao, S., et al. (2009). The analysis of quinolone resistance of the avian *Escherichia coli* and *Salmonella* isolates from 1993 to 2008. *Chinese Journal of Zoonoses*, 25(7), 630–635.
- Pang, J., Chiu, T., Helmuth, R., Schroeter, A., Guerra, B., & Tsen, H. (2007). A pulsed field gel electrophoresis (pfge) study that suggests a major world-wide clone of *Salmonella enterica* serovar enteritidis. *International Journal of Food Microbiology*, 116(3), 305–312.
- Park, C. H., Robicsek, A., Jacoby, G. A., Sahm, D., & Hooper, D. C. (2006). Prevalence in the United States of *aac(6')-Ib-cr* encoding a ciprofloxacin-modifying enzyme. *Antimicrobial Agents and Chemotherapy*, 50, 3953–3955.
- Ribot, E. M., Fair, M. A., Gautam, R., Cameron, D. N., Hunter, S. B., Swaminathan, B., et al. (2006). Standardization of pulsed-field gel electrophoresis protocols for the subtyping of *Escherichia coli* O157: H7, *Salmonella*, and *Shigella* for PulseNet. *Foodborne Pathogens and Disease*, 3(1), 59–67.
- Robicsek, A., Jacoby, G. A., & Hooper, D. C. (2006). The worldwide emergence of plasmid-mediated quinolone resistance. *Lancet Infectious Diseases*, 6(10), 629–640.
- Scallan, E., Hoekstra, R. M., Angulo, F. J., Tauxe, R. V., Widdowson, M. A., Roy, S. L., et al. (2011). Foodborne illness acquired in the United States major pathogens.

- Emerging Infectious Diseases*, 17(1), 7–15.
- Shao, K. (2011). *Study on types and active surveillance of foodborne Salmonella. Spp in Shandong Province*. PA: Shandong.
- Shariat, N., DiMarzio, M. J., Yin, S., Dettinger, L., Sandt, C., Lute, J. R., et al. (2013). The combination of *crispr-mv1st* and *pfge* provides increased discriminatory power for differentiating human clinical isolates of *Salmonella enterica* subsp. *enterica* serovar Enteritidis. *Food Microbiology*, 34(1), 164–173.
- Son, I., Jie, Z., Keys, C. E., Zhao, S., Meng, J., & Brown, E. W. (2013). Analysis of pulsed field gel electrophoresis profiles using multiple enzymes for predicting potential source reservoirs for strains of *Salmonella enteritidis* and *Salmonella typhimurium* isolated from humans. *Infection Genetics and Evolution*, 16(2), 226–233.
- Su, L., Chiu, C. H., Chu, C., & Jonathan, T. (2004). Antimicrobial resistance in non-typhoid *Salmonella* serotypes: a global challenge. *Clinical Infectious Diseases*, 39(4), 546–551.
- Thai, T. H., & Yamaguchi, R. (2012). Molecular characterization of antibiotic-resistant *Salmonella* isolates from retail meat from markets in northern Vietnam. *Journal of Food Protection*, 75(9), 1709–1714 (6).
- Thong, K. L., & Modarressi, S. (2011). Antimicrobial resistant genes associated with *Salmonella* from retail meats and street foods. *Food Research International*, 44(9), 2641–2646.
- U.S. Food and Drug Administration. (2013). *The National Antimicrobial Resistance Monitoring System. Manual of Laboratory Methods*.
- Van, T. T. H., Nguyen, H. N. K., Smooker, P. M., & Coloe, P. J. (2012). The antibiotic resistance characteristics of non-typhoidal *Salmonella enterica* isolated from food-producing animals, retail meat and humans in South East Asia. *International Journal of Food Microbiology*, 154(3), 98–106.
- Wang, X., Cheng, X., Wu, S., Chao, G. X., Jiao, X. A., & Liu, X. F. (2007). Effect of *gyrA* gene mutation in *Salmonella* on the drug susceptibility to fluoroquinolones. *Chinese Journal of Zoonoses*, 23, 891–894.
- Wang, M., Ran, L., Wang, Z., Li, Z., Yao, J., Fu, P., et al. (2004). Study on national active monitoring for food borne pathogens and antimicrobial resistance in China 2001. *Journal of Hygiene Research*, 33(1), 49–54.
- Wang, M., Tran, J., Jacoby, G., Zhang, Y., Wang, F., & Hooper, D. C. (2003). Plasmid-mediated quinolone resistance in clinical isolates of *Escherichia coli* from Shanghai, China. *Antimicrobial Agents and Chemotherapy*, 47(7), 2242–2248.
- Wang, Y., Yang, B., Wu, Y., Zhang, Z., Meng, X., Xi, M., et al. (2015). Molecular characterization of *Salmonella enterica* serovar enteritidis on retail raw poultry in six provinces and two national cities in China. *Food Microbiology*, 46, 74–80.
- White, D. G., Zhao, S., Sudler, R., Ayers, S., Friedman, S., Chen, S., et al. (2001). The isolation of antibiotic-resistant *Salmonella* from retail ground meats. *New England Journal of Medicine*, 345(16), 1147–1154.
- Wouafo, M., Nzouankeu, A., Kinfack, J. A., Fonkoua, M. C., Ejenguele, G., Njine, T., et al. (2010). Prevalence and antimicrobial resistance of *Salmonella* serotypes in chickens from retail markets in Yaounde (Cameroon). *Microbial Drug Resistance*, 16(2), 171–176.
- Wu, W., Hui, W., Jian, L., Wu, J., Chen, M., Xu, Y., et al. (2010). Genetic diversity of *Salmonella enterica* serovar Typhi and Paratyphi in Shenzhen, China from 2002 through 2007. *BMC Microbiology*, 10, 32.
- Yang, B. (2010). *Study on characterization and mechanism of antimicrobial resistance of foodborne Salmonella*. PA: Yangling.
- Yang, B., Qiao, L., Zhang, X., Cui, Y., Xia, X., Cui, S., et al. (2013). Serotyping, antimicrobial susceptibility, pulse field gel electrophoresis analysis of *Salmonella* isolates from retail foods in Henan Province, China. *Food Control*, 32(1), 228–235.
- Yang, B., Dong, Q., Zhang, X., Shen, J., Cui, S., Ying, S., et al. (2010). Prevalence and characterization of *Salmonella* serovars in retail meats of marketplace in Shaanxi, China. *International Journal of Food Microbiology*, 141, 63–72.
- Yang, B., Xi, M., Wang, X., Cui, S., Yue, T., Hao, H., et al. (2011). Prevalence of *Salmonella* on raw poultry at retail markets in China. *Journal of Food Protection*, 74(10), 1724–1728.
- Yin, M., Zhang, X., Ainatula, Duan, L., Gulnazi, Yang, J., et al. (2014). Investigation of *Salmonella* contamination in retail meats in Urumqi, Xinjiang in 2010–2012. *Chinese Journal of Food Hygiene*, 26(2), 172–175.
- Zhang, X., Liao, X., Hao, Z., Hu, D., Chui, H., & Zhang, G. (2009). Pro-active monitoring and establishment of DNA fingerprint database of *Salmonella* in raw meat food in Henan province during 2006–2007. *Chinese Journal of Health Laboratory Technology*, 7, 1545–1548.
- Zhang, Z., Meng, X., Wang, Y., Xia, X., Wang, X., Xi, M., et al. (2014). Presence of *qnr*, *aac(6)-Ib*, *qepA*, *oqxAB*, and mutations in gyrase and topoisomerase in nalidixic acid-resistant *Salmonella* isolates recovered from retail chicken carcasses. *Foodborne Pathogens and Disease*, 11(9), 698–705.
- Zhang, F., Zhu, G., & Li, P. (2008). Surveillance and analysis for food-borne pathogens in retail foods in Shaanxi province, 2002–2006. *Chinese Journal of Public Health*, 24(2), 222.
- Zhu, D., Peng, Z., Liu, S., Lai, H., Han, X., & Zou, L. (2014). Contamination and antimicrobial resistance patterns of *Salmonella* spp. from broiler slaughter and processing chain. *Food Science*, 17, 42.