

Escherichia coli O26 and O113:H21 on Carcasses and Beef from a Slaughterhouse Located in Mato Grosso, Brazil

Elis Caroline Celestina dos Santos,¹ Vinicius Silva Castro,² Adelino Cunha-Neto,^{3,4} Luis Fernando dos Santos,⁵ Deyse Christina Vallim,⁶ Rodrigo de Castro Lisboa,⁶ Ricardo César Tavares Carvalho,³ Carlos Adam Conte Junior,^{2,4,6} and Eduardo Eustáquio de Souza Figueiredo^{1,3}

Abstract

Shiga toxin-producing *Escherichia coli* (STEC) is a group of emerging pathogens that can cause human diseases, including hemolytic uremic syndrome (HUS) and hemorrhagic colitis (HC). Monitoring slaughtering stages and checking contamination points are crucial for the production of safe food. In this context, the aim of this study was to verify contamination by STEC strains, to determine the contamination points and evaluate the resistance profile to 12 antimicrobials used in both veterinary and human medicine. A total of 80 samples were obtained from eight collection points (pen floor, rectum, hide, carcass swabs and esophagus, diaphragm, masseter, and retail beef tissue samples). The isolates were collected by dilution plating on MacConkey agar with sorbitol, cefixime, and tellurite and analyzed by multiplex polymerase chain reaction for virulence genes. Serotyping of non-O157 was performed, and testing for 12 antibiotics by disk diffusion was carried out. A total of 18 STEC strains were isolated, presenting different virulence profiles. Contamination by STEC was observed in the rectum (5/18), carcass surface (5/18), hide (3/18), diaphragm (2/18), retail beef (2/18), and masseter muscle (1/18). Pen floor swabs and esophagus tissues showed no STEC contamination. Moreover, three strains were identified as O26 and three as O113:H21 strains, which have been linked to HUS and HC outbreak cases in Brazil. All STEC isolates were susceptible to all evaluated antimicrobials, except streptomycin. The presence of STEC strains is a direct risk to the consumer, especially when isolated from retail beef, and contamination can occur during different slaughter stages. However, antimicrobial resistance profiles did not identify multidrug-resistant strains, limiting potential antimicrobial resistance transmission to other pathogens.

Keywords: STEC, antimicrobial resistance, foodborne disease, food safety

Introduction

BEEF REPRESENTS ONE OF THE MAIN ANIMAL PROTEIN SOURCES consumed worldwide, with Brazil as the second largest exporter and the third major producer (USDA, 2017). The Brazilian state of Mato Grosso is the main national producer, responsible for ~19.3% of the beef produced in the country (BRASIL, 2016). Considering the high potential for beef production and export, microbiological controls are required to ensure pathogen-free production. Currently, large consumer markets require the monitoring of specific bacterial species involved in meat contamination. For example, the

United States and the European Union require Shiga toxin-producing *Escherichia coli* (STEC) monitoring, which is classified as pathogenic in beef products (USDA, 2011; ISO, 2012). However, for the efficient control of this pathogen during slaughtering, the entire process must be monitored regarding contamination and possible pathogen transmission to retail beef (Buncic *et al.*, 2014; Santos *et al.*, 2017).

E. coli strains reside in the mammal gastrointestinal tract, and most strain serotypes show a symbiotic relationship with hosts (Fouhy *et al.*, 2012). However, *E. coli* comprises six pathogenic groups, with the STEC group of main importance to public health due to its association to severe cases of

¹Programa de Pós graduação em Ciência Animal, Universidade Federal de Mato Grosso, Cuiabá, Brazil.

²Instituto de Química, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.

³Programa de Pós graduação Nutrição, Alimentos e Metabolismo, Universidade Federal de Mato Grosso, Cuiabá, Brazil.

⁴Faculdade de Veterinária, Universidade Federal Fluminense, Rio de Janeiro, Brazil.

⁵Instituto Adolf Lutz, Núcleo de Doenças Entéricas e Infecções por Patógenos Especiais, São Paulo, Brazil.

⁶Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil.

foodborne disease (Clements *et al.*, 2012). The main characteristic of strains belonging to this group is the presence of two immunologically distinct toxins, Stx1 and Stx2, which affect protein transduction in host cells (Castro *et al.*, 2017). The STEC group can be divided into two subgroups: O157, comprising O157: H7, implicated in significant outbreaks, and the non-O157 subgroup, comprising serogroups O26, O45, O103, O111, O121, and O145, most commonly implicated in disease outbreaks (Scallan *et al.*, 2011).

Currently, the STEC prevalence rate in food products in Brazil ranges from 1.9% to 9.5% (Bergamini *et al.*, 2007; Freitas Filho *et al.*, 2014; Cardoso *et al.*, 2017), and cases of hemolytic uremic syndrome (HUS) have been reported in the country for both the O113:H21 and O26 strains (Guth *et al.*, 2002; Bando *et al.*, 2017). In addition to pathogen detection during slaughter, another widely studied factor is STEC resistant to antimicrobials, mainly due to the increase in strains displaying multidrug resistance (Ayaz *et al.*, 2015; Amézquita-López *et al.*, 2016). Therefore, the aim of this study was to evaluate the presence of STEC strains during fresh beef processing and verify their antimicrobial resistance profile.

Materials and Methods

Sample collection

A total of 10 animals were followed from reception at the slaughterhouse lairage to retail beef cutting and wrapping. The samples were collected from eight different locations along the slaughter chain, totaling 80 analyzed samples. The samplings were carried out weekly between May 2015 and July 2015, in a slaughterhouse with an average slaughter capacity of 500 animals per day, under the Brazilian federal inspection system and located in the state of Mato Grosso, Brazil.

The pen floor, anal, and hide swabs (from the gluteobiceps muscle region) were obtained with the use of 3M Swab Samplers[®] kits, inoculated in buffered peptone water (BPW). For esophagus, diaphragm, and masseter samples, tissue fragments (200 g) were collected and packed in sterile plastic bags. The 3M Sponge-Stick kit was used to obtain the swabs from the carcasses, with a sterilized field boundary bracket (10 cm by 10 cm or 100 cm²) from each sampled neck, thorax, abdomen, and hindquarters. The retail beef tissue fragments (100 g) were collected and packed in sterile plastic bags. All samples were refrigerated (1–7°C) and transported to the laboratory for immediate analysis.

Culture enrichment and isolation

The samples were analyzed according to the U.S. Food and Drug Administration guidelines (USDA, 2002), with modifications during the pre-enrichment step. Briefly, a 1 mL aliquot of BPW from the swab samples was inoculated into 9 mL modified tryptone soy broth (mTSB). For tissue fragments, 25 g of each sample was inoculated into 225 mL of mTSB and incubated at 41.5°C. Both samples were homogenized in a stomacher (Marconi[®]) before plating, streaked in agar MacConkey cefixime telluride sorbitol (CT-SMAC), and incubated at 37°C for 24 h. Five typical colonies were striated on trypticase agar with yeast extract (TSA-YE) and incubated at 37°C for 24 h. A colony was inoculated on methylene Levine agar (L-EMB) and subjected to biochemical tests for indole, methyl red, Voges-Proskauer, and citrate

(IMViC) and β -glucuronidase (coli-complete—Biocontrol[®]). The obtained isolates were then stored in a brain heart infusion (BHI) broth containing 10% glycerol at –80°C.

Polymerase chain reaction assay for STEC and major non-O157 strains

A 1 mL aliquot of each BHI-stored isolate with glycerol was reactivated on nutrient agar and the colonies were subjected to DNA extraction and purification using the DNeasy[®] Blood & Tissue kit (Qiagen[®]). DNA quantification was performed using the Qubit 2.0 kit (Invitrogen[®]). Subsequently, two polymerase chain reactions (PCRs) were carried out, based on the primers described by the FDA USDA (2011). In the first reaction, a duplex-PCR was used for the detection of *stx1* and *stx2* genes. In the second reaction, a triplex-PCR was used to amplify the *ehxA* (enterohemolysin), +93 *uidA* (β -glucuronidase mutation characteristic for *E. coli* Enterohemorrhagic (EHEC), and *y-eaeA* (adhesion) genes (Table 1). The *E. coli* O157:H7 strain (ATCC 43895) was used as the positive control, and sterile ultrapure milli-Q water was used as the negative control.

For the m-PCR of the major non-O157 strains, the isolates were analyzed regarding the six major serogroups, known as the “big six.” Two m-PCR assays with the primers described by Bai *et al.* (2012) were carried out, designed for the following genes: *wzxO26*, *wzxO45*, *wzxO103*, *wzxO111*, *wbqEO121*, and *wzxO145* (Table 1). The amplicons were visualized on a photodocumentator (MiniBis-Pro DNT; Bio-Imaging Systems[®]).

Serotyping by agglutination

Serotype determinations of strains negative for the “big six” were carried out by the method described by Guinée *et al.* (1981), employing all available O (O1 to O181) and H (H1 to H56) antisera. Serotyping was carried out at the Center of Bacteriology and Nucleus of Enteric Diseases and Infections at the Special Pathogens Reference Laboratory at the Adolf Lutz Institute.

Antimicrobial resistance

Antimicrobial susceptibility was assessed using the disk diffusion technique described by the Clinical and Laboratory Standards Institute (CLSI, 2017). Each isolate was tested for 12 different antibiotics (Oxoid[®]): ampicillin (10 μ g), imipenem (10 μ g), tetracycline (30 μ g), gentamicin (10 μ g), nitrofurantoin (300 μ g), ciprofloxacin (30 μ g), cefoxitin (30 μ g), streptomycin (10 μ g), chloramphenicol (30 μ g), sulfamethoxazole-trimethoprim (30 μ g), ceftazidime (30 μ g), and nalidixic acid (10 μ g). Each colony was inoculated onto Mueller-Hinton broth 2 (MH; Himedia, India) and incubated between 2 and 4 h. The inoculum was then compared to the standard McFarland scale to 0.5 concentrations, followed by streaking on Mueller-Hinton 2 agar (MH; Himedia). The disks containing the antibiotics were then included and the plates were incubated at 37°C for 18 h. Finally, inhibition zones were measured (CLSI, 2017).

Pulsed-field gel electrophoresis

The pulsed-field gel electrophoresis (PFGE) technique was applied to discriminate between the O26 and O113:H21

TABLE 1. LIST OF PRIMERS, THEIR SEQUENCES, THE AMPLICON SIZE, AND THE REFERENCE USED IN THIS STUDY

Genes	Primer	Primer sequence (5'-3')	Amplicon (bp)	References
<i>stx1</i>	LP30	CAGTTAATGTGGTGGCGAAGG	348	USDA (2011)
	LP31	CACCAGACAATGTAACCGCTG		
<i>stx2</i>	LP43	ATCCTATTCCTCCGGGAGTTTACG	584	USDA (2011)
	LP44	GCGTCATCGTATACACAGGAGC		
+93 <i>uidA</i>	PT-2	GCGAAAACGTGTGGAATTGGG	252	USDA (2011)
	PT-3	TGATGCTCCATCACTTCCTG		
<i>y-eaeA</i>	AE22	ATTACCATCCACACAGACGGT	397	USDA (2011)
	AE20-2	ACAGCGTGGTTGGATCAACC		
<i>ehxA</i>	MFS1Fb	GTTTATTCTGGGGCAGGCTC	166	USDA (2011)
	MFS1R	CTTCACGTCACCATACATAT		
<i>wzxO45</i>	O45-F	GGGCTGTCCAGACAGTTCAT	890	Bai <i>et al.</i> (2012)
	O45-R	TGTACTGCACCAATGCACCT		
<i>wzxO103</i>	O103F2	GCAGAAAATCAAGGTGATTACG	740	Bai <i>et al.</i> (2012)
	O103R2	GGTTAAAGCCATGCTCAACG		
<i>wbqEO121</i>	O121-F2	TCAGCAGAGTGGAATAATTTTGT	587	Bai <i>et al.</i> (2012)
<i>wbqFO121</i>	O121-R2	TGAGCACTAGATGAAAAGTATGGCT		
<i>wzxO145</i>	O145F5	TCAAGTGTGGATTAAGAGGGATT	523	Bai <i>et al.</i> (2012)
	O145R5	CACTCGCGGACACAGTACC		
<i>wzxO26</i>	O26F4	AGGGTGCGAATGCCATATT	417	Bai <i>et al.</i> (2012)
	O26R4	GACATAATGACATACCACGAGCA		
<i>wzxO111</i>	O111F2	TGCATCTTCATTATCACACCAC	230	Bai <i>et al.</i> (2012)
	O111R2	ACCGCAAATGCGATAATAACA		

strains. The analyses were performed according to the CDC PulseNet protocol (Ribot *et al.*, 2006), on a CHEF-DR III (Bio-Rad®) equipment. The *Xba*I enzyme was used for digestion and the PFGE analyses and comparisons were performed using the GelJ software package (Heras *et al.*, 2015). A similarity analysis was performed using the Dice coefficient. Electrophoresis conditions were an initial switch time of 6.76 s, a final switch time of 35.38 s, and a run time of 18 h. The CDC *Salmonella* ser. Branderup isolate H9812 was used as the reference strain. Images of the PFGE gels were taken with the DOC PRINT II (Vilber Lourmat®).

Results

A total of 18 isolates were positive for one or more virulence genes and classified as STEC. The detected virulence gene profiles were the following: *stx2/ehxA* (38.8%), *stx1/stx2/ehxA* (22.2%), *stx2* (16.6%), *stx1/stx2/ehxA/uidA* (5.6%), *stx2/uidA/eaeA* (5.6%), *stx2/uidA* (5.6%), and *stx1/eaeA/uidA* (5.6%). After virulence gene detection, the 18 STEC isolates were subjected to PCR analyses for the six main non-O157 serogroups involved in cases of food outbreaks and serotyping by serum agglutination. Of the total isolates, 3 (16.6%) were positive for the *wzyO26* gene, characterizing *E. coli* O26 strains (Table 2), with one strain isolated from a retail beef sample. Moreover, serum agglutination showed that three isolates belonged to O113:H21, the serotype is involved in HUS and hemorrhagic colitis (HC) cases worldwide.

Furthermore, different STEC virulence profiles were detected in the same samples in some animals. For example, animal 5 had two STEC strains isolated from the carcass swab. However, the S44 and S46 strains were different regarding serogroup determination. In addition, animal 9 was positive for two strains in the hide swab (S101 and S102), but with different virulence genes.

The PFGE profiles indicated that S94 and S95 strains were clones, both obtained from the same sample, while another O113:H21 strain (S97) showed 75% similarity with the clones. In addition, the virulence gene profile was different, with only the presence of the *stx2* gene detected. For the O26 serotype isolated from the same animal (S31 and S35), the PFGE profile showed difference in the strains, and both were different to S101 strain (Fig. 1). Moreover, S35 strain showed the most distant similarity profile and the presence of the *stx2* toxin gene.

All *E. coli* STEC strains were sensitive to the evaluated antimicrobials, except for streptomycin, in which resistance was detected in four isolates (two resistant and two intermediate). Moreover, all *E. coli* O26 were sensitive to the 12 evaluated antimicrobials. The S97 strain showed intermediate resistance to streptomycin (Table 2).

Discussion

Contamination with STEC is a health risk to the consumer due to association with HUS, HC, and purpura thrombocytopenia (Karpman *et al.*, 2017; Castro *et al.*, 2017). The data reported herein indicate a high STEC prevalence in the analyzed samples (22.5%). In addition, all isolates except one presented the *stx2* gene, which encodes the main toxin involved in the most severe disease cases (Fuller *et al.*, 2011).

As expected, most STEC contamination was detected during the first slaughter stages, mainly in the rectum (5/18), hide (3/18), and carcass surface (5/18). This may be due to the fact these areas are more exposed to fecal contamination, with a greater possibility of STEC presence during the first processing stages (Martin and Beutin, 2011).

However, no STEC isolates were detected in swabs obtained from the pen floor, probably due to the high microbial load present in feces, increasing competition before or during enrichment (Vimont *et al.*, 2006). No STEC were detected in

TABLE 2. COLLECTION POINTS, PROFILE OF VIRULENCE GENES, ANTIMICROBIAL RESISTANCE, AND SEROTYPE BY POLYMERASE CHAIN REACTION AND AGGLUTINATION BY SERUM

Strain	Animal	Point	Genes					Antimicrobial resistance	Serotype
			Stx1	Stx2	y-eaeA	+93uidA	ehxA		
S09	2	Masseter muscle	—	+	—	—	+	Streptomycin	O83:H19
S31	4	Rectal swab	—	+	—	—	+	—	O26:HNT
S35	4	Hide swab	—	+	—	—	+	—	O26:HNT
S37	4	Diaphragm muscle	—	+	—	—	+	—	O73:H45
S44	5	Carcass swab	+	+	—	—	+	Streptomycin	O8:H21
S46	5	Carcass swab	—	+	—	—	+	—	O79:H7
S72	6	Diaphragm muscle	+	+	—	+	+	—	O79:H7
S73	6	Carcass swab	—	+	—	—	+	—	O79:H7
S94	7	Retail beef	+	+	—	—	+	—	O113:H21
S95	7	Retail beef	+	+	—	—	+	—	O113:H21
S97	8	Carcass swab	—	+	—	—	—	Intermediary streptomycin	O113:H21
S99	8	Carcass swab	—	+	—	—	+	—	O22:H16
S101	9	Rectal swab	+	+	—	—	+	—	O26:HNT
S102	9	Rectal swab	—	+	—	+	+	—	O117:H7
S103	9	Hide swab	—	+	—	+	—	—	O21:H19
S117	10	Rectal swab	—	+	—	—	—	—	ONT:H10
S119	10	Rectal swab	—	+	—	—	—	Intermediary streptomycin	ONT:H10
S123	10	Hide swab	+	—	+	+	—	—	O132:H21

+93uidA, non-glucuronidase encoding gene; eaeA, intimin encoding gene; ehxA, enterohemolysin-encoding gene; HNT, H nontypeable; stx1, gene encoding Shiga toxin type 1; stx2, gene encoding Shiga toxin type 2.

animals 1 and 3. Moreover, the lack of a “gold standard” methodology with adequate STEC and EHEC strain recoveries (Enterohemorrhagic *E. coli*) and inhibition of accompanying microflora remains a challenge (Conrad *et al.*, 2014).

Furthermore, the absence of strains exhibiting beta-glucuronidase enzyme production was verified in 14 of the 18 strains isolated in this study. The heterogeneity of the STEC group has been reported since the first outbreak in 1999 with a sorbitol-positive strain (Ammon *et al.*, 1999). Some studies report the absence of beta-glucuronidase enzyme production (Yang *et al.*, 2004; Ogura *et al.*, 2007; Perera *et al.*, 2015). Due to rapid cell multiplication and gene transmission, mainly through plasmids, the heterogeneity of *E. coli* metabolic pathways becomes an extra challenge in the development of culture media specific to certain pathogenic groups.

In this study, STEC strains were likely not transferred during processing. Moreover, the presence of pathogenic

O26 was detected in animals 4 and 9, and O113:H21 strains were detected in animals 7 and 8, both linked to certain foodborne illnesses cases (Hoshina *et al.*, 2001; CDC, 2015; Peron *et al.*, 2016).

The presence of *E. coli* O26 strains in central Brazil has been described previously by Salvadori *et al.* (2003) in calves with diarrhea. In addition, *E. coli* O26 was the first non-O157 STEC strain identified in HC and HUS cases in Brazil (Guth *et al.*, 2002). However, to date, this study is the first to report *E. coli* O26 contamination in retail beef produced in Brazil.

The *E. coli* O26 strains isolated herein were initially identified as enteropathogenic (EPEC), and *stx* genes were subsequently detected (Combes *et al.*, 2011). In addition, the presence of the enterohemolysin-encoding gene was verified in all *E. coli* O26 isolates, similar to reports by Lorenz *et al.* (2016), who demonstrated that *E. coli* O26 isolated from food frequently displayed the enterohemolysin gene without the presence of Intimin (*eaeA*). The enterohemolytic toxin can




FIG. 1. PFGE profiles for O26 and O113:H21. Dendrogram of XbaI PFGE for strains S35 (O26); S97 (O113:H21); S95 (O113:H21); S94 (O113:H21); S101 (O26); and S31 (O26) associated with bacterial virulence genes profile. PFGE, pulsed-field gel electrophoresis.

break down red blood cells, causing iron release and leading to HC (Fagan *et al.*, 1999).

E. coli O113:H21 strains have been linked to HUS in a Brazilian patient (Bando *et al.*, 2017). The presence of this strain during the beef processing represents a direct risk to the consumers. In addition, the high production and export of Brazilian beef increase the risk of disease cases or foodborne outbreaks.

The presence of the O group was observed in this study for O26 strains, although the H flagellum was nontypeable, perhaps being the cause of the phylogenetic distance observed between the isolates. Moreover, several studies have indicated genetic heterogeneity between strains with the O26 serotype, and differences in the analyzed virulence genes (Zhang *et al.*, 2007; Miko *et al.*, 2010; Sekse *et al.*, 2011).

Regarding antimicrobial resistance, all evaluated antimicrobials were effective in inhibiting the isolates, except for streptomycin, with only four isolates showing resistance to this agent. Streptomycin belongs to the aminoglycoside class and acts on bacteria through the inhibition of protein synthesis by binding to prokaryotic ribosomes (Mingeot-Leclercq *et al.*, 1999; Kester *et al.*, 2012). Bacterial resistance to streptomycin is primarily due to alteration of the binding site of the ribosome to the agent molecule (Murray, 1991).

In Brazil, the government instituted the National Program for the Prevention and Control of Antimicrobial Resistance in Agriculture in 2017, with the aim of preventing and monitoring antimicrobial resistance throughout the production chain (BRASIL, 2017). In addition, the use of antimicrobials as growth promoters has been controlled in the country since 1999 (BRASIL, 1999).

However, the use of antimicrobials in herds is common in clinical treatment, and the use of streptomycin is related to the treatment of diseases such as *Dermatophilosis* in the state of Mato Grosso (Bacha *et al.*, 2014). In addition, other cattle-related diseases are commonly treated with streptomycin, such as *Leptospira* (Martins and Lilenbaum, 2017).

Resistance to streptomycin has been reported by Park *et al.* (2015) as being prevalent in STEC from slaughterhouse beef and supermarkets. In a Turkish study, 11 STEC isolates from cattle were tested against several antimicrobial agents and all were resistant to streptomycin (Goncuoglu *et al.*, 2010). In Brazil, Cergole-Novella *et al.* (2006) reported that 75% of STEC isolated in the state of São Paulo from different origins exhibited resistance to streptomycin. However, in the study carried out by Cergole-Novella *et al.* (2006), resistance to tetracycline was most prevalent, while all of the STEC isolates in this study were sensitive to this antibiotic.

Conclusions

Contamination by Shiga toxin-producing bacteria can occur throughout the slaughter chain, with transference from hide to carcass and beef cuts through cross-contamination during the process. The presence of *E. coli* O26 and O113:H21 is a public health risk, due to association with cases of foodborne disease, including HC and HUS in Brazil. In addition, antimicrobial resistance for streptomycin detected in this study would only slightly increase the risk in treating bacterial infections, as, unlike other studies available in the literature, isolates were not resistant to multiple antimicrobials.

Acknowledgments

The authors would like to thank the Enterobacteria Laboratory (LABENT) at the Fundação Oswaldo Cruz—FIOCRUZ, the Fundação de Amparo à Pesquisa do Estado de Mato Grosso—FAPEMAT, and the Coordenação de Aperfeiçoamento de Pessoal do Ensino Superior (CAPES).

Disclosure Statement

No competing financial interests exist.

References

- Amézquita-López BA, Quiñones B, Soto-Beltrán M, Lee BG, Yambao JC, Lugo-Melchor OF, Chaidez C. Antimicrobial resistance profiles of Shiga toxin-producing *Escherichia coli* O157 and Non-O157 recovered from domestic farm animals in rural communities in Northwestern Mexico. *Antimicrob Resist Infect Control* 2016;5:1.
- Ammon A, Petersen LR, Karch H. A large outbreak of hemolytic uremic syndrome caused by an unusual sorbitol-fermenting strain of *Escherichia coli* O157:H-. *J Infect Dis* 1999;179:1274–1277.
- Ayaz ND, Gencay YE, Erol I. Phenotypic and genotypic antibiotic resistance profiles of *Escherichia coli* O157 from cattle and slaughterhouse wastewater isolates. *Ann Microbiol* 2015; 65:1137–1144.
- Bacha FB, Faccin TC, Lima SC, Leal CRB, Lemos RAA. *Dermatophytosis in Nelore calves in Mato Grosso do Sul*. Semina: Ciências Agrárias, 2014;1947–1954.
- Bai J, Paddock ZD, Shi X, Li S, An B, Nagaraja TG. Applicability of a multiplex PCR to detect the seven major Shiga toxin-producing *Escherichia coli* based on genes that code for serogroup-specific O-antigens and major virulence factors in cattle feces. *Foodborne Pathog Dis* 2012;9:541–548.
- Bando SY, Iamashita P, Guth BE, dos Santos LF, Fujita A, Abe CM, Ferreira LR, Moreira-Filho CA. A hemolytic-uremic syndrome-associated strain O113:H21 Shiga toxin-producing *Escherichia coli* specifically expresses a transcriptional module containing *dicA* and is related to gene network dysregulation in Caco-2 cells. *PLoS One* 2017;12:0189613.
- Bergamini AMM, Simões M, Irino K, Gomes TAT, Guth BEC. Prevalence and characteristics of Shiga Toxin-producing *Escherichia coli* (STEC) strains in ground beef in São Paulo, Brazil. *Braz J Microbiol* 2007;38:553–556.
- BRASIL. Ministério da Agricultura Quantidade de Abate Estadual for Ano/Espécie. 2016. Available at: http://sigsif.agricultura.gov.br/sigsif_cons/ap_abate_estaduais_cons?p_select=SIM&p_ano=2016&p_id_especie=9, accessed November 27, 2017.
- BRASIL. Ministério da Agricultura, Pecuária e abastecimento. 1999. Available at: www.agricultura.gov.br/assuntos/inspecao/produtos-animal/plano-de-nacional-de-controle-de-residuos-e-contaminantes/documentos-da-pncrc/instrucao-normativa-sda-n-o-42-de-20-de-dezembro-de-1999.pdf, Accessed March 27, 2018.
- BRASIL. Ministério da Agricultura, Pecuária e abastecimento. Plano de controle de antimicrobianos utilizados na produção animal. 2017. Available at: www.lex.com.br/legis_27542328_INSTRUCAO_NORMATIVA_N_41_DE_23_DE_OUTUBRO_DE_2017.aspx, accessed March 27, 2018.
- Buncic S, Nychas G-J, Lee MRF, Koutsoumanis K, Hébraud M, Desvaux M, Chorianopoulos N, Bolton D, Blagojevic B,

- Antic D. Microbial pathogen control in the beef chain: Recent research advances. *Meat Sci* 2014;97:288–297.
- Cardoso P, Marin JM, Cardoso P, Marin JM. Occurrence of non-O157 Shiga toxin-encoding *Escherichia coli* in artisanal mozzarella cheese in Brazil: Risk factor associated with food workers. *Food Sci Technol* 2017;37:41–44.
- Castro VS, Carvalho RCT, Conte-Junior CA, Figueiredo EES. Shiga-toxin producing *Escherichia coli*: Pathogenicity, supershedding, diagnostic methods, occurrence, and foodborne outbreaks. *Compr Rev Food Sci Food Saf* 2017; DOI: 10.1111/1541-4337.12302.
- CDC. Centre Diseases Control. *Escherichia coli* O26 infections linked to chipotle Mexican grill restaurants. | *E. coli* | CDC. 2015. Available at: <https://www.cdc.gov/ecoli/2015/o26-11-15/index.html> 2015, accessed November 1, 2017.
- Cergole-Novella MC, Nishimura LS, Irino K, Vaz TM, de Castro AF, Leomil L, Guth BE. *Stx* genotypes and antimicrobial resistance profiles of Shiga toxin-producing *Escherichia coli* strains isolated from human infections, cattle, and foods in Brazil. *FEMS Microbiol Lett* 2006;259:234–239.
- Clements A, Young JC, Constatinou N, Frankel G. Infection strategies of enteric pathogenic *Escherichia coli*. *Gut Microbes* 2012;3:71–87.
- CLSI. *Performance Standards for Antimicrobial Susceptibility Testing an Informational Supplement for Global Application Developed Through the Clinical and Laboratory Standards Institute*. 2017. Available at: https://clsi.org/media/1469/m100s27_sample.pdf, accessed March 12, 2018.
- Conrad CC, Stanford K, McAllister TA, Thomas J, Reuter T. Further development of sample preparation and detection methods for O157 and the top 6 non-O157 STEC serogroups in cattle feces. *J Microbiol Methods* 2014;105:22–30.
- Coombes BK, Gilmour MW, Goodman CD. The evolution of virulence in non-o157 shiga toxin-producing *Escherichia coli*. *Front Microbiol* 2011;90:1–3.
- Fagan PK, Hornitzky MA, Bettelheim KA, Djordjevic SP. Detection of shiga-like toxin (*stx1* and *stx2*), intimin (*eaeA*), and enterohemorrhagic *Escherichia coli* (EHEC) hemolysin (EHEC *hlyA*) genes in animal feces by multiplex PCR. *Appl Environ Microbiol* 1999;65:868–872.
- Fouhy F, Ross RP, Fitzgerald GF, Stanton C, Cotter PD. Composition of the early intestinal microbiota: Knowledge, knowledge gaps and the use of high-throughput sequencing to address these gaps. *Gut Microbes* 2012;3:203–220.
- Freitas Filho EG, Ferreira MRA, Pinto JFN, Conceição FR, Moreira CN. Enterohemorrhagic *Escherichia coli* O157:H7 from healthy dairy cattle in Mid-West Brazil: Occurrence and molecular characterization. *Pesqui Vet Bras* 2014;34:24–28.
- Fuller CA, Pellino CA, Flagler MJ, Strasser JE, Weiss AA. Shiga toxin subtypes display dramatic differences in potency. *Infect Immun* 2011;79:1329–1337.
- Goncuoglu M, Ormanci FSB, Ayaz ND, Erol I. Antibiotic resistance of *Escherichia coli* O157:H7 isolated from cattle and sheep. *Ann Microbiol* 2010;60:489–494.
- Guinée PAM, Jansen WH, Wadström T, Sellwood R. *Escherichia coli* associated with neonatal diarrhoea in piglets and calves, in Laboratory diagnosis in neonatal calf and pig diarrhoea. *Curr Top Vet Animal Sci* 1981;13:126–162.
- Guth BEC, Lopes de Souza R, Vaz TMI, Irino K. First Shiga toxin-producing *Escherichia coli* isolate from a patient with hemolytic uremic syndrome, Brazil. *Emerg Infect Dis* 2002;8:535–536.
- Heras J, Domínguez C, Mata E, Pascual V, Lozano C, Torres C, Zarazaga M. GelJ—A tool for analyzing DNA fingerprint gel images. *BMC Bioinform* 2015;26:16:270.
- Hoshina K, Itagaki A, Seki R, Yamamoto K, Masuda S, Muku T, Okada N. Enterohemorrhagic *Escherichia coli* O26 outbreak caused by contaminated natural water supplied by facility owned by local community. *Jpn J Infect Dis* 2001;54:247–248.
- ISO. International Organization for Standardization. Microbiology of food and animal feed—Real-time polymerase chain reaction (PCR) -based method for the detection of food-borne pathogens, Horizontal method for the detection of Shiga toxin-producing *Escherichia coli* (STEC) and the determination of O157, O111, O26, O103 and O145 serogroups. 2012. Available at: <https://www.iso.org/standard/53328.html>, accessed July 12, 2017.
- Karpman D, Loos S, Tati R, Arvidsson I. Haemolytic uraemic syndrome. *J Intern Med* 2017;281:123–148.
- Kester M, Karpa KD, Vrana KE. Treatment of infectious diseases. In: Vrana KE, ed. *Elsevier's Integrated Review Pharmacology*. Philadelphia, PA: Elsevier, 2012.
- Lorenz SC, Monday SR, Hoffmann M, Fischer M, Kase JA. Plasmids from Shiga toxin-producing *Escherichia coli* strains with rare enterohemolysin gene (*ehxA*) subtypes reveal pathogenicity potential and display a novel evolutionary path. *Appl Environ Microbiol* 2016;82:6367–6377.
- Martin A, Beutin L. Characteristics of Shiga toxin-producing *Escherichia coli* from meat and milk products of different origins and association with food producing animals as main contamination sources. *Int J Food Microbiol* 2011;146:99–104.
- Martins G, Lilenbaum W. Control of bovine leptospirosis: Aspects for consideration in a tropical environment. *Res Vet Sci* 2017;112:156–160.
- Miko A, Lindstedt BA, Brandal LT, Lobersli I, Beutin L. Evaluation of multiple-locus variable number of tandem-repeats analysis (MLVA) as a method for identification of clonal groups among enteropathogenic, enterohaemorrhagic and avirulent *Escherichia coli* O26 strains. *FEMS Microbiol Lett* 2010;303:137–146.
- Mingeot-Leclercq MP, Glupczynski Y, Tulkens PM. Aminoglycosides: Activity and resistance. *Antimicrob Agents Chemother* 1999;43:727–737.
- Murray BE. New aspects of antimicrobial resistance and the resulting therapeutic dilemmas. *J Infect Dis* 1991;163:1185–1194.
- Ogura Y, Ooka T, Whale A, Garmendia J, Beutin L, Tennant S, Krause G, Morabito S, Chinen I, Tobe T, Abe H, Tozzoli R, Caprioli A, Rivas M, Robins-Browne R, Hayashi T, Frankel G. TccP2 of O157:H7 and non-O157 enterohemorrhagic *Escherichia coli* (EHEC): Challenging the dogma of EHEC-induced actin polymerization. *Infect Immun* 2007;75:604–612.
- Park H, Yoon JW, Heo E-J, Ko EK, Kim KY, Kim YJ, Yoon HJ, Wee SH, Park YH, Moon JS. Antibiotic resistance and virulence potentials of Shiga toxin-producing *Escherichia coli* isolates from raw meats of slaughterhouses and retail markets in Korea. *J Microbiol Biotechnol* 2015;25:1460–1466.
- Perera A, Clarke CM, Dykes GA, Fegan N. Characterization of Shiga toxigenic *Escherichia coli* O157 and non-O157 isolates from ruminant feces in Malaysia. *Biomed Res Int* 2015;2015:382403.

- Peron E, Zaharia A, Zota LC, Severi E, Mardh O, Usein C, Balgradean M, Espinosa L, Jansa J, Scavia G, Rafila A, Serban A, Pistol A. Early findings in outbreak of haemolytic uraemic syndrome among young children caused by Shiga toxin-producing *Escherichia coli*, Romania, January to February 2016. *EuroSurveill* 2016;21:30170.
- Ribot EM, Fair MA, Gautom R, Cameron DN, Hunter SB, Swaminathan B, Barrett TJ. Standardization of pulsed-field gel electrophoresis protocols for the subtyping of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* for PulseNet. *Foodborne Pathog Dis* 2006;3:59–67.
- Salvadori MR, Valadares GF, Leite DS, Blanco J, Yano T. Virulence factors of *Escherichia coli* isolated from calves with diarrhea in Brazil. *Braz J Microbiol* 2003;34:230–235.
- Santos ECC, Cunha-Neto A, Castro VS, Carvalho RCT, Figueiredo EES. Evaluation of the sanitary conditions of head meat, esophagus, diaphragm meat, and boning scrap processing. *J Food Qual* 2017;48:1–4.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 2011;17:7–15.
- Sekse C, Sunde M, Lindstedt B-A, Hopp P, Bruheim T, Cudjoe KS, Kvitle B, Urdahl AM. Potentially human-pathogenic *Escherichia coli* O26 in Norwegian sheep flocks. *Appl Environ Microbiol* 2011;77:4949–4958.
- USDA. United States Department of Agriculture. Food and Drug Administration—Bacteriological Analytical Manual Chapter 4^a Diarrheagenic *Escherichia coli*. 2011. Available at: www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm070080.htm#STECPRO, accessed September 28, 2017.
- USDA. United States Department of Agriculture. Food and Drug Administration. Enumeration of *Escherichia coli* and the Coliform Bacteria. Bacteriological Analytical Manual (BAM). 2002. Available at: www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm064948.htm, accessed September 28, 2017.
- USDA. United States Department of Agriculture. Foreign Agricultural Service. 2017. Available at: [https://www.fas.usda.gov/data/search?f\[0\]=field_commodities%3A23&f\[1\]=field_topics%3A43](https://www.fas.usda.gov/data/search?f[0]=field_commodities%3A23&f[1]=field_topics%3A43), accessed September 28, 2017.
- Vimont A, Vernozy-Rozand C, Delignette-Muller ML. Isolation of *E. coli* O157:H7 and non-O157 STEC in different matrices: Review of the most commonly used enrichment protocols. *Lett Appl Microbiol* 2006;42:102–108.
- Yang Z, Kovar J, Kim J, Nietfeldt J, Smith DR, Moxley RA, Olson M, Fey P, Benson A. Identification of common subpopulations of non-sorbitol-fermenting, beta-glucuronidase-negative *Escherichia coli* O157:H7 from bovine production environments and human clinical samples. *Appl Environ Microbiol* 2004;70:6846–6854.
- Zhang Y, Laing C, Steele M, Ziebell K, Johnson R, Benson A, Taboada E, Gannon V. Genome evolution in major *Escherichia coli* O157:H7 lineages. *BMC Genomics* 2007;8:121.

Address correspondence to:

Eduardo Eustáquio de Souza Figueiredo, DVM, MSc, PhD
 Programa de Pós graduação em Ciência Animal
 Universidade Federal de Mato Grosso
 Fernando Correa da Costa Avenue, Boa Esperança
 Cuiabá CEP 78060-900
 Mato Grosso
 Brazil

E-mail: figueiredoeduardo@ufmt.br