

Prevalence, Molecular Characterization, and Antimicrobial Susceptibility of Methicillin-Resistant *Staphylococcus aureus* (MRSA) Isolated from Milk and Dairy Products

Maha Abdou Al-Ashmawy,¹ Khalid Ibrahim Sallam,¹ Samir Mohammed Abd-Elghany,¹ Mohamed Elhadidy,² and Tomohiro Tamura³

Abstract

The present work was undertaken to study the prevalence, molecular characterization, virulence factors, and antimicrobial susceptibility of methicillin-resistant *Staphylococcus aureus* (MRSA) in raw milk and dairy products in Mansoura City, Egypt. MRSA was detected in 53% (106/200) among all milk and dairy products with prevalence rates of 75%, 65%, 40%, 50%, and 35% in raw milk, Damietta cheese, Kareish cheese, ice cream, and yogurt samples, respectively. The mean *S. aureus* counts were 3.49, 3.71, 2.93, 3.40, and 3.23 log₁₀ colony-forming units (CFU)/g among tested raw milk, Damietta cheese, Kareish cheese, ice cream and yogurt, respectively, with an overall count of 3.41 log₁₀ CFU/g. Interestingly, all recovered *S. aureus* isolates were genetically verified as MRSA strains by molecular detection of the *mecA* gene. Furthermore, genes encoding α -hemolysin (*hla*) and staphylococcal enterotoxins (*sea*, *seb*, *sec*) were detected in all isolates. The antimicrobial susceptibility pattern of recovered MRSA isolates against 13 tested antimicrobials revealed that the least effective drugs were penicillin G, cloxacillin, tetracycline, and amoxicillin with bacterial resistance percentages of 87.9%, 75.9%, 65.2%, and 55.6%, respectively. These findings suggested that milk and dairy products represent a potential infection risk threat of multidrug-resistant and toxigenic *S. aureus* in Egypt due to neglected hygienic practices during production, retail, or storage stages. These findings highlighted the crucial importance of applying more restrictive hygienic measures in dairy production in Egypt for food safety.

Introduction

STAPHYLOCOCCUS AUREUS IS A MAJOR foodborne pathogen causing outbreaks of food poisoning worldwide (Meyrand *et al.*, 1998). Furthermore, *S. aureus* represents a main cause of mastitis in dairy cattle (Virgin *et al.*, 2009; Lee *et al.*, 2014; Xue *et al.*, 2014). Methicillin-resistant *S. aureus* (MRSA) represents those *S. aureus* strains that have acquired the *mecA* gene that encodes penicillin-binding protein 2a (PBP2a) mediating resistance to methicillin and all other β -lactam antibiotics and therefore represent a global health concern and public health threat (Arsic *et al.*, 2012).

Traditional dairy products, especially those produced from raw milk under neglected hygienic conditions, are potential vehicles for the transmission of different foodborne pathogens including toxigenic *S. aureus* (Kadariya *et al.*, 2014).

MRSA, including animal-associated strains, have been frequently recovered from different dairy products such as raw milk or raw-milk cheeses worldwide (Peton and Le Loir, 2014). Kareish cheese (acid curd skim-milk cheese) is a kind of homemade soft cheese manufactured in Egyptian villages that is considered a major protein supplement consumed by most Egyptian farmers, while Damietta cheese is a famous soft, white pickled cheese industrially processed and distributed in Egypt under different market names. Since these types of cheese are manufactured from unpasteurized milk under low hygienic standards, they may serve as a potential vehicle for the transmission of MRSA.

Several virulence factors have been reported to be correlated with the symptoms and severity of infections caused by *S. aureus* and representing major concerns for the food-processing industry (Xing *et al.*, 2016). These factors include

¹Departments of Food Hygiene and Control, and ²Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt.

³Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Sapporo, Japan.

hemolysins (alpha, beta, gamma, and delta), leukocidin, toxic shock syndrome toxin-1, and staphylococcal enterotoxins (SEs) (Dinges *et al.*, 2000). Staphylococcal enterotoxins are a group of single-chain, low molecular mass proteins that are produced during all phases of growth (Naffa *et al.*, 2006) and are responsible for gastrointestinal symptoms such as nausea, emesis, abdominal cramps, and diarrhea (McLauchlin *et al.*, 2000; Le Loir *et al.*, 2003). Many types of *S. aureus* enterotoxins have been reported but the major five serological types are SEA, SEB, SEC, SED, and SEE (Omoe *et al.*, 2003; Orwin *et al.*, 2003). SEs are resistant to inactivation by gastrointestinal proteases as well as to heat and therefore may retain their biological and immunological activities even following pasteurization, food processing, and exposure to gastrointestinal proteases (Asao *et al.*, 2003). Therefore, the detection of these SEs is proposed to be a reliable method for the confirmation of staphylococcal outbreaks and determination of the enterotoxigenicity of strains (da Cunha *et al.*, 2007). The enterotoxigenic *S. aureus* had been implicated in causing contamination of raw milk (Heidinger *et al.*, 2009; Fusco *et al.*, 2011), cheeses (Ertas *et al.*, 2010; Rosengren *et al.*, 2010), ice cream (Gücükoğlu *et al.*, 2013), and yogurt (Zakary *et al.*, 2011).

The present study was planned to gain better insight into the prevalence of MRSA in raw milk, Damietta cheese, Kareish cheese, ice cream, and yogurt marketed in different locations in Mansoura City, Egypt and to characterize the strains recovered using molecular analysis of marker genes (*nuc*, *coa*, and *mecA*), virulence genes (*hla*, *sea*, *seb*, *sec*, and *tst*), and antimicrobial resistance patterns.

Materials and Methods

Collection of samples

A total of 200 samples (40 samples of raw milk, Damietta cheese, Kareish cheese, ice cream, and yogurt) were collected on 12 occasions, during the period of June–November 2012 from different retail outlets, different shops, and supermarkets in Mansoura City, Egypt. All samples were transported in an ice box to Food Hygiene and Control Department Laboratory, Mansoura University, Egypt, for conventional bacteriological analysis.

Isolation, enumeration, and identification of *S. aureus*

The count of *S. aureus* was determined according to the International Commission on Microbiological Specifications for Foods (ICMSF, 1996) by using a surface plate technique onto Baird Parker agar (BP) with Egg Yolk Tellurite Emulsion (Baird Parker; Oxoid Ltd., Hampshire, UK, CM0257; Egg Yolk Tellurite Emulsion; Oxoid Ltd., Hampshire, UK, SR0054) followed by incubation at 37°C for 48 h. Suspected colonies were selected and picked up for further identification as previously reported by Bennett and Lancette (2001). One *S. aureus* isolate per positive sample was used to calculate the prevalence rate among different samples of raw milk and dairy products.

Molecular characterization of MRSA

Molecular identification and characterization of isolated *S. aureus* strains was carried out in the Bioproduction Research Institute, National Institute of Advanced Industrial

Science and Technology, Hokkaido, Japan. All *S. aureus* strains isolated from the sampled milk and dairy products were screened by polymerase chain reaction (PCR) for nuclease (*nuc*) and coagulase (*coa*) genes (*S. aureus* species-specific determinants) and *mecA* (methicillin resistance determinant). Simultaneous amplification of the three genes was used for MRSA confirmation. MRSA-confirmed isolates were used as targets for detection of five selected virulence genes including *hla*, *sea*, *seb*, *sec*, and *tst* genes encoding α -hemolysin, staphylococcal enterotoxins A, B, C, and TSS-T1, respectively. Genomic DNA of *S. aureus* isolates were prepared according to the method described by Sallam *et al.* (2015) using the Maxwell 16-cell DNA purification kit (Promega Corporation, Madison, WI). Genomic DNA from *E. coli* K12DH5 α strain was similarly prepared and used as a negative control template for PCR analyses. PCR for target genes was conducted using primers sets and cycling conditions previously described (Sallam *et al.*, 2015). The oligonucleotide primers were synthesized by Hokkaido System Science Co. Ltd. (Hokkaido, Sapporo, Japan). GeneAmp PCR system 2700 thermal cycler (Applied Biosystems, Foster City, CA) was used for detection of various target genes.

PCR was carried out in a 20- μ L reaction mixture containing 1.6 μ L *S. aureus* genomic DNA template, 1 μ L (6 pmol) for each of forward and reverse primers, 4 μ L dNTPs (2 mM), 0.4 μ L KOD FX Neo Polymerase enzyme (1.0U/ μ L), 10 μ L of 2 \times PCR Buffer for KOD FX Neo (Toyobo Co., Ltd., Osaka, Japan) and 2 μ L PCR grade water. Amplified genes of each PCR reaction mixture were separated by subjecting 3 μ L aliquots to agarose (1.2%) gel electrophoresis for 30 min at 100 V followed by a 20-min staining in ethidium bromide solution. The separated PCR products were then visualized under ultraviolet light and photographed.

DNA sequencing

For confirmation of the amplified genes, DNA sequencing was conducted using the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied BioSystems) following the manufacturer's instructions on an ABI Prism 3100 automated sequencer (Applied Biosystems). The resultant sequences were subjected to the GenBank database for homology search using BLAST, Basic Local Alignment Search Tool (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Primers used for PCR for sequencing were the same used for DNA amplification. Nucleotide sequence data were then analyzed by the GENETYXMAC software, version 12 (GENETYX Corp., Tokyo, Japan).

Antimicrobial susceptibility testing

Antimicrobial susceptibility patterns for recovered *S. aureus* isolates were determined by the agar disk diffusion method as published (Jorgensen and Turnidge, 2007). The antimicrobials tested were penicillin (10 IU), amoxicillin (30 μ g), tetracycline (30 μ g), streptomycin (10 μ g), cloxacillin (5 μ g), rifampicin (5 μ g), chloramphenicol (30 μ g), netilmicin (30 μ g), ciprofloxacin (5 μ g), amikacin (30 μ g), gentamicin (10 μ g), vancomycin (30 μ g), and sulfamethoxazole/trimethoprim (25 μ g). Results were recorded and interpreted after 24-h incubation at 35°C according to the guidelines of the National Committee of Clinical Laboratory Standards guidelines (Kiehlbauch *et al.*, 2000).

Results and Discussion

In this study, the mean *S. aureus* counts were 3.49, 3.71, 2.93, 3.40, and 3.23 log₁₀ CFU/g among tested raw milk, Damietta cheese, Kareish cheese, ice cream, and yogurt, respectively, with an overall count of 3.41 log₁₀ CFU/g (Table 1). Our results showed that 75% (30/40) of raw milk samples exceeded the maximal limits of 100 CFU/mL proposed by both Egyptian standards (ES, 2005) and European regulations (EEC, 1992). On the other hand, 65% (26/40), 40% (16/40), 40% (16/40), and 30% (12/40) of Damietta cheese, Kareish cheese, ice cream, and yogurt samples, respectively, exceeded the maximal limit of 100 CFU/g set for *S. aureus* by European Economic Community food legislation for soft and fresh cheeses, frozen milk-based products, and fermented dairy products (EEC, 1992) (Table 1). According to the Egyptian standards, soft cheeses (including Damietta and Kareish cheeses), ice cream, and yogurt must be free from pathogenic organisms and their toxins. Therefore, all cheese, ice cream, and yogurt samples in this study exceeded the permissible limits of *S. aureus* counts reported in Egyptian Standards.

From each positive sample identified through conventional culturing on BP agar, five characteristic colonies or all characteristic colonies of *S. aureus* were picked from BP plates and then subjected to biochemical identification and molecular characterization by positive amplification of *nuc*, *coa*, and *mecA* genes. The *nuc* gene encodes the thermonuclease enzyme and has been used as a valuable genetic marker for rapid identification of *S. aureus*. The staphylocoagulase is used as a genetic marker for discriminating coagulase-positive *S. aureus* from other less pathogenic coagulase-negative staphylococci (Watanabe *et al.*, 2009). All coagulase-positive *S. aureus* were screened for MRSA using positive detection of the *mecA* gene. This gene is located on a mobile genetic element (staphylococcal cassette chromosome *mec* [SCC*mec*]) and encodes for PBP2a responsible for resistance against β -lactam antibiotics by interfering with their binding to cell wall proteins (Sakoulas *et al.*, 2001). By simultaneous amplification *nuc*, *coa*, and *mecA* genes, a total of 53% (106/200) of tested samples were confirmed as MRSA positive. Among 106 positive samples, a total of 414 colonies were confirmed to be MRSA. This molecular screening of MRSA is reported to be more efficient than other phenotypic methods that are considered time consuming, expensive, and less accurate (Oliveira and Len-

castre, 2002; Normanno *et al.*, 2007; Aras *et al.*, 2012). PCR analyses verified the existence of *nuc*, *coa*, and *mecA* genes at the expected molecular size of 660 bp (Fig. 1A), 1000 bp (Fig. 1B), and 1200 bp (Fig. 1C), respectively, in all isolates. Furthermore, sequence analyses of the amplified genes showed 100% identity with the corresponding gene sequences of *S. aureus* accessible in the GenBank. Among dairy samples tested, MRSA was detected in raw milk, Damietta cheese, Kareish cheese, ice cream, and yogurt samples with prevalence rates of 75% (30/40), 65% (26/40), 40% (16/40), 50% (20/40), and 35% (14/40), respectively. The relatively lower prevalence rate in both Kareish cheese and yogurt might be attributed, in part, to the acidity of these dairy products. On the other hand, the higher contamination rate in Damietta cheese samples suggested lack of appropriate hygienic measures during production, handling, and/or distribution of cheese. A significantly lower prevalence rate from raw milk and dairy products was reported during a mini-survey analysis in Dakahlia province, Egypt where only 5 samples out of 95 tested yielded positive results for the *mecA* gene (3 raw milk samples and 1 each from Kareish cheese and ice cream samples). Similarly, very low contamination rates for MRSA in milk and dairy products were recorded in different countries. Gücükoğlu *et al.* (2013) revealed that 2.8% (1/35) of ice cream samples tested in Turkey were positive for MRSA. In another study in Turkey, 7.5% (3/40) *S. aureus* strains isolated from Urfa cheese samples were MRSA positive (Kav *et al.*, 2011). MRSA was not detected among any of the *S. aureus*-positive samples associated with foodborne illness in Alberta, Canada (Crago *et al.*, 2012). In Italy, Normanno *et al.* (2007) reported that 3.75% (6/160) of dairy products were positive for MRSA. In Korea, Lee (2003) revealed the occurrence of MRSA in 1.34% (12/894) of bovine milk samples. Another study in Iran revealed that 4% (2/50) of both pasteurized milk and traditional soft cheese samples were positive for MRSA while it could not be detected in raw milk (Mirzaei *et al.*, 2011). A study in the United States identified MRSA in 21.8% (29/133) of bovine milk samples (Matyi *et al.*, 2013). This diversity in prevalence rates among different studies might be related to different attributes including source of samples, geographical origin, sensitivity of the identification methods, the quantity of samples, inappropriate antimicrobial administration, prevention practices, production techniques, use of either pasteurized or raw milk in production, storage, and handling of different samples (Sader *et al.*, 2006; Chen *et al.*, 2010).

TABLE 1. *STAPHYLOCOCCUS AUREUS* COUNTS (LOG₁₀ CFU/G OR mL) FROM MILK AND DAIRY PRODUCTS ON BAIRD PARKER AGAR

| Product | No. of samples tested | Minimum | Maximum | Mean | Samples that exceeded the maximal limit ^a |
|-----------------|-----------------------|---------|---------|-------|--|
| Raw milk | 40 | 2.0 | 4.45 | 3.49* | 30 (75%) |
| Damietta cheese | 40 | 2.3 | 5.04 | 3.71* | 26 (65%) |
| Kareish cheese | 40 | 2.0 | 3.69 | 2.93* | 16 (40%) |
| Ice cream | 40 | 2.3 | 4.08 | 3.40* | 16 (40%) |
| Yogurt | 40 | 2.0 | 4.65 | 3.23* | 12 (30%) |
| All products | 200 | 2.0 | 5.04 | 3.41* | 100 (50%) |

*No significant difference at $p \leq 0.05$.

^aMaximum permissible limit is 2 log₁₀ colony-forming units (CFU)/g or mL according to the European Economic Community (EEC, 1992).




FIG. 1. Representative agarose gel electrophoresis for polymerase chain reaction (PCR) products of the marker genes identified in the *Staphylococcus aureus* strains isolated from milk and different dairy products. Three microliters from the PCR products were separated by electrophoresis on 1.2% agarose gel. Amplified DNA of the expected molecular size of 660 bp for *nuc* gene (A), 1000 bp for *coa* gene (B), and 1200 bp for *mecA* gene (C), were visualized under ultraviolet light. M, DNA marker (Gene Ladder Wide 1) used as a reference for fragment size; Lane C, *Escherichia coli* K12 DH5 α as a negative control strain; Lanes with the key numbers from 1 to 14 are representative of positive strains.

All MRSA confirmed isolates were further tested for five additional virulence genes, including *hla*, *sea*, *seb*, *sec*, and *tst* genes encoding α -hemolysin, *S. aureus* enterotoxins *sea*, *seb*, *sec*, and toxic shock syndrome toxin, respectively. The *hla* gene, which encodes the α -hemolysin (Hla) toxin is considered one of the most important virulence factors of *S. aureus* and is formed by most *S. aureus* strains. Interestingly, all MRSA recovered isolates (100%) were positive for the *hla* gene, which was detected at the expected molecular size of 744 bp (Fig. 2A). Likewise, all MRSA isolates (100%) were positive for the three enterotoxin genes, *sea*, *seb*, and *sec*, which were detected by PCR at the expected molecular size of 500 bp (Fig. 2B), 600 bp (Fig. 2C), and 300 bp (Fig. 2D), respectively. Nonetheless, no amplification of the *tst* gene was observed among recovered isolates (0%). Reports of the existence of SEs in *S. aureus* isolated from milk and dairy products vary from one study to another.

Normanno *et al.* (2005) reported that 55.5% of isolated *S. aureus* from several food products including raw milk, cheeses, ice cream, and other dairy products produced one or more SEs. Of these strains, 33.9% produced SEC, 26.5% produced SEA, 2.7% produced SEB, 1.7% produced SEA+SEB, 0.3% produced SEA + SEC, and 0.3% SEB + SEC. In another study, Morandi *et al.* (2007) revealed that 36 (51%) of 71 isolates from cow's milk were positive for the *sea* gene. In contrast to our high findings for the existence of *sea*, *seb*, and *sec* in all MRSA




FIG. 2. Representative agarose gel electrophoresis for polymerase chain reaction (PCR) products of the virulence genes identified in the *Staphylococcus aureus* strains isolated from different dairy products. Three microliters from the PCR products were separated by electrophoresis on 1.2% agarose gel. Amplified DNA of the expected molecular size of 744 bp for *hla* gene (A), 500 bp for *sea* gene (B), 600 bp for *seb* gene (C), and 300 bp for *sec* gene (D), were visualized under ultraviolet light. M, DNA marker (Gene Ladder Wide 1) used as a reference for fragment size; Lane C, *Escherichia coli* K12 DH5 α as a negative control strain; Lanes from 1 to 14 are representative positive strains for the target genes.

TABLE 2. ANTIMICROBIAL SUSCEPTIBILITY PATTERN OF MRSA ISOLATED FROM MILK AND DAIRY PRODUCTS

| Type of antimicrobials | No. of MRSA isolates (n=414) | | |
|--|------------------------------|-----------|------------|
| | S (%) | I (%) | R (%) |
| Tetracycline (30 μ g) | 80 (19.3) | 64 (15.5) | 270 (65.2) |
| Netilmicin (30 μ g) | 286 (69.1) | 50 (12.1) | 78 (18.8) |
| Amoxicillin (10 μ g) | 114 (27.5) | 70 (16.9) | 230 (55.6) |
| Cloxacillin (5 μ g) | 4 (0.97) | 96 (23.2) | 314 (75.9) |
| Streptomycin (10 μ g) | 134 (32.4) | 94 (22.7) | 186 (44.9) |
| Sulfamethoxazole/ trimethoprim (25 μ g) | 312 (75.4) | 44 (10.6) | 58 (14.0) |
| Gentamicin (10 μ g) | 262 (63.3) | 38 (9.2) | 114 (27.5) |
| Penicillin G (10 IU) | 26 (6.3) | 24 (5.8) | 364 (87.9) |
| Rifampicin (5 μ g) | 222 (53.6) | 40 (9.7) | 152 (36.7) |
| Chloramphenicol (30 μ g) | 206 (49.8) | 80 (38.3) | 128 (30.9) |
| Ciprofloxacin (5 μ g) | 290 (70.1) | 60 (14.5) | 64 (15.5) |
| Amikacin (30 μ g) | 238 (57.5) | 30 (7.3) | 146 (35.3) |
| Vancomycin (30 μ g) | 316 (76.3) | 62 (15.0) | 36 (8.7) |

MRSA, methicillin-resistant *Staphylococcus aureus*; S, susceptible; I, intermediate; R, resistant.

isolates recovered from milk and dairy products, Zouharova and Rysanek (2008) detected *sea* in 27.1% of isolates while the *seb* and *sec* genes were observed in 10% and 1.4% of the isolates, respectively. Jørgensen *et al.* (2005) reported that the *sea* gene was not detected in any of the isolates of *S. aureus* from bulk tank cow's milk samples, but found the *sec* gene in 17.2% of isolates and *seb* in 1.2% of the isolates. Low detection rates for SEs were also reported by Oliveira *et al.* (2011) who detected the *sec* gene in 6% (5/83) of isolated *S. aureus* strains from milk obtained from cows with subclinical mastitis, while they could not identify enterotoxin genes, *sea* or *seb* in any of the strains tested.

The antimicrobial susceptibility patterns for MRSA isolates are shown in Table 2. Of the 13 tested antimicrobials, the least effective drugs were penicillin G, cloxacillin, tetracycline, and amoxicillin with bacterial resistance percentages of 87.9%, 75.9%, 65.2%, and 55.6%, respectively. The most effective antimicrobials against MRSA isolates were vancomycin, sulfamethoxazole/trimethoprim, ciprofloxacin, netilmicin, and gentamicin, which exhibited bacterial sensitivity percentages of 76.3%, 75.45%, 70.1%, 69.1%, and 63.3%, respectively. Of the MRSA isolates tested, 348 (84.1%) isolates were resistant to 3 or more antimicrobials. In agreement with our study, MRSA strains isolated from Turkish Tulum cheese were found to have resistances to multiple antibiotics (Yesim and Çelik, 2012). Furthermore, several studies reported different antibiotic resistance patterns by *S. aureus* strains recovered from different raw milk and dairy products (Aarestrup *et al.*, 1995; Lange *et al.*, 1999; Tondo *et al.*, 2000; King *et al.*, 2016). The heterogeneity in antimicrobial susceptibility patterns observed among MRSA isolates may reflect the microbial adaptive response to the use and overuse of antimicrobials, geographical location, drugs approved, and farm-level management. One idea is that multidrug-resistant bacteria, particularly staphylococci associated with milk samples and its related products, might be related to human contamination rather than contamination with animal origins (Spanu *et al.*, 2012).

In summary, the present study suggested that raw milk and dairy products marketed in Mansoura, Egypt might be considered as potential vehicles for the transmission of multidrug-resistant MRSA that might have the potential to cause severe infection in humans due to the presence of a wide array of virulence factors. The high prevalence rate in this study shows that the application of more strict hygienic measures in the production of raw milk and dairy products is crucial to prevent the further spread of MRSA. As the implementation of a food safety system in a dairy processing plant included applying good manufacturing practices, sanitation standard operating procedures, and hazard analysis and critical control point (HACCP) (Cusato *et al.*, 2013), proper diagnosis of indicator microorganisms including MRSA from dairy samples is crucial in assessing the system's implementation performance. Future studies should be directed toward using a larger number of strains and samples in parallel with studying the epidemiological and pathogenic potential of these strains to monitor distribution, pathogenesis, and epidemiology of enterotoxigenic MRSA.

Disclosure Statement

No competing financial interests exist.

References

- Aarestrup FM, Wegener HC, Rosdahl VT. Evaluation of phenotypic and genotypic methods for epidemiological typing of *Staphylococcus aureus* isolates from bovine mastitis in Denmark. *Vet Microbiol* 1995;45:139–150.
- Asao T, Kumeda Y, Kawai T, Shibata T, Oda H, Haruki K, Nakazawa H, Kozaki S. An extensive outbreak of *Staphylococcus* food poisoning due to the low-fat milk in Japan: Estimation of enterotoxin A in the incriminated milk and powdered skim milk. *Epidemiol Infect* 2003;130:33–40.
- Aras Z, Aydinb I, Kav K. Isolation of methicillin-resistant *Staphylococcus aureus* from caprine mastitis cases. *Small Ruminant Res* 2012;102:68e73.
- Arsic B, Zhu Y, Heinrichs DE, McGavin MJ. Induction of the staphylococcal proteolytic cascade by antimicrobial fatty acids in community acquired methicillin resistant *Staphylococcus aureus*. *PLoS One* 2012;7:e45952.
- Bennett RW, Lancette GA. *Staphylococcus aureus*. In: *BAM Bacteriological Analytical Manual*. U.S. Food and Drug Administration, Centre for Food Safety & Applied Nutrition, AOAC International, Gaithersburg, MD; 2001, pp. 222–228.
- Cusato S, Gameiro AH, Corassin CH, Sant'ana AS, Cruz AG, Faria Jde A, de Oliveira CA. Food safety systems in a small dairy factory: Implementation, major challenges, and assessment of systems' performances. *Foodborne Pathog Dis* 2013;10:6–12.
- Chen CB, Chang HC, Huang YC. Nasal methicillin-resistant *Staphylococcus aureus* carriage among intensive care unit hospitalised adult patients in a Taiwanese medical centre: One time-point prevalence, molecular characteristics and risk factors for carriage. *J Hosp Infect* 2010;74:238–244.
- Crago B, Ferrato C, Drews SJ, Severson LW, Tyrrel G, Louie M. Prevalence of *Staphylococcus aureus* and methicillin-resistant *S. aureus* (MRSA) in food samples associated with foodborne illness in Alberta, Canada from 2007 to 2010. *Food Microbiol* 2012;32:202–205.
- da Cunha ML, Calsolari RA, Junior JP. Detection of enterotoxin and toxic shock syndrome toxin 1 genes in *Staphylococcus*, with emphasis on coagulase-negative staphylococci. *Microbiol Immunol* 2007;51:381–390.
- Dinges MM, Orwin PM, Schlievert PM. Enterotoxins of *Staphylococcus aureus*. *Clin Microbiol Rev* 2000;13:16–34.
- [EEC] European Economic Community. Council Directive No 92/46/EEC, 1992. Laying down the health rules for the production and placing on the market of raw milk, heat treated milk and milk based products. *Offic J Eur Commun* 1992; L268:1–32.
- [ES] Egyptian Standards. Milk and Dairy Products, Part 1: Raw milk, Egyptian organization for standardization and quality control, Directive No. 154-1, Arab Republic of Egypt, 2005.
- Ertas N, Gonulalan Z, Yildirim Y, Kum E. Detection of *Staphylococcus aureus* enterotoxins in sheep cheese and dairy desserts by multiplex PCR technique. *Int J Food Microbiol* 2010;142:74–77.
- Fusco V, Quero G, Morea M, Blaiotta G, Visconti A. Rapid and reliable identification of *Staphylococcus aureus* harbouring the enterotoxin gene cluster (*egc*) and quantitative detection in raw milk by real time PCR. *Int J Food Microbiol* 2011; 144:528–537.
- Gücükoğlu A, CadirciÖ, Terzi G, Kevenk T, Alisarli M. Determination of enterotoxigenic and methicillin resistant *Staphylococcus aureus* in ice cream. *J Food Sci* 2013;78:738–741.
- Heidinger J, Winter C, Cullor J. Quantitative microbial risk assessment for *Staphylococcus aureus* and *Staphylococcus enterotoxin A* in raw milk. *J Food Prot* 2009;72:1641–1653.

- [ICMSF] International Commission on Microbiological Specifications for Foods. *Staphylococcus aureus*. In: *Microorganisms in Food 5: Characteristics of Microbial Pathogens*. Roberts TA, Baird-Parker AC, Tompkin RB (eds.). London: Blackie Academic and Professional, 1996, pp. 299–333.
- Jørgensen H, Mørk T, Hogasen H, Røvik L. Enterotoxigenic *Staphylococcus aureus* in bulk milk in Norway. *J Appl Microbiol* 2005;99:158–166.
- Jørgensen JH, Turnidge JD. Susceptibility test methods: Dilution and disc diffusion methods. In: Murray PR, Baron EJ, Jørgensen JH, Landry ML, Pfaller MA (eds.). *Manual of Clinical Microbiology*, 9th ed. Washington, DC: ASM Press, 2007, pp. 1152–1172.
- Kadariya J, Smith TC, Thapaliya D. *Staphylococcus aureus* and staphylococcal food-borne disease: an ongoing challenge in public health. *Biomed Res Int* 2014;2014:827965.
- Kav K, Col R, Ardic M. Characterization of *Staphylococcus aureus* isolates from white-brined Urfa cheese. *J Food Prot* 2011;74:1788–1796.
- Kiehlbauch JA, Hannett GE, Salfinger M, Archinal W, Monserrat C, Carlyn C. Use of the National Committee for Clinical Laboratory Standards Guidelines for disk diffusion susceptibility testing in New York State laboratories *J Clin Microbiol* 2000;38:3341–3348.
- Lange C, Cardoso M, Senczek D, Schwarz S. Molecular subtyping of *Staphylococcus aureus* isolates from cases of bovine mastitis in Brazil. *Vet Microbiol* 1999;67:127–141.
- Le Loir Y, Baron F, Gautier M. *Staphylococcus aureus* and food poisoning. *Genet Mol Res* 2003;2:6–76.
- Lee JH. Methicillin (Oxacillin)-resistant *Staphylococcus aureus* strains isolated from major food animals and their potential transmission to humans. *Appl Environ Microbiol* 2003;69:6489–6494.
- Lee SH, Mangolin BL, Gonçalves JL, Neeff DV, Silva MP, Cruz AG, Oliveira CA. Biofilm-producing ability of *Staphylococcus aureus* isolates from Brazilian dairy farms. *J Dairy Sci* 2014;97:1812–1816.
- Matyi S, Dupre J, Johnson W, Hoyt P, White D, Brody T, Odenwald W, Gustafson J. Isolation and characterization of *Staphylococcus aureus* strains from a Paso del Norte dairy. *J Dairy Sci* 2013;96:3535–3542.
- McLauchlin J, Narayanan G, Mithani V, O'Neill G. The detection of enterotoxins and toxic shock syndrome toxin genes in *Staphylococcus aureus* by polymerase chain reaction. *J Food Prot* 2000;63:479–488.
- Meyrand A, Boutrand-Loei S, Ray-Gueniot S, Mazuy C, Gaspard CE, Jaubert G, Perrin G, Lapeyre C, Vernozy-Rozand C. Growth and enterotoxin production of *Staphylococcus aureus* during the manufacture and ripening of Camembert-type cheeses from raw goats' milk. *J Appl Microbiol* 1998;85:537–544.
- Mirzaei H, Tofighi A, Sarabi HK, Farajli M. Prevalence of methicillin-resistant *Staphylococcus aureus* in raw milk and dairy products in Sarab by culture and PCR techniques. *J Anim Vet Adv* 2011;10:3107–3111.
- Morandi S, Brasca M, Lodi R, Cremonesi PB, Castiglioni B. Detection of classical enterotoxins and identification of enterotoxin genes in *Staphylococcus aureus* from milk and dairy products. *Vet Microbiol* 2007;124:66–72.
- Naffa RG, Bdour SM, Migdadi HM, Shehabi AA. Enterotoxigenicity and genetic variation among clinical *Staphylococcus aureus* isolates in Jordan. *J Med Microbiol* 2006;55:183–187.
- Normanno G, Firinu A, Virgilio S, Mula G, Dambrosio A, Poggiu A, Decastelli L, Mioni R, Scuota S, Bolzoni G, Di Giannatale E, Salinetti AP, La Salandra G, Bartoli M, Zuccon F, Pirino T, Sias S, Parisi A, Quaglia NC, Celano GV. Coagulase-positive *Staphylococci* and *Staphylococcus aureus* in food products marketed in Italy. *Int J Food Microbiol* 2005;98:73–79.
- Normanno G, La Salandra G, Dambrosio A, Quaglia N, Corrente M, Parisi A, Santagada G, Firinu A, Crisetti E, Celano G. Occurrence, characterization and antimicrobial resistance of enterotoxigenic *Staphylococcus aureus* isolated from meat and dairy products. *Int J Food Microbiol* 2007;115:290–294.
- Oliveira DC, Lencastre H. Multiplex PCR strategy for rapid identification of structural types and variants of the *mec* element in methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2002;46:2155–2161.
- Oliveira L, Rodrigues AC, Hulland C, Ruegg PL. Enterotoxin production, enterotoxin gene distribution, and genetic diversity of *Staphylococcus aureus* recovered from milk of cows with subclinical mastitis. *Am J Vet Res* 2011;72:1361–1368.
- Omoe K, Hu D, Takahashi-Omoe H, Nakane A, Shinagawa K. Identification and characterization of a new staphylococcal enterotoxin-related putative toxin encoded by two kinds of plasmids. *Infect Immun* 2003;71:6088–6094.
- Orwin P, Fitzgerald J, Leung D, Gutierrez J, Bohach G, Schlievert P. Characterization of *Staphylococcus aureus* enterotoxin. *Infect Immun* 2003;71:2916–2919.
- Peton V, Le Loir Y. *Staphylococcus aureus* in veterinary medicine. *Infect Genet Evol* 2014;21:602–615.
- Rosengren A, Fabricius A, Guss B, Sylvén S, Lindqvist R. Occurrence of foodborne pathogens and characterization of *Staphylococcus aureus* in cheese produced on farm-dairies. *Int J Food Microbiol* 2010;144:263–269.
- Sader HS, Streit JM, Fritsche TR, Jones RN. Antimicrobial susceptibility of gram-positive bacteria isolated from European medical centres: Results of the Daptomycin Surveillance Programme (2002–2004). *Clin Microbiol Infect* 2006;12:844–852.
- Sakoulas G., Gold HS, Venkataraman L, DeGirolami PC, Eliopoulos GM, Qian Q. Methicillin-resistant *Staphylococcus aureus*: Comparison of susceptibility testing methods and analysis of *mecA*-positive susceptible strains. *J Clin Microbiol* 2001;39:3946–3951.
- Sallam KI, Abd-Elghany SM, Elhadidy M, Tamura T. Prevalence, molecular characterization and antimicrobial resistance profile of methicillin-resistant *Staphylococcus aureus* (MRSA) in retail chicken products. *J Food Prot* 2015;78:1879–84.
- Spanu V, Spanu C, Virids S, Cossu F, Scarano C, De Santis E. Virulence factors and genetic variability of *Staphylococcus aureus* strains isolated from raw sheep's milk cheese. *Int J Food Microbiol* 2012;34:587–590.
- Tondo EC, Guimarães MC, Henriques JA, Ayub MA. Assessing and analyzing contamination of a dairy products processing plant by *Staphylococcus aureus* using antibiotic resistance and PFGE. *Can J Microbiol* 2000;46:1108–1114.
- Virgin JE, Van Slyke TM, Lombard JE, Zadoks RN. Methicillin resistant *Staphylococcus aureus* detection in US bulk tank milk. Short communication. *J Dairy Sci* 2009;92:4988–4991.
- Watanabe S, Ito T, Sasaki T, Li S, Uchiyama I, Kishii K, Kikuchi K, Skov RL, Hiramatsu K. Genetic diversity of staphylocoagulase genes (*coa*): Insight into the evolution of variable chromosomal virulence factors in *Staphylococcus aureus*. *PLoS One* 2009;4:e5714.
- Xing X, Zhang Y, Wu Q, Wang X, Ge W, Wu C. Prevalence and characterization of *Staphylococcus aureus* isolated from goat milk powder processing plants. *Food Control* 2016;59:644–650.
- Xue T, Chen X, Shang F. Effects of lactose and milk on the expression of biofilm-associated genes in *Staphylococcus aureus* strains isolated from a dairy cow with mastitis. *J Dairy Sci* 2014;97:6129–6134.

- Yesim CH, Çelik TH. Detection of enterotoxigenic and antimicrobial resistant *S. aureus* in Turkish cheeses. *Food Control* 2012;24:100–103.
- Zakary E, Nassif M, Mohammed G. Detection of *Staphylococcus aureus* in bovine milk and its product by real time PCR assay. *Global J Biotechnol Biochem* 2011;6:171–177.
- Zouharova M, Rysanek D. Multiplex PCR and RPLA identification of *Staphylococcus aureus* enterotoxigenic strains from bulk tank milk. *Zoonoses Public Hlth* 2008;55:313–319.

Address correspondence to:
Maha Abdou Al-Ashmawy, PhD
Department of Food Hygiene and Control
Faculty of Veterinary Medicine
Mansoura University
Mansoura 35516, Egypt

E-mail: mahaalashmawy@yahoo.com