



Characterization of *Salmonella* isolated from apparently healthy slaughtered cattle and retail beef in Hawassa, southern Ethiopia



Kokeb Kore^a, Biruhtesfa Asrade^b, Kassa Demissie^c, Kassaye Aragaw^{b,*}

^a Gamo Gofa Zone Livestock and Fishery Resources Development Department, Arba Minch, Ethiopia

^b Hawassa University, School of Veterinary Medicine, P.O. Box 05, Hawassa, Ethiopia

^c Debre Berhan University, Department of Animal Sciences, P.O. Box 445, Debre Berhan, Ethiopia

ARTICLE INFO

Keywords:

Beef
Cattle
Salmonella
Serotypes
Ethiopia

ABSTRACT

The present study was conducted to determine the occurrence, serotype distribution and antimicrobial resistance of *Salmonella* serotypes in slaughtered cattle, slaughterhouse environment and retail beef. Cecal content (CC), mesenteric lymph nodes (MLN), spleen and carcass swab (CS) samples (each, n = 150) were collected from 150 cattle slaughtered at Hawassa municipality slaughterhouse. Floor swab specimens (SHFS) were collected on 11 occasions from the slaughterhouse, and 100 beef samples were collected from 100 butcher shops. The samples were cultured for *Salmonella*, following standard procedures. A total of 14 *Salmonella* isolates belonging to 3 serotypes namely *Salmonella enterica* serotype Muenchen (4 isolates), *S. enterica* serotype 1,4,5,12:i:- (5) and *S. enterica* serotype Korovi (5) were recovered. All of the 5 *S. enterica* serotype 1,4,5,12:i:- isolates belonged to phage type 120. Four (2.7%) of the slaughtered cattle carried *Salmonella* in their CC and/or MLN, while none of the spleen samples were positive for *Salmonella*. *Salmonella* was isolated from four (2.7%) CC and two (1.3%) MLN samples. Out of the total of 150 CS samples, two (1.3%) were found contaminated with *Salmonella*, while 4 (4%) of the 100 beef samples obtained from butcher shops yielded *Salmonella*. Two of the 11 (18%) SHFS were positive for *Salmonella*. All the four isolates from beef were *S. enterica* serotype Muenchen, while both of the isolates from MLN were *S. enterica* serotype 1,4,5,12:i:-. Both *S. enterica* serotype 1,4,5,12:i:- and *S. enterica* serotype Korovi were isolated from CC, CS and slaughterhouse environment. All the 14 isolates recovered during the study were tested and found pan-susceptible to a panel of 14 antimicrobials. The present study helped to update the information on the occurrence, serotype distribution and antibiogram of *Salmonella* in slaughter cattle and beef in Ethiopia.

1. Introduction

Salmonellosis is among the main foodborne diseases in humans with worldwide distribution. The global burden of non-typhoidal *Salmonella* gastroenteritis has been estimated to be 93.8 million cases each year with 155,000 deaths (Majowicz et al., 2010). Animals are the principal reservoirs of *Salmonella* infection to humans (Acha and Szyfres, 2001; EFSA/ECDC, 2014a) although foods of vegetable origin contaminated by animal products, human excreta or contaminated utensils have been implicated as the vehicle of human salmonellosis (Acha and Szyfres, 2001). Foods containing products from farm animals, including cattle, are important sources of human *Salmonella* infections (Smerdon et al., 2001; van Duynhoven et al., 2002). Beef has been implicated as a vehicle of human infection in several outbreaks (CDC, 1995; Davies et al., 1996; Roels et al., 1997). Contamination of carcass by *Salmonella* may occur at abattoirs from carrier animals excreting/harboring the

organism (Samuel et al., 1980; Nabbut and Al-Nakhli, 1982; Morgan et al., 1987; Berends et al., 1997; Arguello et al., 2012) and by contaminated abattoir equipment and utensils (Smeltzer et al., 1980; Mannion et al., 2012). Meat available at retail outlets comes through a long chain of slaughtering and transportation, where each step may pose a risk of microbial contamination including salmonellae. The sanitary conditions of abattoir and meat shop environments play important roles in the spreading of *Salmonella* contamination (Acha and Szyfres, 2001; Mannion et al., 2012; Garedew et al., 2015).

Contamination of food with antibiotic-resistant *Salmonella* can be a major threat to public health, due partly to increased virulence and treatment failures (Helms et al., 2002; Angulo et al., 2004; Varma et al., 2005). The use of antimicrobial agents in food animals may promote development of antimicrobial resistance in *Salmonella*, with potential for transmission to humans via the food supply or other routes (Angulo et al., 2000; Threlfall, 2002).

* Corresponding author.

E-mail address: kassaye.a@gmail.com (K. Aragaw).

Even if data regarding meat-borne diseases in Ethiopia are scarce, several studies conducted in different parts of the country have shown the prevalence of *Salmonella* in cattle, cattle carcasses and beef (Nyeleti et al., 2000; Alemayehu et al., 2003; Ejeta et al., 2004; Addis et al., 2011; Sibhat et al., 2011; Alemu and Zewde, 2012). The prevalence of non-typhoidal salmonellae in humans has also been described (Aseffa et al., 1994; Mache et al., 1997; Addis et al., 2011; Eguale et al., 2015). A high level of antimicrobial resistance in *Salmonella* isolated from food animals has also been reported in Ethiopia (Alemayehu et al., 2003; Molla et al., 2003; Molla et al., 2004; Aragaw et al., 2007).

Knowledge on the occurrence and identification of sources and points of contamination are imperative for implementation of effective strategies for prevention of contamination of meat with *Salmonella*. It is also important to monitor antimicrobial resistance profile of *Salmonella* isolates for proper treatment of patients. Therefore, this study was conducted with the aim of describing the occurrence and antimicrobial resistance of *Salmonella* serotypes in slaughter cattle, slaughterhouse environment and retail beef in Hawassa, southern Ethiopia.

2. Materials and methods

2.1. Study area and animals

The study was undertaken from November 2015 to April 2016 in Hawassa town and involved the Hawassa Municipality abattoir and 100 butcher shops in the town. Hawassa is the capital city of Southern Nations, Nationalities and Peoples Regional State (SNNPRS) located about 275 km South of Addis Ababa. It is geographically positioned at 7°5′ latitude N and 38°29′ longitude E at an altitude of 1708 m a.s.l. The mean annual rainfall and temperature of the area vary from 800 to 1000 mm and 20.1–25 °C respectively. According to the Central Statistical Agency (CSA) population census report, Hawassa town had 259,803 residents in 2007 (CSA, 2008). Hawassa is frequently visited by national and international tourists. The municipality of Hawassa town has its abattoir that gives service to the town's community.

On average, 100–150 adult beef cattle are slaughtered per day at Hawassa Municipality abattoir, except on Wednesdays and Fridays. After arriving at the slaughterhouse, the animals are inspected, and those fit to be slaughtered are moved into pens where they spend an average of 8–10 h without feed before slaughter.

2.2. Sampling and sample collection

The study involved bacteriological examination of samples collected from 150 apparently healthy slaughtered cattle, 11 slaughterhouse floor swabs (SHFS) and 100 beef samples from butcher shops for *Salmonella*. The sample size for the slaughter cattle was calculated with an expected 7% estimated prevalence (Alemu and Zewde, 2012), desired 95% confidence interval and 5% precision (Thrusfield, 2005) which gives a sample size of 100 animals; the sample size was increased to 150 to improve precision. The study animals were selected with systematic random sampling technique from a batch of cattle destined to be slaughtered on the sample collection day. On each sampling day, which was held weekly, four types of samples were collected from 10 to 15 selected cattle. The chosen animals were followed along the slaughtering process to collect the appropriate samples. Accordingly, carcass swabs (CS), mesenteric lymph nodes (MLN), spleen and cecal content (CC) samples, 150 each, were collected aseptically in separate sterile containers making the total number of samples collected from slaughtered animals 600. Simultaneously, one weekly swab sample was collected for a total of 11 weeks (representing sample collection dates) from slaughterhouse floor on the same date of sample collection from slaughtered animals by rubbing a 1 m² (2 m²) floor area using a gauze swab. Additionally, 100 beef samples were collected from 100 randomly selected butcher shops in the town. The butcher shops were selected from a list of butcher shops registered in the town with a

simple random sampling technique. A total of 711 samples from various sources were collected during the study, for isolation and characterization of *Salmonella*.

For CS sample, sterile gauze moistened with buffered peptone water (BPW) was rubbed over different parts of the carcass (100 cm² area each on the rump, flank, brisket and neck) and was then placed into wide-mouthed, screw-capped, 30 ml, universal glass bottles. Cecal content samples were collected directly from the cecum into sterile universal bottles through a puncture made with sterile scalpel blades. The samples were transported to the Microbiology Laboratory of the School of Veterinary Medicine of Hawassa University in ice boxes with ice packs. Upon arrival at the laboratory the samples were stored at 4 °C in the refrigerator for immediate microbiological analysis within 24 h of sampling. Samples were identified by date of collection, sources and sample type.

2.3. Isolation and identification

For the isolation and identification of *Salmonella*, the technique recommended by the International Organization for Standardization (ISO-6579) (ISO, 2002) and Quinn et al. (1999) were employed as previously described (Aragaw et al., 2007) with minor modification. For pre-enrichment, 25 g of MLN, spleen and meat samples each were weighed, passed over flame to disinfect the surface (MLN and spleen), cut into small pieces on sterile Petri dish using sterile scalpel blade and placed in screw capped wide mouth bottles containing 225 ml buffered peptone water (BPW) (HiMedia, Mumbai, India), thoroughly mixed by shaking and incubated for 16–20 h at 37 °C. Similarly, 25 g of CC was placed in a wide mouth container containing 225 ml of BPW shaken and incubated as described above. The swab samples (CS and SHFS) were also pre-enriched in 225 ml BPW. Rappaport-Vassiliadis (RV) medium (HiMedia, Mumbai, India) was used for selective enrichment of the samples. After vortexing the pre-enrichment broth culture 0.1 ml was transferred to a tube containing 10 ml of RV medium and incubated at 42 °C for 24 h.

Xylose lysine desoxycholate (XLD) agar and *Salmonella*-Shigella (SS) agar (both from Oxoid, Hampshire, England) plates were used for plating out and identification. A loop-full of inoculum from RV broth culture was plated onto XLD and SS plates each and incubated at 37 °C for 24 h. After incubation, the plates were examined for the presence of typical and suspect colonies. Up to five of these colonies were selected from the selective plating media, streaked onto the surface of nutrient agar (HiMedia, Mumbai, India) plates and incubated at 37 °C for 24 h. Presumptive *Salmonella* colonies were characterized using biochemical tests following standard methods (Quinn et al., 1999; ISO, 2002).

Biochemically typical *Salmonella* isolates were stab-cultured in tryptone soya agar (Titan Biotech Ltd., Rajasthan, India) and shipped to the Public Health Agency of Canada, Office International des Epizooties (OIE) Reference Laboratory for Salmonellosis, Guelph, Ontario, Canada for serotyping and phage typing.

2.4. Serotyping and phage typing

For serotyping, the somatic (O) antigens were determined by slide agglutination tests (Ewing, 1986) and flagellar antigens were determined using a microplate agglutination technique (Shipp and Rowe, 1980). The antigenic formulae of Grimont and Weill (2007) were used to identify and assign the serotypes of the isolates.

Phage typing of *S. enterica* serotype 1:4,5,12:i- isolates was performed by the methods developed by Callow (1959) and extended by Anderson et al. (1977) with reference phages obtained from the Public Health England, Gastrointestinal Bacteria Reference Unit, Colindale, England and the Public Health Agency of Canada, National Laboratory for Enteric Pathogens, Winnipeg, Canada.

Table 1Resistance breakpoints for a panel of antimicrobials used to determine antimicrobial susceptibility among *Salmonella* isolates.

Antimicrobial	Range (µg/ml)	Susceptible (µg/ml)	Resistant (µg/ml)	Reference
Amoxicillin/Clavulanic Acid (AMC)	1–32	< = 8	> = 32	CLSI M100-S24 ^a
Ampicillin (AMP)	1–32	< = 8	> = 32	CLSI M100-S24 ^a
Azithromycin (AZM)	0.12–16	< = 16	> = 32	No CLSI Interpretive Criteria ^b
Cefoxitin (FOX)	0.5–32	< = 8	> = 32	CLSI M100-S24 ^a
Ceftiofur (TIO)	0.12–8	< = 2	> = 8	CLSI VET01-S2 ^c
Ceftriaxone (CRO)	0.25–64	< = 1	> = 4	CLSI M100-S24 ^a
Chloramphenicol (CHL)	2–32	< = 8	> = 32	CLSI M100-S24 ^a
Ciprofloxacin (CIP)	0.015–4	< = 0.06	> = 1	CLSI M100-S24 ^a
Gentamicin (GEN)	0.25–16	< = 4	> = 16	CLSI M100-S24 ^a
Nalidixic Acid (NAL)	0.5–32	< = 16	> = 32	CLSI M100-S24 ^a
Streptomycin (STR)	2–64	< = 32	> = 64	No CLSI Interpretive Criteria ^b
Sulfisoxazole (SOX)	16–256	< = 256	> = 512	CLSI M100-S24 ^a
Tetracycline (TCY)	4–32	< = 4	> = 16	CLSI M100-S24 ^a
Trimethoprim/Sulphamethoxazole (SXT)	0.12–4	< = 2	> = 4	CLSI M100-S24 ^a

^a CLSI M100-S24.^b No CLSI breakpoints for *Enterobacteriaceae* available for this antimicrobial. Breakpoints based on distribution of MIC's and were harmonized with NARMS.^c CLSI VET01-S2.

2.5. Antimicrobial susceptibility testing

The antimicrobial susceptibility testing was carried out at Public Health Agency of Canada, National Microbiology Laboratory at Guelph, Canadian Integrated Program for Antimicrobial Resistance Surveillance (CIPARS), Guelph AMR Testing Laboratory, Guelph, Ontario, Canada. All the 14 isolates, obtained during the study, were tested for a panel of 14 antimicrobials (Panel Type: CMV3AGNF) using the Sensititre® Automated Microbiology System (Trek Diagnostics, Inc., Weslake, OH, USA) microtitre plates to determine the minimal inhibitory concentration (MIC) as per the manufacturer's instructions. Antimicrobial susceptibility results were interpreted according to Clinical and Laboratory Standards Institute (CLSI) (2013, 2014) guideline. The list of a panel of antimicrobials utilized, their symbols, and concentrations and breakpoints are shown in Table 1.

2.6. Data management and analysis

The data collected for each slaughtered animal included sex, age, breed, body condition, origin and results of the bacteriological examination of the samples as *Salmonella* negative or positive. An animal was considered *Salmonella* positive when it was positive for at least one of CC, MLN or spleen samples. Carcass swab sample was used as a measure of contamination. The data were entered in Microsoft Excel and were analyzed with Stata statistical software version 13 (Stata Corp., College Station, TX).

3. Results

3.1. Occurrence of *Salmonella*

Out of the total of 711 samples examined 14 (2.0%, 95% CI: 1.2, 3.3) were found positive for *Salmonella*. At an animal level (if *Salmonella* was isolated from at least one of the CC, MLN or spleen samples) *Salmonella* was isolated from four (2.7%, 95% CI: 1.0, 7.0) of the 150 slaughtered cattle examined. Two (1.3%, 95% CI: 0.3, 5.3) of the 150 carcasses were found contaminated with *Salmonella*. The occurrence at sample level for the slaughtered cattle was 2.7% (95% CI: 1.0, 7.0) for CC and 1.3% (95% CI: 0.3, 5.3) for MLN. Four of the 100 beef samples (4.0%, 95% CI: 1.5, 10.3) collected from butcher shops and two of the 11 (18.2%, 95% CI: 3.5, 58.0) SHFS samples yielded *Salmonella* (Table 2).

Table 2Occurrence of *Salmonella* by sample type.

Sample type	Number of samples		Proportion (%)	95% CI
	Examined	Positive		
CC	150	4	2.7	1.0–7.0
CS	150	2	1.3	0.3–5.3
MLN	150	2	1.3	0.3–5.3
Spleen	150	0	0	
SHFS	11	2	18	3.5–58
Beef	100	4	4.0	1.5–10.3
Total	711	14	2.0	1.2–3.3

CC, cecal content; CS, carcass swab; MLN, mesenteric lymph nodes; SHFS, slaughterhouse floor swab.

3.2. Serotype distribution

A total of 14 isolates belonging to the *Salmonella enterica* subsp. *enterica* group were recovered during the study. These isolates belonged to 3 serotypes namely *Salmonella enterica* serotype Muenchen (4 isolates), *S. enterica* serotype Korovi (5) and *S. enterica* serotype 1,4,5,12:i:- (5). All of the *S. enterica* serotype 1:4,5,12:i:- isolates were found to belong to phage type 120.

Salmonella enterica serotype Korovi was isolated from CC (n = 3), CS (1) and SHFS (1), and *S. enterica* serotype 1:4,5,12:i:- was isolated from CC (n = 1), MLN (2), CS (1) and SHFS (1). *Salmonella enterica* serotype Muenchen, however, was isolated only from beef samples collected from butcher shops (Table 3).

Only one individual animal was detected having two different serotypes in various samples: *S. enterica* serotype Korovi and *S. enterica* serotype 1,4,5,12:i:- in CC and MLN respectively.

3.3. Antimicrobial resistance

All the 14 isolates of *Salmonella* obtained in the study were tested for susceptibility to 14 antimicrobials (Table 1) and all of them were found susceptible to all of the antimicrobials used.

4. Discussion

The occurrence of *Salmonella* in slaughtered cattle, CS and retail beef observed in this study was generally low compared to previous reports from different parts of the county (Ashenafi, 1994; Nyeleti et al., 2000; Alemayehu et al., 2003; Sibhat et al., 2011; Alemu and Zewde, 2012; Garedew et al., 2015). However, we acknowledge that our

Table 3

Serotype distribution of *Salmonella* isolated from slaughtered cattle, SHFS and retail beef with sample type.

Sample type	No. examined	Serotype	No. of isolates	<i>Salmonella</i> occurrence
CC	150	S. Korovi	3	2.7%
		S. 1:4,5,12:i:-	1	
MLN	150	S. 1:4,5,12:i:-	2	1.3%
Spleen	150	–	0	0%
CS	150	S. Korovi	1	1.3%
		S. 1:4,5,12:i:-	1	
Beef	100	S. Muenchen	4	4.0%
SHFS	11	S. Korovi	1	18%
		S. 1:4,5,12:i:-	1	
Total	711		14	2.0%

CC, cecal content; MLN, mesenteric lymph nodes; CS, carcass swab; SHFS, slaughterhouse floor swab.

carcass sample size was too small to identify any trends and possible comparison with other studies. The reason for this was that the sample size formula used was not adjusted for precision down from the default 5% used. Thus, a more proper precision of 7% \pm 2% would have given a sample size of 625.

Salmonella occurrence in slaughtered animals, tissues and carcasses may be affected by hygienic conditions of holding pens, transportation stress, length of stay in lairage, hygienic status of the slaughterhouse environment and even the prevalence of *Salmonella* in beef cattle population supplying the slaughterhouses (Smeltzer et al., 1980; Morgan et al., 1987; Berends et al., 1997; Mannion et al., 2012).

We isolated three serotypes of *Salmonella* belonging to the *S. enterica* subsp. *enterica* group. Both *Salmonella enterica* serotype Korovi and serotype 1,4,5,12:i:- were isolated from CC, CS and slaughterhouse floor, while serotype 1,4,5,12:i:- and serotype Muenchen were isolated only from MLN and beef, respectively. Earlier works in Ethiopia have reported the isolation of *S. enterica* serotype Muenchen from cattle (Nyeleti et al., 2000), camels (Pegram et al., 1981; Molla et al., 2004) and pigs (Aragaw et al., 2007). The serotype was among the most frequently isolated serotypes from meat or poultry in the US (FSIS, 2016) and the ten most commonly identified serotypes causing human salmonellosis in 2012 in the US (CDC, 2014). *Salmonella enterica* serotype Muenchen has been associated with several disease outbreaks in people in different parts of the world (Proctor et al., 2001; Buchholz et al., 2005; OzFoodNet, 2005; CDC, 2016).

Salmonella enterica serotype 1,4,5,12:i:- and serotype Korovi, the other two serotypes isolated in the present study, are to the best of our knowledge reported here for the first time in Ethiopia, while none of the common previously reported serotypes in cattle (Alemayehu et al., 2003; Sibhat et al., 2011; Alemu and Zewde, 2012) were detected. However, the isolation of *S. enterica* serotype 1,4,5,12:i:- is in agreement with the rise of the serotype throughout the world. The recently emerged serotype has become one of the most frequently isolated serotypes in many countries (Echeita et al., 1999, 2001; CDC, 2008; Switt et al., 2009; Hauser et al., 2010; Mandilara et al., 2013; FSIS, 2016; Yang et al., 2015). Several molecular studies conducted on this serotype in different parts of the world (Echeita et al., 2001; Zamperini et al., 2007; Bugarel et al., 2012; Mandilara et al., 2013; Yang et al., 2015) demonstrated that it is a monophasic variant of *S. enterica* serotype Typhimurium unable to express the phase 2 flagellar antigens (lacking the second phase of H-antigen).

During 2011 *Salmonella enterica* serotype 1,4,5,12:i:- was the third frequently isolated serotype in Ireland next only to *S. enterica* serotype Typhimurium and *S. enterica* serotype Enteritidis (NSSLRL, 2011). During 2011 and 2012, it was the 3rd most frequently reported serotype from confirmed human cases in the EU/EEA countries (EFSA/ECDC, 2014a). It was also the 5th most commonly isolated serotype among

laboratory-confirmed human salmonellosis infections in the US in 2012 (CDC, 2014). This bacterium has also ranked among the four most frequent serotypes causing human salmonellosis in China (Yang et al., 2015). The identification of this serotype, which emerged in Europe in the mid-1990s and growing fast in proportional prevalence, in Ethiopia indicates that the serotype might have been imported to the country through contaminated foods or infected animals or human beings. The antimicrobial profiles of our *S. enterica* serotype 1,4,5,12:i:- isolates, however, differ from the most commonly encountered multidrug-resistant isolates (clones) reported elsewhere (CDC, 2008; EFSA/ECDC, 2014b; Yang et al., 2015).

Although many studies consider pigs as the reservoir of *S. enterica* serotype 1,4,5,12:i:- (De la Torre et al., 2003; Hauser et al., 2010), outbreaks associated with beef have been reported in Denmark (ECDC, 2012). Beef, along with pork, was reported to be one of the most important reservoirs of this serotype in China (Yang et al., 2015).

There are only a few published reports of isolation of *S. enterica* serotype Korovi from around the world. It was first reported from a serpent from the then Belgian Congo in 1955 (Kauffmann and van Oye, 1955). It was also isolated from 'lizards and snakes' in Sweden in 1994 for the first time in that country (only one isolate) (Boqvist et al., 2003). Our findings are the first report of the serotype from Ethiopia. The isolation of this rare serotype, which has been reported mainly from reptiles, in a relatively high frequency ($n = 3$) in the CC of slaughter cattle on different sample collection dates and its detection from CS and slaughterhouse floor samples needs further study.

Contrary to our finding of pan-susceptibility in all 14 isolates obtained in our study, *S. enterica* serotype 1,4, [5],12:i:- was reported to be characterized by multidrug resistance (; Echeita et al., 1999, 2001; Mandilara et al., 2013; EFSA/ECDC, 2014b, Yang et al., 2015). Extremely high levels of resistance were observed for ampicillin, streptomycin, sulfonamides and tetracycline (resistance [R] type-ASSuT) in EU (EFSA/ECDC, 2014b) and China (Yang et al., 2015). *Salmonella enterica* serotype Muenchen has also been found in some studies to demonstrate multiple drug resistance for up to seven antimicrobials: ampicillin, chloramphenicol, streptomycin, sulfamethoxazole, tetracycline, amoxicillin/clavulanic acid, and kanamycin (R-type ACSSuT-AxK) (Gebreyes and Thakur, 2005). Our finding was, however, consistent with previous reports in Ethiopia which indicated susceptibility of the serotype to all antimicrobials tested (Molla et al., 2004; Aragaw, 2005). As the lack of antimicrobial resistance of the isolates observed in this study was based on a small number of observations ($n = 14$), there is a need for a study with larger number of isolates to substantiate it.

The results of this study indicate that the occurrence of *Salmonella* in cattle slaughtered at Hawassa and beef sold in butcher shops in the town are relatively low, with 2.3% slaughter cattle and 4.0% fresh retail beef positive for culture. The study also demonstrated that serotypes which are characterized by multiple drug resistance somewhere may be pan-susceptible elsewhere. This study contributes to the global information on the occurrence and antibiogram of *Salmonella* in general and the expanding *S. enterica* serotype 1,4,5,12:i:- and the rare and usually reptile-associated *S. enterica* serotype Korovi in particular as the first report of their isolation in Ethiopia.

The contamination of carcasses and beef with *Salmonella* serotypes known to cause human infections, although at a low level, is a significant public health risk in Ethiopia where a considerable proportion of the population consumes raw or undercooked beef. Efforts to keep the prevalence low through control in animals, prevention of contamination of animal products at slaughter and butcher shops, and adequate cooking of meat are recommended.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgements

The authors are grateful for the cooperation they got from the staff of Hawassa Municipality abattoir during sample collection. The kind collaboration of Public Health Agency of Canada, Office International des Epizooties (OIE) Reference Laboratory for Salmonellosis, Guelph, Ontario, Canada for serotyping and phage typing and Public Health Agency of Canada, National Microbiology Laboratory at Guelph, Canadian Integrated Program for Antimicrobial Resistance Surveillance (CIPARS), Guelph AMR Testing Laboratory, Guelph, Ontario, Canada for antimicrobial resistance testing of our isolates is highly appreciated. This study was partially financed by School of Graduate Studies of Hawassa University, in the form of an MSc research fund provided to K.K.

References

- Acha, P., Szyfres, B., 2001. 3rd ed. Zoonoses and Communicable Diseases Common to Man and Animals: Bacterioses and Mycoses, vol. 1. P an American Health Organization, Washington, D.C, pp. 233–246.
- Addis, Z., Kebede, N., Sisay, Z., Alemayehu, H., Wubetie, A., Kassa, T., 2011. Prevalence and antimicrobial resistance of *Salmonella* isolated from lactating cows and in contact humans in dairy farms of Addis Ababa: a cross sectional study. *BMC Infect. Dis.* 11 (1), 222.
- Alemayehu, D., Molla, B., Muckle, A., 2003. Prevalence and antimicrobial resistance pattern of *Salmonella* isolates from apparently healthy slaughtered cattle in Ethiopia. *Trop. Anim. Health Prod.* 35, 309–319.
- Alemu, S., Zewde, B.M., 2012. Prevalence and antimicrobial resistance profiles of *Salmonella enterica* serovars isolated from slaughtered cattle in Bahir Dar, Ethiopia. *Trop. Anim. Health Prod.* 44 (3), 595–600.
- Anderson, E.S., Ward, L.R., Saxe, M.J., de Sa, J.D., 1977. Bacteriophage-typing designations of *Salmonella typhimurium*. *J. Hyg. (Lond.)* 78 (2), 297–300.
- Angulo, F.J., Johnson, K.R., Tauxe, R.V., Cohen, M.L., 2000. Origins and consequences of antimicrobial-resistant nontyphoidal *Salmonella*: implications for the use of fluoroquinolones in food animals. *Microb. Drug Resist.* 6 (1), 77–83.
- Angulo, F.J., Nargund, V.N., Chiller, T.C., 2004. Evidence of an association between use of anti-microbial agents in food animals and anti-microbial resistance among bacteria isolated from humans and the human health consequences of such resistance. *J. Vet. Med. B* 51 (8–9), 374–379.
- Aragaw, K., Molla, B., Muckle, A., Cole, L., Wilkie, E., Poppe, C., Kleer, J., Hildebrandt, G., 2007. The characterization of *Salmonella* serovars isolated from apparently healthy slaughtered pigs at Addis Ababa abattoir, Ethiopia. *Prev. Vet. Med.* 82, 252–261.
- Aragaw, K., 2005. *Salmonella* in Apparently Healthy Slaughtered Swine in Addis Ababa Ethiopia. Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia (Unpublished MSc thesis).
- Arguello, H., Carvajal, A., Collazos, J.A., García-Feliz, C., Rubio, P., 2012. Prevalence and serovars of *Salmonella enterica* on pig carcasses, slaughtered pigs and the environment of four Spanish slaughterhouses. *Food Res. Int.* 45 (2), 905–912.
- Aseffa, A., Mengistu, G., Tiruneh, M., 1994. *Salmonella* newport: outbreak of food poisoning among college students due to contaminated undercooked eggs. *Ethiop. Med. J.* 32 (1), 1–6.
- Ashenafi, M., 1994. Microbial flora and incidence of some foodborne pathogens on fresh raw beef from butcher's shops in Awassa, Ethiopia. *Bull. Anim. Hlth. Prod. Afr.* 42, 273–277.
- Berends, B.R., Van Knapen, F., Snijders, J.M., Mossel, D.A., 1997. Identification and quantification of risk factors regarding *Salmonella* spp. on pork carcasses. *Int. J. Food Microbiol.* 36, 199–206.
- Boqvist, S., Hansson, I., Bjerselius, U.N., Hamilton, C., Wahlström, H., Noll, B., Tysen, E., Engvall, A., 2003. *Salmonella* isolated from animals and feed production in Sweden between 1993 and 1997. *Acta Vet. Scand.* 44, 181–197.
- Buchholz, U., Brodhun, B., Brockmann, S.O., Dreweck, C.M., Prager, R., Tschäpe, H., Ammon, A., 2005. An outbreak of *Salmonella* München in Germany associated with raw pork meat. *J. Food Prot.* 68 (2), 273–276.
- Bugarel, M., Granier, S.A., Bonin, E., Vignaud, M.L., Roussel, S., Fach, P., Brisabois, A., 2012. Genetic diversity in monophasic (1,4,[5],12:i:- and 1,4,[5],12:-1,2) and in non-motile (1,4,[5],12:-) variants of *Salmonella enterica* S. *Typhimurium*. *Food Res. Int.* 45, 1016–1024.
- CDC, 1995. Outbreak of salmonellosis associated with beef jerkey-New Mexico. *MMWR Morb. Mortal. Wkl. Rep.* 44 (42), 785–788.
- CDC, 2008. *Salmonella: Annual Summary*. (<http://www.cdc.gov/ncidod/dbmd/phlhisdata/salmtab/2006/SalmonellaAnnualSummary2006.pdf>). Accessed 15 June 2016).
- CDC, 2014. National Enteric Disease Surveillance: *Salmonella* Annual Report. (<http://www.cdc.gov/nceid/dwef/pdfs/salmonella-annual-report-2012-508c.pdf>). Accessed 25 May 2016).
- CDC, 2016. Multistate Outbreak of *Salmonella* Infections Linked to Alfalfa Sprouts from One Contaminated Seed Lot (Final Update). (<http://www.cdc.gov/salmonella/muenchen-02-16/>). Accessed 25 May 2016).
- CLSI, 2013. Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals, 2nd Informational Supplement VET01-S2. Clinical and Laboratory Standards Institute, Wayne, PA.
- CLSI, 2014. Performance Standards for Antimicrobial Susceptibility Testing: 24th Informational Supplement M100-S24. Clinical and Laboratory Standards Institute, Wayne, PA.
- CSA, 2008. Summary and Statistical Report of the Population and Housing Census. Population Size by Age and Sex. Population Census Commission. Federal Democratic Republic of Ethiopia, Addis Ababa.
- Callow, B.R., 1959. A new phage-typing scheme for *Salmonella typhimurium*. *J. Hyg. (Lond.)* 57, 346–359.
- Davies, A., O'Neill, P., Towers, L., Cooke, M., 1996. An outbreak of *Salmonella typhimurium* DT104 food poisoning associated with eating beef. Communicable disease report. *CDR Rev.* 6 (11), R159–62.
- De la Torre, E., Zapata, D., Tello, M., Mejia, W., Frias, N., Pena, F.G., Mateu, E.M., Torre, E., 2003. Several *Salmonella enterica* subsp. *enterica* serotype 4,5,12:i:- phage types isolated from swine samples originate from serotype Typhimurium DT U302. *J. Clin. Microbiol.* 41 (6), 2395–2400.
- ECDC, 2012. Urgent Inquiries Outbreak of Monophasic *Salmonella* Typhimurium in Denmark. European Center for Disease Prevention and Control, Stockholm.
- EFSA/ECDC, 2014a. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2012. *EFSA J.* 12 (2), 3547.
- EFSA/ECDC, 2014b. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2012. *EFSA J.* 12 (3), 3590.
- Echeita, M.A., Aladueno, A., Cruchaga, S., Usera, M.A., 1999. Emergence and spread of an atypical *Salmonella enterica* subsp. *enterica* serotype 4,5,12:i:2 strain in Spain. *J. Clin. Microbiol.* 37, 3425.
- Echeita, M.A., Herrera, S., Usera, M.A., 2001. Atypical fljB-negative *Salmonella enterica* subsp. *enterica* strain of serovar 4,[5],12:i:- appears to be a monophasic variant of *Salmonella Typhimurium*. *J. Clin. Microbiol.* 39, 2981–2983.
- Egualde, T., Gebreyes, W.A., Asrat, D., Alemayehu, H., Gunn, J.S., Engidawor, E., 2015. Non-typhoidal *Salmonella* serotypes, antimicrobial resistance and co-infection with parasites among patients with diarrhea and other gastrointestinal complaints in Addis Ababa, Ethiopia. *BMC Infect. Dis.* 15, 497.
- Ejeta, G., Molla, B., Alemayehu, D., Muckle, A., 2004. *Salmonella* serotypes isolated from minced meat beef, mutton and pork in Addis Ababa, Ethiopia. *Revue Med. Vet.* 155 (11), 547–551.
- Ewing, W.H., 1986. Serologic identification of *Salmonella*. In: Ewing, W.H. (Ed.), *Edwards and Ewing's Identification of Enterobacteriaceae*, 4th ed. Elsevier Science Publishing Co., New York.
- FSIS, 2016. Serotypes Profile of *Salmonella* Isolates from Meat and Poultry Products January 1998 Through. (<https://www.fsis.usda.gov/wps/wcm/connect/3866026a-582d-4f0e-a8ce-851b39c7390f/Salmonella-Serotype-Annual-2014.pdf?MOD=AJPERES>). Accessed 25 May 2016).
- Garedew, L., Hagos, Z., Addis, Z., Tesfaye, R., Zegeye, B., 2015. Prevalence and antimicrobial susceptibility patterns of *Salmonella* isolates in association with hygienic status from butcher shops in Gondar town, Ethiopia. *Antimicrob. Resist. Infect. Control.* 4 (1), 21.
- Gebreyes, W.A., Thakur, S., 2005. Multidrug-resistant *Salmonella enterica* serovar Muenchen from pigs and humans and potential interserovar transfer of antimicrobial resistance. *Antimicrob. Agents Chemother.* 49 (2), 503–511.
- Grimont, P.A.D., Weill, F.-X., 2007. Antigenic Formulae of the *Salmonella* Serovars, 9th ed. WHO Collaborating Centre for Reference and Research on *Salmonella*. Institut Pasteur, Paris, France.
- Hauser, E., Tietze, E., Helmuth, R., Junker, E., Blank, K., Prager, R., Rabsch, W., Appel, B., Fruth, A., Malorny, B., 2010. Pork contaminated with *Salmonella enterica* serovar 4, [5], 12: i:-, an emerging health risk for humans. *Appl. Environ. Microbiol.* 76 (14), 4601–4610.
- Helms, M., Vastrup, P., Gerner-Smidt, P., Mølbak, K., 2002. Excess mortality associated with antimicrobial drug-resistant *Salmonella Typhimurium*. *Emerg. Infect. Dis.* 8 (5), 490–495.
- ISO, 2002. Microbiology of Food and Animal Feeding Stuff: Horizontal Method for the Detection of *Salmonella* Spp. ISO-6579, Geneva, pp. 511–525.
- Kauffmann, F., van Oye, E., 1955. A new *Salmonella* type (*Salmonella korovi*) from the Belgian Congo. *Acta Pathol. Microbiol. Scand.* 36 (4), 352.
- Mache, A., Mengistu, Y., Cowley, C., 1997. *Salmonella* serogroups identified from adult diarrhoeal out-patients in Addis Ababa, Ethiopia: antibiotic resistance and plasmid profile analysis. *East Afr. Med. J.* 74, 183–186.
- Majowicz, S.E., Musto, J., Scallan, E., Angulo, F.J., Kirk, M., O'Brien, S.J., Jones, T.F., 2010. The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin. Infect. Dis.* 50, 882–889.
- Mandilara, G., Lambiri, M., Polemis, M., Passiotou, M., Vatopoulos, A., 2013. Phenotypic and molecular characterisation of multiresistant monophasic *Salmonella Typhimurium* (1,4,[5],12:i:-) in Greece, 2006–2011. *Euro Surveill.* 18 (22), 1–8.
- Mannion, C., Fanning, J., McLernon, J., Lendrum, L., Gutierrez, M., Duggan, S., Egan, J., 2012. The role of transport, lairage and slaughter processes in the dissemination of *Salmonella* spp in pigs in Ireland. *Food Res. Int.* 45 (2), 871–879.
- Molla, B., Mesfin, A., Alemayehu, D., 2003. Multiple antimicrobial resistant *Salmonella* serotype isolated from chicken carcass and giblets in Debre-zeit and Addis Ababa, Ethiopia. *J. Health Dev.* 17, 131–149.
- Molla, B., Salah, W., Alemayehu, D., Mohammed, A., 2004. Antimicrobial resistance pattern of *Salmonella* serotypes isolated from apparently healthy slaughtered camels (*Camelus dromedarius*) in eastern Ethiopia. *Berl. Munch. Tierarztl. Wochenschr.* 117, 39–45.
- Morgan, I.R., Krautill, F.L., Craven, J.A., 1987. Effect of time in lairage on caecal and carcass *Salmonella* contamination of slaughter pigs. *Epidemiol. Infect.* 98, 323–330.
- NSLRL, 2011. National *Salmonella*, *Shigella* and *Listeria* Reference Laboratory of Ireland

- (human Health): Annual Report for 2011. (http://www.nuigalway.ie/research/salmonella_lab/downloads/nssrlr_annual_report_2011.pdf. Accessed 20 May 2016).
- Nabbut, N.H., Al-Nakhli, H.M., 1982. *Salmonella* in lymph nodes, spleens and feces of sheep and goats slaughtered in Riyadh Public Abattoir. *J. Food Protect.* 45, 1314–1317.
- Nyeleti, C., Hildebrandt, G., Kleer, J., Molla, B., 2000. Prevalence of *Salmonella* in Ethiopian cattle and minced beef. *Berl. Munch. Tierarztl. Wochenschr.* 113, 431–434.
- OzFoodNet, 2005. *Salmonella* Muenchen in Queensland, Pathogen Fact Sheet. (<https://www.health.qld.gov.au/foodsafety/documents/muenchenfs.pdf>. Accessed 25 May 2016).
- Pegram, R.G., Ropeder, P.L., Hall, M.L.M., Rowe, B., 1981. *Salmonella* in livestock and animal by-products in Ethiopia. *Trop. Anim. Health Prod.* 13, 203–207.
- Proctor, M.E., Hamacher, M., Tortorello, M.L., Archer, J.R., Davis, J.P., 2001. Multistate outbreak of *Salmonella* serovar Muenchen infections associated with alfalfa sprouts grown from seeds pretreated with calcium hypochlorite. *J. Clin. Microbiol.* 39 (10), 3461–3465.
- Quinn, P.J., Carter, M.E., Markey, B.K., Carter, G., 1999. *Clinical Veterinary Microbiology*, Mosby Yearbook. Europe Ltd., pp. 226–234.
- Roels, T.H., Frazak, P.A., Kazmierczak, J.J., Mackenzie, W.R., Proctor, M.E., Kurzynski, T.A., Davis, J.P., 1997. Incomplete sanitation of a meat grinder and ingestion of raw ground beef: contributing factors to a large outbreak of *Salmonella* Typhimurium infection. *Epidemiol. Infect.* 119 (02), 127–134.
- Samuel, J.L., O'Boyle, D.A., Mathers, W.J., Frost, A.J., 1980. Isolation of *Salmonella* from mesenteric lymph nodes of healthy cattle at slaughter. *Res. Vet. Sci.* 28 (2), 238–241.
- Shipp, C.R., Rowe, B., 1980. A mechanised microtechnique for *Salmonella* serotyping. *J. Clin. Pathol.* 33 (6), 595–597.
- Sibhat, B., Molla Zewde, B., Zerihun, A., Muckle, A., Cole, L., Boerlin, P., Wilkie, E., Perets, A., Mistry, K., Gebreyes, W.A., 2011. *Salmonella* serovars and antimicrobial resistance profiles in beef cattle, slaughterhouse personnel and slaughterhouse environment in Ethiopia. *Zoonoses Public Health* 58 (2), 102–109.
- Smeltzer, T., Thomas, R., Collins, G., 1980. The role of equipment having accidental or indirect contact with the carcass in the spread of *Salmonella* in an abattoir. *Aust. Vet. J.* 56, 14–17.
- Smerdon, W.J., Adak, G.K., O'Brien, S.J., Gillespie, I.A., Reacher, M., 2001. General outbreaks of infectious intestinal disease linked with red meat, England and Wales, 1992–1999. *Commun. Dis. Public Health* 4, 259–267.
- Switt, A.L., Soyer, Y., Warnick, L.D., Wiedmann, M., 2009. Emergence, distribution, and molecular and phenotypic characteristics of *Salmonella enterica* serotype 4,5,12:i. *Foodborne Pathog. Dis.* 6 (4), 407–415.
- Threlfall, E.J., 2002. Antimicrobial drug resistance in *Salmonella*: problems and perspectives in food- and water-borne infections. *FEMS Microbiol. Rev.* 26, 141–148.
- Thrusfield, M., 2005. *Veterinary Epidemiology*, 3rd ed. Blackwell science Ltd., London, pp. 228–246.
- Varma, J.K., Mølbak, K., Barrett, T.J., Beebe, J.L., Jones, T.F., Rabatsky-Ehr, T., Smith, K.E., Vugia, D.J., Chang, H.-G.H., Angulo, F.J., 2005. Antimicrobial-resistant nontyphoidal *Salmonella* is associated with excess bloodstream infections and hospitalizations. *J. Infect. Dis.* 191, 554–561.
- Yang, X., Wu, Q., Zhang, J., Huang, J., Guo, W., Cai, S., 2015. Prevalence and characterization of monophasic *Salmonella* serovar 1,4,[5],12:i:- of food origin in China. *PLoS One* 10 (9), e0137967. <http://dx.doi.org/10.1371/journal.pone.0137967>.
- van Duijkeren, E., Wannet, W.J.B., Houwers, D.J., Van Pelt, W., 2002. Serotype and phage type distribution of *Salmonella* strains isolated from humans, cattle, pigs, and chickens in The Netherlands from 1984 to 2001. *J. Clin. Microbiol.* 40, 3980–3985.
- Zamperini, K., Soni, V., Waltman, D., Sanchez, S., Theriault, E.C., Bray, J., Maurer, J.J., 2007. Molecular characterization reveals *Salmonella enterica* serovar 4,[5],12:i:- from poultry is a variant Typhimurium serovar. *Avian Dis.* 51 (4), 958–964.