

# Antimicrobial Resistance of *Campylobacter* Species Isolated from Broilers in Live Bird Markets in Shanghai, China

Beibei Li<sup>1,\*</sup>, Licai Ma<sup>2,\*</sup>, Yingli Li<sup>1</sup>, Haiyan Jia<sup>1</sup>, Jianchao Wei<sup>1</sup>, Donghua Shao<sup>1</sup>, Ke Liu<sup>1</sup>,  
Yuanyuan Shi<sup>1</sup>, Yafeng Qiu<sup>1</sup>, and Zhiyong Ma<sup>1</sup>

## Abstract

This study was conducted to determine the prevalence of antimicrobial resistance in *Campylobacter* spp. isolates from broilers in live bird markets (LBMs). A total of 209 *Campylobacter* spp. isolates (84 *Campylobacter jejuni*; 125 *Campylobacter coli*) were recovered from 364 broiler cecum samples collected from five LBMs in Shanghai, China. Minimum inhibitory concentrations of 13 antimicrobials were determined using agar dilution method. More than 96% of the *Campylobacter* spp. isolates were resistant to quinolones and tetracyclines. A high prevalence of macrolide resistance (erythromycin, 84.0%; azithromycin, 80.8%) was observed in *C. coli*, but not in *C. jejuni* (erythromycin, 6.0%; azithromycin, 2.4%). *C. coli* also showed significantly higher resistance than *C. jejuni* to clindamycin, gentamicin, and kanamycin. In contrast, *C. coli* isolates had lower resistance to florfenicol than the *C. jejuni* isolates. The majority of the *C. jejuni* (88.1%) and *C. coli* (97.6%) isolates exhibited multidrug resistance (MDR) to three or more classes of antimicrobials. All of the 208 ciprofloxacin-resistant *Campylobacter* spp. isolates were positive for the C257T mutation of the *gyrA* gene. In addition, the *tet(O)* gene was identified in all of the 202 doxycycline-resistant *Campylobacter* spp. isolates. Furthermore, 75.7% and 20.4% of the 103 azithromycin-resistant *Campylobacter* spp. isolates were positive for the A2075G mutation of the 23S rRNA gene and the presence of the *erm(B)* gene, respectively. Moreover, the *cat* gene was found in 14.3% (8/56) and 76.8% (73/95) of the chloramphenicol-resistant *C. jejuni* and *C. coli* isolates, respectively. To the best of our knowledge, this is the first report of the prevalence of antimicrobial resistance among *Campylobacter* spp. isolates originating from LBMs. The high prevalence of MDR *Campylobacter* spp. isolates in LBMs highlights the need to implement efficient intervention measures to control not only *Campylobacter* contamination in LBMs but also dissemination of antimicrobial resistance among *Campylobacter* spp. in poultry production.

**Keywords:** *Campylobacter*, antimicrobial resistance, broilers, live bird markets

## Introduction

*CAMPYLOBACTER* spp., ESPECIALLY *Campylobacter jejuni* and *Campylobacter coli*, are leading causes of bacterial gastroenteritis globally (Koluman and Dikici, 2013). Although usually self-limiting, severe or long-lasting *Campylobacter* spp. infections, especially of patients with compromised immune systems, often require antimicrobial therapy (McGill *et al.*, 2009). *Campylobacter* spp. are prevalent in food-producing animals, and the use of antimicrobials in food animal production has led to an increasing prevalence of antimicrobial-resistant *Campylobacter* spp. infection transmitted to humans

through the food chain, thereby presenting a major threat to public health (Mäsaar *et al.*, 2016).

The handling and consumption of contaminated poultry, particularly broilers, seem to be the most common risk factor (Damjanova *et al.*, 2011). Broilers are regarded as a main reservoir of *Campylobacter* spp., and the colonization level of *Campylobacter* spp. in broiler ceca can reach  $10^9$  CFU/g of cecal content (Stern *et al.*, 2008). *Campylobacter* contamination often occurs during the slaughtering process and consumption of meat from a contaminated carcass is a significant source of human campylobacteriosis (Melero *et al.*, 2012).

<sup>1</sup>Shanghai Veterinary Research Institute, Chinese Academy of Agricultural Science, Shanghai, China.

<sup>2</sup>East China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Shanghai, China.

\*These two authors contributed equally to this work.

In China and other Asian countries, live bird markets (LBMs) are among the most important termini of the poultry industry, as people in these countries prefer to purchase live or freshly slaughtered poultry. Birds sold in LBMs are continually introduced from different sources and caged at high densities with those collected from other areas. These conditions provide an optimal environment for amplification and persistence of infectious agents. Hence, LBMs are potential reservoirs of poultry-associated pathogens, such as various avian influenza viruses that continue to cause human infection and death (Zhou *et al.*, 2009; Cowling *et al.*, 2013; He *et al.*, 2014). In addition, unlike other consummation termini of the poultry industry, consumers at LBMs are in direct and close contact with live or freshly slaughtered poultry. Together, the conditions of LBMs likely increase the risk for human *Campylobacter* spp. infection. However, studies concerning *Campylobacter* spp. from LBMs are very limited (Mansouri-najand *et al.*, 2012). Therefore, in this study, we investigated the prevalence and antimicrobial resistance of *C. jejuni* and *C. coli* isolates collected from five LBMs in Shanghai, one of the largest cities in China.

## Materials and Methods

### Sample collection

From November 2012 to April 2013, a total of 364 cecum samples from broilers were randomly collected from five LBMs located in five geographically separated districts (Minhang, Fengxian, Pudong, Songjiang, and Yangpu) of Shanghai (Table 1). For each LBM, 71–77 samples were collected at three to four time points between November 2012 and April 2013. Each cecum sample was collected from a different chicken during evisceration and put into a stomach bag, and then rapidly transported on ice to our laboratory within 2 h.

### Isolation and identification of *Campylobacter* spp.

A loopful of fecal content from each sample was directly streaked onto a *Campylobacter* spp. Selective Agar (Base) plate (Sigma, St. Louis, MO) containing 5% fresh sterile defibrinated sheep blood and *Campylobacter* supplement III (Sigma) (Ma *et al.*, 2014). The plates were incubated under microaerobic conditions (CO<sub>2</sub>, 10%; O<sub>2</sub>, 5%; N<sub>2</sub>, 85%) at 42°C for 48 h. One presumptive colony from each plate was

subcultured and identified by PCR strategies (Keramas *et al.*, 2003) and API-Campy kits (BioMérieux, Marcy l'Etoile, France). All the identified isolates were stored at –80°C in brain-heart infusion broth with 20% (v/v) glycerol.

### Antibiotic susceptibility testing

The standard agar dilution method, as described by the Clinical Laboratory Standards Institute (CLSI, 2010), was employed to determine the susceptibility of *Campylobacter* spp. isolates to 13 antibiotic agents: nalidixic acid, ciprofloxacin, enrofloxacin, levofloxacin, erythromycin, tetracycline, doxycycline, azithromycin, clindamycin, chloramphenicol, florfenicol, gentamicin, and kanamycin. These antimicrobials, with the exception of nalidixic acid (Sigma), were all obtained from the China Institute of Veterinary Drug Control (Beijing, China). Breakpoints for each antimicrobial agent and minimum inhibitory concentration (MIC) ranges for quality control strains *C. jejuni* ATCC 33560 and *Escherichia coli* ATCC25922 are shown in Table 2. A *Campylobacter* spp. isolate simultaneously resistant to three or more classes of antimicrobials was defined as multidrug resistant.

### Detection of resistance determinants

The presence of genes involved in macrolide, tetracycline, and phenicol resistance was detected by PCR. Genomic DNA of *Campylobacter* spp. isolates was extracted using the TIANamp Bacterial DNA Kit (TIANGEN, Beijing, China), according to the manufacturer's directions, and used as templates for subsequent PCR reactions. The detection of the *tet*(O) (Qin *et al.*, 2011), *cat* (Wang and Taylor, 1990), *cfr* (Kehrenberg and Schwarz, 2004), *cmlA* & *floR* (Dai *et al.*, 2008), and *erm*(B) (Spiliopoulou *et al.*, 2004) genes was conducted by PCR analysis, as previously described. In addition, the mismatch amplification mutation assay (MAMA) PCR was used to detect the C257T mutation in the quinolone resistance-determining region (QRDR), which confers high-level quinolone resistance (Zirnstein *et al.*, 2000), and the mutations A2074C and A2075G in the 23S rRNA gene, which are associated with macrolide resistance (Alonso *et al.*, 2005). Moreover, the macrolide-resistant isolates were also screened for mutations in the L4 and L22 ribosomal protein genes (Cagliero *et al.*, 2006).

TABLE 1. THE PREVALENCE OF *CAMPYLOBACTER* SPECIES IN CECUM SAMPLES COLLECTED FROM LBMs IN SHANGHAI, CHINA

District	Nos. of cecum samples	No. of <i>Campylobacter jejuni</i> <sup>a</sup>	No. of <i>Campylobacter coli</i> <sup>a</sup>	No. of total isolates from each district <sup>a</sup>
Minhang	71	20 (28.2)	47 (66.2)	67 (94.4)
Fengxian	77	16 (20.8)	22 (28.6)	38 (49.4)
Songjiang	73	12 (16.4)	34 (46.6)	46 (63)
Putuo	71	20 (28.2)	8 (11.3)	28 (39.4)
Yangpu	72	16 (22.2)	14 (19.4)	30 (41.7)
Total (%)	364	84 (23.1)	125 (34.3)	209 (57.4)

<sup>a</sup>Numbers in parentheses indicate percentages.  
LBM, live bird market.

TABLE 2. ANTIMICROBIAL SUSCEPTIBILITY OF 84 *C. JEJUNI* AND 125 *C. COLI* STRAINS ISOLATED FROM LBMs IN SHANGHAI, CHINA

Antimicrobial agent	<i>MIC QC range</i> (μg/mL) <sup>a</sup>		<i>MIC breakpoint</i> (μg/mL) <sup>b</sup>			<i>C. jejuni</i>		<i>C. coli</i>	
	ATCC33560	ATCC25922	S	I	R	<i>MIC</i> <sub>50</sub> / <i>MIC</i> <sub>90</sub> (μg/mL)	Resistant isolates (%) <sup>c</sup>	<i>MIC</i> <sub>50</sub> / <i>MIC</i> <sub>90</sub> (μg/mL)	Resistant isolates (%) <sup>c</sup>
Nalidixic acid	4–16	—	≤16	32	≥64	256/256	84 (100)	256/256	125 (100)
Ciprofloxacin	0.06–0.5	—	≤1	2	≥4	128/256	83 (98.8)	32/128	125 (100)
Enrofloxacin	—	0.008–0.03	≤0.5	1–2	≥4	16/128	83 (98.8)	8/16	122 (97.6)
Levofloxacin	0.03–0.25	—	≤2	4	≥8	32/256	83 (98.8)	16/32	122 (97.6)
Erythromycin	1–4	—	≤8	16	≥32	2/8	5 (6.0)	256/256	105 (84.0)
Azithromycin	0.03–0.12	—	≤2	4	≥8	0.125/1	2 (2.4)	256/256	101 (80.8)
Clindamycin	0.12–0.5	—	≤2	4	≥8	2/8	14 (16.7)	64/256	106 (84.8)
Chloramphenicol	1–4	—	≤8	16	≥32	32/64	56 (66.7)	32/64	95 (76.0)
Florfenicol	0.5–2	—	≤4	8	≥16	32/64	67 (79.8)	4/8	8 (6.4)
Tetracycline	0.25–1	—	≤4	8	≥16	256/256	83 (98.8)	256/256	122 (97.6)
Doxycycline	0.25–2	—	≤2	4	≥8	64/128	81 (96.4)	64/128	121 (96.8)
Gentamicin	0.5–4	—	≤2	4	≥8	0.25/256	15 (17.9)	256/256	117 (93.6)
Kanamycin	—	1–4	≤16	32	≥64	8/256	23 (27.4)	256/256	116 (92.8)

<sup>a</sup>The QC ranges of *C. jejuni* ATCC 33560 were directly adopted from CLSI (2010). Due to the lack of QC ranges of *C. jejuni* ATCC 33560 for enrofloxacin and kanamycin, we used *E. coli* ATCC 25922 as QC strain for these two antimicrobial agents (CLSI, 2010).

<sup>b</sup>MIC breakpoints for nalidixic acid, ciprofloxacin, erythromycin, azithromycin, clindamycin, florfenicol, tetracycline, doxycycline, and gentamicin are those recommended by the CLSI (2010). Since standardized MIC breakpoints for enrofloxacin, levofloxacin, chloramphenicol, and kanamycin are not available for *Campylobacter* spp., we used the breakpoints of *Enterobacteriaceae* for these four antimicrobial agents as recommended by CLSI (2010).

<sup>c</sup>Numbers in parentheses indicate percentages.

I, intermediate; QC, quality control; R, resistant; S, susceptible.

### Statistical analysis

The resistance frequencies for each class of antimicrobial agents and multidrug resistance (MDR) rates between *C. jejuni* and *C. coli* isolates from LBMs were compared with the chi-square test using SPSS 17.0 statistical software (SPSS Inc., Chicago, IL). Differences were considered significant at  $p < 0.05$ .

## Results and Discussion

### Incidence of *Campylobacter* spp.

Overall, *Campylobacter* spp. were isolated in 209 (57.4%) of the 364 broiler cecum samples. *Campylobacter* spp. isolation rates varied from 39.4% to 94.4% across the five surveyed LBMs (Table 1). Among these samples, the prevalence of *C. jejuni* and *C. coli* was 23.1% and 34.3%, respectively, which corresponds well with previous reports on the greater proportion of *C. coli* than *C. jejuni* in broilers (Henry *et al.*, 2011; Ma *et al.*, 2014). However, the observation that *C. coli* was the predominant *Campylobacter* spp. in broilers was contrary to other reports (Chen *et al.*, 2010; Anderson *et al.*, 2012). It was consistent with the speculation that *C. coli*, instead of *C. jejuni*, is becoming the predominant *Campylobacter* spp. in chickens in China, which might be attributed to the fact that *C. coli* is intrinsically more adaptable to antimicrobial treatment, and the extensive use of antimicrobials in broiler production may have favorably selected *C. coli* over *C. jejuni* (Ma *et al.*, 2014; Wang *et al.*, 2016).

### Quinolone resistance

High prevalence of quinolone resistance was observed among the 209 *Campylobacter* spp. isolates (nalidixic acid,

100%; ciprofloxacin, 99.5%; enrofloxacin, 98.1%; levofloxacin, 98.1%) (Table 2). These results are similar to previous observations in China that almost 100% of *Campylobacter* spp. strains isolated from chickens were resistant to quinolones (Chen *et al.*, 2010; Ma *et al.*, 2014), which might be caused by the extended use of these antimicrobials in chicken production. In contrast, a very low incidence of quinolone-resistant *Campylobacter* spp. has been reported in Canada (Agunos *et al.*, 2013), the United States (Ge *et al.*, 2013), and France (Guyard-Nicodème *et al.*, 2015), which may be benefited from prudent use policy of antimicrobials in these countries.

Quinolone resistance in *Campylobacter* spp. is usually conferred by point mutations in the QRDR of DNA gyrase A (GyrA), and the C257T mutation, leading to the Thr-86-Ile substitution in the gyrase, is the most frequently detected (Hormeño *et al.*, 2016). All of the 208 *Campylobacter* spp. (*C. jejuni* = 83; *C. coli* = 125) isolates resistant to ciprofloxacin were positive for the C257T mutation in the QRDR of gyrA. However, some *Campylobacter* spp. isolates examined in this study showed a high level of fluoroquinolone resistance. For example, 21 of 84 *C. jejuni* strains exhibited ciprofloxacin MICs of ≥256 μg/mL. However, the C257T mutation is not sufficient to confer such a high level of fluoroquinolone resistance in *Campylobacter* spp.; thus other mechanisms, such as additional mutations the QRDR of gyrA and enhanced activity of the MDR efflux pump CmeABC, may be involved (Guo *et al.*, 2010; Dai *et al.*, 2015).

### Tetracycline resistance

High tetracycline (tetracycline, 98.1%; doxycycline, 96.7%) resistance rate was also present in the *Campylobacter*

spp. isolates of this study, which was consistent with other investigations performed in China (Chen *et al.*, 2010; Ma *et al.*, 2014). Tetracycline resistance in *Campylobacter* spp. is mainly conferred by the ribosomal protection protein gene *tet(O)*, which is located either on the chromosome or a transmissible plasmid (Gibree *et al.*, 2004; Abdi-Hachesoo *et al.*, 2014). In this study, all the 202 doxycycline-resistant *Campylobacter* spp. strains (81 *C. jejuni* and 121 *C. coli*) were positive for the *tet(O)* gene. Interestingly, three doxycycline-sensitive and *tet(O)*-negative *Campylobacter* spp. isolates (2 *C. jejuni* and 1 *C. coli*) exhibited high-level resistance to tetracycline (MIC,  $\geq 256 \mu\text{g/mL}$ ), indicating the possible existence of an unknown tetracycline resistance mechanism in these strains.

#### Macrolide and lincosamide resistance

The prevalence of macrolide and lincosamide resistance in *C. coli* was significantly higher than in *C. jejuni* (erythromycin, 84.0% vs. 6.0%; azithromycin, 80.8% vs. 2.4%; clindamycin, 84.8% vs. 16.7%, respectively) (Table 2) ( $p < 0.01$ ). Resistance rates of *C. coli* (84.0%) to erythromycin were lower than that reported in previous studies performed in China (Chen *et al.*, 2010; Ma *et al.*, 2014), but higher than in Australia (Obeng *et al.*, 2012) and Poland (Wieczorek *et al.*, 2013). Furthermore, the incidence of erythromycin-resistant *C. jejuni* (6.0%) was comparable to some other findings (Gblossi Bernadette *et al.*, 2012; Torralbo *et al.*, 2015), but higher than that reported in most previous studies (Obeng *et al.*, 2012; Shin *et al.*, 2012).

In *Campylobacter* spp., macrolide resistance is often mediated by point mutations in the 23S rRNA gene, especially A2074C and A2075G (Lehtopolku *et al.*, 2011). In this study, no A2074C mutation was detected among the 103 azithromycin-resistant *Campylobacter* spp. isolates (2 *C. jejuni* and 101 *C. coli*), while 75.7% (78/103) of these strains, except 25 *C. coli*, harbored the A2075G point mutation as determined by MAMA PCR. Screening of the *erm(B)* gene and mutations in the L4 and L22 genes in the 25 *C. coli* isolates without the 23S rRNA mutations revealed that 21 harbored the *erm(B)* gene, but no L4 or L22 mutations were found. The *erm(B)*-positive rate in LBM-origin *Campylobacter* spp. isolates (10.0%, 21/209) was significantly higher than that in *Campylobacter* spp. strains from other sources (3.7%, 58/1554) (Wang *et al.*, 2014). The high density and continued introduction of broilers in LBMs may provide optimal conditions for the spread and persistence of *erm(B)*-carrying *Campylobacter* spp. isolates, finally resulting in the high incidence of the *erm(B)* gene. Notably, there were four *C. coli* isolates demonstrating high-level macrolide resistance that did not contain the 23S rRNA mutations or the *erm(B)* gene. This may be attributed to the enhanced activity of the CmeABC efflux pump and/or other unknown mechanisms, thus further studies are warranted (Guo *et al.*, 2010; Qin *et al.*, 2014).

#### Phenicol resistance

Interestingly, the florfenicol resistance rate of *C. jejuni* was significantly higher compared with *C. coli* (79.8% vs. 6.4%, respectively) (Table 2) ( $p < 0.01$ ), but the *Campylobacter* spp. with the higher chloramphenicol resistance rate was *C. coli*, rather than *C. jejuni* (76.0% vs. 66.7%, respectively).

29.2% *Campylobacter* spp. strains (*C. jejuni*, 54; *C. coli*, 7) were resistant to both phenicols. Low chloramphenicol resistance rates in *Campylobacter* spp. isolates were found in investigations conducted in Spain (Perez-Boto *et al.*, 2013), Japan (Ozawa *et al.*, 2012), and Iran (Chakeri *et al.*, 2012). However, it is worth noting that the incidence of chloramphenicol resistance among *Campylobacter* spp. isolates was very high in China (Chen *et al.*, 2010) and Brazil (de Moura *et al.*, 2013).

In this study, of the 151 (*C. jejuni*, 56; *C. coli*, 95) chloramphenicol-resistant *Campylobacter* spp. isolates, 81 (*C. jejuni*, 8; *C. coli*, 73) were positive for the *cat* gene, but not *cmlA*. In addition, the *floR* and *cfr* genes, which confer florfenicol resistance, were not detected in any of the 75 (*C. jejuni*, 67; *C. coli*, 8) florfenicol-resistant *Campylobacter* spp. isolates.

#### Aminoglycoside resistance

A very high prevalence of gentamicin (93.6%) and kanamycin (92.8%) resistance was observed among the *C. coli* isolates, but the corresponding resistance rates of *C. jejuni* were only 17.9% and 27.4%, respectively (Table 2). In addition, as mentioned above, the resistance rates of the *C. coli* isolates to azithromycin and clindamycin were also significantly higher than those of the *C. jejuni* isolates. Some previous studies also reported that resistance to some antimicrobial classes (i.e.,

TABLE 3. ANTIMICROBIAL RESISTANCE PATTERNS OF *CAMPYLOBACTER* ISOLATES FROM FIVE LBMS IN SHANGHAI, CHINA

Antimicrobial resistance patterns <sup>a</sup>	No. (%) of <i>Campylobacter</i> strains	
	<i>C. jejuni</i> (n = 84) <sup>b</sup>	<i>C. coli</i> (n = 125) <sup>b</sup>
Q	1 (1.2)	0
Q-A	0	1 (0.8)
Q-T	9 (10.7)	2 (1.6)
Q-M-T	0	1 (0.8)
Q-P-A	0	1 (0.8)
<b>Q-P-T</b>	<b>41 (48.8)<sup>c</sup></b>	1 (0.8)
Q-T-A	4 (4.8)	6 (4.8)
Q-L-P-T	3 (3.6)	0
Q-L-T-A	1 (1.2)	3 (2.4)
Q-M-L-A	0	1 (0.8)
Q-M-P-T	1 (1.2)	0
Q-M-T-A	0	1 (0.8)
Q-P-T-A	12 (14.3)	6 (4.8)
Q-L-P-T-A	8 (9.5)	0
Q-M-L-P-T	1 (1.2)	1 (0.8)
Q-M-L-T-A	0	14 (11.2)
Q-M-P-T-A	2 (2.4)	0
<b>Q-M-L-P-T-A</b>	<b>1 (1.2)</b>	<b>87 (69.6)<sup>c</sup></b>
MDR (%) <sup>d</sup>	74 (88.1)	122 (97.6)

<sup>a</sup>Abbreviations of antimicrobial agents: Q, quinolones (nalidixic acid, ciprofloxacin, enrofloxacin, and levofloxacin); P, phenicols (florfenicol and chloramphenicol); M, macrolides (erythromycin and azithromycin); T, tetracyclines (tetracycline and doxycycline); L, clindamycin; A, aminoglycosides (gentamicin and kanamycin).

<sup>b</sup>n, the number of *Campylobacter* isolates.

<sup>c</sup>The bold text indicates the major resistance patterns.

<sup>d</sup>The number (%) of multidrug-resistant *Campylobacter*.

aminoglycosides, macrolides, and lincosamides) was much higher in *C. coli* than *C. jejuni* (Ma *et al.*, 2014; Torralbo *et al.*, 2015). Further research is needed to better elucidate the molecular mechanism leading to the differences in antimicrobial resistance between *C. jejuni* and *C. coli*.

### Multidrug resistance

As is shown in Table 3, all, but one, of the *Campylobacter* spp. isolates were resistant to two or more classes of antimicrobials. Overall, 196 (93.8%) of the 209 *Campylobacter* spp. isolates exhibited MDR. In addition, the MDR rate was significantly higher ( $p < 0.05$ ) among *C. coli* isolates than *C. jejuni* (97.6%, 122/125 vs. 88.1%, 74/84, respectively). The most common antimicrobial resistance pattern in *C. jejuni* (48.8%, 41/84) was the combination of quinolones, phenicols, and tetracyclines. In *C. coli*, 87 of 125 (69.6%) isolates exhibited resistance to all of the antimicrobial classes tested. The difference in dominant MDR patterns between the *Campylobacter* spp. might be due to the intrinsic ability of *C. coli* to develop resistance to these antimicrobials (Chen *et al.*, 2010). Interestingly, a similar high MDR prevalence was also observed in the surveillance studies concerning human-origin *Campylobacter* spp. in China (Zhang *et al.*, 2014; Zhou *et al.*, 2016). The extremely high rates of antimicrobial resistance in *Campylobacter* spp. isolates from human and LBM origin suggested that some of the antimicrobial classes, such as quinolones and tetracyclines, may no longer be suitable for the treatment of human *Campylobacter* infection in China.

### Conclusions

This is the first report addressing antimicrobial resistance of *Campylobacter* spp. isolated from LBMs in China. An extremely high rate of antimicrobial resistance was observed in the *Campylobacter* spp. isolates, which was likely due to the extensive use of various antimicrobials in poultry production. Furthermore, given the fact that consumers purchasing poultry from LBMs have direct contact with live or freshly slaughtered birds, the high prevalence of MDR *Campylobacter* spp. isolates in LBMs may pose a high risk factor for human *Campylobacter* spp. infection and treatment. The findings of this study highlight the need for efficient measures to control *Campylobacter* spp. contamination in LBMs, while the extensive antimicrobial resistance among *Campylobacter* spp. isolates examined in this study underlines the need for prudent use of antimicrobials in poultry production to minimize the emergence and spread of antibiotic-resistant *Campylobacter* spp. strains.

### Acknowledgment

This work was funded by the National Natural Science Foundation of China (grant nos. 31302145 and 31672606).

### Disclosure Statement

No competing financial interests exist.

### References

Abdi-Hachesoo B, Khoshbakht R, Sharifiyazdi H, Tabatabaei M, Hosseinzadeh S, Asasi K. Tetracycline resistance genes in

- Campylobacter jejuni* and *C. coli* isolated from poultry carcasses. Jundishapur J Microbiol 2014;7:e12129.
- Agunos A, Léger D, Avery BP, Parmley EJ, Deckert A, Carson CA, Dutil L. Ciprofloxacin-resistant *Campylobacter* spp. in retail chicken, western Canada. Emerg Infect Dis 2013;19:1121–1124.
- Alonso R, Mateo E, Churrua E, Martinez I, Girbau C, Fernandez-Astorga A. MAMA-PCR assay for the detection of point mutations associated with high-level erythromycin resistance in *Campylobacter jejuni* and *Campylobacter coli* strains. J Microbiol Methods 2005;63:99–103.
- Anderson J, Horn BJ, Gilpin BJ. The prevalence and genetic diversity of *Campylobacter* spp. in domestic 'Backyard' poultry in Canterbury, New Zealand. Zoonoses Public Health 2012;59:52–60.
- Cagliero C, Mouline C, Cloeckert A, Payot S. Synergy between efflux pump CmeABC and modifications in ribosomal proteins L4 and L22 in conferring macrolide resistance in *Campylobacter jejuni* and *Campylobacter coli*. Antimicrob Agents Chemother 2006; 50:3893–3896.
- Chakeri A, Foroushani MSH, Torki Z, Rahimi E, Ebadi AG. Antimicrobial resistance of *Campylobacter* species isolated from fecal samples from cats and dogs in Iran. J Pure Appl Microbiol 2012;6:1823–1827.
- Chen X, Naren G-W, Wu, C-M, Wang Y, Dai L, Xia L-N, Luo P-J, Zhang Q, Shen J-Z. Prevalence and antimicrobial resistance of *Campylobacter* isolates in broilers from China. Vet Microbiol 2010;144:133–139.
- [CLSI] Clinical and Laboratory Standards Institute. *Methods for Antimicrobial Dilution and Disk Susceptibility Testing of Infrequently Isolated or Fastidious Bacteria; Approved Guideline* (2<sup>nd</sup> ed.), Wayne, PA: CLSI, 2010.
- Cowling BJ, Jin LM, Lau EH, Liao QH, Wu P, Jiang H, Tsang TK, Zheng JD, Fang VJ, Chang ZR, Ni MY, Zhang Q, Ip DK, Yu JX, Li Y, Wang LP, Tu WX, Meng L, Wu JT, Luo HM, Li Q, Shu YL, Li ZJ, Feng ZJ, Yang WZ, Wang Y, Leung GM, Yu HJ. Comparative epidemiology of human infections with avian influenza A H7N9 and H5N1 viruses in China: A population-based study of laboratory-confirmed cases. Lancet 2013;382:129–137.
- Dai L, Lu LM, Wu CM, Li BB, Huang SY, Wang SC, Qi YH, Shen JZ. Characterization of antimicrobial resistance among *Escherichia coli* isolates from chickens in China between 2001 and 2006. FEMS Microbiol Lett 2008; 286:178–183.
- Dai L, Muraoka WT, Wu Z, Sahin O, Zhang Q. A single nucleotide change in mutY increases the emergence of antibiotic-resistant *Campylobacter jejuni* mutants J Antimicrob Chemother 2015;70:2739–2748.
- Damjanova I, Jakab M, Farkas T, Mészáros J, Galántai Z, Turcsányi I, Bistyák A, Juhász A, Pásztai J, Kiss I, Kardos G. From farm to fork follow-up of thermotolerant *Campylobacters* throughout the broiler production chain and in human cases in a Hungarian county during a ten-months period. Int J Food Microbiol 2011;150:95–102.
- de Moura HM, Silva PR, da Silva PH, Souza NR, Racanicci AM, Santana AP. Antimicrobial resistance of *Campylobacter jejuni* isolated from chicken carcasses in the Federal District, Brazil. J Food Prot 2013;76:691–693.
- Gblossi Bernadette G, Eric Essoh A, Elise Solange KN, Natalie G, Souleymane B, Lamine Sebastien N, Mireille D. Prevalence and antimicrobial resistance of thermophilic *Campylobacter* isolated from chicken in Cote d'Ivoire. Int J Microbiol 2012. DOI:10.1155/2012/150612

- Ge B, Wang F, Sjölund-Karlsson M, McDermott PF. Antimicrobial resistance in *Campylobacter*: Susceptibility testing methods and resistance trends. *J Microbiol Methods* 2013; 95:57–67.
- Gibreel A, Tracz DM, Nonaka L, Ngo TM, Connell SR, Taylor DE. Incidence of antibiotic resistance in *Campylobacter jejuni* isolated in Alberta, Canada, from 1999 to 2002, with special reference to *tet*(O)-mediated tetracycline resistance. *Antimicrob Agents Chemother* 2004;48:3442–3450.
- Guo B, Lin J, Reynolds DL, Zhang Q. Contribution of the multidrug efflux transporter CmeABC to antibiotic resistance in different *Campylobacter* species. *Foodborne Pathog Dis* 2010;7:77–83.
- Guyard-Nicodème M, Rivoal K, Houard E, Rose V, Quesne S, Mourand G, Rouxel S, Kempf I, Guillier L, Gauchard F. Prevalence and characterization of *Campylobacter jejuni* from chicken meat sold in French retail outlets. *Int J Food Microbiol* 2015;203:8–14.
- He Y, Liu PH, Tang SZ, Chen Y, Pei E, Zhao BH, Ren H, Li J, Zhu YY, Zhao HJ, Pan QC, Gu BK, Yuan ZG, Wu F. Live poultry market closure and control of avian influenza A(H7N9), Shanghai, China. *Emerg Infect Dis* 2014;20:1565–1566.
- Henry I, Reichardt J, Denis M, Cardinale E. Prevalence and risk factors for *Campylobacter* spp. in chicken broiler flocks in Reunion Island (Indian Ocean). *Prev Vet Med* 2011;100: 64–70.
- Hormeño L, Palomo G, Ugarte-Ruiz M, Porrero MC, Borge C, Vadillo S, Piriz S, Domínguez L, Campos MJ, Quesada A. Identification of the main quinolone resistance determinant in *Campylobacter jejuni* and *Campylobacter coli* by MAMA-DEG PCR. *Diagn Microbiol Infect Dis* 2016;84:236–239.
- Kehrenberg C, Schwarz S. *fexA*, a novel *Staphylococcus lentus* gene encoding resistance to florfenicol and chloramphenicol. *Antimicrob Agents Chemother* 2004;48:615–618.
- Keramas G, Bang DD, Lund M, Madsen M, Rasmussen SE, Bunkenborg H, Telleman P, Christensen CBV. Development of a sensitive DNA microarray suitable for rapid detection of *Campylobacter* spp. *Mol Cell Probes* 2003;17:187–196.
- Koluman A, Dikici A. Antimicrobial resistance of emerging foodborne pathogens: Status quo and global trends. *Crit Rev Microbiol* 2013;39:57–69.
- Lehtopolku M, Kotilainen P, Haanperä-Heikkinen M, Nakari UM, Hänninen ML, Huovinen P, Siitonen A, Eerola E, Jalava J, Hakanen AJ. Ribosomal mutations as the main cause of macrolide resistance in *Campylobacter jejuni* and *Campylobacter coli*. *Antimicrob Agents Chemother* 2011;55:5939–5941.
- Ma LC, Wang Y, Shen JZ, Zhang QJ, Wu CM. Tracking *Campylobacter* contamination along a broiler chicken production chain from the farm level to retail in China. *Int J Food Microbiol* 2014;181:77–84.
- Mäesaar M, Kramarenko T, Meremäe K, Sögel J, Lillenberg M, Häkkinen L, Ivanova M, Kovalenko K, Hörman A, Hänninen ML, Roasto M. Antimicrobial resistance profiles of *Campylobacter* spp. isolated from broiler chicken meat of Estonian, Latvian and Lithuanian Origin at Estonian retail level and from patients with severe enteric infections in Estonia. *Zoonoses Public Health* 2016;63:89–96.
- Mansouri-najand L, Saleha AA, Wai SS. Prevalence of multidrug resistance *Campylobacter jejuni* and *Campylobacter coli* in chickens slaughtered in selected markets, Malaysia. *Trop Biomed* 2012;29:231–238.
- McGill K, Kelly L, Madden RH, Moran L, Carroll C, O'Leary A, Moore JE, McNamara E, O'Mahony M, Fanning S, Whyte P. Comparison of disc diffusion and epsilometer (E-test) testing techniques to determine antimicrobial susceptibility of *Campylobacter* isolates of food and human clinical origin. *J Microbiol Methods* 2009;79:238–241.
- Melero B, Juntunen P, Hänninen ML, Jaime I, Rovira J. Tracing *Campylobacter jejuni* strains along the poultry meat production chain from farm to retail by pulsed-field gel electrophoresis, and the antimicrobial resistance of isolates. *Food Microbiol* 2012;32:124–128.
- Obeng AS, Rickard H, Sexton M, Pang Y, Peng H, Barton M. Antimicrobial susceptibilities and resistance genes in *Campylobacter* strains isolated from poultry and pigs in Australia. *J Appl Microbiol* 2012;113:294–307.
- Ozawa M, Makita K, Tamura Y, Asai T. Associations of antimicrobial use with antimicrobial resistance in *Campylobacter coli* from grow-finish pigs in Japan. *Prev Vet Med* 2012; 10:295–300.
- Perez-Boto D, Garcia-Pena FJ, Abad-Moreno JC, Echeita MA. Antimicrobial susceptibilities of *Campylobacter jejuni* and *Campylobacter coli* strains isolated from two early stages of poultry production. *Microb Drug Resist* 2013;19: 323–330.
- Qin SS, Wang Y, Zhang QJ, Deng FR, Shen ZQ, Wu CM, Wang SL, Zhang JZ, Shen JZ. Report of ribosomal RNA methylase gene *erm*(B) in multidrug resistant *Campylobacter coli*. *J Antimicrob Chemother* 2014;69:964–968.
- Qin SS, Wu CM, Wang Y, Jeon B, Shen ZQ, Zhang QJ, Shen JZ. Antimicrobial resistance in *Campylobacter coli* isolated from pigs in two provinces of China. *Int J Food Microbiol* 2011;146:94–98.
- Shin E, Oh Y, Kim M, Jung J, Lee Y. Antimicrobial resistance patterns and corresponding multilocus sequence types of the *Campylobacter jejuni* isolates from human diarrheal samples. *Microb Drug Resist* 2012;19:110–116.
- Spiliopoulou I, Petinaki E, Papandreou P, Dimitracopoulos G. *erm*(C) is the predominant genetic determinant for the expression of resistance to macrolides among methicillin-resistant *Staphylococcus aureus* clinical isolates in Greece. *J Antimicrob Chemother* 2004;53:814–817.
- Stern NJ, Eruslanov BV, Pokhilenko VD, Kovalev YN, Volodina LL, Pereygin VV, Mitsevich EV, Mitsevich IP, Borzenkov VN, Levchuk VP, Svetoch OE, Stepanshin YG, Svetoch EA. Bacteriocins reduce *Campylobacter jejuni* colonization while bacteria producing bacteriocins are ineffective. *Microb Ecol Health Dis* 2008;20:74–79.
- Torralbo A, Borge C, Garcia-Bocanegra I, Meric G, Perea A, Carbonero A. Higher resistance of *Campylobacter coli* compared to *Campylobacter jejuni* at chicken slaughterhouse. *Comp Immunol Microbiol Infect Dis* 2015;39:47–52.
- Wang Y, Dong Y, Deng F, Liu D, Yao H, Zhang Q, Shen J, Liu Z, Gao Y, Wu C, Shen Z. Species shift and multidrug resistance of *Campylobacter* from chicken and swine, China, 2008–14. *J Antimicrob Chemother* 2016;71:666–669.
- Wang Y, Taylor D E. Chloramphenicol resistance in *Campylobacter coli*: Nucleotide sequence, expression, and cloning vector construction. *Gene* 1990;94:23–28.
- Wang Y, Zhang MJ, Deng FR, Shen ZQ, Wu CM, Zhang JQ, Zhang Q, Shen JZ. Emergence of multidrug-resistant *Campylobacter* species isolates with a horizontally acquired rRNA methylase. *Antimicrob Agents Chemother* 2014;58: 5405–5412.
- Wieczorek K, Kania I, Osek J. Erythromycin-resistant *Campylobacter coli* from slaughtered animals as a potential public health risk. *Vet Med (Praha)* 2013;58:352–358.

- Zhang M, Liu X, Xu X, Gu, Y, Tao, X, Yang, X, Yan, G, Zhang, J. Molecular subtyping and antimicrobial susceptibilities of *Campylobacter coli* isolates from diarrheal patients and food-producing animals in China. *Foodborne Pathog Dis* 2014;11:610–619.
- Zhou L, Liao Q, Dong L, Huai Y, Bai T, Xiang N, Shu Y, Liu W, Wang S, Qin P, Wang M, Xing X, Lv J, Chen RY, Feng Z, Yang W, Uyeki TM, Yu H. Risk factors for human illness with avian influenza A (H5 N1) virus infection in China. *J Infect Dis* 2009;199:1726–1734.
- Zhou J, Zhang M, Yang W, Fang Y, Wang G, Hou F. A seventeen-year observation of the antimicrobial susceptibility of clinical *Campylobacter jejuni* and the molecular mechanisms of erythromycin-resistant isolates in Beijing, China. *Int J Infect Dis* 2016;42:28–33.
- Zirnstein G, Helsel L, Li Y, Swaminathan B, Besser J. Characterization of *gyrA* mutations associated with fluoroquinolone resistance in *Campylobacter coli* by DNA sequence analysis and MAMA PCR. *FEMS Microbiol Lett* 2000;190:1–7.

Address correspondence to:

Zhiyong Ma, PhD

Shanghai Veterinary Research Institute

Chinese Academy of Agricultural Science

518 Ziyue Road

Shanghai 200241

China

E-mail: zhiyongma@shvri.ac.cn