

Evaluating Dissemination Mechanisms of Antibiotic-Resistant Bacteria in Rural Environments in China by Using CTX-M-Producing *Escherichia coli* as an Indicator

Huiyun Zou,^{1,*} Beiwen Zheng,^{2,*} Mingli Sun,³ Jakob Ottoson,⁴ Yubo Li,³ Björn Berglund,^{2,5} Xiaohui Chi,¹ Xiang Ji,¹ Xuewen Li,¹ Cecilia Stålsby Lundborg,⁶ and Lennart E. Nilsson⁵

It is becoming increasingly recognized that the environment plays an important role both in the emergence and in dissemination of antibiotic-resistant bacteria (ARB). Mechanisms and factors facilitating this development are, however, not yet well understood. The high detection rate of CTX-M genes in environmental sources provides an opportunity to explore this issue. In this study, 88 CTX-M-producing *Escherichia coli* were isolated from 30 pig feces samples from 30 pig farms and 201 environmental samples. CTX-M-producing *E. coli* was detected with the following frequencies in the different types of samples: pig feces, 73%; river water, 64%; river sediment, 52%; wastewater, 31%; drinking water, 23%; outlet sediment, 21%; soil, 17%; and vegetables, 4.4%. Dissemination of CTX-M-producing *E. coli* to different environmental matrices was evaluated by analyzing the genetic relatedness of isolates from different environmental sources, and putative transmission routes through bird feces, pig feces, drinking water, river sediment, river water, and wastewater were hypothesized. Dissemination through these routes is likely facilitated by anthropogenic activities and environmental factors. Wild birds as potential vectors for dissemination of CTX-M-producing *E. coli* have the capacity to spread ARB across long distances. Regional dissemination between different environmental matrices of CTX-M-producing *E. coli* increases the exposure risk of humans and animals in the area.

Keywords: ESBL-producing *E. coli*, dissemination mechanism, multiple environmental matrices, animal

Introduction

AT PRESENT, antibiotic-resistant bacteria (ARB) constitute a great threat to human health worldwide. If the situation is left unchecked, it has been estimated that the global costs up until 2050 will be 100 trillion USD and an increase in deaths per year up to a staggering 10 million in 2050.¹ To prevent this development, and tackle the global challenge of antibiotic resistance, a “One Health” perspective, which takes into account the connections between human and animal health and the environment, is required.²

It is becoming increasingly recognized that the environment functions as a reservoir of ARB and antibiotic resistance genes (ARGs), and plays an important role for the dissemination and evolution of these environmental contaminants. ARGs can be readily transferred to humans through contaminated environmental matrices, which contribute to the dissemination and persistence of ARB in the general population.³ For example, environmental exposure to antibiotic-resistant *Escherichia coli* has been associated with fecal carriage in humans, including carriage of pathogenic *E. coli*, which further indicates the risk to humans

¹Department of Environment and Health, School of Public Health, Shandong University, Jinan, China.

²State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, China.

³Center of Disease Prevention and Control, Zhucheng City, Shandong Province, China.

⁴National Food Agency, Department of Risk and Benefit Assessment, Uppsala, Sweden.

⁵Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden.

⁶Global Health—Health Systems and Policy: Medicines, Focusing Antibiotics, Department of Public Health Sciences, Karolinska Institutet, Stockholm, Sweden.

*These two authors contributed equally to this work.

posed by dissemination of ARB between different environmental matrices.⁴

The dissemination mechanisms and risk factors pertaining to dissemination of ARB and ARGs are not well understood and need to be further elucidated to enable risk assessment and design of suitable interventions.⁵ To date, most studies focus on ARB and ARGs associated with clinical settings or animal breeding, whereas the environmental sector is usually neglected. Furthermore, environmental studies on ARB and ARGs are usually limited by focusing on a single type of environmental matrix; studies simultaneously investigating multiple environmental sources are rarely undertaken.^{6,7} Lastly, the detection rates of ARB and ARGs for resistance to last-resort antibiotics are relatively low, which increases the difficulty of drawing conclusions from the gathered data. For example, one study reported that the detection rate of total carbapenem-resistant gram-negative bacilli from a Swedish river downstream of a wastewater treatment plant was 4.5% and 2.5% in influent sewage water and environmental waters, respectively.⁸ Other studies have reported on a prevalence rate of 1.3–3.6% of the plasmid-mediated colistin resistance gene *mcr-1* among isolates of *E. coli* from environmental samples.^{9,10} Therefore, the mentioned ARB are not suitable as biomarkers for environmental dissemination of ARB and ARGs across different environmental matrices.

Extended-spectrum beta-lactamases (ESBLs) are enzymes that hydrolyze the β -lactam ring of penicillins and third-generation cephalosporins, such as cefotaxime and ceftazidime.¹¹ Since the 2000s, CTX-M genes have become the predominant ESBL type.¹² Among Enterobacteriaceae, *E. coli* is most frequently associated with production of ESBLs.¹³ *E. coli* is a commensal micro-organism and opportunistic pathogen of the gastrointestinal tract in humans and many animals, and is particularly suited to disseminate ARGs in the environment through the fecal route.¹⁴ *E. coli* is also the most prevalently isolated Enterobacteriaceae from environmental sources, and is well adapted for survival outside of animal hosts.¹⁵ CTX-M-producing *E. coli* is widely disseminated and has been reported among various environmental matrices including river water, well water, municipal wastewater, soil, and vegetables.^{16–20} Accordingly, CTX-M-producing *E. coli* may be a suitable indicator for studying the mechanisms of environmental dissemination of antibiotic resistance through fecal pollution.²¹ In this study, the prevalence and characteristics of CTX-M-producing *E. coli* among various environmental sources in a rural area of China were investigated to elucidate the dissemination of ARB and ARGs among different animal and environmental matrices.

Materials and Methods

Sampling

The study site is located in a region of rural Shandong province in eastern China. We chose to conduct the research in a rural area because household pigs live in proximity to humans and many residents work as farmers (*i.e.*, they have extended periods of contact with animals and the environment). Twelve villages that hold a number of available backyard pig farms were included in our study. In each village two to three households with backyard farms (de-

fined as harboring <50 pigs) with at least 1 pig were randomly selected. The pig farms chosen usually did not harbor other animals aside from pigs. Additional information regarding the scale of the pig farming in the sampled villages is presented in Supplementary Table S1. During July and September 2015, 30 pig feces samples and 201 environmental samples were collected (Fig. 1) from 12 villages: villages A ($n=27$), B ($n=19$), C ($n=15$), D ($n=22$), E ($n=19$), F ($n=17$), G ($n=19$), H ($n=21$), I ($n=19$), J ($n=18$), K ($n=16$), and L ($n=19$). The environmental samples collected were of wastewater ($n=36$), outlet sediment ($n=14$), river water ($n=25$), river sediment ($n=25$), drinking water ($n=43$), soil ($n=23$), vegetables ($n=23$), bird feces ($n=6$), and human waste ($n=6$). The number of samples and types of samples are summarized in Table 1. The pig feces samples were collected from 30 pig farms (a single sample per farm) and were collected by pooling fresh pig droppings on the ground. Wastewater directly discharged from the pig farms (*i.e.*, without treatment) were collected by using sterilized bottles. Samples of bird feces were only collected from village A; samples were collected twice daily for 3 days from the ground of a villager's yard, which was frequented by swallows. Human waste samples were collected from public toilets, and since some villages lacked public toilets, only six samples of human waste from six villages were collected. Waste samples of animals and humans were collected using the ESwab collection and transport system (Copan, Brescia, Italy). Water samples (from rivers, streams, and wells) were collected by gathering 1 L of water in sterile glass bottles. The sediment samples were taken by a sediment grabber and the sediment cores were sliced into 5 cm layers. At least 30 grams of river sediment and soil samples from vegetable gardens were collected in sterile plastic bottles. Edible parts of different vegetables were collected in plastic bags, including beans ($n=10$), cucumbers ($n=6$), peppers ($n=3$), celery ($n=2$), and leeks ($n=2$).

All samples were put in cool boxes with ice packs (4–8°C) upon collection, and transported to the local CDC laboratory at the end of the day. Water samples were subjected to filtration through 0.45 μ m sterile membrane filters (Millipore, Billerica). After filtration, membranes were transferred into ESwabs and stored at –80°C until cultivation. Sediments (2.0 g) were homogenized in a fivefold volume of sterile brain heart infusion broth (Oxoid, Basingstoke, UK). The vegetable samples were not washed, and the inedible or damaged parts were removed using a sterilized knife. The surface was sampled using the ESwab collection and transport system, and stored until analysis as described hereunder.

Screening of samples for ESBL-producing *E. coli*

Each sample was inoculated in brain heart infusion broth (Oxoid) and cultured at 37°C overnight. The enriched solutions were plated on ChromID ESBL agar (bioMérieux, Marcy l'Etoile, France) for 18–24 hours at 37°C to isolate potential ESBL-producing isolates. Screening was performed as previously described.²² Species confirmation of presumptive *E. coli* isolates was performed by using matrix-assisted laser desorption ionization time-of-flight mass spectrometry. ESBL production was confirmed by using the




FIG. 1. The sampling distribution of CTX-M-producing *Escherichia coli* among different environmental samples from a rural area in Shandong province, China.

double disk diffusion method, using cefotaxime, ceftazidime, each alone and in combination with clavulanic acid according to the manufacturer's protocol (Oxoid). *E. coli* ATCC 25922 was used as control strain.

Genetic characterization of CTX-M genes

DNA was extracted from the ESBL isolates using Gentra Puregene Yeast/Bact. Kit (QIAGEN, Germany), following the manufacturer's instructions. The presence of CTX-M genes was confirmed by using a multiplex PCR as previ-

ously described.²³ Four *bla*_{CTX-M} subgroups (group 1, 2, 8, and 9) were discriminated by multiplex PCR assay as previously described.²⁴ PCR products were verified by sequencing (BGI Company, Beijing, China).

Antibiotic susceptibility testing

The confirmed CTX-M-producing isolates were cultured on blood agar, incubated overnight at 37°C, and then subjected to antibiotic susceptibility testing using agar dilution with Müller-Hinton agar (Oxoid), according to the Clinical

TABLE 1. SAMPLES FROM 10 DIFFERENT SOURCES WERE COLLECTED FROM 12 VILLAGES IN SHANDONG PROVINCE, CHINA, IN 2015

Village	Human waste	Bird feces	Pig feces	Wastewater	Outlet sediment	River water	River sediment	Drinking water	Soil	Vegetables	Total
A	1 (1)	5 (6)	2 (1)	0 (4)	0 (0)	3 (3)	2 (3)	3 (5)	1 (2)	0 (2)	17 (27)
B	0 (0)	0 (0)	5 (3)	1 (4)	0 (0)	2 (3)	2 (3)	2 (3)	0 (2)	0 (1)	12 (19)
C	1 (1)	0 (0)	3 (3)	2 (3)	1 (1)	0 (1)	0 (1)	1 (3)	0 (1)	1 (1)	9 (15)
D	0 (0)	0 (0)	1 (3)	2 (4)	0 (0)	2 (3)	2 (3)	2 (5)	0 (2)	0 (2)	9 (22)
E	0 (0)	0 (0)	2 (3)	0 (3)	0 (1)	2 (2)	1 (2)	1 (4)	1 (2)	0 (2)	7 (19)
F	0 (0)	0 (0)	0 (2)	1 (3)	0 (1)	0 (2)	1 (2)	0 (3)	1 (2)	0 (2)	3 (17)
G	0 (0)	0 (0)	3 (3)	0 (3)	0 (1)	2 (2)	1 (2)	0 (4)	0 (2)	0 (2)	6 (19)
H	0 (1)	0 (0)	1 (2)	1 (2)	0 (5)	2 (2)	1 (2)	0 (4)	0 (1)	0 (2)	5 (21)
I	0 (1)	0 (0)	1 (3)	1 (3)	0 (0)	1 (2)	0 (2)	1 (4)	0 (2)	0 (2)	4 (19)
J	0 (1)	0 (0)	3 (2)	0 (2)	2 (2)	2 (2)	3 (2)	0 (3)	0 (2)	0 (2)	10 (18)
K	1 (1)	0 (0)	0 (2)	0 (2)	0 (1)	0 (2)	0 (2)	0 (2)	0 (2)	0 (2)	1 (16)
L	0 (0)	0 (0)	1 (3)	3 (3)	0 (2)	0 (1)	0 (1)	0 (3)	1 (3)	0 (3)	5 (19)
Total	3 (6)	5 (6)	22 (30)	11 (36)	3 (14)	16 (25)	13 (25)	10 (43)	4 (23)	1 (23)	88 (231)

The number of samples positive for CTX-M-producing *Escherichia coli* are denoted in the table with the total number of samples collected given in parentheses.




FIG. 2. Distribution of CTX-M gene types among CTX-M-producing *E. coli* from a rural area in Shandong province, China. **(a)** Distribution of CTX-M gene types in different matrices; **(b)** distribution of CTX-M gene types in different villages; *n* represents the number of CTX-M-producing *E. coli* in different matrices and villages, respectively. (BF, bird feces; HW, human waste; F, pig feces; WW, wastewater; OS, outlet sediment; RW, river water; RS, river sediment; DW, drinking water; S, soil; V, vegetables).

& Laboratory Standards Institute (CLSI, 2016), except for colistin, for which broth microdilution was used. The results were evaluated in accordance with the European Committee on Antimicrobial Susceptibility Testing (EUCAST) breakpoints version 8.1,^a except for tetracycline and nitrofurantoin, for which the CLSI breakpoints version 2017 was used, and florfenicol, for which the EUCAST epidemiological cutoff value was used. Multidrug resistance was defined as acquired nonsusceptibility to at least one agent in three or more antimicrobial categories.²⁵

Molecular typing and determination of putative transmission routes

For multilocus sequence typing of *E. coli* isolates, internal fragments of the seven housekeeping genes (*adhA*, *fumC*, *gyrB*, *icd*, *mdh*, *purA*, and *recA*) were amplified by PCR from DNA, as previously described,²⁶ and subsequently sequenced (BGI Company). Sequences were imported into the *E. coli* Multilocus sequence typing (MLST) database website^b to determine sequence types (STs). A phylogenetic tree based on the concatenated MLST sequences was constructed by using Molecular Evolutionary Genetics Analysis 5. Annotation of the phylogenetic tree was performed with Interactive Tree of Life.^c

Preparation of DNA for pulsed-field gel electrophoresis (PFGE) was performed as previously described.^{27,28} A dendrogram with the PFGE results was generated using BioNumerics 6.1 (Applied Maths, Sint-Martens-Latem, Belgium). According to the Unweighted Pair group Method using Arithmetic average (UPGMA) cluster analysis, isolates were defined as being of a related clone if dice similarity indices of the PFGE fingerprints were >80%. Putative transmission routes of CTX-M-producing *E. coli* were determined by identifying related clones isolated from different sources.

Results

A total of 88 CTX-M-producing *E. coli* were isolated from 231 samples (38%); CTX-M-producing *E. coli* was detected among all sample types. The detection rate was highest in pig feces (73%), followed by river water (64%), river sediment (52%), human waste (50%), wastewater (31%), drinking water (23%), outlet sediment (21%), soil (17%), and vegetables (4.4%) (Table 1). In addition, five CTX-M-producing *E. coli* were isolated from three samples of bird feces (two samples were isolated with four different CTX-M-producing *E. coli*).

Genes of the CTX-M-9 group were most frequently detected (*n*=64), and *bla*_{CTX-M-14} (*n*=35) was the predominant genotype, followed by *bla*_{CTX-M-65} (*n*=20), *bla*_{CTX-M-55} (*n*=10), *bla*_{CTX-M-15} (*n*=9), *bla*_{CTX-M-27} (*n*=9), and *bla*_{CTX-M-3} (*n*=5). *bla*_{CTX-M-14} was the most common CTX-M gene found in pig feces as well as some of the environmental sources sampled (human waste, bird feces, river water, river sediment, and vegetables), whereas *bla*_{CTX-M-65} was the most common found in drinking water, soil, and outlet sediment, and *bla*_{CTX-M-55} was the most common CTX-M gene in *E. coli* from wastewater samples. The distribution of CTX-M genes in different matrices and different villages is shown in Fig. 2. *bla*_{CTX-M-14} was found in all isolates except for those from villages F and L. *bla*_{CTX-M-65} was predominant in villages E, F, and J, whereas *bla*_{CTX-M-27} was the main genotype in villages D and L.

All CTX-M-producing *E. coli* isolates were cefotaxime resistant, and the majority (>80%) of these isolates were also resistant to tetracycline, trimethoprim-sulfamethoxazole, and florfenicol (Table 2). Fifty percent to 75% of the isolates were resistant to fosfomycin, piperacillin-tazobactam, ciprofloxacin, gentamicin, and amoxicillin-clavulanic acid; 43% was resistant to nitrofurantoin, and <25% of the isolates were resistant to colistin, ceftazidime, ceftiofur, and amikacin. Only one isolate was resistant to meropenem, whereas none of the isolates were resistant to tigecycline. All isolates were multidrug resistant.

^awww.eucast.org/

^b<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>

^c<http://itol.embl.de/>

TABLE 2. ANTIBIOTIC RESISTANCE RATES OF CTX-M-PRODUCING *E. COLI* ISOLATED FROM DIFFERENT SOURCES IN A RURAL AREA OF SHANDONG PROVINCE, CHINA, IN 2015

Antibiotic	Rate of resistance (%) by source (No. of isolates)										Total
	Human waste (3)	Bird feces (5)	Pig feces (22)	Wastewater (11)	Outlet sediment (3)	River water (16)	River sediment (13)	Drinking water (10)	Soil (4)	Vegetables (1)	
CTX	100	100	100	100	100	100	100	100	100	100	100
TE	67	80	100	100	100	88	100	100	75	100	94
SXT	100	60	86	100	67	56	92	90	75	100	82
FFC	33	60	82	91	100	81	100	60	75	100	81
FOF	33	60	68	73	100	88	69	90	75	100	75
TZP	67	60	64	91	67	81	69	70	75	100	73
CIP	67	80	77	46	33	50	92	60	75	100	66
GEN	100	80	64	46	0	50	85	30	75	100	59
AMC	67	80	55	64	33	44	62	80	50	100	59
F	0	40	36	18	33	63	46	60	50	100	43
CL	0	20	18	36	67	31	15	20	25	100	25
CAZ	33	40	9.1	18	33	19	18	50	25	0	22
FOX	0	20	9.1	0	33	38	31	30	25	0	20
AMK	0	20	0	0	0	13	0	10	0	0	4.5
MEM	0	0	0	0	0	6.3	0	0	0	0	1.1
TGC	0	0	0	0	0	0	0	0	0	0	0

CTX, cefotaxime; TE, tetracycline; SXT, trimethoprim/sulfamethoxazole; FOF, fosfomicin; FFC, florfenicol; TZP, piperacillin/tazobactam; CIP, ciprofloxacin; AMC, amoxicillin/clavulanate; GEN, gentamicin; F, nitrofurantoin; CL, collistin; CAZ, ceftazidime; FOX, ceftaxime; AMK, amikacin; MEM, meropenem; TGC, tigecycline.

A total of 57 STs were detected, including 2 new STs (Supplementary Fig. S1). The CTX-M-9 producers were distributed among 43 ST types ($n=64$), whereas the CTX-M-1-producers were distributed among 18 ST types ($n=24$). The most prevalent STs were ST10 ($n=10$), ST48 ($n=7$), ST3489 ($n=4$), ST34 ($n=3$), and ST1638 ($n=3$). ST10 was the most common ST overall, and was found in pig feces, wastewater, drinking water, river water, and river sediment. Five isolates from wild birds belonged to ST216, ST542, ST1011, ST2807, and ST4633. In addition, two isolates (both belonging to ST2614) from pig waste belonged to phylogroup B2, whereas three isolates (belonging to ST38, ST174, and ST2309, respectively) from river sediment, wastewater, and drinking water, respectively, belonged to phylogroup D. The predominant STs were distributed over several villages; ST10 isolates were found in villages D ($n=4$), L ($n=3$), H ($n=1$), I ($n=1$), and J ($n=1$); ST48 isolates were found in villages C ($n=3$), G ($n=2$), F ($n=1$), and H ($n=1$); and ST3489-isolates were distributed in villages B ($n=3$) and G ($n=1$). Isolates sharing STs were found in villages A, B, C, D, G, J, and L (Supplementary Fig. S2).

The PFGE analysis revealed a high clonal diversity among the isolates, nevertheless, genetically related isolates could be determined based on the PFGE dendrogram (Supplementary Fig. S3). Twenty-three groups (consisting of two or more isolates) of related isolates were identified, with 18 of

the groups containing isolates originating from different environmental sources (Table 3). From all the villages, isolates genetically related to isolates from other villages could be determined.

Discussion

The highest detection rate of CTX-M-producing *E. coli* among all sample types was among pig fecal samples (73%), indicating that pig feces can be a source of wider dissemination of CTX-M-producing *E. coli*. A previous study on CTX-M-producing *E. coli* isolated from fecal samples in 21 pig farms in China reported a prevalence of 43%.²⁹ The detection rate of CTX-M-producing *E. coli* in river water in Taian city, China, was 53% in a previous study.³⁰ Compared with these studies, the prevalence of CTX-M-producing *E. coli* in pig feces and river water (64%) was higher in the study area of this study. In the rural area of Laiwu, in the Shandong province, China, the prevalence of CTX-M-producing *E. coli* in well water has been reported as 23%,³¹ which was similar to the results in this study. In rural areas, well water is essential in daily life for activities such as tooth brushing, bathing, and as drinking water for humans (usually after boiling) and pigs. Such activities could potentially increase the risk of dissemination of CTX-M-producing *E. coli* from the environment to humans and animals. Notably, CTX-M-producing *E. coli* was also detected in vegetables (4.4%). The vegetable samples were collected from edible parts that can be eaten fresh without preparation, such as cucumbers and pepper. Through vegetables, humans and animals can be exposed to CTX-M-producing *E. coli* in daily life. Three isolates of CTX-M-producing *E. coli* (50%) originated from six human waste samples, and five isolates of CTX-M-producing *E. coli* originated from three samples of bird feces (two samples were isolated with four different CTX-M-producing *E. coli*). Results similar to ours have been reported in a study in Sweden in which CTX-M-producing *E. coli* could be isolated from 47% of fecal samples from wild birds.³² Birds often live close to human settings and are exposed to our bacteria and antibiotic residues through, for example, garbage dumps, river, and agricultural areas. Thus, wild birds may be an important source and facilitator of ARB transmission.

Notably, all of the CTX-M-producing *E. coli* isolated in this study ($n=88$) displayed the multidrug resistance phenotype, with rates of resistance to tetracycline, trimethoprim-sulfamethoxazole, and florfenicol being particularly high (94%, 82%, and 81% respectively). *bla*_{CTX-M} genes may also be selected by environmental exposure to non- β -lactam antibiotics, either by being physically linked on the same plasmid as other ARGs or by being carried by ARB resistant to other antibiotics.³³ A previous study in the sampling area, conducted during the same period, showed that the various environmental compartments had levels of antibiotic contamination that may pose a risk for selection of ARB and ARGs, further emphasizing the risk of dissemination of CTX-M-producing *E. coli* in the local area.³⁴

CTX-M enzymes are the most widely disseminated ESBLs worldwide, and among clinical isolates, CTX-M-14 is the most common.³⁵⁻³⁷ CTX-M-65 has been frequently reported in *E. coli* isolated from humans and food-producing animals in China.^{38,39} In this study, 20 isolates carrying the

TABLE 3. BASED ON GENETIC RELATEDNESS AS DETERMINED BY THE PULSED-FIELD GEL ELECTROPHORESIS ANALYSIS ($\geq 80\%$ HOMOGENEITY), 23 GROUPS OF GENETICALLY RELATED ISOLATES WERE DETERMINED, 18 OUT OF WHICH (VI–XXIII) CONSISTED OF TWO OR MORE ISOLATES ORIGINATING FROM DIFFERENT ENVIRONMENTAL SOURCES

Group	Isolates (n)	Isolation source	Source village
I	2	WW	E, I
II	2	RS	B, F
III	2	RW	B
IV	3	F	G, J
V	2	F	B
VI	3	RW, RS	E
VII	3	BF, DW, WW	A, D, L
VIII	2	RS, WW	A, L
IX	2	DW, F	E, I
X	2	HW, RW	A, J
XI	6	DW, F, RW, WW	A, B, D, H
XII	4	BF, RS, S, WW	A, C, I, L
XIII	4	BF, HW, RW	A, C, K
XIV	8	BF, F, RS, RW	A, B, C, I
XV	6	F, RW, WW	D, G, H, I, J
XVI	3	DW, F, RW	A, C, J
XVII	3	RS, RW, S	A, D
XVIII	2	OS, RW	C, D
IX	3	DW, F, WW	B, C, D
XX	2	DW, F	B, D
XXI	2	DW, F	E, G
XXII	2	F, RS	D, G
XXIII	2	DW, RS	B, L

BF, bird feces; DW, drinking water; F, pig feces; HW, human waste; OS, outlet sediment; RS, river sediment; RW, river water; S, soil; WW, wastewater.

*bla*_{CTX-M-65} gene were detected in samples from pig feces, wastewater, soil, drinking water, and river water, implying that *bla*_{CTX-M-65} has been disseminated in pigs as well as in various environmental matrices inside and outside pig farms. *bla*_{CTX-M-55} is not frequently reported in environmental isolates from China, nevertheless, the gene was detected in this study among isolates belonging to six different STs (ST88, ST206, ST1844, ST2526, ST2973, and ST5020), showing that *bla*_{CTX-M-55} has disseminated among multiple *E. coli* lineages in this region.

ST10 has been described from different sources worldwide, including river water, wastewater, pigs, chicken, and migratory birds.^{35,40–42} One study reported on the zoonotic potential of avian pathogenic isolates of *E. coli* belonging to the ST10 clonal complex (includes ST10 and ST48), which were associated with poultry and human cases of urinary tract infections and sepsis.⁴³ In this study, one isolate from wild birds belonged to ST1011. *E. coli* belonging to ST1011 have previously been reported both as extraintestinal pathogenic *E. coli* isolated from humans and as avian pathogenic *E. coli* isolated from birds, suggesting the zoonotic potential through wild birds of CTX-M-producing *E. coli*.⁴⁴ Isolates belonging to phylogroup B2 (ST2614) and phylogroup D (ST38, ST174, and ST2309) were also detected in this study. *E. coli* of phylogroups B2 and D can possess toxin functions and are commonly causative agents of urinary tract infections.⁴⁵ The presence of *E. coli* strains capable of causing urinary tract infections in the environment increases the exposure risk for humans.

CTX-M genes are a form of environmental pollution resulting from anthropogenic activities, either by direct exposure of the environment to ARGs and ARB through waste or through emergence through environmental exposure to antibiotics of subtherapeutic levels.⁴⁶ Thus, CTX-M-producing *E. coli* is potentially of use as an indicator to track contamination sources and dissemination of ARB and ARGs. In this study, genetic relatedness of isolates was determined by using the PFGE dendrogram (Supplementary Fig. S2), and putative dissemination routes were explored based on this phylogeny. The results showed that from all the sampled villages, isolates could be determined that were genetically related to isolates from other villages, indicating dissemination in the local geographical region (Table 3). Furthermore, 18 groups consisting of genetically related isolates of CTX-M-producing *E. coli* from different environmental sources were identified (Table 3). A large number of isolates with close relatives isolated from different environmental sources were identified, suggesting a widespread dissemination of CTX-M-producing *E. coli* between environmental matrices in the local area. Pig feces and environmental sources such as bird feces, drinking water, river sediment, river water, and wastewater had a particularly high number of isolates, which could be determined to have close relatives in other environmental matrices; from all these sources, isolates genetically related to isolates from each other of the aforementioned sources were identified (excepting that no isolates from bird feces was found that were closely related to any isolate from drinking water). These results suggest that these environmental matrices may serve as nodes through which dissemination of ARGs and ARB can occur. Although it is likely that transmission can occur in any direction from any of these environmental

matrices, the methods employed in this study cannot establish the direction of transmission. Nevertheless, it is possible to speculate as to the likely routes and mechanisms of transmission (Supplementary Fig. S4).

Anthropogenic activities might modulate the environmental resistance gene pool and promote antibiotic resistance dissemination.⁴⁶ Pig farms are often flush cleaned with water, and pig fecal material can be discharged into the environment through the run-off water from the cleaning.⁴⁷ In addition, owing to the lack of management measures for livestock manure, the animal manure was directly piled up by the river in the study area. During rain, wastewater from the manure piles can be leached directly into the river, introducing large number of bacteria carrying different types of ARGs into the river water.⁴⁸ For example, a high degree of homology was observed between isolates EC6 (river water, village A), EC8 (river water, village A), and EC41 (feces, village C). These results indicated that ARB from animal feces can spread not only to environmental matrices in the surrounding aquatic environment in the same village but also may spread through the aquatic route across villages.

Rivers can likely function as vectors for ARGs and ARB transmission, either allowing the dissemination along the river to villages along its length, or to drinking water (well water) through leaching into groundwater. Previous studies have shown that environmental factors such as river direction and aerosols can affect the dissemination of ESBL genes.^{49–52} Another study found that presence of CTX-M-producing *E. coli* in river water was found to be higher in regions with large number of animals being raised than in regions with lower number of animals being raised.⁵³ In this study, genetically homologous isolates from river water were also observed in drinking water and wastewater of different villages, suggesting the possibility of dissemination through the river to other aquatic environments in the local area. For example, EC1 (drinking water, village A), EC22 (drinking water, village B), and EC76 (river water, village H) showed high homology in the PFGE analysis, indicating that dissemination may have occurred between river water and drinking water (well water) between different villages.

ARGs and ARB can also disseminate from river water to soil. For example, CTX-M-producing *E. coli* isolated from soil shared high homology with isolates collected from aquatic environments, which indicates that irrigation may facilitate dissemination of ARB between different environmental matrices (Supplementary Fig. S5). ARGs may migrate to deeper soil layers and contaminate groundwater through vertical transport.¹⁴ Animal manure is often applied to soil as fertilizer, thus contaminating the soil with ARB present in the animal manure.⁵⁴ However, in this study, no dissemination of CTX-M-producing *E. coli* from pig feces to soil was indicated by the homology analysis. A possible explanation for this is that the soil and pig manure samples were collected during autumn, whereas the fields are generally fertilized during spring. A failure to observe homologous isolates in the two sample types could indicate either that the microbiota of the pigs had changed or that contaminating soil through animal manure only gives rise to a transient presence of the ARB. A previous study investigated ARGs in soils from areas along the Pearl River irrigated with wastewater, and concluded that wastewater was the primary source of antibiotics in the soil.⁵⁵

Wild birds have been hypothesized to be reservoirs for ARB, and facilitators of antibiotic resistance dissemination.^{35,56} In this study, CTX-M-producing *E. coli* isolated from wild bird feces were found to be related to isolates from animal waste, human waste, aquatic environments, and the soil, indicating that wild birds may be responsible for disseminating CTX-M-producing *E. coli* to these environmental matrices. Furthermore, isolates genetically related to isolates from wild birds were observed in different villages. For example, EC18, isolated from bird feces in village A, was closely related to EC36 (wastewater, village C) and EC87 (river sediment, village I). Village I is remotely located from both villages A and C, and all three villages lack a connection through river (Fig. 1). A possibility is that isolates may have spread between villages A, C, and I through birds; it had been previously shown that birds can carry CTX-M-producing *E. coli* from long distances while migrating and that an extensive dissemination between individual birds can allow ARB to persist in dense bird populations.⁵⁷ In addition, the isolate EC17 carrying *bla*_{CTX-M-14} gene from wild bird showed a significant genetic homology to seven isolates carrying *bla*_{CTX-M-14} gene from different matrices (Supplementary Fig. S3). This indicates that both plasmid and clonal expansion may be important modes of dissemination for CTX-M genes in the area.

Conclusion

In this study, CTX-M-producing *E. coli* were detected in all sampled environmental matrices, including drinking water and vegetables. The prevalence of CTX-M-producing *E. coli* was relatively higher in pig feces and river water than shown in previous studies. Through drinking water and vegetables from edible parts, humans and animals can be exposed to CTX-M-producing *E. coli* in daily life. Positive samples were distributed over several villages. Dissemination of CTX-M-producing *E. coli* to different environmental matrices was evaluated by analyzing the genetic relatedness of isolates from different environmental sources, and putative transmission routes through bird feces, pig feces, drinking water, river sediment, river water, and wastewater were hypothesized. Dissemination through these routes is likely facilitated by anthropogenic activities and environmental factors. Wild birds as potential vectors for dissemination of CTX-M-producing *E. coli* have the capacity to spread ARB across long distances. The PFGE analysis indicated that both plasmid and clonal expansion may be important modes of dissemination for CTX-M genes in the area. Regional dissemination between different environmental matrices of CTX-M-producing *E. coli* increases the exposure risk of humans and animals in the area.

Acknowledgments

The authors thank the IMPACT Consortium for its efforts in this project. This project was supported by the National Natural Science Foundation of China (Grant Nos. 41771499, 81361138021), the Fundamental Research Funds of Shandong University (Grant No. 2018JC102), the Swedish Research Council, Public Health Agency of Sweden (Grant No. D0879801), and the Swedish Research Council Formas (Grant No. 2016-00640).

Disclosure Statement

No competing financial interests exist.

Supplementary Material

Supplementary Figure S1
Supplementary Figure S2
Supplementary Figure S3
Supplementary Figure S4
Supplementary Table S1

References

- O'Neill, J. 2014. Antimicrobial Resistance: Tackling a Crisis for the Health and Wealth of Nations. Review on Antimicrobial Resistance 2014, UK. <http://amr-review.org>.
- Robinson, T.P., D.P. Bu, J. Carrique-Mas, E.M. Fevre, M. Gilbert, D. Grace, S.I. Hay, J. Jiwakanon, M. Kakkar, S. Kariuki, R. Laxminarayan, J. Lubroth, U. Magnusson, P. Thi Ngoc, T.P. Van Boeckel, and M.E.J. Woolhouse. 2016. Antibiotic resistance is the quintessential One Health issue. *Trans. R. Soc. Trop. Med. Hyg.* 110:377–380.
- Huijbers, P.M.C., E.A.M. Graat, A. van Hoek, and C. Veenman. 2016. Transmission dynamics of extended-spectrum beta-lactamase and AmpC beta-lactamase-producing *Escherichia coli* in a broiler flock without antibiotic use. *Prev. Vet. Med.* 131:12–19.
- Leonard, A.F.C., L. Zhang, A.J. Balfour, R. Garside, P.M. Hawkey, A.K. Murray, O.C. Ukoumunne, and W.H. Gaze. 2018. Exposure to and colonisation by antibiotic-resistant *E. coli* in UK coastal water users: environmental surveillance, exposure assessment, and epidemiological study (Beach Bum Survey). *Environ. Int.* 114:326–333.
- Larsson, D.G.J., A. Andremon, J. Bengtsson-Palme, K.K. Brandt, A.M. de Roda Husman, P. Fagerstedt, J. Fick, C.F. Flach, W.H. Gaze, M. Kuroda, K. Kvint, R. Laxminarayan, C.M. Manaia, K.M. Nielsen, L. Plant, M.C. Ploy, C. Segovia, P. Simonet, K. Smalla, J. Snape, E. Topp, A.J. van Hengel, D.W. Verner-Jeffreys, M.P.J. Virta, E.M. Wellington, and A.S. Wernersson. 2018. Critical knowledge gaps and research needs related to the environmental dimensions of antibiotic resistance. *Environ. Int.* 117:132–138.
- Daoud, Z., J. Farah, E.S. Sokhn, K.E. Kfoury, E. Dahdouh, K. Masri, C. Afif, R.M. Abdel-Massih, and G.M. Matar. 2018. Multidrug-resistant Enterobacteriaceae in lebanese hospital wastewater: Implication in the one health concept. *Microb. Drug. Resist.* 24:166–174.
- Purohit, M.R., S. Chandran, H. Shah, V. Diwan, A.J. Tamhankaret, and C. Stålsby Lundborg. 2017. Antibiotic Resistance in an Indian Rural Community: A “One-Health” Observational Study on Commensal Coliform from Humans, Animals, and Water. *Int. J. Environ. Res. Public Health* 14:386.
- Khan, F.A., B. Hellmark, R. Ehricht, B. Söderquist, and J. Jass. 2018. Related carbapenemase-producing Klebsiella, isolates detected in both a hospital and associated aquatic environment in Sweden. *Eur. J. Clin. Microbiol. Infect. Dis.* 37:2241–2251.
- Runchaon, C., K.E. Raven, S. Reuter, T. Kallonen, S. Paksanont, J. Thammachote, S. Anun, B. Blane, J. Parkhill, S.J. Peacock, and N. Chantratita. 2017. Whole genome sequencing of ESBL-producing *Escherichia coli* isolated from patients, farm waste and canals in Thailand. *Genome. Med.* 9:81.

10. Zurfluh, K., K. Kieffer, L. Poirel, P. Nordmann, and R. Stephan. 2016. Features of the mcr-1 cassette related to colistin resistance. *Antimicrob. Agents Chemother.* 60: 6438–6439.
11. Knothe, H., P. Shah, V. Krcmery, M. Antal, and S. Mitsuhashi. 1983. Transferable resistance to cefotaxime, ceftiofur, cefamandole and cefuroxime in clinical isolates of *Klebsiella pneumoniae* and *Serratia marcescens*. *Infection* 11:315–317.
12. Woerther, P.L., C. Burdet, E. Chachaty, and A. Andremont. 2013. Trends in human fecal carriage of extended-spectrum β -lactamases in the community: toward the globalization of ctx-m. *Clin. Microbiol. Rev.* 26:744–758.
13. Willemsen, I., S. Oomeet, C. Verhulst, A. Pettersson, K. Verduin, and J. Kluytmans. 2015. Trends in extended spectrum beta-lactamase (ESBL) producing Enterobacteriaceae and ESBL genes in a Dutch teaching hospital, measured in 5 yearly point prevalence surveys (2010–2014). *PLoS One*. 10:e0141765.
14. Huang, X., C. Liu, K. Li, F. Liu, D. Liao, L. Liu, G. Zhu, and J. Liao. 2013. Occurrence and distribution of veterinary antibiotics and tetracycline resistance genes in farmland soils around swine feedlots in Fujian Province, China. *Environ. Sci. Pollut. Res. Int.* 20:9066–9074.
15. Haque, A., A. Yoshizumi, T. Sagaet, Y. Ishii, and Y. Tateda. 2014. ESBL-producing Enterobacteriaceae in environmental water in Dhaka, Bangladesh. *J. Infect. Chemother.* 20:735–737.
16. Yamashita, N., Y. Katakawa, and H. Tanaka. 2017. Occurrence of antimicrobial resistance bacteria in the Yodo River basin, Japan and determination of beta-lactamases producing bacteria. *Ecotoxicol. Environ. Saf.* 143:38–45.
17. Sun, P., Z. Bi, M. Nilsson, B. Zheng, B. Berglund, C. Stålsby Lundborg, S. Börjesson, X. Li, B. Chen, H. Yin, and L.E. Nilsson. 2017. Occurrence of bla_{KPC-2}, bla_{CTX-M}, and mcr-1 in enterobacteriaceae from well water in rural China. *Antimicrob. Agents Chemother.* 61:02569–16.
18. Lamba, M., and S.Z. Ahammad. 2017. Sewage treatment effluents in Delhi: a key contributor of beta-lactam resistant bacteria and genes to the environment. *Chemosphere*. 188: 249–256.
19. Brower, C.H., S. Mandal, S. Hayer, M. Sran, A. Zehra, S.J. Patel, R. Kaur, L. Chatterjee, S. Mishra, B.R. Das, P. Singh, R. Singh, J.P.S. Gill, and R. Laxminarayan. 2017. The prevalence of extended-spectrum beta-lactamase-producing multidrug-resistant *Escherichia coli* in poultry chickens and variation according to farming practices in Punjab, India. *Environ. Health. Perspect.* 125:077015.
20. Kraemer, J.G., J. Pires, M. Kueffer, E. Semaani, A. Endimiani, M. Hilty, and A. Oppliger. 2017. Prevalence of extended-spectrum beta-lactamase-producing Enterobacteriaceae and Methicillin-Resistant *Staphylococcus aureus* in pig farms in Switzerland. *Sci. Total Environ.* 603–604: 401–405.
21. Guenther, S., C. Ewers, and L.H. Wieler. 2011. Extended-spectrum beta-lactamases producing *E. coli* in wildlife, yet another form of environmental pollution? *Front. Microbiol.* 2:246.
22. Sun, Q., Y. Wang, A. Hulth, Y. Xiao, L.E. Nilsson, X. Li, Z. Bi, Y. Liu, H. Yin, Y. Luo, M. Nilsson, C. Sun, Y. Zhu, B. Zheng, B. Chen, P. Sun, L. Ding, X. Xia, J. Ottosson, S. Löfmark, O.J. Dyar, S. Börjesson, and C. Stålsby Lundborg. 2018. Study protocol for One Health data collections, analyses and intervention of the Sino-Swedish integrated multisectoral partnership for antibiotic resistance containment (IMPACT). *BMJ. Open*. 8:e017832.
23. Monstein, H.J., A. Östholm-Balkhed, M.V. Nilsson, M. Nilsson, K. Dornbusch, and L.E. Nilsson. 2010. Multiplex PCR amplification assay for the detection of bla_{SHV}, bla_{TEM} and bla_{CTX-M} genes in Enterobacteriaceae. *APMIS*. 115:1400–1408.
24. Xu, L., V. Ensor, S. Gossain, K. Nye, and P. Hawkey. 2005. Rapid and simple detection of bla_{CTX-M} genes by multiplex PCR assay. *J. Med. Microbiol.* 54:1183.
25. Magiorakos, A.P., A. Srinivasan, R.B. Carey, Y. Carmeli, M.E. Falagas, and C.G. Giske, S. Harbarth, J.F. Hindler, G. Kahlmeter, B. Olsson-Liljequist, D.L. Paterson, L.B. Rice, J. Stelling, M.J. Struelens, A. Vatopoulos, J.T. Weber, and D.L. Monnet. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin. Microbiol. Infect.* 18:268–281.
26. Wirth, T., D. Falush, R. Lan, F. Colles, P. Mensa, L.H. Wieler, H. Karch, P.R. Reeves, M.C. Maiden, H. Ochman, and M. Achtman. 2006. Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol. Microbiol.* 60: 1136–1151.
27. Versalovic, J., T. Koeuth, and J.R. Lupski. 1991. Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic. Acids. Res.* 19:6823–6831.
28. Zheng, B., J. Zhang, J. Ji, Y. Fang, P. Shen, C. Ying, J. Lv, Y. Xiao, and L. Li. 2015. Emergence of Raoultella ornithinolytica coproducing IMP-4 and KPC-2 carbapenemases in China. *Antimicrob. Agents Chemother.* 59:7086–7089.
29. Xu, G., W. An, H. Wang, and X. Zhang. 2015. Prevalence and characteristics of extended-spectrum beta-lactamase genes in *Escherichia coli* isolated from piglets with post-weaning diarrhea in Heilongjiang province, China. *Front. Microbiol.* 6:1103.
30. Li, S., W. Song, Y. Zhou, Y. Tang, Y. Gao, and Z. Miao. 2015. Spread of extended-spectrum beta-lactamase-producing *Escherichia coli* from a swine farm to the receiving river. *Environ. Sci. Pollut. Res. Int.* 22:13033–13037.
31. Zhang, H., Y. Gao, and W. Chang. 2016. Comparison of extended-spectrum beta-lactamase-producing *Escherichia coli* isolates from drinking well water and pit latrine wastewater in a rural area of China. *Biomed. Res. Int.* 2016: 4343564.
32. Hessman, J., C. Atterby, B. Olsen, and J.D. Jarhult. 2018. High prevalence and temporal variation of extended Spectrum beta-lactamase-producing bacteria in Urban Swedish Mallards. *Microb. Drug Resist.* 24:822–829.
33. Zurfluh, K., H. Hachler, N. Nuesch-Inderbinen, and R. Stephan. 2013. Characteristics of extended-spectrum beta-lactamase- and carbapenemase-producing Enterobacteriaceae isolates from rivers and lakes in Switzerland. *Appl. Environ. Microb.* 79:3021–3026.
34. Hanna, N., P. Sun, Q. Sun, X. Li, X. Yang, X. Ji, H. Zou, J. Ottosson, L.E. Nilsson, B. Berglund, O. Dyar, A.J. Tamhankar, and C. Stålsby Lundborg. 2018. Presence of antibiotic residues in various environmental compartments of Shandong province in eastern China: its potential for resistance development and ecological and human risk. *Environ. Int.* 114:131–142.
35. Mohsin, M., S. Raza, K. Schaeffer, N. Roschanski, F. Sarwar, T. Semmler, P. Schierack, and S. Guenther. 2017.

- High prevalence of CTX-M-15-type ESBL-producing *E. coli* from migratory avian species in Pakistan. *Front. Microbiol.* 8:2476.
36. Rao, L., L. Lv, Z. Zeng, S. Chen, D. He, X. Chen, C. Wu, Y. Wang, T. Yang, P. Wu, Y. Liu, and L.H. Liu. 2015. Increasing prevalence of extended-spectrum cephalosporin-resistant *Escherichia coli* in food animals and the diversity of CTX-M genotypes during 2003-2012. *Vet. Microbiol.* 172:534–541.
 37. Bevan, E.R., A.M. Jones, and P.M. Hawkey. 2017. Global epidemiology of CTX-M beta-lactamases: temporal and geographical shifts in genotype. *J. Antimicrob. Chemother.* 72:2145–2155.
 38. Wang, J., Z.L. Zeng, X.Y. Huang, Z.B. Ma, Z.W. Guo, L.C. Lv, Y.B. Xia, L. Zeng, Q.H. Song, and J.H. Liu. 2018. Evolution and comparative genomics of F33:A:B plasmids carrying blaCTX-M-55 or blaCTX-M-65 in *Escherichia coli* and *Klebsiella pneumoniae* isolated from animals, food products, and humans in China. *Mosphere.* 3:e00137–18.
 39. Yang, X., W. Liu, Y. Liu, J. Wang, L. Lv, X. Chen, D. He, T. Yang, J. Hou, Y. Tan, L. Xing, Z. Zeng, and J.H. Liu. 2014. F33: A- B-, IncHI2/ST3, and IncI1/ST71 plasmids drive the dissemination of fosA3 and blaCTX-M-55/-14/-65 in *Escherichia coli* from chickens in China. *Front. Microbiol.* 5:688.
 40. Jorgensen, S.B., A.V. Soraas, L.S. Arnesen, T.M. Leegaard, A. Sundsfjord, and P.A. Jenum. 2017. A comparison of extended spectrum beta-lactamase producing *Escherichia coli* from clinical, recreational water and wastewater samples associated in time and location. *PLoS One.* 12: e0186576.
 41. Moremi, N., E.V. Manda, L. Falgenhauer, H. Ghosh, C. Imirzalioglu, M. Matee, T. Chakraborty, and S.E. Mshana. 2016. Predominance of CTX-M-15 among ESBL producers from environment and fish gut from the shores of lake victoria in Mwanza, Tanzania. *Front. Microbiol.* 7:1862.
 42. Nirupama, K.R., O.R. V.K., B.S. Pruthvishree, D.K. Sinha, M.S. Murugan, N. Krishnaswamy, and B.R. Singh. 2018. Molecular characterisation of bla_{oxa}-48 carbapenemase-, extended-spectrum β -lactamase- and shiga toxin-producing *Escherichia coli* isolated from farm piglets in India. *J. Glob. Antimicrob. Resist.* 13:201–205.
 43. Giufre, M., C. Graziani, M. Accogli, I. Luzzi, L. Busani, and M. Cerquetti. 2012. *Escherichia coli* of human and avian origin: detection of clonal groups associated with fluoroquinolone and multidrug resistance in Italy. *J. Antimicrob. Chemother.* 67:860–867.
 44. Maluta, R.P., C.M. Logue, M.R. Casas, T. Meng, E.A. Guastalli, T.C. Rojas, A.C. Montelli, T. Sadatsune, M. de Carvalho Ramos, L.K. Nolan, and W.D. da Silveira. 2014. Overlapped sequence types (STs) and serogroups of avian pathogenic (APEC) and human extra-intestinal. *PLoS One.* 9:e105016.
 45. Farajzadah, A., S. Aslani, Hosseini, S.A., M. Amin, H. Goodarzi, N. Jomehzadeh, R. Ranjbar, M. Moradzadeh, S. Azarpira, M. Moradi, and M. Hashemzadeh. 2018. Phylo-typing based virulence associated genes and drug susceptibility of Uropathogenic *Escherichia coli* in urinary tract infection patients. *J. Microbiol. Biotechnol.* [Epub ahead of print]; DOI: 10.4014/jmb.1801.01031.
 46. Tacão, M., A. Correia, and I. Henriques. 2012. Resistance to broad-spectrum antibiotics in aquatic systems: anthropogenic activities modulate the dissemination of bla(ctx-m)-like genes. *Appl. Environ. Microbiol.* 78:4134–4140.
 47. Zhang, Q.Q., G.G. Ying, C.G. Pan, Y.S. Liu, and J.L. Zhao. 2015. Comprehensive evaluation of antibiotics emission and fate in the river basins of China: source analysis, multimedia modeling, and linkage to bacterial resistance. *Environ. Sci. Technol.* 49:6772–6782.
 48. Amos, G.C., P.M. Hawkey, W.H. Gaze, and E.M. Wellington. 2014. Waste water effluent contributes to the dissemination of CTX-M-15 in the natural environment. *J. Antimicrob. Chemother.* 69:1785–1791.
 49. Song, Q., D. Zhang, H. Gao, and J. Wu. 2018. Salmonella species' persistence and their high level of antimicrobial resistance in flooded man-made rivers in China. *Microb. Drug. Resist.* 00:1–8.
 50. Dohmen, W., H. Schmitt, M. Bonten, and D. Heederik. 2017. Air exposure as a possible route for ESBL in pig farmers. *Environ. Res.* 155:359–364.
 51. von Salviati, C., H. Laube, B. Guerra, U. Roesler, and A. Friese. 2015. Emission of ESBL/AmpC-producing *Escherichia coli* from pig fattening farms to surrounding areas. *Vet. Microbiol.* 175:77–84.
 52. Schmithausen, R.M., S.V. Schulze-Geisthoevel, F. Stemmer, M. El-Jade, M. Reif, S. Hack, A. Meilaender, G. Montabauer, R. Fimmers, M. Parcina, A. Hoerauf, M. Exner, B. Petersen, G. Bierbaum, and I. Bekeredjian-Ding. 2015. Analysis of transmission of MRSA and ESBL-E among pigs and farm personnel. *PLoS One.* 10:e0138173.
 53. Chen, P., C. Hung, P. Huang, J. Chen, I. Huang, W. Chen, Y. Chiou, W. Hung, J. Wang, and M. Cheng. 2016. Characteristics of CTX-M extended-spectrum beta-lactamase producing *Escherichia coli* strains isolated from multiple rivers in southern taiwan. *Appl. Environ. Microbiol.* 82: 1889–1897.
 54. Cheng, W., J. Li, Y. Wu, L. Xu, C. Su, Y. Qian, Y.G. Zhu, and H. Chen. 2016. Behavior of antibiotics and antibiotic resistance genes in eco-agricultural system: a case study. *J. Hazard. Mater.* 304:18–25.
 55. Pan, M., and L.M. Chu. 2017. Occurrence of antibiotics and antibiotic resistance genes in soils from wastewater irrigation areas in the Pearl River Delta region, southern China. *Sci. Total Environ.* 624:145.
 56. Atterby, C., S. Borjesson, S. Ny, J.D. Jarhult, S. Byfors, and J. Bonnedahl. 2017. ESBL-producing *Escherichia coli* in Swedish gulls—a case of environmental pollution from humans? *PLoS. One.* 12:e0190380.
 57. Sandegren, L., J. Stedt, U. Lustig, J. Bonnedahl, D.I. Andersson, and J.D. Jarhult. 2018. Long-term carriage and rapid transmission of extended spectrum beta-lactamase-producing *E. coli* within a flock of Mallards in the absence of antibiotic selection. *Environ. Microbiol. Rep.* 10:576–582.

Address correspondence to:

Xuwen Li, PhD

Department of Environment and Health

School of Public Health

Shandong University

Jinan 250012

China

E-mail: lxw@sdu.edu.cn