

Research Note

Characterization of Extended-Spectrum β -Lactamase–Producing *Escherichia coli* Isolates from Pigs and Farm Workers

HONGNA ZHANG,^{1*} ZHENZHEN ZHAI,^{2†} QING LI,^{1†} LINGHONG LIU,^{1†} SHUYUAN GUO,^{1†} QIMENG LI,¹
LINGLING YANG,¹ CHAOQUN YE,¹ WEISHAN CHANG,^{1*} AND JING ZHAI^{3*}

¹College of Animal Science and Technology, Shandong Agricultural University, Tai'an 271000, People's Republic of China; ²Digestive Disease Institute, Central Hospital of Tai'an City, Tai'an 271000, People's Republic of China; and ³College of Basic Medicine, Taishan Medical University, Tai'an 271000, People's Republic of China

MS 16-093: Received 2 March 2016/Accepted 27 May 2016

ABSTRACT

Food-producing animals can serve as reservoirs for extended-spectrum β -lactamase (ESBL)–producing *Escherichia coli*. The present study aimed to characterize and compare ESBL-carrying *E. coli* isolates from both pigs and farm workers. Rectal swabs were obtained from 60 pigs on four pig-fattening farms (15 samples per farm), and rectal swabs were taken from 40 farm workers on these farms (10 samples per farm). ESBL-carrying *E. coli* isolates from the workers and pigs were characterized by ESBL genotype, antibiotic susceptibility, enterobacterial repetitive intergenic consensus type, and multilocus sequence type. ESBL-producing *E. coli* was detected in 34 (56.7%) of 60 pigs, and 20.0% (8 of 40) of the farm workers were positive for ESBL-producing *E. coli*. More importantly, ESBL-producing *E. coli* isolates with the same β -lactamase genes, antibiotic resistance profiles, enterobacterial repetitive intergenic consensus types, and multilocus sequence types were detected in both pigs and workers on the same pig farm. These findings were suggestive for transfer of ESBL-producing *E. coli* between animals and humans.

Key words: *Escherichia coli*; Extended-spectrum β -lactamase; Farm workers; Pigs; Transfer

Extended-spectrum cephalosporins are frequently used to treat bacterial infections and are listed as important antimicrobials by the World Health Organization (6). However, reports about extended-spectrum β -lactamase (ESBL)–producing bacteria isolated from both humans and animals have become increasingly frequent (21). ESBLs are mainly plasmid-encoded enzymes that can hydrolyze the third- and fourth-generation cephalosporins and can be produced by a variety of bacteria, such as *Escherichia coli* and *Klebsiella pneumoniae*, the most commonly found ESBL-producing bacteria (4, 10).

In recent years, ESBL-producing *E. coli* have frequently been isolated from food-producing animals across the world (3, 7, 8, 11, 16, 22, 30). In China, an increasing number of ESBL-producing *E. coli* isolates have been recovered from food-producing animals since 2005 (16, 22, 27, 30). Therefore, food animals have been regarded as a potential source for ESBL-producing *E. coli* and have attracted considerable attention (23).

Of note, possible transmission of ESBL-producing *E. coli* from animal meat products to humans has been suggested (13, 15, 20, 25, 26). ESBL-producing *E. coli* can also be transmitted from live animals to humans (or vice

versa). Direct contact with live animals may be a potential risk factor for human ESBL carriage (8). The present study was therefore designed to characterize and compare ESBL-producing *E. coli* isolates from both pigs and pig caretakers.

MATERIALS AND METHODS

Collection of rectal swab samples. During May to July 2015, on four pig-fattening farms (A, B, C, and D) in Shandong Province, People's Republic of China, rectal swab samples from pigs and farm workers were obtained and immediately transported into our laboratory for analysis within 6 h. Briefly, on each farm, rectal swabs from 15 pigs were randomly obtained by the farm veterinarians with sterile swabs, and rectal swab samples from 10 farm workers were collected by themselves using sterile swabs according to instructions by trained veterinarians. All participants gave written informed consent. The study protocol was approved by the Bio-medical Research Ethics Committee of Shandong Agricultural University (no. 2015-5-16).

Isolation and confirmation of ESBL-producing *E. coli*. All rectal swab samples were spread on MacConkey agar plates with added ceftriaxone (1 mg/liter) and were incubated at 37°C for 24 h. One presumptive colony with typical *E. coli* morphology was selected from each rectal swab sample and was identified using the API 20E system (bioMérieux, Marcy-l'Étoile, France). Suspected ESBL-producing *E. coli* isolates were confirmed by double-disk synergy test using both cefotaxime and ceftazidime in the presence or absence of clavulanic acid as recommended by the Clinical and Laboratory Standards Institute (5).

* Authors for correspondence. Tel and Fax: +00865388248213; E-mail: zhang186982@126.com (H.Z.), sdauwschang@sina.com (W.C.). Tel and Fax: +008605386221301; E-mail: tsmczhaijing@sina.com (J.Z.).

† These authors contributed equally to this work.

TABLE 1. *Primer pairs used for PCR amplification in this study*

Primer pair	Target gene or genetic element	Sequence (5'–3')	Amplicon size (bp)	Annealing temp (°C)
CTX-M group 1 F	CTX-M group 1	TTA GGA ART GTG CCG CTG YA	415	57
CTX-M group 1 R		CGA TAT CGT TGG TGG TRC CAT		
CTX-M group 2 F	CTX-M group 2	CGT TAA CGG CAC GAT GAC	552	57
CTX-M group 2 R		CGA TAT CGT TGG TGG TRC CAT		
CTX-M group 9 F	CTX-M group 9	TCA AGC CTG CCG ATC TGG T	205	57
CTX-M group 9 R		TGA TTC TCG CCG CTG AAG		
TEM F	TEM	CAT TTC CGT GTC GCC CTT ATT C	840	56
TEM R		CGT TCA TCC ATA GTT GCC TGA C		
SHV F	SHV	AGC CGC TTG AGC AAA TTA AAC	1,051	56
SHV R		ATC CCG CAG ATA AAT CAC CAC		
ERIC F	ERIC	GTAAGCTCCTGGGGATTAC		50
ERIC R		AAGTAAGTGACTGGGGTGAGCG		

Antimicrobial susceptibility testing. The disk diffusion method was used according to Clinical and Laboratory Standards Institute guidelines (5) to carry out antimicrobial susceptibility testing. A panel of 13 antimicrobial drugs was tested in this study, including ampicillin, cephalothin, cefotaxime, cefoxitin, imipenem, chloramphenicol, gentamicin, neomycin, streptomycin, tetracycline, ciprofloxacin, nalidixic acid, and trimethoprim-sulfamethoxazole. In addition, the MICs of four selected antimicrobials (cefotaxime, ceftazidime, cefepime, and imipenem), which are important for the scientific community to know, were tested using Etest strips (AB Biodisk), and the results were interpreted according to the Clinical and Laboratory Standards Institute guidelines (5). *E. coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853 were used as quality control strains.

Detection of β -lactamase gene. ESBL-producing *E. coli* isolates were analyzed for the presence of genes encoding TEM, SHV, and CTX-M β -lactamases by PCR as described previously (Table 1) (1, 18, 30). The amplification conditions included an initial denaturation at 94°C for 5 min, 35 cycles of 94°C for 1 min, 56 or 57°C (depending on the primers) for 1 min, and 72°C for 1 min, with a final extension at 72°C for 7 min. The PCR products were sequenced using an ABI 3730 automated sequencer (BaseClear, Leiden, The Netherlands). Forward and reverse sequences were aligned with reference sequences in the BLAST database (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) to confirm the β -lactamase type (http://www.ncbi.nlm.nih.gov/projects/pathogens/submit_beta_lactamase).

ERIC PCR. The genetic similarity of ESBL-producing *E. coli* isolates from both pigs and farm workers was analyzed using enterobacterial repetitive intergenic consensus (ERIC)-based primers for PCR, as previously described (Table 1) (17, 19). The PCR protocol included an initial denaturation (3 min at 94°C), 30 denaturation cycles (1 min at 94°C), annealing (1 min at 50°C), chain extension (3 min at 72°C), and a final extension (5 min at 72°C). The PCR products were separated in 2% agarose gels by electrophoresis. NTSYSpc software (version 2.10) was used to produce a dendrogram, and clustering analysis was performed by the unweighted pair group method with arithmetic averages according to the Dice similarity index (17).

MLST. Forty-two ESBL-producing *E. coli* isolates recovered from both pigs and farm workers were subjected to multilocus sequence type (MLST) analysis (8). Briefly, MLST was carried out

using seven conserved housekeeping genes (*adk*, *fumC*, *gyrB*, *icd*, *mdh*, *purA*, and *recA*) (<http://mlst.warwick.ac.uk/mlst/>).

RESULTS

ESBL-producing *E. coli* isolates from pigs and farm workers. A total of 42 ESBL-producing *E. coli* isolates were obtained in this study. Thirty-four ESBL-producing *E. coli* isolates (34 of 60, 56.7%) were obtained from four pig farms, including 10 isolates from farm A (10 of 15, 66.7%), 10 from farm B (10 of 15, 66.7%), 9 from farm C (9 of 15, 60.0%), and 5 from farm D (5 of 15, 33.3%). Eight ESBL-producing *E. coli* isolates were isolated from 40 farm workers (8 of 40, 20.0%), including 3 workers from farm A (3 of 10, 30.0%), 2 from B (2 of 10, 20.0%), 2 from C (2 of 10, 20.0%), and 1 from D (1 of 10, 10.0%) (Fig. 1).

Antimicrobial susceptibility. All 42 ESBL-producing *E. coli* isolates from both pigs and farm workers were resistant to ampicillin, cephalothin, and cefotaxime. Fortunately, these isolates were all susceptible to imipenem and cefoxitin. Among the 34 ESBL-producing *E. coli* isolates from pigs, most of the isolates showed resistance to gentamicin (30 of 34, 88.2%), tetracycline (30 of 34, 88.2%), streptomycin (24 of 34, 70.6%), and trimethoprim-sulfamethoxazole (24 of 34, 70.6%); moderate resistance rates were observed for nalidixic acid (20 of 34, 58.8%), chloramphenicol (18 of 34, 52.9%), and neomycin (14 of 34, 41.2%); and relatively low resistance rates were observed for ciprofloxacin (7 of 34, 20.6%). Among the eight ESBL-producing *E. coli* isolates from farm workers, high resistance rates were observed for tetracycline (6 of 8, 75.0%) and nalidixic acid (6 of 8, 75.0%); moderate resistance rates were observed for gentamicin (4 of 8, 50.0%), neomycin (4 of 8, 50.0%), streptomycin (4 of 8, 50.0%), and trimethoprim-sulfamethoxazole (4 of 8, 50.0%); and relatively low resistance rates were observed for chloramphenicol (3 of 8, 37.5%) and ciprofloxacin (3 of 8, 37.5%) (Fig. 1). The MIC ranges for cefotaxime, ceftazidime, cefepime, and imipenem were 4 to ≥ 16 μ g/ml, ≤ 0.5 to 32 μ g/ml, 0.5 to 16 μ g/ml, and 0.19 to 0.25 μ g/ml, respectively.




FIGURE 1. Characteristics of ESBL-producing *E. coli* isolated in this study, including origins, antimicrobial resistances, β -lactamase genes, MLST, and ERIC-PCR fingerprints. A-, B-, C-, and D-P, ESBL-producing *E. coli* isolates from pigs on farms A to D; A-, B-, C-, and D-H, ESBL-producing *E. coli* isolates from farm workers on farms A to D; black, resistant; white, susceptible; AMP, ampicillin; CF, cephalothin; CTX, cefotaxime; FOX, ceftiofur; GEN, gentamicin; N, neomycin; STR, streptomycin; TET, tetracycline; CIP, ciprofloxacin; NA, nalidixic acid; SXT, trimethoprim-sulfamethoxazole; C, chloramphenicol; IMP, imipenem.

Characterization of β -lactamase genes. All 42 ESBL-producing *E. coli* isolates from both pigs and workers carried *bla*_{CTX-M} genes. However, no *bla*_{SHV} genes were detected in this study. Among the 34 ESBL-producing *E. coli* isolates from pigs, the dominant *bla*_{CTX-M} type was *bla*_{CTX-M-55} (11 of 34, 32.4%), followed by *bla*_{CTX-M-14} (10 of 34, 29.4%), *bla*_{CTX-M-15} (7 of 34, 20.6%), and *bla*_{CTX-M-65} (5 of 34, 14.7%). Of note, 25 ESBL-producing *E. coli* isolates (73.5%) from pigs contained both *bla*_{CTX-M} and *bla*_{TEM-1}. Among the eight ESBL-producing *E. coli* isolates from farm workers, the dominant *bla*_{CTX-M} type was *bla*_{CTX-M-14} (6 of 8, 75.0%), followed by *bla*_{CTX-M-55} (2 of 8, 25.0%). Additionally, six of the eight ESBL-producing *E. coli* isolates (75.0%) from farm workers contained both *bla*_{CTX-M} and *bla*_{TEM-1} (Fig. 1).

ERIC and MLST. Thirteen different MLSTs were detected in isolates from both farm workers and pigs. Twenty-two of 42 ESBL-producing *E. coli* isolates belonged

to sequence type 10 (ST-10). More importantly, four sets of ESBL-producing *E. coli* isolates from both farm workers and pigs with the same β -lactamase genes and antibiotic resistance profiles were identical in their MLSTs and ERIC types. Specifically, at pig farm A, two farm worker isolates and one pig isolate were identified as *E. coli* ST-10; at pig farm B, one farm worker isolate and one pig isolate were identified as *E. coli* ST-453; at pig farm C, one farm worker isolate and one pig isolate were identified as *E. coli* ST-10; and at pig farm D, one farm worker isolate and one pig isolate were identified as *E. coli* ST-4056 (Fig. 1).

DISCUSSION

In this study, ESBL-producing *E. coli* isolates with the same β -lactamase genes, antibiotic resistance profiles, ERIC types, and MLSTs were detected in pigs and farm workers on the same farms, indicating possible transfer of ESBL-producing *E. coli* between pigs and farm workers. Results from other studies are also suggestive for transmission of

ESBL-producing *E. coli* between animals and humans (8, 11). Additionally, the ERIC PCR results revealed genetic heterogeneity of ESBL-producing *E. coli* isolates among the different pig farms. However, a few clusters with high similarity ($\geq 90\%$) composed of isolates from different pig farms were also detected. The isolates within these clusters were subjected to MLST analysis, which confirmed strain relatedness in some but not all cases. Therefore, ERIC PCR in combination with MLST can be considered as an appropriate molecular method to analyze genetic relatedness among ESBL-producing *E. coli* isolates.

All 42 isolates from both pigs and farm workers in our study were resistant to three or more of the antibiotics tested, which defines them as multidrug resistant. Of note, the 42 isolates were all resistant to ampicillin, cephalothin, and cefotaxime. This result may be mainly associated with the fact that ampicillin, cephalothin, and cefotaxime have been frequently used in local human and animal clinical practices. Fortunately, these isolates were all susceptible to imipenem and ceftiofloxacin.

Food-producing animals have been recognized as important carriers of ESBL-producing *E. coli* (2). A connection between ESBL-producing *E. coli* in food production animals, chicken meat, and humans has been suggested (14), and several studies revealed the presence of community-associated ESBL-producing *E. coli* infection among healthy humans (9, 28). Thus, the high rate of isolation of ESBL-producing *E. coli* from pigs (56.7%) found in this study is of great concern. Of note, extended-spectrum cephalosporins, such as cephalothin and cefotaxime, have frequently been used on these pig farms, which may contribute to the high rate of isolation of ESBL-producing *E. coli*.

All 34 ESBL-producing *E. coli* isolates from pigs harbored *bla* genes, and *bla*_{CTX-M} was the dominant ESBL gene, which is similar to findings reported previously in China (22, 29). Among the 34 ESBL-producing *E. coli* isolates recovered from pigs, a high diversity of CTX-M variants was observed. *bla*_{CTX-M-55} (32.4%) and *bla*_{CTX-M-14} (29.4%) were the most common genes detected that confer resistance to cephalosporins, in agreement with several reports from China (22, 29). Additionally, *bla*_{CTX-M-15} and *bla*_{CTX-M-65} were detected in ESBL-producing *E. coli* isolates from pigs.

In the current study, the isolation rate for ESBL-producing *E. coli* in farm workers was 20.0%, higher than the reported isolation rate of 7.0% in rural inhabitants in Shandong Province, China (28). This result suggested that frequent contact with animals carrying ESBL-producing *E. coli* may be a risk factor for human carriage of ESBL. Among the eight ESBL-producing *E. coli* isolates from farm workers, the dominant *bla*_{CTX-M} type was *bla*_{CTX-M-14} (75.0%), followed by *bla*_{CTX-M-55} (25.0%). *bla*_{CTX-M-14} is distributed in both human and animal ESBL-producing *E. coli* isolates worldwide. In addition, several studies have shown that *bla*_{CTX-M-55} in human ESBL-producing *E. coli* isolates has become the second most common CTX-M enzyme in China (12, 24).

In summary, food animals can become reservoirs for ESBL-producing bacteria. Farm workers seem to have a

higher risk for carrying ESBLs than the general population due to the fact that they have more chances to contact animals carrying ESBL-producing bacteria.

ACKNOWLEDGMENT

We acknowledge the financial support from the Special Grant of Innovation Team of Shandong Province (SDAIT-13-011-11).

REFERENCES

1. Batchelor, M., K. Hopkins, E. J. Threlfall, F. A. Clifton-Hadley, A. D. Stallwood, R. H. Davies, and E. Liebana. 2005. *bla*_(CTX-M) genes in clinical *Salmonella* isolates recovered from humans in England and Wales from 1992 to 2003. *Antimicrob. Agents Chemother.* 49:1319–1322.
2. Carattoli, A. 2008. Animal reservoirs for extended spectrum beta-lactamase producers. *Clin. Microbiol. Infect.* 14:117–123.
3. Carmen, D., H. Nils-Olaf, K. Annelene, M. Alexander, D. Kathleen, and K. Axel. 2015. Occurrence of ESBL-producing *Escherichia coli* in livestock and farm workers in Mecklenburg-Western Pomerania, Germany. *PLOS One* 10:e0143326.
4. Chong, Y., Y. Ito, and T. Kamimura. 2011. Genetic evolution and clinical impact in extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*. *Infect. Genet. Evol.* 11:1499–1504.
5. Clinical and Laboratory Standards Institute. 2012. Performance standards for antimicrobial susceptibility testing: 22nd informational supplement. CLSI document M100-S22. Clinical and Laboratory Standards Institute, Wayne, PA.
6. Collignon, P., J. H. Powers, T. M. Chiller, A. Aidara-Kane, and F. M. Aarestrup. 2009. World Health Organization ranking of antimicrobials according to their importance in human medicine: a critical step for developing risk management strategies for the use of antimicrobials in food production animals. *Clin. Infect. Dis.* 49:132–141.
7. Dierikx, C., J. Van Der Goot, T. Fabri, A. van Essen-Zandbergen, H. Smith, and D. Mevius. 2013. Extended-spectrum- β -lactamase- and AmpC- β -lactamase-producing *Escherichia coli* in Dutch broilers and broiler farmers. *J. Antimicrob. Chemother.* 68:60–67.
8. Dohmen, W., M. J. Bonten, M. E. Bos, S. van Marm, J. Scharringa, J. A. Wagenaar, and D. J. Heederik. 2015. Carriage of extended-spectrum β -lactamases in pig farmers is associated with occurrence in pigs. *Clin. Microbiol. Infect.* 21:917–923.
9. Doi, Y., Y. S. Park, J. I. Rivera, J. M. Adams-Haduch, A. Hingwe, E. M. Sordillo, J. S. Lewis II, W. J. Howard, L. E. Johnson, B. Polsky, J. H. Jorgensen, S. S. Richter, K. A. Shutt, and D. L. Paterson. 2013. Community-associated extended-spectrum beta-lactamase-producing *Escherichia coli* infection in the United States. *Clin. Infect. Dis.* 56:641–648.
10. Falagas, M. E., and D. E. Karageorgopoulos. 2009. Extended-spectrum beta-lactamase-producing organisms. *J. Hosp. Infect.* 73:345–354.
11. Hammerum, A. M., J. Larsen, V. D. Andersen, C. H. Lester, T. S. Skovgaard Skytte, F. Hansen, S. S. Olsen, H. Mordhorst, R. L. Skov, F. M. Aarestrup, and Y. Agersø. 2014. Characterization of extended-spectrum β -lactamase (ESBL)-producing *Escherichia coli* obtained from Danish pigs, pig farmers and their families from farms with high or no consumption of third- or fourth-generation cephalosporins. *J. Antimicrob. Chemother.* 69:2650–2657.
12. Hu, Y. Y., J. C. Cai, H. W. Zhou, D. Chi, X. F. Zhang, W. L. Chen, R. Zhang, and G. X. Chen. 2013. Molecular typing of CTX-M-producing *Escherichia coli* isolates from environmental water, swine feces, specimens from healthy humans, and human patients. *Appl. Environ. Microbiol.* 79:5988–5996.
13. Jouini, A., L. Vinué, K. B. Slama, Y. Sáenz, N. Klibi, S. Hammami, A. Boudabous, and C. Torres. 2007. Characterization of CTX-M and SHV extended-spectrum β -lactamases and associated resistance genes in *Escherichia coli* strains of food samples in Tunisia. *J. Antimicrob. Chemother.* 60:1137–1141.
14. Kluytmans, J. A., I. T. Overdeest, I. Willemsen, M. F. Kluytmans-van den Bergh, K. van der Zwaluw, M. Heck, M. Rijnsburger, C. M.

- Vandenbroucke-Grauls, P. H. Savelkoul, B. D. Johnston, D. Gordon, and J. R. Johnson. 2013. Extended-spectrum beta-lactamase-producing *Escherichia coli* from retail chicken meat and humans: comparison of strains, plasmids, resistance genes, and virulence factors. *Clin. Infect. Dis.* 56:478–487.
15. Kola, A., C. Kohler, Y. Pfeifer, F. Schwab, K. Kühn, K. Schulz, V. Balau, K. Breitbach, A. Bast, W. Witte, P. Gastmeier, and I. Steinmetz. 2012. High prevalence of extended-spectrum beta-lactamase-producing Enterobacteriaceae in organic and conventional retail chicken meat, Germany. *J. Antimicrob. Chemother.* 67:2631–2634.
16. Li, L., Z. G. Jiang, L. N. Xia, J. Z. Shen, L. Dai, Y. Wang, S. Y. Huang, and C. M. Wu. 2010. Characterization of antimicrobial resistance and molecular determinants of beta-lactamase in *Escherichia coli* isolated from chickens in China during 1970–2007. *Vet. Microbiol.* 144:505–510.
17. Li, S., W. Song, Y. Zhou, Y. Tang, Y. Gao, and Z. Miao. 2015. Spread of extended-spectrum beta-lactamase-producing *Escherichia coli* from a swine farm to the receiving river. *Environ. Sci. Pollut. Res. Int.* 22:13033–13037.
18. Monstein, H. J., A. Ostholm-Balkhed, M. V. Nilsson, M. Nilsson, K. Dornbusch, and L. E. Nilsson. 2007. Multiplex PCR amplification assay for the detection of *bla*_{SHV}, *bla*_{TEM} and *bla*_{CTX-M} genes in Enterobacteriaceae. *APMIS* 115:1400–1408.
19. Moraru, R., A. M. Pourcher, A. Jadas-Hecart, I. Kempf, C. Ziebal, M. Kervarrec, P. Y. Comunal, M. Mares, and P. Dabert. 2012. Changes in concentrations of fluoroquinolones and of ciprofloxacin-resistant Enterobacteriaceae in chicken feces and manure stored in a heap. *J. Environ. Qual.* 41:754–763.
20. Overdevest, I., I. Willemsen, M. Rijnsburger, A. Eustace, L. Xu, P. Hawkey, M. Heck, P. Savelkoul, C. Vandenbroucke-Grauls, K. van der Zwaluw, X. Huijsdens, and J. Kluytmans. 2011. Extended-spectrum beta-lactamase genes of *Escherichia coli* in chicken meat and humans, the Netherlands. *Emerg. Infect. Dis.* 7:1216–1222.
21. Pitout, J. D., and K. B. Laupland. 2008. Extended-spectrum beta-lactamase-producing Enterobacteriaceae: an emerging public-health concern. *Lancet Infect. Dis.* 8:159–166.
22. Rao, L., L. Lv, Z. Zeng, S. Chen, D. He, X. Chen, C. Wu, Y. Wang, T. Yang, P. Wu, Y. Liu, and J. H. Liu. 2014. Increasing prevalence of extended-spectrum cephalosporin-resistant *Escherichia coli* in food animals and the diversity of CTX-M genotypes during 2003–2012. *Vet. Microbiol.* 172:534–541.
23. Smet, A., A. Martel, D. Persoons, J. Dewulf, M. Heyndrickx, L. Herman, F. Haesebrouck, and P. Butaye. 2010. Broad-spectrum beta-lactamases among Enterobacteriaceae of animal origin: molecular aspects, mobility and impact on public health. *FEMS Microbiol. Rev.* 34:295–316.
24. Wang, X., X. Wang, and S. Zhao. 2013. Detection and genotyping of extended-spectrum beta-lactamases in *Escherichia coli* and *Klebsiella pneumoniae* isolated from a hospital. *Chin. J. Infect. Control* 12:339–343.
25. Xi, M., Q. Wu, X. Wang, B. Yang, X. Xia, and D. Li. 2015. Characterization of extended-spectrum beta-lactamase-producing *Escherichia coli* strains isolated from retail foods in Shaanxi Province, China. *J. Food Prot.* 78:1018–1023.
26. Yu, T., X. Jiang, K. Fu, B. Liu, D. Xu, S. Ji, and L. Zhou. 2013. Detection of extended-spectrum β -lactamase and plasmid-mediated quinolone resistance determinants in *Escherichia coli* isolates from retail meat in China. *J. Food Prot.* 76:2040–2044.
27. Yuan, L., J. H. Liu, G. Z. Hu, Y. S. Pan, Z. M. Liu, J. Mo, and Y. J. Wei. 2009. Molecular characterization of extended-spectrum beta-lactamase-producing *Escherichia coli* isolates from chickens in Henan Province, China. *J. Med. Microbiol.* 58:1449–1453.
28. Zhang, H., Y. Zhou, S. Guo, and W. Chang. 2015. High prevalence and risk factors of fecal carriage of CTX-M type extended-spectrum beta-lactamase-producing Enterobacteriaceae from healthy rural residents of Taian, China. *Front. Microbiol.* 6:239.
29. Zhao, W. H., and Z. Q. Hu. 2013. Epidemiology and genetics of CTX-M extended-spectrum beta-lactamases in Gram-negative bacteria. *Crit. Rev. Microbiol.* 39:79–101.
30. Zheng, H., Z. Zeng, S. Chen, Y. Liu, Q. Yao, Y. Deng, X. Chen, L. Lv, C. Zhuo, Z. Chen, and J. H. Liu. 2012. Prevalence and characterisation of CTX-M beta-lactamases amongst *Escherichia coli* isolates from healthy food animals in China. *Int. J. Antimicrob. Agents* 39:305–310.