

# Estimation by direct maximization of the likelihood

---

## 3.1 Introduction

We saw in Equation (2.12) that the likelihood of an HMM is given by

$$L_T = \Pr(\mathbf{X}^{(T)} = \mathbf{x}^{(T)}) = \boldsymbol{\delta} \mathbf{P}(x_1) \boldsymbol{\Gamma} \mathbf{P}(x_2) \cdots \boldsymbol{\Gamma} \mathbf{P}(x_T) \mathbf{1}',$$

where  $\boldsymbol{\delta}$  is the initial distribution (that of  $C_1$ ) and  $\mathbf{P}(x)$  the  $m \times m$  diagonal matrix with  $i$ th diagonal element the state-dependent probability or density  $p_i(x)$ . In principle we can therefore compute  $L_T = \boldsymbol{\alpha}_T \mathbf{1}'$  recursively via

$$\boldsymbol{\alpha}_1 = \boldsymbol{\delta} \mathbf{P}(x_1)$$

and

$$\boldsymbol{\alpha}_t = \boldsymbol{\alpha}_{t-1} \boldsymbol{\Gamma} \mathbf{P}(x_t) \quad \text{for } t = 2, 3, \dots, T.$$

If the Markov chain is assumed to be stationary (in which case  $\boldsymbol{\delta} = \boldsymbol{\delta} \boldsymbol{\Gamma}$ ), we can choose to use instead

$$\boldsymbol{\alpha}_0 = \boldsymbol{\delta}$$

and

$$\boldsymbol{\alpha}_t = \boldsymbol{\alpha}_{t-1} \boldsymbol{\Gamma} \mathbf{P}(x_t) \quad \text{for } t = 1, 2, \dots, T.$$

We shall first consider the stationary case.

The number of operations involved is of order  $Tm^2$ , making the evaluation of the likelihood quite feasible even for large  $T$ . Parameter estimation can therefore be performed by numerical maximization of the likelihood with respect to the parameters.

But there are several problems that need to be addressed when the likelihood is computed in this way and maximized numerically in order to estimate parameters. The main problems are numerical underflow, constraints on the parameters, and multiple local maxima in the likelihood function. In this chapter we first discuss how to overcome these problems, in order to arrive at a general strategy for computing MLEs which is easy to implement. Then we discuss the estimation of standard errors for parameters. We defer to the next chapter the EM algorithm, which necessitates some discussion of the forward and backward probabilities.

### 3.2 Scaling the likelihood computation

In the case of discrete state-dependent distributions, the elements of  $\alpha_t$ , being made up of products of probabilities, become progressively smaller as  $t$  increases, and are eventually rounded to zero. In fact, with probability 1 the likelihood approaches 0 or  $\infty$  exponentially fast; see Leroux and Puterman (1992). The problem is therefore not confined to the discrete case and underflow; overflow may occur in a continuous case. The remedy is, however, the same for over- and underflow, and we confine our attention to underflow.

Since the likelihood is a product of matrices, not of scalars, it is not possible to circumvent numerical underflow simply by computing the log of the likelihood as the sum of logs of its factors. In this respect the computation of the likelihood of an independent mixture model is simpler than that of an HMM.

To solve the problem, Durbin *et al.* (1998, p. 78) suggest (*inter alia*) a method of computation that relies on the following approximation. Suppose we wish to compute  $\log(p+q)$ , where  $p > q$ . Write  $\log(p+q)$  as

$$\log p + \log(1 + q/p) = \log p + \log(1 + \exp(\tilde{q} - \tilde{p})),$$

where  $\tilde{p} = \log p$  and  $\tilde{q} = \log q$ . The function  $\log(1 + e^x)$  is then approximated by interpolation from a table of its values; apparently quite a small table will give a reasonable degree of accuracy.

We prefer to compute the logarithm of  $L_T$  using a strategy of scaling the vector of forward probabilities  $\alpha_t$ . Define, for  $t = 0, 1, \dots, T$ , the vector

$$\phi_t = \alpha_t / w_t,$$

where  $w_t = \sum_i \alpha_t(i) = \alpha_t \mathbf{1}'$ .

First we note certain immediate consequences of the definitions of  $\phi_t$  and  $w_t$ :

$$\begin{aligned} w_0 = \alpha_0 \mathbf{1}' &= \delta \mathbf{1}' = 1; \\ \phi_0 &= \delta; \\ w_t \phi_t &= w_{t-1} \phi_{t-1} \mathbf{B}_t; \\ L_T = \alpha_T \mathbf{1}' &= w_T (\phi_T \mathbf{1}') = w_T. \end{aligned} \tag{3.1}$$

Hence  $L_T = w_T = \prod_{t=1}^T (w_t / w_{t-1})$ . From (3.1) it follows that

$$w_t = w_{t-1} (\phi_{t-1} \mathbf{B}_t \mathbf{1}'),$$

and so we conclude that

$$\log L_T = \sum_{t=1}^T \log(w_t / w_{t-1}) = \sum_{t=1}^T \log(\phi_{t-1} \mathbf{B}_t \mathbf{1}').$$

The computation of the log-likelihood is summarized below in the form

of an algorithm. Note that  $\mathbf{\Gamma}$  and  $\mathbf{P}(x_t)$  are  $m \times m$  matrices,  $\mathbf{v}$  and  $\phi_t$  are vectors of length  $m$ ,  $u$  is a scalar, and  $l$  is the scalar in which the log-likelihood is accumulated.

```

set  $\phi_0 \leftarrow \delta$  and  $l \leftarrow 0$ 
for  $t = 1, 2, \dots, T$ 
     $\mathbf{v} \leftarrow \phi_{t-1} \mathbf{\Gamma} \mathbf{P}(x_t)$ 
     $u \leftarrow \mathbf{v} \mathbf{1}'$ 
     $l \leftarrow l + \log u$ 
     $\phi_t \leftarrow \mathbf{v}/u$ 
return  $l$ 

```

The required log-likelihood,  $\log L_T$ , is then given by the final value of  $l$ . This procedure will avoid underflow in many cases. Clearly, variations of the technique are possible: for instance, the scale factor  $w_t$  could be chosen instead to be the largest element of the vector being scaled, or the mean of its elements (as opposed to the sum). See A.1.3 (in Appendix A) for an implementation of the above algorithm.

The algorithm is easily modified to compute the log-likelihood without assuming stationarity of the Markov chain. If  $\delta$  is the initial distribution, replace the first two lines above by

```

set  $w_1 \leftarrow \delta \mathbf{P}(x_1) \mathbf{1}'$ ,  $\phi_1 \leftarrow \delta \mathbf{P}(x_1)/w_1$  and  $l \leftarrow \log w_1$ 
for  $t = 2, 3, \dots, T$ 

```

Of course, if the initial distribution happens to be the stationary distribution, the more general algorithm still applies.

### 3.3 Maximization of the likelihood subject to constraints

#### 3.3.1 Reparametrization to avoid constraints

The elements of  $\mathbf{\Gamma}$  and those of  $\boldsymbol{\lambda}$ , the vector of state-dependent means in a Poisson–HMM, do not range over the whole of  $\mathbb{R}$ , the set of all real numbers. Neither therefore should any sensible estimates of the parameters. In particular, the row sums of  $\mathbf{\Gamma}$ , and any estimate thereof, should equal one. Thus when maximizing the likelihood we have a constrained optimization problem to solve, not an unconstrained one.

Special-purpose software, e.g. NPSOL (Gill *et al.*, 1986) or the corresponding NAG routine E04UCF, can be used to maximize a function of several variables which are subject to constraints. The advice of Gill, Murray and Wright (1981, p. 267) is that it is ‘rarely appropriate to alter linearly constrained problems’. However — depending on the implementation and the nature of the data — constrained optimization can be slow. For example, the constrained optimizer `constrOptim` available in **R** is acknowledged to be slow if the optimum lies on the boundary of the parameter space. We shall focus on the use of the unconstrained op-

imizer `nlm`. Exercise 3 and A.4 explore the use of `constrOptim`, which can minimize a function subject to linear inequality constraints.

In general, there are two groups of constraints: those that apply to the parameters of the state-dependent distributions and those that apply to the parameters of the Markov chain. The first group of constraints depends on which state-dependent distribution(s) are chosen; e.g. the ‘success probability’ of a binomial distribution lies between 0 and 1.

In the case of a Poisson–HMM the relevant constraints are:

- the means  $\lambda_i$  of the state-dependent distributions must be nonnegative, for  $i = 1, \dots, m$ ;
- the rows of the transition probability matrix  $\mathbf{\Gamma}$  must add to 1, and all the parameters  $\gamma_{ij}$  must be nonnegative.

Here the constraints can be circumvented by making certain transformations. The transformation of the parameters  $\lambda_i$  is relatively easy. Define  $\eta_i = \log \lambda_i$ , for  $i = 1, \dots, m$ . Then  $\eta_i \in \mathbb{R}$ . After we have maximized the likelihood with respect to the unconstrained parameters, the constrained parameter estimates can be obtained by transforming back:  $\hat{\lambda}_i = \exp \hat{\eta}_i$ .

The reparametrization of the matrix  $\mathbf{\Gamma}$  requires more work, but can be accomplished quite elegantly. Note that  $\mathbf{\Gamma}$  has  $m^2$  entries but only  $m(m-1)$  free parameters, as there are  $m$  row sum constraints

$$\gamma_{i1} + \gamma_{i2} + \dots + \gamma_{im} = 1 \quad (i = 1, \dots, m).$$

We shall show one possible transformation between the  $m^2$  constrained probabilities  $\gamma_{ij}$  and  $m(m-1)$  unconstrained real numbers  $\tau_{ij}, i \neq j$ .

For the sake of readability we show the case  $m = 3$ . We begin by defining the matrix

$$\mathbf{T} = \begin{pmatrix} - & \tau_{12} & \tau_{13} \\ \tau_{21} & - & \tau_{23} \\ \tau_{31} & \tau_{32} & - \end{pmatrix}, \text{ a matrix with } m(m-1) \text{ entries } \tau_{ij} \in \mathbb{R}.$$

Now let  $g : \mathbb{R} \rightarrow \mathbb{R}^+$  be a strictly increasing function, e.g.

$$g(x) = e^x \quad \text{or} \quad g(x) = \begin{cases} e^x & x \leq 0 \\ x + 1 & x \geq 0. \end{cases}$$

Define

$$\varrho_{ij} = \begin{cases} g(\tau_{ij}) & \text{for } i \neq j \\ 1 & \text{for } i = j. \end{cases}$$

We then set  $\gamma_{ij} = \varrho_{ij} / \sum_{k=1}^3 \varrho_{ik}$  (for  $i, j = 1, 2, 3$ ) and  $\mathbf{\Gamma} = (\gamma_{ij})$ . It is left to the reader as an exercise to verify that the resulting matrix  $\mathbf{\Gamma}$  satisfies the constraints of a transition probability matrix. We shall refer to the parameters  $\eta_i$  and  $\tau_{ij}$  as **working parameters**, and to the parameters  $\lambda_i$  and  $\gamma_{ij}$  as **natural parameters**.

Using the above transformations of  $\mathbf{\Gamma}$  and  $\boldsymbol{\lambda}$ , we can perform the calculation of the likelihood-maximizing parameters in two steps.

1. Maximize  $L_T$  with respect to the working parameters  $\mathbf{T} = \{\tau_{ij}\}$  and  $\boldsymbol{\eta} = (\eta_1, \dots, \eta_m)$ . These are all unconstrained.
2. Transform the estimates of the working parameters to estimates of the natural parameters:

$$\hat{\mathbf{T}} \rightarrow \hat{\mathbf{\Gamma}}, \quad \hat{\boldsymbol{\eta}} \rightarrow \hat{\boldsymbol{\lambda}}.$$

See A.1.1 and A.1.2 for  $\mathbf{R}$  functions that transform natural parameters to working and vice versa.

As an illustration we consider the first row of  $\mathbf{\Gamma}$  for the case  $g(x) = e^x$  and  $m = 3$ . We have

$$\begin{aligned} \gamma_{11} &= 1 / (1 + \exp(\tau_{12}) + \exp(\tau_{13})), \\ \gamma_{12} &= \exp(\tau_{12}) / (1 + \exp(\tau_{12}) + \exp(\tau_{13})), \\ \gamma_{13} &= \exp(\tau_{13}) / (1 + \exp(\tau_{12}) + \exp(\tau_{13})). \end{aligned}$$

The transformation in the opposite direction is

$$\begin{aligned} \tau_{12} &= \log(\gamma_{12} / (1 - \gamma_{12} - \gamma_{13})) = \log(\gamma_{12} / \gamma_{11}), \\ \tau_{13} &= \log(\gamma_{13} / (1 - \gamma_{12} - \gamma_{13})) = \log(\gamma_{13} / \gamma_{11}). \end{aligned}$$

This generalization of the logit and inverse logit transforms has long been used in (for example) the context of compositional data: see Aitchison (1982), where several other transforms are described as well.

### 3.3.2 Embedding in a continuous-time Markov chain

A different reparametrization is described by Zucchini and MacDonald (1998). In a continuous-time Markov chain on a finite state-space, the transition probability matrix  $\mathbf{P}_t$  is given by  $\mathbf{P}_t = \exp(t\mathbf{Q})$ , where  $\mathbf{Q}$  is the matrix of transition intensities. The row sums of  $\mathbf{Q}$  are zero, but the only constraint on the off-diagonal elements of  $\mathbf{Q}$  is that they be non-negative. The one-step transition probabilities in a discrete-time Markov chain can therefore be parametrized via  $\mathbf{\Gamma} = \exp(\mathbf{Q})$ . This is effectively the parametrization used in the  $\mathbf{R}$  package `msm` (Jackson *et al.*, 2003).

## 3.4 Other problems

### 3.4.1 Multiple maxima in the likelihood

The likelihood of an HMM is a complicated function of the parameters and frequently has several local maxima. The goal of course is to find the global maximum, but there is no simple method of determining in

general whether a numerical maximization algorithm has reached the global maximum. Depending on the starting values, it can easily happen that the algorithm identifies a local, but not the global, maximum. This applies also to the main alternative method of estimation, the EM algorithm, which is discussed in Chapter 4. A sensible strategy is therefore to use a range of starting values for the maximization, and to see whether the same maximum is identified in each case.

### 3.4.2 Starting values for the iterations

It is often easy to find plausible starting values for some of the parameters of an HMM: for instance, if one seeks to fit a Poisson–HMM with two states, and the sample mean is 10, one could try 8 and 12, or 5 and 15, for the values of the two state-dependent means. More systematic strategies based on the quantiles of the observations are possible, however: e.g. if the model has three states, use as the starting values of the state-dependent means the lower quartile, median and upper quartile of the observed counts.

It is less easy to guess values of the transition probabilities  $\gamma_{ij}$ . One strategy is to assign a common starting value (e.g. 0.01 or 0.05) to all the off-diagonal transition probabilities. A consequence of such a choice, perhaps convenient, is that the corresponding stationary distribution is uniform over the states; this follows by symmetry. Choosing good starting values for parameters tends to steer one away from numerical instability.

### 3.4.3 Unbounded likelihood

In the case of HMMs with continuous state-dependent distributions, just as in the case of independent mixtures (see Section 1.2.3), it may happen that the likelihood is unbounded in the vicinity of certain parameter combinations. As before, we suggest that, if this creates difficulties, one maximizes the discrete likelihood instead of the joint density. This has the advantage in any case that it applies more generally to data known only up to some interval. Applications of this kind are described in Sections 10.3 and 10.4; code for the latter is in A.3.

## 3.5 Example: earthquakes

Figure 3.1 shows the result of fitting (stationary) Poisson–hidden Markov models with two and three states to the earthquakes series by means of the unconstrained optimizer `nlm`. The relevant code appears in A.1. The




Figure 3.1 *Earthquakes series. Left: marginal distributions of Poisson-HMMs with 2 and 3 states, and their components, compared with a histogram of the observations. Right: the state-dependent means (horizontal lines) compared to the observations.*

two-state model is

$$\mathbf{\Gamma} = \begin{pmatrix} 0.9340 & 0.0660 \\ 0.1285 & 0.8715 \end{pmatrix},$$

with  $\boldsymbol{\delta} = (0.6608, 0.3392)$ ,  $\boldsymbol{\lambda} = (15.472, 26.125)$ , and log-likelihood given by  $l = -342.3183$ . It is clear that the fitted (Markov-dependent) mixture of two Poisson distributions provides a much better fit to the marginal distribution of the observations than does a single Poisson distribution, but the fit can be further improved by using a mixture of three or four Poisson distributions.

The three-state model is

$$\mathbf{\Gamma} = \begin{pmatrix} 0.955 & 0.024 & 0.021 \\ 0.050 & 0.899 & 0.051 \\ 0.000 & 0.197 & 0.803 \end{pmatrix},$$

with  $\boldsymbol{\delta} = (0.4436, 0.4045, 0.1519)$ ,  $\boldsymbol{\lambda} = (13.146, 19.721, 29.714)$  and  $l =$

−329.4603, and the four-state is as follows:

$$\Gamma = \begin{pmatrix} 0.805 & 0.102 & 0.093 & 0.000 \\ 0.000 & 0.976 & 0.000 & 0.024 \\ 0.050 & 0.000 & 0.902 & 0.048 \\ 0.000 & 0.000 & 0.188 & 0.812 \end{pmatrix},$$

with  $\delta = (0.0936, 0.3983, 0.3643, 0.1439)$ ,  $\lambda = (11.283, 13.853, 19.695, 29.700)$ , and  $l = -327.8316$ .

The means and variances of the marginal distributions of the four models compare as follows with those of the observations. By a one-state Poisson–HMM we mean a model that assumes that the observations are realizations of independent Poisson random variables with common mean.

	mean	variance
observations:	19.364	51.573
‘one-state HMM’:	19.364	19.364
two-state HMM:	19.086	44.523
three-state HMM:	18.322	50.709
four-state HMM:	18.021	49.837

As regards the autocorrelation functions of the models, i.e.  $\rho(k) = \text{Corr}(X_{t+k}, X_t)$ , we have the following results, valid for all  $k \in \mathbb{N}$ , based on the conclusions of Exercise 4 of Chapter 2:

- two states:  $\rho(k) = 0.5713 \times 0.8055^k$ ;
- three states:  $\rho(k) = 0.4447 \times 0.9141^k + 0.1940 \times 0.7433^k$ ;
- four states:  $\rho(k) = 0.2332 \times 0.9519^k + 0.3682 \times 0.8174^k + 0.0369 \times 0.7252^k$ .

In all these cases the ACF is just a linear combination of the  $k$ th powers of the eigenvalues other than 1 of the transition probability matrix.

For model selection, e.g. choosing between competing models such as HMMs and independent mixtures, or choosing the number of components in either, see Section 6.1.

A phenomenon that is noticeable when one fits models with three or more states to relatively short series is that the estimates of one or more of the transition probabilities turn out to be very close to zero; see the three-state model above (one such probability,  $\gamma_{13}$ ) and the four-state model (six of the twelve off-diagonal transition probabilities).

This phenomenon can be explained as follows. In a stationary Markov chain, the expected number of transitions from state  $i$  to state  $j$  in a series of  $T$  observations is  $(T-1)\delta_i\gamma_{ij}$ . For  $\delta_3 = 0.152$  and  $T = 107$  (as in our three-state model), this expectation will be less than 1 if  $\gamma_{31} < 0.062$ . In such a series, therefore, it is likely that if  $\gamma_{31}$  is fairly small there will be no transitions from state 3 to state 1, and so when we



seek to estimate  $\gamma_{31}$  in an HMM the estimate is likely to be effectively zero. As  $m$  increases, the probabilities  $\delta_i$  and  $\gamma_{ij}$  get smaller on average; this makes it increasingly likely that at least one estimated transition probability is effectively zero.

### 3.6 Standard errors and confidence intervals

Relatively little is known about the properties of the maximum likelihood estimators of HMMs; only asymptotic results are available. To exploit these results one requires estimates of the variance-covariance matrix of the estimators of the parameters. One can estimate the standard errors from the Hessian of the log-likelihood at the maximum, but this approach runs into difficulties when some of the parameters are on the boundary of their parameter space, which occurs quite often when HMMs are fitted. An alternative here is the parametric bootstrap, for which see [Section 3.6.2](#). The algorithm is easy to code (see A.2.1), but the computations are time-consuming.

#### 3.6.1 Standard errors via the Hessian

Although the point estimates  $\hat{\Theta} = (\hat{\Gamma}, \hat{\lambda})$  are easy to compute, exact interval estimates are not available. Cappé *et al.* (2005, Chapter 12) show that, under certain regularity conditions, the MLEs of HMM parameters are consistent, asymptotically normal and efficient. Thus, if we can estimate the standard errors of the MLEs, then, using the asymptotic normality, we can also compute approximate confidence intervals. However, as pointed out by Frühwirth-Schnatter (2006, p. 53) in the context of independent mixture models, ‘The regularity conditions are often violated, including cases of great practical concern, among them small data sets, mixtures with small component weights, and overfitting mixtures with too many components.’ Furthermore, McLachlan and Peel (2000, p. 68) warn: ‘In particular for mixture models, it is well known that the sample size  $n$  has to be very large before the asymptotic theory of maximum likelihood applies.’

With the above caveats in mind we can, to estimate the standard errors of the MLEs of an HMM, use the approximate Hessian of minus the log-likelihood at the minimum, e.g. as supplied by `nlm`. We can invert it and so estimate the asymptotic variance-covariance matrix of the estimators of the parameters. A problem with this suggestion is that, if the parameters have been transformed, the Hessian available will be that which refers to the working parameters  $\phi_i$ , not the original, more readily interpretable, natural parameters  $\theta_i$  ( $\Gamma$  and  $\lambda$  in the case of a Poisson-HMM).

The situation is therefore that we have the Hessian

$$\mathbf{H} = - \left( \frac{\partial^2 l}{\partial \phi_i \partial \phi_j} \right)$$

available at the minimum of  $-l$ , and what we really need is the Hessian with respect to the natural parameters:

$$\mathbf{G} = - \left( \frac{\partial^2 l}{\partial \theta_i \partial \theta_j} \right).$$

There is, however, the following relationship between the two Hessians at the minimum:

$$\mathbf{H} = \mathbf{M} \mathbf{G} \mathbf{M}' \quad \text{and} \quad \mathbf{G}^{-1} = \mathbf{M}' \mathbf{H}^{-1} \mathbf{M}, \quad (3.2)$$

where  $\mathbf{M}$  is defined by  $m_{ij} = \partial \theta_j / \partial \phi_i$ . (Note that all the derivatives appearing here are as evaluated at the minimum.) In the case of a Poisson-HMM, the elements of  $\mathbf{M}$  are quite simple: see [Exercise 7](#) for details.

With  $\mathbf{M}$  at our disposal, we can use (3.2) to deduce  $\mathbf{G}^{-1}$  from  $\mathbf{H}^{-1}$ , and use  $\mathbf{G}^{-1}$  to find standard errors for the natural parameters, provided such parameters are not on the boundary of the parameter space. It is however true in many applications that some of the estimated parameters lie on or very close to the boundary; this limits the usefulness of the above results.

As already pointed out on p. 52, for series of moderate length the estimates of some transition probabilities are expected to be close to zero. This is true of  $\hat{\gamma}_{13}$  in the three-state model for the earthquakes series. An additional example of this type can be found in Section 10.2.2. In Section 12.2.1, several of the estimates of the parameters in the state-dependent distributions are practically zero, their lower bound; see Table 12.1. The same phenomenon is apparent in Section 16.9.2; see Table 16.1.

### *Recursive computation of the Hessian*

An alternative method of computing the Hessian is that of Lystig and Hughes (2002). They present the forward algorithm  $\boldsymbol{\alpha}_t = \boldsymbol{\alpha}_{t-1} \mathbf{G} \mathbf{P}(x_t)$  in a form which incorporates automatic or ‘natural’ scaling, and then extend that approach in order to compute (in a single pass, along with the log-likelihood) its Hessian and gradient with respect to the natural parameters, those we have denoted above by  $\theta_i$ . Turner (2008) has used this approach in order to find the analytical derivatives needed to maximize HMM likelihoods directly by the Levenberg–Marquardt algorithm.

While this may be a more efficient and more accurate method of computing the Hessian than that outlined above, it does not solve the fundamental problem that the use of the Hessian to compute standard errors

(and thence confidence intervals) is unreliable if some of the parameters are on or near the boundary of their parameter space.

### 3.6.2 Bootstrap standard errors and confidence intervals

As an alternative to the technique described in Section 3.6.1 one may use the **parametric bootstrap** (Efron and Tibshirani, 1993). Roughly speaking, the idea of the parametric bootstrap is to assess the properties of the model with parameters  $\Theta$  by using those of the model with parameters  $\hat{\Theta}$ . The following steps are performed to estimate the variance-covariance matrix of  $\hat{\Theta}$ .

1. Fit the model, i.e. compute  $\hat{\Theta}$ .
- 2.(a) Generate a sample, called a bootstrap sample, of observations from the fitted model, i.e. the model with parameters  $\hat{\Theta}$ . The length should be the same as the original number of observations.
- (b) Estimate the parameters  $\Theta$  by  $\hat{\Theta}^*$  for the bootstrap sample.
- (c) Repeat steps (a) and (b)  $B$  times (with  $B$  ‘large’) and record the values  $\hat{\Theta}^*$ .

The variance-covariance matrix of  $\hat{\Theta}$  is then estimated by the sample variance-covariance matrix of the bootstrap estimates  $\hat{\Theta}^*(b)$ ,  $b = 1, 2, \dots, B$ :

$$\widehat{\text{Var-Cov}}(\hat{\Theta}) = \frac{1}{B-1} \sum_{b=1}^B \left( \hat{\Theta}^*(b) - \hat{\Theta}^*(.) \right)' \left( \hat{\Theta}^*(b) - \hat{\Theta}^*(.) \right),$$

where  $\hat{\Theta}^*(.) = B^{-1} \sum_{b=1}^B \hat{\Theta}^*(b)$ .

The parametric bootstrap requires code to generate realizations from a fitted model; for a Poisson–HMM this is given in A.2.1. Since code to fit models is available, that same code can be used to fit models to the bootstrap sample.

The bootstrap method can be used to estimate confidence intervals directly. In the example given in the next section we use the well-known ‘percentile method’ (Efron and Tibshirani, 1993); other options are available.

## 3.7 Example: the parametric bootstrap applied to the three-state model for the earthquakes data

A bootstrap sample of size 500 was generated from the three-state model for the earthquakes data, which appears on p. 51. In fitting models to the bootstrap samples, we noticed that, in two cases out of the 500, the starting values we were in general using caused numerical instability or

Table 3.1 *Earthquakes data: bootstrap confidence intervals for the parameters of the three-state HMM.*

parameter	MLE	90% conf. limits	
$\lambda_1$	13.146	11.463	14.253
$\lambda_2$	19.721	13.708	21.142
$\lambda_3$	29.714	20.929	33.160
$\gamma_{11}$	0.954	0.750	0.988
$\gamma_{12}$	0.024	0.000	0.195
$\gamma_{13}$	0.021	0.000	0.145
$\gamma_{21}$	0.050	0.000	0.179
$\gamma_{22}$	0.899	0.646	0.974
$\gamma_{23}$	0.051	0.000	0.228
$\gamma_{31}$	0.000	0.000	0.101
$\gamma_{32}$	0.197	0.000	0.513
$\gamma_{33}$	0.803	0.481	0.947
$\delta_1$	0.444	0.109	0.716
$\delta_2$	0.405	0.139	0.685
$\delta_3$	0.152	0.042	0.393

Table 3.2 *Earthquakes data: bootstrap estimates of the correlations of the estimators of  $\lambda_i$ , for  $i = 1, 2, 3$ .*

	$\lambda_1$	$\lambda_2$	$\lambda_3$
$\lambda_1$	1.000	0.483	0.270
$\lambda_2$		1.000	0.688
$\lambda_3$			1.000

convergence problems. By choosing better starting values for these two cases we were able to fit models successfully and complete the exercise. The resulting sample of parameter values then produced the 90% confidence intervals for the parameters that are displayed in Table 3.1, and the estimated parameter correlations that are displayed in Table 3.2. What is noticeable is that the intervals for the state-dependent means  $\lambda_i$  overlap, the intervals for the stationary probabilities  $\delta_i$  are very wide, and the estimators  $\hat{\lambda}_i$  are quite strongly correlated.

These results, in particular the correlations shown in Table 3.2, should make one wary of over-interpreting a model with nine parameters based on only 107 (dependent) observations. In particular, they suggest that the states are not well defined, and one should be cautious of attaching a substantive interpretation to them.

## Exercises

- Consider the following parametrization of the t.p.m. of an  $m$ -state Markov chain. Let  $\tau_{ij} \in \mathbb{R}$  ( $i, j = 1, 2, \dots, m; i \neq j$ ) be  $m(m-1)$  arbitrary real numbers. Let  $g : \mathbb{R} \rightarrow \mathbb{R}^+$  be some strictly increasing function, e.g.  $g(x) = e^x$ . Define  $\varrho_{ij}$  and  $\gamma_{ij}$  as on p. 48.
  - Show that the matrix  $\mathbf{\Gamma}$  with entries  $\gamma_{ij}$  that are constructed in this way is a t.p.m., i.e. show that  $0 \leq \gamma_{ij} \leq 1$  for all  $i$  and  $j$ , and that the row sums of  $\mathbf{\Gamma}$  are equal to one.
  - Given a t.p.m.  $\mathbf{\Gamma} = \{\gamma_{ij} : i, j = 1, 2, \dots, m\}$ , derive an expression for the parameters  $\tau_{ij}$ , for  $i, j = 1, 2, \dots, m; i \neq j$ .
- The purpose of this exercise is to investigate the numerical behaviour of an ‘unscaled’ evaluation of the likelihood of an HMM, and to compare this with the behaviour of an alternative algorithm that applies scaling.

Consider the stationary two-state Poisson–HMM with parameters

$$\mathbf{\Gamma} = \begin{pmatrix} 0.9 & 0.1 \\ 0.2 & 0.8 \end{pmatrix}, \quad (\lambda_1, \lambda_2) = (1, 5).$$

Compute the likelihood,  $L_{10}$ , of the following sequence of ten observations in two ways: 2, 8, 6, 3, 6, 1, 0, 0, 4, 7.

- Use the unscaled method  $L_{10} = \boldsymbol{\alpha}_{10} \mathbf{1}'$ , where  $\boldsymbol{\alpha}_0 = \boldsymbol{\delta}$  and  $\boldsymbol{\alpha}_t = \boldsymbol{\alpha}_{t-1} \mathbf{B}_t$ ;

$$\mathbf{B}_t = \mathbf{\Gamma} \begin{pmatrix} p_1(x_t) & 0 \\ 0 & p_2(x_t) \end{pmatrix};$$

and

$$p_i(x_t) = \lambda_i^{x_t} e^{-\lambda_i} / x_t!, \quad i = 1, 2; \quad t = 1, 2, \dots, 10.$$

Examine the numerical values of the vectors  $\boldsymbol{\alpha}_0, \boldsymbol{\alpha}_1, \dots, \boldsymbol{\alpha}_{10}$ .

- Use the algorithm given in Section 3.2 to compute  $\log L_{10}$ .

Examine the numerical values of the vectors  $\boldsymbol{\phi}_0, \boldsymbol{\phi}_1, \dots, \boldsymbol{\phi}_{10}$ . (It is easiest to store these vectors as rows in an  $11 \times 2$  matrix.)

- Use the **R** function `constrOptim` to fit HMMs with two to four states to the earthquakes data, and compare your models with those given in Section 3.5.
- Consider the following transformation:

$$\begin{aligned} w_1 &= \sin^2 \theta_1 \\ w_i &= \left( \prod_{j=1}^{i-1} \cos^2 \theta_j \right) \sin^2 \theta_i \quad i = 2, \dots, m-1 \\ w_m &= \prod_{i=1}^{m-1} \cos^2 \theta_i. \end{aligned}$$

Show how this transformation can be used to convert the constraints

$$\sum_{i=1}^m w_i = 1, \quad w_i \geq 0 \quad i = 1, \dots, m$$

into simple ‘box constraints’, i.e. constraints of the form  $a \leq \theta_i \leq b$ . How could this be used in the context of estimation in HMMs?

- 5.(a) Consider a stationary Markov chain, with t.p.m.  $\mathbf{\Gamma}$  and stationary distribution  $\boldsymbol{\delta}$ . Show that the expected number of transitions from state  $i$  to state  $j$  in a series of  $T$  observations (i.e. in  $T - 1$  transitions) is  $(T - 1)\delta_i\gamma_{ij}$ .

Hint: this expectation is  $\sum_{t=2}^T \Pr(X_{t-1} = i, X_t = j)$ .

- (b) Show that, for  $\delta_3 = 0.152$  and  $T = 107$ , this expectation is less than 1 if  $\gamma_{31} < 0.062$ .
6. Prove the relation (3.2) between the Hessian  $\mathbf{H}$  of  $-l$  with respect to the working parameters and the Hessian  $\mathbf{G}$  of  $-l$  with respect to the natural parameters, both being evaluated at the minimum of  $-l$ .
7. (See [Section 3.6.1](#).) Consider an  $m$ -state Poisson–HMM, with natural parameters  $\gamma_{ij}$  and  $\lambda_i$ , and working parameters  $\tau_{ij}$  and  $\eta_i$  defined as in [Section 3.3.1](#), with  $g(x) = e^x$ .
- (a) Show that
- $$\begin{aligned} \partial\gamma_{ij}/\partial\tau_{ij} &= \gamma_{ij}(1 - \gamma_{ij}), \text{ for all } i, j; & \partial\gamma_{ij}/\partial\tau_{il} &= -\gamma_{ij}\gamma_{il}, \text{ for } j \neq l; \\ \partial\gamma_{ij}/\partial\tau_{kl} &= 0, \text{ for } i \neq k; \text{ and} & \partial\lambda_i/\partial\eta_i &= e^{\eta_i} = \lambda_i, \text{ for all } i. \end{aligned}$$
- (b) Hence find the matrix  $\mathbf{M}$  in this case.
8. Modify the **R** code in [A.1](#) in order to fit a Poisson–HMM to interval-censored observations. (Assume that the observations are available as a  $T \times 2$  matrix of which the first column contains the lower bound of the observation and the second the upper bound, possibly **Inf**.)
9. Verify the autocorrelation functions given on p. 52 for the two-, three- and four-state models for the earthquakes data. (Hint: use the **R** function **eigen** to find the eigenvalues and -vectors of the relevant transition probability matrices.)
10. Consider again the soap sales series introduced in Exercise 5 of Chapter 1.
- (a) Fit stationary Poisson–HMMs with two, three and four states to these data.
- (b) Find the marginal means and variances, and the ACFs, of these models, and compare them with their sample equivalents.