

Here we provide short scripts to check the coincidence between the factors in the class number formula for biquadratic fields $\mathbb{Q}(\sqrt{-m}, \zeta_3)$ and the index of the reduction morphism $\mathcal{O}_{-m}[\zeta_3]^\times \rightarrow \mathcal{O}_m[\zeta_3]/(\zeta_3 - 1)^\times \cong \mathcal{O}_{-m}/(3)^\times$.

The following short SAGE-script goes through all square-free $m \leq 1000$ with $m \equiv 2 \pmod{3}$, computes the class numbers of the fields $K = \mathbb{Q}(\sqrt{-m})$, $L = \mathbb{Q}(\sqrt{-m}, \zeta_3)$ and $T = \mathbb{Q}(\sqrt{3m})$ and the factor $\frac{h_{\mathbb{Q}(\sqrt{-m}, \sqrt{-3})}}{h_{\mathbb{Q}(\sqrt{-m})}h_{\mathbb{Q}(\sqrt{3m})}}$. Then it computes a fundamental unit and checks if the unit is ± 1 modulo the ideal $(\zeta_3 - 1)$. The final line checks that the reduction of the fundamental unit is ± 1 if and only if the class number formula factor is $\frac{1}{2}$. For more discussion and a table of the first few outputs, see Example 6.3 of the paper “ON FARRELL–TATE COHOMOLOGY OF GL₃ OVER RINGS OF QUADRATIC INTEGERS” by BUI ANH TUAN, ALEXANDER D. RAHM AND MATTHIAS WENDT.

```
for m in range(1000):
    if mod(m,3) == 2 and Integer(m).is_squarefree():
        print('m='+str(m))
        K.<sqnm> = NumberField(x^2+m)
        R.<y> = PolynomialRing(K)
        L.<zeta_3> = K.extension(y^2+y+1)
        S.<c> = PolynomialRing(QQ)
        T.<b> = NumberField(c^2-3*m)
#     print('class number='+str(L.class_number()))
    factor = L.class_number()/(K.class_number()*T.class_number())
#     print('class number formula factor='+str(factor))
    fu = L.units()[0]
    ideal = L.ideal(zeta_3-1)
    if not ((fu - 1 in ideal) or (fu + 1 in ideal)) == (factor == 1/2):
        print('conjecture false')
```

For the case $m \equiv 1 \pmod{3}$, only few modifications have to be made. Obviously the change to $\text{mod}(m,3) == 1$ in the second line, and then the last line of the previous script has to be replaced by

```
index = mod(ideal.ideallog(fu)[0],4)
print('unit index='+str(index))
if mod(index,2) == 1:
    print('exceptional case')
if not ((index == 2) == (factor == 1)):
    print('conjecture false')
```

Here, the function `ideallog(fu)` is used to compute the index of the reduction of the fundamental unit in $\mathcal{O}_{-m}/(3)^\times \cong \mathbb{Z}/8\mathbb{Z}$. According to our expectation in Example 6.5 of the paper “ON FARRELL–TATE COHOMOLOGY OF GL₃ OVER RINGS OF QUADRATIC INTEGERS” by BUI ANH TUAN, ALEXANDER D. RAHM AND MATTHIAS WENDT, the reduction of the fundamental unit should only be a generator in the case $m = 1$, otherwise it should have reduction equal to ± 1 (and thus yield a number of 5 orbits) if and only if the class number formula factor is $\frac{1}{2}$. For more discussion and a table of outcomes, cf. the quoted Example 6.5.

BUI ANH TUAN, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF SCIENCE, VNU-HCM, 227 NGUYEN VAN CU STR., DIST. 5, HO CHI MINH CITY, VIETNAM
Email address: `batuan@hcmus.edu.vn`

ALEXANDER D. RAHM, LABORATOIRE DE MATHÉMATIQUES GAATI, UNIVERSITÉ DE LA POLYNÉSIE FRANÇAISE, BP 6570, 98702 FAAA, FRENCH POLYNESIA
Email address: `Alexander.Rahm@upf.pf`

MATTHIAS WENDT, FACHGRUPPE MATHEMATIK/INFORMATIK, BERGISCHE UNIVERSITÄT WUPPERTAL, GAUSSSTRASSE 20, 42119 WUPPERTAL, GERMANY
Email address: `wendt@math.uni-wuppertal.de`