

## Project 5

CODE:

```
import math

#parameters in the question
seed=1234567
a=7**5
c=0
m=(2**31)-1

randomNumbers=[]

#function to create random numbers
def generateRandom():
    global seed
    seed=(a*seed+c)%m
    return round(seed/m,3)

for i in range(100):

    randomNumbers.append(generateRandom())

copyrandomNumbers=randomNumbers.copy()
randomNumbers.sort()

print(randomNumbers)

#this part here is for kolmogrov-smirnov test

dpluses=[] #array to sto
re all d+ values
dnegatives=[] #array to sto
re all d- values
for i in range(len(randomNumbers)):
    dpluses.append(((i+1)/len(randomNumbers))-
randomNumbers[i]) #i in the for loop starts from 0 thats why i need +1 here
    dnegatives.append(randomNumbers[i]-
((i)/len(randomNumbers))) #i in the for loop starts from 0 thats why i dont ne
ed -1 here
```

```

dpluses.sort()                                #sorting the
array so we can get the max
dnegatives.sort()                            #sorting the
array so we can get the max
dplus = dpluses[len(randomNumbers)-
1]                                           #getting the max
dnegative = dnegatives[len(randomNumbers)-
1]                                           #getting the max
d=max(dplus,dnegative)                      #getting the
max

dalpha=1.36/math.sqrt(len(randomNumbers))    #alpha value

if(dalpha>d):
    print("numbers are uniform for alpha level of 0.05")
else:
    print("numbers are not uniform for alpha level of 0.05")

print("d $\alpha$  = ",dalpha)
print("d = ",d)

#this part is for runs up and runs down test
currentRunType=None
runCount=0

#loop for counting runs
for i in range(1,len(copyrandomNumbers)):
    num=copyrandomNumbers[i]
    if currentRunType!=None:                #if it is not the first 2 elements th
is block runs
        if (num>copyrandomNumbers[i-
1]):    #checks if the current element is greater than the previous element
            if(currentRunType=="-
"):    #then checks if it runs down if it runs down it means now its running
up
                currentRunType="+"          #thats why we change the run type

                runCount+=1                  #and increment the run count
                #there is not a else block because

```

```

        #if current run type is up we are continuing the run and there is no
change in the run type

    else:                                #if previous number is greater than t
he current element

        if(currentRunType=="+"):        #if it is running up that means now i
t started to run down
            currentRunType="-"
"            #so we change the run type and increment the counter
            runCount+=1
            #there is not a else block because
            #if current run type is down we are continuing the run and there is n
o change in the run type

    else:                                # if we are checking the first 2 elem
ents then this else block runs
        if(num>copyrandomNumbers[i-1]):
            currentRunType="+"
        else:
            currentRunType="-"
        runCount+=1

meana=((2*len(randomNumbers))-1)/3      #mean of a
vara=((16*len(randomNumbers))-29)/90   #variance of a
z=1.96                                  #z value that we need to compare with
z0=(runCount-meana)/math.sqrt(vara)

if (z0>-z and z0<z):
    print("numbers are independent for alpha level of 0.05")
else:
    print("numbers are not independent for alpha level of 0.05")
print("z = ",z)
print("z0 = ",z0)

```

random numbers generated with these parameters( $X_0 = 123457$ ,  $a = 75$ ,  $c = 0$ ,  $m = 231 - 1$ ):

0.662, 0.212, 0.396, 0.317, 0.85, 0.786, 0.731, 0.977, 0.926, 0.004, 0.009, 0.312, 0.394, 0.587, 0.97,  
0.379, 0.882, 0.927, 0.31, 0.221, 0.407, 0.927, 0.127, 0.192, 0.72, 0.88, 0.819, 0.772, 0.078, 0.769, 0.057,  
0.547, 0.805, 0.122, 0.404, 0.144, 0.699, 0.424, 0.052, 0.067, 0.056, 0.159, 0.683, 0.253, 0.332, 0.592,  
0.379, 0.226, 0.726, 0.12, 0.158, 0.872, 0.469, 0.913, 0.688, 0.295, 0.306, 0.829, 0.318, 0.72, 0.854,  
0.317, 0.021, 0.541, 0.792, 0.314, 0.226, 0.838, 0.467, 0.175, 0.456, 0.219, 0.047, 0.858, 0.651, 0.489,  
0.799, 0.034, 0.378, 0.117, 0.921, 0.956, 0.74, 0.196, 0.257, 0.529, 0.49, 0.715, 0.715, 0.932, 0.701,  
0.824, 0.842, 0.775, 0.322, 0.917, 0.321, 0.266, 0.217, 0.484

3)

numbers are uniform for alpha level of 0.05

$d\alpha = 0.136$

$d = 0.073000000000000006$

```
numbers are uniform for alpha level of 0.05
dα = 0.136
d = 0.073000000000000006
```

4)

numbers are independent for alpha level of 0.05

$z = 1.96$

$z_0 = -0.5584829866163423$

```
numbers are independent for alpha level of 0.05
z = 1.96
z0 = -0.5584829866163423
```