

Half-spherical twists on derived categories of coherent sheaves

based on arXiv:2302.12501

Hayato Arai

Graduate School of Mathematical Sciences, The University of Tokyo

1 Introduction

1.1 Mirror symmetry for Kodaira fibers of type I_n

Let $\pi: S \rightarrow C$ be a relatively minimal, smooth projective elliptic surface. The possible singular fibers of π are classified by Kodaira and Néron, and they are called *Kodaira fibers*. Among them, the type I_n Kodaira fiber is the cyclic configuration of n smooth rational curves.

Lekili and Polishchuk [LP17] established mirror symmetry between the type I_n Kodaira fiber Y_n and the n -punctured torus T_n , i.e. they showed that the derived category $D^b(Y_n)$ of coherent sheaves on Y_n and the wrapped Fukaya category $D^\pi(\mathcal{W}(T_n))$ of T_n are equivalent.

For example, the equivalence includes the following correspondence of objects, where G_i 's are irreducible components and γ_F denotes the Lagrangian submanifold corresponding to the object $F \in D^b(Y_n)$:




Figure 1: The correspondence of objects via $D^b(Y_n) \simeq D^\pi(\mathcal{W}(T_n))$, $n = 3$.

Combining this with the theory of topological Fukaya categories of surfaces (such as Haiden, Katzarkov, and Kontsevich [HKK17]), Oppen [Opp20] described the autoequivalence group of $D^b(Y_n)$ with the following exact sequence:

$$1 \rightarrow (\mathbb{C}^\times)^n \times \mathbb{Z}[1] \times \text{Pic}^0(Y_n) \rightarrow \text{Auteq } D^b(Y_n) \xrightarrow{\Upsilon} \text{MCG}(T_n) \rightarrow 1.$$

Here $\text{MCG}(T_n)$ denotes the mapping class group of T_n and the morphism Υ is induced by the equivalence $D^b(Y_n) \simeq D^\pi(\mathcal{W}(T_n))$.

1.2 Autoequivalences of elliptic surfaces

Uehara [Ueh16] gave the following description of the autoequivalence group $\text{Auteq } D^b(S)$.

Suppose that

- S has non-zero Kodaira dimension,
- all singular fibers of π are non-multiple and of type I_n , $n \geq 2$, and
- $B = \langle T_{\mathcal{O}_G(a)} \mid G \subset S : \text{an irreducible component of a singular fiber, } a \in \mathbb{Z} \rangle$: the subgroup of $\text{Auteq } D^b(S)$ generated by twist functors $T_{\mathcal{O}_G(a)}$, where $\mathcal{O}_G(a)$ is the line bundle of degree a on $G \simeq \mathbb{P}^1$.

Then there is the exact sequence

$$1 \rightarrow \langle B, (-) \otimes \mathcal{O}_S(D) \mid D.F = 0, F \text{ is a fiber} \rangle \rtimes \text{Aut } S \times \mathbb{Z}[2] \rightarrow \text{Auteq } D^b(S) \rightarrow \text{SL}(2, \mathbb{Z}).$$

This result implies that the study of the structure of $\text{Auteq } D^b(S)$ reduces to that of B .

1.3 Main results

We study the group B in terms of the mapping class group of the n -punctured torus so that we reveal the whole structure of $\text{Auteq } D^b(S)$.

- (1) We construct a natural “restriction” morphism $\text{res}: B \rightarrow \text{Auteq } D^b(F)$ for each fiber F , which is nontrivial if F is a reducible fiber.
- (2) Combining with mirror symmetry for the Kodaira fibers of type I_n , we have the exact sequences

$$1 \rightarrow \langle (-) \otimes \mathcal{O}_S(Y_{n_j}) \mid 1 \leq j \leq m \rangle \rightarrow B \xrightarrow{\Upsilon \circ \text{res}} \prod_{j=1}^m \text{MCG}(T_{n_j}),$$

where Y_{n_j} is the Kodaira fiber of type I_{n_j} and $\{Y_{n_j}\}_{j=1}^m$ is the set of all singular fibers of S .

- (3) For $G \subset Y_n$, $a \in \mathbb{Z}$, and the curve $\gamma_{\mathcal{O}_G(a)}$ on T_n corresponding to $\mathcal{O}_G(a)$ under the equivalence $D^b(Y_n) \simeq D^\pi(\mathcal{W}(T_n))$, the twist functor $T_{\mathcal{O}_G(a)} \in B$ is mapped to the half twist along $\gamma_{\mathcal{O}_G(a)}$ in $\text{MCG}(T_n) \subset \prod_{j=1}^m \text{MCG}(T_{n_j})$.
- (4) The image of $\Upsilon \circ \text{res}$ is generated by the half twists along the finite number of curves $\{\gamma_{\mathcal{O}_G}, \gamma_{\mathcal{O}_G(-1)} \mid G \subset Y_{n_j} : \text{an irreducible component, } 1 \leq j \leq m\}$.




Figure 2: The half twist along the curve γ .

2 Sketch of the proof

2.1 Result (1)

We prepare the following theorem to prove the result (1). This is a generalization of the relation between twist functors and \mathbb{P} -twists by Huybrechts and Thomas [HT06].

Theorem 2.1.

- $\pi: X \rightarrow T$: a flat morphism between smooth quasi-projective varieties
- $i: Y = \pi^{-1}(0) \hookrightarrow X$: the fiber at $0 \in T$

- $E \in D^b(Y)$: an object such that $i_*E \in D^b(X)$ is a spherical object
- $T_{i_*E} \in \text{Auteq } D^b(X)$: the corresponding twist functor

Then there is a unique $H_E \in \text{Auteq } D^b(Y)$ which makes the following diagram commutative:

$$\begin{array}{ccc} D^b(Y) & \xrightarrow{i_*} & D^b(X) \\ H_E \downarrow & & \downarrow T_{i_*E} \\ D^b(Y) & \xrightarrow{i_*} & D^b(X). \end{array} \quad (1)$$

Proof of the result (1). The uniqueness and the construction of H_E ensure that we have a well-defined morphism $\text{res}: B \rightarrow \text{Auteq } D^b(F)$ such that $\text{res}(T_{\mathcal{O}_G(a)}) = H_{\mathcal{O}_G(a)}$ if $G \subset F$ and $\text{res}(T_{\mathcal{O}_G(a)}) = \text{id}$ otherwise. \square

2.2 Result (3)

For the proof of the result (3), we use the following three facts.

- The group $\pi_1(T_n)$ is generated by the curves $\gamma_{\mathcal{O}_{Y_n}}, \gamma_{\mathcal{O}_{x_1}}, \dots, \gamma_{\mathcal{O}_{x_n}}$ (see Figure 1).
- An element of $\text{MCG}(T_n)$ is determined by its action on $\pi_1(T_n)$ (Dehn–Nielsen–Baer theorem).
- [HKK17, Opp20] There are correspondences between indecomposable objects of $D^b(Y_n)$ and homotopy classes of curves on T_n (+ some data), and between dimensions of Hom-spaces in $D^b(Y_n)$ and intersection numbers of curves on T_n .

Proof of the result (3). For simplicity, we assume that $x_1 \in G$ and $a = 0$. We denote the image under the morphism $\Upsilon \circ \text{res}$ of the twist functor $T_{\mathcal{O}_G}$ by H , and the half twist along $\gamma_{\mathcal{O}_G}$ by H' .

Given the first and second facts, we only need to check the identities $H(\gamma_{\mathcal{O}_{Y_n}}) = H'(\gamma_{\mathcal{O}_{Y_n}})$, $H(\gamma_{\mathcal{O}_{x_1}}) = H'(\gamma_{\mathcal{O}_{x_1}})$, \dots , $H(\gamma_{\mathcal{O}_{x_n}}) = H'(\gamma_{\mathcal{O}_{x_n}})$ of curves.

Next, due to the construction of Υ and res , the curve $H(\gamma_{\mathcal{O}_{Y_n}})$ on T_n (resp. $H(\gamma_{\mathcal{O}_{x_i}})$) corresponds to the object $H_{\mathcal{O}_G}(\mathcal{O}_{Y_n}) \in D^b(Y_n)$ (resp. $H_{\mathcal{O}_G}(\mathcal{O}_{x_i})$) under the third fact.

Finally, these correspondences and the third fact enable us to compute some intersection numbers between $H(\gamma_{\mathcal{O}_{Y_n}})$ (or $H(\gamma_{\mathcal{O}_{x_i}})$) and other curves using basic computations in homological algebra. We can collect enough intersection numbers to determine the curves $H(\gamma_{\mathcal{O}_{Y_n}})$ and $H(\gamma_{\mathcal{O}_{x_i}})$, and this finishes the proof. \square

References

- [HKK17] F. Haiden, L. Katzarkov, and M. Kontsevich, *Flat surfaces and stability structures*, Publ. Math. Inst. Hautes Études Sci. **126** (2017), 247–318. MR 3735868
- [HT06] Daniel Huybrechts and Richard Thomas, *\mathbb{P} -objects and autoequivalences of derived categories*, Math. Res. Lett. **13** (2006), no. 1, 87–98. MR 2200048
- [LP17] Yankı Lekili and Alexander Polishchuk, *Arithmetic mirror symmetry for genus 1 curves with n marked points*, Selecta Math. (N.S.) **23** (2017), no. 3, 1851–1907. MR 3663596
- [Opp20] Sebastian Oppen, *Spherical objects, transitivity and auto-equivalences of Kodaira cycles via gentle algebras*, arXiv e-prints (2020), arXiv:2011.08288.
- [Ueh16] Hokuto Uehara, *Autoequivalences of derived categories of elliptic surfaces with non-zero Kodaira dimension*, Algebr. Geom. **3** (2016), no. 5, 543–577. MR 3568337