

# Half-spherical twists on derived categories of coherent sheaves

based on arXiv:2302.12501

Hayato Arai

Graduate School of Mathematical Sciences, The University of Tokyo

email : hayato@ms.u-tokyo.ac.jp


## 1 Introduction

### 1.1 Mirror symmetry for singular fibers of type $I_n$ of elliptic surfaces

Let  $\pi: S \rightarrow C$  be a relatively minimal, smooth projective elliptic surface. The possible singular fibers of  $\pi$  are classified by Kodaira and Néron. Among them, the singular fiber of type  $I_n$  is the cyclic configuration of  $n$  smooth rational curves.

Lekili and Polishchuk [LP17] established mirror symmetry between type  $I_n$  singular fiber  $Y_n$  and the  $n$ -punctured torus  $T_n$ , i.e. they showed that the derived category  $D^b(Y_n)$  of coherent sheaves on  $Y_n$  and the wrapped Fukaya category  $D^\pi(\mathcal{W}(T_n))$  of  $T_n$  are equivalent.

For example, the equivalence includes the following correspondence of objects:



Combining this with the theory of topological Fukaya categories of surfaces by Haiden, Katzarkov, and Kontsevich [HKK17], Opper [Opp20] described the autoequivalence group of  $D^b(Y_n)$  with the following exact sequence:

$$1 \rightarrow (\mathbb{C}^\times)^n \times \mathbb{Z}[1] \times \text{Pic}^0(Y_n) \rightarrow \text{Auteq } D^b(Y_n) \xrightarrow{\Upsilon} \text{MCG}(T_n) \rightarrow 1. \quad (1)$$

Here  $\text{MCG}(T_n)$  denotes the mapping class group of  $T_n$  and the morphism  $\Upsilon$  is induced by the equivalence  $D^b(C_n) \simeq D^\pi(\mathcal{W}(T_n))$ .

### 1.2 Autoequivalences of elliptic surfaces

Uehara [Ueh16] gave the following description of the autoequivalence group  $\text{Auteq } D^b(S)$  of  $D^b(S)$ :

**Theorem 1.1.**

- $S$  has non-zero Kodaira dimension
- all singular fibers of  $\pi$  are non-multiple and of type  $I_n$ ,  $n \geq 2$
- $B = \langle T_{\mathcal{O}_G(a)} \mid G \subset S : \text{an irreducible component of a singular fiber, } a \in \mathbb{Z} \rangle$  : the subgroup of  $\text{Auteq } D^b(S)$  generated by twist functors  $T_{\mathcal{O}_G(a)}$ , where  $\mathcal{O}_G(a)$  is the line bundle of degree  $a$  on  $G \simeq \mathbb{P}^1$

Then there is the exact sequence

$$1 \rightarrow \langle B, (-) \otimes \mathcal{O}_S(D) \mid D.F = 0, F \text{ is a fiber} \rangle \rtimes \text{Aut } S \times \mathbb{Z}[2] \quad (2)$$

$$\rightarrow \text{Auteq } D^b(S) \rightarrow \text{SL}(2, \mathbb{Z}). \quad (3)$$

### 1.3 Main results

- (1) In a general setting, we construct a “restriction” morphism  $B \rightarrow \text{Auteq } D^b(F)$  for each fiber  $F$ , which is nontrivial if  $F$  is reducible.
- (2) Combining with mirror symmetry for the singular fibers of type  $I_n$ , we described the group  $B$  in terms of the mapping class group of the  $n$ -punctured torus. Then there exists the exact sequence

$$1 \rightarrow \langle (-) \otimes \mathcal{O}_S(Y_{n_j}) \mid j = 1, \dots, m \rangle \rightarrow B \xrightarrow{r} \prod_{j=1}^m \text{MCG}(T_{n_j}), \quad (4)$$

where  $Y_{n_j}$  is the singular fiber of type  $I_{n_j}$ .

- (3) For  $G \subset Y_{n_j}$ ,  $a \in \mathbb{Z}$ , and the curve  $\gamma_{\mathcal{O}_G(a)}$  on  $T_{n_j}$  corresponding to  $\mathcal{O}_G(a)$  under the equivalence  $D^b(C_n) \simeq D^\pi(\mathcal{W}(T_n))$ , the twist functor  $T_{\mathcal{O}_G(a)}$  is mapped to the half twist along  $\gamma_{\mathcal{O}_G(a)}$ .
- (4) The image of  $r$  is generated by the half twists along the finite number of curves  $\{\gamma_{\mathcal{O}_G}, \gamma_{\mathcal{O}_G(-1)} \mid G \subset Y_{n_j} : \text{an irreducible component, } 1 \leq j \leq m\}$ .




Figure 1: The half twist along the curve  $\gamma$ .

## 2 Sketch of the proof

### 2.1 Result (1)

The result (1) is a consequence of the following theorem which generalizes the result of [HT06].

**Theorem 2.1.**

- $\pi: X \rightarrow T$  : a flat morphism between smooth quasi-projective varieties
- $i: Y = \pi^{-1}(0) \hookrightarrow X$  : the fiber at  $0 \in T$
- $E \in D^b(Y)$  : an object such that  $i_*E \in D^b(X)$  is a spherical object
- $T_{i_*E} \in \text{Auteq } D^b(X)$  : the corresponding twist functor

Then there is a unique  $H_E \in \text{Auteq } D^b(Y)$  which makes the following diagram commutative:

$$\begin{array}{ccc} D^b(Y) & \xrightarrow{i_*} & D^b(X) \\ H_E \downarrow & & \downarrow T_{i_*E} \\ D^b(Y) & \xrightarrow{i_*} & D^b(X). \end{array} \quad (5)$$

### 2.2 Result (3)

- (a) There are correspondences between indecomposable objects of  $D^b(Y_n)$  and homotopy classes of curves on  $T_n$  (+ additional data), and between dimensions of Hom-spaces in  $D^b(Y_n)$  and intersection numbers of curves on  $T_n$ .
- (b) An element of  $\text{MCG}(T_n)$  is determined by its action on  $\pi_1(T_n)$  (Dehn–Nielsen–Baer theorem).

## References

- [HKK17] F. Haiden, L. Katzarkov, and M. Kontsevich, *Flat surfaces and stability structures*, Publ. Math. Inst. Hautes Études Sci. **126** (2017), 247–318. MR 3735868
- [HT06] Daniel Huybrechts and Richard Thomas,  *$\mathbb{P}$ -objects and autoequivalences of derived categories*, Math. Res. Lett. **13** (2006), no. 1, 87–98. MR 2200048
- [LP17] Yankı Lekili and Alexander Polishchuk, *Arithmetic mirror symmetry for genus 1 curves with  $n$  marked points*, Selecta Math. (N.S.) **23** (2017), no. 3, 1851–1907. MR 3663596
- [Opp20] Sebastian Opper, *Spherical objects, transitivity and auto-equivalences of Kodaira cycles via gentle algebras*, arXiv e-prints (2020), arXiv:2011.08288.
- [Ueh16] Hokuto Uehara, *Autoequivalences of derived categories of elliptic surfaces with non-zero Kodaira dimension*, Algebr. Geom. **3** (2016), no. 5, 543–577. MR 3568337