Document 0932

```
nt of blood surgical patients can donate and store before surgery can be ev-w1037f1 ev-w3190f1
                                                                                                                                                                                                                                                                                 before surgery can be increased by the new genetically engineered drug , EPO .
     APPS.member

APPS.member

APPS.member

APPS.member

APPS.member

APPS.member

ACT-arg

ACT-ar
   [20932003]
A genetically engineered version of the human protein developed ev-w1152f1

A genetically engineered version of the human protein developed ev-w918f3

A genetically engineered version of the human protein developed ev-w918f3

A genetically engineered version of the human protein developed ev-w918f3

A genetically engineered version of the human protein developed ev-w918f3
                                                                                                                                                                                                                                                                                                                                                                                 _ _ _ ev-w1943f1 _ _ _
A competing version of EPO is being developed by Genetics Institute Inc. in Cambridge , Mass .
   The drug is being used primarily to treat anemias .
 [20932006]
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  PAT-arg
                                                                                                                                                                                                                                                                                                                               ne , involved giving injections of Amgen 's EPO to 23 patients who wanted to store units ev-w1756f1 ev-w1465f6
     The patients began receiving ev-w2542f1 ev-w2542f1  

EPO injections about a month before their scheduled surgery .
                                                                                                             ACT-arg
                                                                                                                                       (TWHEN)
     They then began donating blood twice a week , receiving an EPO injection each time .
 The EPO-treated patients donated an average of 5.4 units of blood
                                                                                                                                                                                                                                                                                                                                   d with only 4.1 units donated by a similar group of surgical patients who received a placebo injection .
                                                                                                                                                                                                                                                                                                                                                                                                                      REG (RAT.urg) (MAT) (CONJ.member) (REG) (R
                                                                                                                                                                                                                                                                  was 41 % higher per doi
```

1		