


Übungsblatt 3

Montag, 16. November 2020

16:20

①

4 neighborhood:



only edges

(17, 42) and (289, 68)

• you need 2 squares to move diagonal on x and y axis



• y-axis: $68 - 42 = 26$ } we need to move 26 upwards in y

• x-axis: $289 - 17 = 272$ } and 272 in x direction

• (1) We have 26 diagonal steps, which will be put together with $272 - 26$ steps in x axis (order doesn't matter)

$\Rightarrow (272 - 26) + 2 \cdot 26 = 298$ steps or pixels in distance ✓

8 neighborhood:



• We need only 1 pixel for diagonal movement

$\Rightarrow (272 - 26) + 26 = 272$ steps/pixels in distance ✓

euclidean distance

$$d((17, 42), (289, 68)) = \sqrt{(17 - 289)^2 + (42 - 68)^2} \approx 273,23 \checkmark$$

3

②

	0	1	2	3
0	R = 40	G = 80	R = 60	G = 100
1	G = 80	r = ? g = ? B = 100	r = ? G = 100 b = ?	B = 40
2	R = 40	r = ? G = 100 b = ?	R = 100 g = ? b = ?	G = 25
3	G = 25	B = 20	G = 75	B = 20

$$r_{11} = (R_{00} + R_{02} + R_{20} + R_{22}) / 4 = 60 \checkmark$$

$$g_{11} = (G_{01} + G_{12} + G_{21} + G_{10}) / 4 = 90 \checkmark$$

$$r_{12} = (R_{02} + R_{22}) / 2 = 80 \checkmark$$

$$b_{12} = (B_{11} + B_{13}) / 2 = 70 \checkmark$$

$$r_{21} = (R_{20} + R_{22}) / 2 = 70 \checkmark$$

$$b_{21} = (B_{11} + B_{31}) / 2 = 60 \checkmark$$

$$g_{22} = (G_{12} + G_{23} + G_{32} + G_{21})/4 = 15 //$$

$$b_{22} = (B_{11} + B_{13} + B_{31} + B_{33})/4 = 45 //$$

③

$$I^E = [1 \ 2 \ 1] \quad \text{and} \quad I^{\tilde{F}_1} = I^{\tilde{F}_2} [-1 \ 0 \ 1]$$

Widerspruchsbeweis:

Beh: Cross correlation is associative

$$\text{Bew: } I^{E^T} \otimes (I^{\tilde{F}_1^T} \otimes I^{\tilde{F}_2^T}) = (I^{E^T} \otimes I^{\tilde{F}_1^T}) \otimes I^{\tilde{F}_2^T}$$

$$\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \otimes \left(\begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right) = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} -1 \cdot -1 + 0 \cdot 0 + 1 \cdot 1 \\ -1 \cdot -1 + 0 \cdot 0 + 1 \cdot 1 \\ -1 \cdot -1 + 0 \cdot 0 + 1 \cdot 1 \end{pmatrix} \quad \text{f}$$

$$= \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \cdot 2 + 2 \cdot 2 + 1 \cdot 2 \\ 1 \cdot 2 + 2 \cdot 2 + 1 \cdot 2 \\ 1 \cdot 2 + 2 \cdot 2 + 1 \cdot 2 \end{pmatrix} = \begin{pmatrix} 8 \\ 8 \\ 8 \end{pmatrix} \quad \text{f}$$

$$\left(\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right) \otimes \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} = 1 \cdot -1 + 2 \cdot 0 + 1 \cdot 1 \quad \approx \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$= \begin{pmatrix} 2 \cdot -1 + 2 \cdot 0 + 2 \cdot 1 \\ 2 \cdot -1 + 2 \cdot 0 + 2 \cdot 1 \\ 2 \cdot -1 + 2 \cdot 0 + 2 \cdot 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \quad \text{f}$$

$$\Rightarrow \begin{pmatrix} 8 \\ 8 \\ 8 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \quad \text{⚡}$$

□

0