

Contents

1	Introduction	1
1.1	Fermions, Fermi surfaces and quasiparticles	2
1.2	The SYK model.	5
1.2.1	The SYK self consistent equations	6
1.3	Other versions of the SYK used in this thesis	8
1.3.1	Complex SYK	8
1.3.2	Yukawa-SYK model	9
1.3.3	b-SYK	12
1.4	Holographic dual of the SYK model and $N - AdS_2$ spacetime	13
1.5	Graphene and its bilayers	19
1.6	The concept of a higher order van Hove singularity	22
1.7	Enhanced interactions near van hove points - towards Twisted Bilayer graphene	23
1.8	This thesis	28
1.8.1	Chapter 2 - Chaos in the bipartite Sachdev-Ye-Kitaev model	28
1.8.2	Chapter 3 - Wormholes in the Yukawa-Sachdev-Ye-Kitaev model	29
1.8.3	Chapter 4 - The Kondo effect in Twisted bilayer graphene	29
2	Lyapunov Exponents in the bipartite SYK model	31
2.1	Introduction.	32
2.2	Model and methods.	33
2.2.1	The bipartite SYK model	33
2.2.2	Schwinger-Dyson equations	34
2.2.3	Real time formalism	37
2.3	The four-point function.	38
2.3.1	Conformal limit	41
2.3.2	Numerical analysis for weak and intermediate coupling	43

2.4	Results	43
2.4.1	Analytics and numerics	43
2.4.2	Discussion and Conclusion	46
2.5	Appendix	48
2.5.1	Mathematical consistency of the Lyapunov ansatz	48
2.5.2	Recovery of the maximal Lyapunov exponent of the regular SYK	49
2.5.3	Finite-size dependence of two-point functions and Lyapunov exponents	50
3	A BCS-Josephson Wormhole in coupled superconducting Yukawa-SYK metals	51
3.1	Introduction.	52
3.2	Two Yukawa-SYK models with a tunneling contact with shared disorder.	56
3.3	The YSYK-TFD/wormhole for the metallic normal state.	58
3.4	Inclusion of superconductivity: the BCS Josephson wormhole	65
3.5	A standard Josephson current	70
3.6	Conclusion	72
3.7	Appendix	75
3.7.1	Derivation of the effective action	75
3.7.2	Conformal limit of the one sided Schwinger-Dyson equations	79
3.7.3	Numerical details	80
4	Kondo effect in Twisted Bilayer graphene	81
4.1	Introduction.	82
4.2	Van Hove singularities in the Bistritzer-MacDonald model	84
4.2.1	Particle-hole symmetric Bistritzer-MacDonald model	85
4.2.2	Characterization of van Hove singularities in the particle-hole symmetric BM model	86
4.3	The Kondo problem.	89
4.3.1	Poor man's scaling approach	91
4.3.2	Numerical Renormalization Group	92
4.3.3	Observables	94
4.3.4	Limiting cases of the Kondo problem	95

4.4	Results and discussion	99
4.5	Conclusions	104
4.6	Appendix	106
4.6.1	Locating saddle point positions in the particle-hole symmetric BM model	106
4.6.2	Mapping from TBG density of states to the NRG Wilson chain.	107
4.6.3	NRG calculations for an impurity coupled to a conventional log-vHs host	108
	References	109
	Samenvatting	129
	Summary	131
	List of Publications	133
	Curriculum Vitae	135
	Acknowledgements	137

Samenvatting

Vaste stoffen bestaan uit elektronen en ionen. De meeste vaste stoffen kunnen effectief worden gemodelleerd met een theorie waarin de beweging van de elektronen allemaal als onafhankelijk van elkaar kan worden beschouwd. Dit proefschrift houdt zich bezig met enkele gevallen waarin deze beschrijving onvermijdelijk faalt, en meerdere-deeltjes-interacties de lage-energiefysica domineren. Dergelijke systemen zijn veel moeilijker te bestuderen. Soms kan vooruitgang worden geboekt met behulp van geavanceerde numerieke technieken, of door creatief geconstrueerde, precies oplosbare speelgoedmodellen.

Dit laatste is de benadering van het Sachdev-Ye-Kitaev (SYK)-model, dat een kwantumdot beschrijft met een groot aantal fermionen erop, die allemaal met elkaar interacteren door middel van een gerandomiseerde interactie met vier fermionen. De eenvoud van het model verhult een overvloed aan exotische verschijnselen die voortkomen uit een emergente conforme symmetrie in het infrarood, met eclectische toepassingen van vreemde metalen tot zwarte gaten. Door zogenaamde holografische dualiteit kan de lage-energietheorie van het model, die voortkomt uit het verbreken van de bovengenoemde conforme symmetrie, in kaart worden gebracht via verstoringen van een bijna AdS_2 ruimtetijd.

Eén van die verschijnselen staat bekend als maximale chaos. Het SYK-model verzadigt een grens aan de snelheid waarmee in een kwantumsysteem verstoringen kunnen dissiperen in zijn vrijheidsgraden. Dit kenmerk blijft bestaan, zelfs als men bepaalde gewijzigde versies van het SYK-model construeert, zoals het bipartiete SYK-model. De mate van chaos verandert niet, zelfs niet als de schaaldimensies van de samenstellende fermionen worden veranderd, zoals beschreven in hoofdstuk 2. De eerste subleidende correctie van de chaos exponent blijft ook onafhankelijk te zijn van de schaaldimensie.

Wanneer twee identieke kopieën van het SYK-model aan elkaar worden gekoppeld door een tunnelinteractie, bij lage temperatuur beschrijft het holografische dual de ruimtetijd van een wormgat. Een op een dergelijke manier gemaakt Josephson-contact kan worden bestudeerd met behulp van de SYK-versie van het elektron-fonon-gekoppelde systeem, dat bekend staat als het Yukawa SYK-model. Supergeleiding in het Yukawa SYK-model vindt plaats bij lage temperaturen als het systeem tijdinversie niet schendt. In hoofdstuk 3 wordt aangetoond dat de niet-supergeleidende

toestand de correcte niet-analytische afhankelijkheid van de "gap" vertoont van de sterkte van de tunnelinteractie, kenmerkend voor de wormgattoestand. Er wordt aangetoond dat de supergeleidende toestand effectief kan worden beschreven door een twee-vloeistoffenmodel, waarbij de Cooperparen onafhankelijk aan beide zijden leven, terwijl excitaties van ongebonden elektronen nog steeds de wormgattoestand vormen.

Als tegenpool van de hyperbolische geometrie van de AdS_2 ruimte, hebben we in het tweede deel van dit proefschrift een systeem beschouwd waarvan het niet-interagerende fermi-oppervlak is samengesteld uit een familie van hyperbolen. In gedraaid dubbellaags grafeen voor kleine draaihoeken is dit te zien in de vorm van van Hove-singulariteiten, die precies onder de magische hoek een hogere orde krijgen. In hoofdstuk 4 wordt aangetoond dat een kenmerk voor het waarnemen van het effect ervan in een realistische fysieke omgeving de reactie is op een magnetische onzuiverheid. De onzuiverheidsentropie vertoont duidelijke kenmerken nabij de magische hoek, en toont stromingen naar de verschillende soorten vaste punten die beschikbaar zijn in de faseruimte van het systeem. Totdat de magische hoek is bereikt, werkt het bestaan van de Dirac-kegel bij de laagste energieën als een afschermingsmiddel voor Kondo-screening, wat zich manifesteert als een entropie van $\log 2$ per onzuiverheid. Bij de magische hoek domineert de verschijning van de hogere orde Van Hove-singulariteit de bandstructuur, en dit leidt tot een versterkt Kondo-effect tot de laagst mogelijke energieschalen.

Summary

Solids are composed of electrons and ions. Most solids can be effectively modeled by a theory in which the motion of the electrons can all be considered independent to each other. This thesis is concerned with some cases where that description fails, and many body interactions dominate the low energy physics. Such systems are far more difficult to study, and sometimes progress can be made either with sophisticated numerical tools, or by creatively constructed exactly solvable toy models.

The latter is the approach taken by the Sachdev-Ye-Kitaev (SYK) model, which describes a quantum dot with a large number of fermions on it, all interacting with each other by means of a randomized four-fermion interaction. The simplicity of the model belies a plethora of exotic phenomena that arise from an emergent conformal symmetry in the infrared, with eclectic applications from strange metals to black holes. Through the holographic duality, the low energy theory of the model arising from the breaking of the aforementioned conformal symmetry can be mapped to perturbations of a nearly AdS_2 spacetime.

One such phenomenon is known as maximal chaos. The SYK model saturates a bound on the rate at which a quantum system can dissipate perturbations into its degrees of freedom. This feature persists even when one constructs certain modified versions of the SYK model, such as the bipartite SYK model, and the rate of chaos doesn't change even when the scaling dimensions of the constituent fermions are changed as is described in chapter 2. The first subleading correction to the chaos exponent is also found to be independent of the scaling dimension.

When two identical copies of the SYK model are coupled together by a tunneling interaction, its holographic dual describes a wormhole spacetime at low temperature. A Josephson contact made in such a way can be studied using the SYK version of the electron-phonon coupled system, which is known as the Yukawa SYK model. Superconductivity in the Yukawa SYK model occurs at low temperatures if time reversal symmetry is present. In chapter 3, it was shown that the non-superconducting state shows the correct non-analytic dependence of the gap on the strength of the tunneling interaction, characteristic of the wormhole state. The superconducting state was shown to be effectively described by a two-fluid model, where the Cooper pairs live independently on the two sides, while single electron excitations still form the wormhole state.

In juxtaposition to the hyperbolic geometry of the AdS_2 space, we consider a system whose non-interacting fermi surface was composed of a family of hyperbolae in the second part of this thesis. In twisted bilayer graphene for small twists, this is seen in the form of van Hove singularities, which turn higher order at precisely the magic angle. In chapter 4, a probe for sensing its effect in a realistic physical setting was shown to be the response of a magnetic impurity. The impurity entropy shows distinct signatures near the magic angle, showing flows to the different kinds of fixed points available in the phase space of the system. Until the magic angle is reached, the existence of the Dirac cone at the lowest energies acts as a deterrent for Kondo screening, which manifests itself as a $\log 2$ entropy per impurity. At the magic angle, the appearance of the higher order van Hove singularity dominates the band structure, and it leads to an enhanced Kondo effect until the lowest energy scales possible.

List of Publications

1. “ Lyapunov exponents in a Sachdev-Ye-Kitaev-type model with population imbalance in the conformal limit and beyond ”,
A. S. Shankar, M.Fremling, S. Plugge and L. Fritz
Phys. Rev. D, 108, 094039 (2023).
2. “ Kondo effect in twisted bilayer graphene ”,
A. S. Shankar, D. O. Oriekhov, Andrew K. Mitchell, and L. Fritz
Phys. Rev. B, 107, 245102 (2023).

Curriculum Vitae

I was born on September 26th, 1996 in Chennai, a big city in the south of India. After finishing high school, I went to West Bengal for a five-year integrated Bachelor-Master program in Physics at the Indian Institute of Technology, Kharagpur.

During this period, I took courses in several areas of physics and engineering, including high energy physics and gravitational physics, with my favorite being condensed matter and statistical physics. I was also fortunate to get an early start at doing my own research by being selected for several internships throughout my bachelor and master studies. With Prof. Pinaki Sengupta at NTU Singapore, I worked on writing my first quantum Monte Carlo code to investigate the Heisenberg XXZ model on ladder-like systems. I then worked briefly with Prof. Barbara Terhal and Prof. Fabian Hassler on DMRG simulations at RTWH aachen, and then on numerical and analytical studies on the Anderson model on the Bethe lattice with Prof. Antonello Scardicchio at the ICTP, Trieste, looking for signatures of anomalous diffusion. I concluded my master studies with a thesis titled "Disordered Superconductors", working with Prof. Sudhansu Sekhar Mandal.

I started my PhD at Leiden university in the Quantum matter group of Prof. Jan Zaanen and Prof. Koenraad Schalm in August 2019. After a short stint there, I moved to the group of Prof. Vadim Cheianov in April 2021. It was also at this time that I started working very closely with Prof. Lars Fritz and his group at Utrecht University.

As the early years of my PhD programme were marred by the coronavirus pandemic, I attended several condensed matter schools online, including editions of the Princeton PSSC summer school and the Maglab winter school. After the pandemic ended, I was able to present my work at several schools and conferences, including poster presentations in Trieste, Italy, at the ICTP conference "From quantum criticality to flat bands", at the school "Emergence of quantum phases in novel materials", at CSIC, Madrid, Spain, at the Veldhoven Physics conference in the Netherlands, and an online talk at the APS March meeting in 2023, among others. I also had the opportunity to TA several undergraduate and graduate courses, in particular to write a set of extensive lecture notes on Information Geometry with Prof. Subodh Patil.

After the defense of my PhD, I will move on to begin a Postdoctoral position at the ICTP in Trieste.

Acknowledgements

First and foremost, I would like to thank my promotor Prof. Vadim Cheianov and my co-promotor Prof. Koenraad Schalm for giving me the independence to carve my own path, even though it might have been slightly unconventional. I'm very thankful to Vadim for his infinite patience with my questions and for addressing them sincerely even if they might have been trivial at times. To Koenraad I'm thankful for teaching me to improve my scientific rigor and for the importance he taught me to place on communicating my results well. Thank you very much.

I'm most greatly indebted to Prof. Lars Fritz, his contribution to this work goes far beyond his supervision of the two published papers that form parts of this thesis. I'm exceedingly grateful to him for welcoming me into his group, and for his wonderful mentorship over the years. I'm also thankful to Lars for introducing me to Prof. Andrew K. Mitchell, who I greatly enjoyed collaborating with. I hope this collaboration will last for quite some time to come.

I'm also very grateful to the PhD students and postdocs that I worked with in Leiden and in Utrecht - Dima Oriekhov, Mikael Fremling and Stephan Plugge. I am also particularly grateful to the several intense discussions I had with Nicolas Chagnet over physics and on numerics in general. I would also like to acknowledge master students Alina Melissakis and Nina Doerfler whose theses I helped co-supervise- I learned quite a lot from this experience.

I extend thanks next to the several friends I made along the way. I will always have a special place for my covid-housemate Alicja Dutkiewicz, and my post-covid housemate Dimitrios Krommydas, with whom I felt that family was always close by.

I also thank Floris Balm, Luca Barbera, Alexander Barnaveli, Norman Bluemel, Meike Bos, Aleksandar Bukva, Zhenya Cheipesh, Jose Collado, Tjeerd de Leede, David Dechant, Emanuele di Salvio, Alvaro Donis-Vela, Patrick Emonts, Inci Erbilen, Matthias Flor, Ireth Garcia-Aguilar, Alice Garoffolo, Peter George, Joren Harms, Yaroslav Herasymenko, Ludwig Hoffman, Stef Koenis, Ilse Kuijf, Gal Lemut, Samadarshi Maity, Savvas Malikis, Simon Marshall, Kirsten Martens, Kamil Matuszelski, Anna Negro, Vladimir Ohanesjan, Maialen Ortego-Larrazabal, Michal Pacholski, Adrian Perez, Stefano Polla, Kshiti Sneha Rai, Midas Ratsma, Philippe Sabella-Garnier, Katerina Saiti, Kolya Semenyakin, Fabian Sies, Tereza Vakhtel, Vova Zakharov and Masha Zakharova for their warm company, and for discussions both

physics and otherwise. I hope I didn't miss anyone out!

I'm also very thankful to my friends from India - Mohit Chajjed, Mohit Dhariwal, Muthu Ganesh, Philip Jacob, Viraj Jangam, Daksh Khanter, Pranai Kumar, Praveen PR, Saloni Priya, Suvrat Rao, Darvin Sai Teja Reddy, Baibhaw Kumar Singh, Tulja Varun for keeping me company with frequent phone calls, especially during the pandemic.

Lastly, I thank my family - Amma, Appa, Aparna and Paati-Thatha for your support, and for being understanding of my absence from home.