NEKHOROSHEV STABILITY FOR QUASI-CONVEX HAMILTONIANS: BLUEPRINT (TRACK N)

SCOTT ARMSTRONG AND ZOHAR KOMARGODSKI

Abstract. A proof plan, step by step, for the three frozen statements of track N of the KAM formalization: the exponential stability estimate for strictly quasi-convex nearly integrable Hamiltonians with a complex Hessian bound (N-HEAD-C, after Pöschel 1993, Theorem 1), the exact integrable dynamics for zero perturbation (N-ZERO) and the nonsingular isoenergetic derivative (N-QC). Every result is labelled and cites its pinned source. This version contains Part 0 only; the plan of the remaining parts is in `PLAN.md`.

Contents

1. Conventions, definitions and the frozen statements

This part fixes notation and restates, in mathematical form, the three frozen theorems of track N (owner approval D20, 27 October 2026) and the eleven frozen definitions of `KAM/Frozen/Nekhoroshev/Defs/` together with the five track-K definitions they use. The Lean declarations are authoritative; the statements below are transcriptions of them, with the Lean names recorded under each. N-HEAD with real bounds only is *not* frozen (source gap F-N1, D17/D18/D20); the frozen headline is N-HEAD-C. The additions of D22 (N-PARTIAL-CONVEX, N-AVERAGING) are not yet frozen and are not part of this part. Results of track K's blueprint are cited as [K:`bp:p1:...`].

1.1. Conventions.

- $n \in \mathbb{N}$ is the number of degrees of freedom; N-HEAD-C assumes $n \geq 3$ (D17), N-ZERO and N-QC are stated for every n. A phase point is $z = (\theta, I)$ with angles θ and actions I, in this order.
- Ambient spaces are those of the Lean text: $\mathbb{R}^n = (\texttt{Fin}\ n \to \mathbb{R})$, $\mathbb{C}^n = (\texttt{Fin}\ n \to \mathbb{C})$, with the max norms $|v| = \max_i |v_i|$ on both. The embedding $\mathbb{R}^n \subset \mathbb{C}^n$ is componentwise. The norm $\|\cdot\|$ in every frozen statement is the max norm; on the phase space $\mathbb{R}^n \times \mathbb{R}^n$ it is $\|(\theta, I)\| = \max(|\theta|, |I|)$ (Lean product norm). The only Euclidean structure is the dot product $\sum_i \omega_i v_i$ used for orthogonality in quasi-convexity; Euclidean norms used later for comparison with the sources come with explicit inequalities ($|v| \leq \|v\|_2 \leq \sqrt{n}|v|$). For $n = 0$ all these spaces are $\{0\}$ and the norm is 0.
- Operator norms of linear and multilinear maps are the induced norms of these max norms. In particular $\|D^2 h(z)\|$ for complex h is the norm of $D^2 h(z) \in \mathcal{L}(\mathbb{C}^n, \mathcal{L}(\mathbb{C}^n, \mathbb{C}))$ (Lean `fderiv` applied twice), i.e. $\sup_{|u|,|v|\leq 1} |D^2 h(z)[u, v]|$.
- Angles are *not* reduced modulo $\mathbb{Z}^n$ in the statements: $\theta \in \mathbb{R}^n$ or $\theta \in T_s \subset \mathbb{C}^n$, and 1-periodicity is a hypothesis on f (Definition **??**). So the torus is $\mathbb{R}^n/\mathbb{Z}^n$ and the Fourier characters are $e^{2\pi i k \cdot \theta}$ as in track K ([K:`bp:p1:decay`], [K:`bp:p1:summation`]). Pöschel's torus is $\mathbb{R}^n/2\pi\mathbb{Z}^n$ and his norms are Euclidean; the conversion is a separate step of the proof (Part 1 of `PLAN.md`).
- The complex strip is $T_s = \{\theta \in \mathbb{C}^n : |\operatorname{Im} \theta_i| < s\ \forall i\}$ (Lean `KAM.complexStrip`), as in track K.

1.2. **Track-K definitions used.**

Definition 1.1 (Complex action neighbourhood)**.** For $B \subset \mathbb{R}^n$ and $r \in \mathbb{R}$, $B_r = \{I \in \mathbb{C}^n : \exists J \in B,\ |I - J| < r\}$ (max norm, J viewed in $\mathbb{C}^n$). Thus $B_r = \emptyset$ for $r \le 0$, and for B open B_r is open.
Frozen Lean declaration: `KAM.actionNbhd`
Source: BM11 p. 4 (D_s); track K, [K:`bp:p0:domains`].

Definition 1.2 (Frequency map)**.** For $h : \mathbb{C}^n \to \mathbb{C}$ and $I \in \mathbb{R}^n$, $\omega_h(I) = \nabla h(I) \in \mathbb{R}^n$ is the vector with $\omega_h(I)_i = D\big(J \mapsto \operatorname{Re} h(J)\big)(I)[e_i]$, the real partial derivatives of $\operatorname{Re} h$ on $\mathbb{R}^n$ at I. It is 0 if $\operatorname{Re} h|_{\mathbb{R}^n}$ is not differentiable at I (Lean totality, Remark **??**).
Frozen Lean declaration: `KAM.frequencyMap`
Source: BM11 p. 1; Pos93 p. 190 ($\omega(I) = \partial_I h(I)$).

Definition 1.3 (Hamiltonian vector field)**.** For $H : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ and $z = (\theta, I)$, $X_H(z) = \big((\partial_{I_i} H(z))_i,\ (-\partial_{\theta_i} H(z))_i\big)$, where $\partial_{I_i} H(z) = DH(z)[(0, e_i)]$ and $\partial_{\theta_i} H(z) = DH(z)[(e_i, 0)]$ are the Fréchet derivatives in the Lean sense (0 if H is not differentiable at z).
Frozen Lean declaration: `KAM.hamiltonianVectorField`
Source: BM11 p. 1; track K, [K:`bp:p0:hamiltonian`].

Definition 1.4 (Real Hamiltonian)**.** For $h : \mathbb{C}^n \to \mathbb{C}$, $f : \mathbb{C}^n \times \mathbb{C}^n \to \mathbb{C}$ and real $z = (\theta, I)$, $H_{h,f}(z) = \operatorname{Re}\big(h(I) + f(\theta, I)\big)$, with θ, I viewed in $\mathbb{C}^n$.
Frozen Lean declaration: `KAM.realHamiltonian`
Source: BM11 p. 5 ($H = h + f$).

Remark 1.5 (Strip and neighbourhood notation). `KAM.complexStrip` $n\, s = T_s$ as above; for $n > 0$ it is empty when $s \le 0$, and for $n = 0$ it is the whole (one-point) space $\mathbb{C}^0$ for every s. No hypothesis $n > 0$ is added to N-ZERO or N-QC.
Frozen Lean declaration: `KAM.complexStrip`

1.3. **Frozen track-N definitions.**

Definition 1.6 (Action ball)**.** $B(0, R) = \{I \in \mathbb{R}^n : |I| < R\}$, the open ball of the max norm; written `actionBall` $n\, R$.
Frozen Lean declaration: `KAM.Nekhoroshev.actionBall`
Source: BM11 p. 4.

Definition 1.7 (Complex domain)**.** $D_s(R) = T_s \times B(0, R)_s \subset \mathbb{C}^n \times \mathbb{C}^n$; `complexDomain` $n\, R\, s$. It is the period-one lift of BM11's D_s, with a single width s for angles and actions. It is open.
Frozen Lean declaration: `KAM.Nekhoroshev.complexDomain`
Source: BM11 p. 4 (D_s); ours for the lift.

Definition 1.8 (Real action domain)**.** $\mathcal{B}_s(R) = \{I \in \mathbb{R}^n : I \in B(0, R)_s\}$; `realActionDomain` $n\, R\, s$.
Frozen Lean declaration: `KAM.Nekhoroshev.realActionDomain`
Source: BM11 p. 4 (B_s).

Remark 1.9 (Computing $\mathcal{B}_s(R)$). Since $|I - J|$ for real I, J is the real max norm, $\mathcal{B}_s(R) = \{I \in \mathbb{R}^n : \exists J,\ |J| < R,\ |I - J| < s\} = B(0, R + s)$ for $R, s > 0$. This identity is ours and is proved from the definitions (it is not stated in the source); it shows that $B(0, R) \subset \mathcal{B}_s(R)$ and that $\mathcal{B}_s(R)$ is open.

Definition 1.10 (Real part of the integrable part)**.** For $h : \mathbb{C}^n \to \mathbb{C}$ and $I \in \mathbb{R}^n$, $h_{\mathbb{R}}(I) = \operatorname{Re} h(I)$; `realPart` h.
Frozen Lean declaration: `KAM.Nekhoroshev.realPart`
Source: BM11 p. 4 ($h : B_s \to \mathbb{R}$).

Definition 1.11 (Analytic integrable part, $h \in A_s(D)$)**.** `IsAnalyticIntegrable` $R\, s\, h$ holds if

(i) h is complex differentiable on $B(0, R)_s$;
(ii) h is continuous on $\overline{B(0, R)_s}$ (closure in $\mathbb{C}^n$);

(iii) there is $A \geq 0$ with $|h(I)| \leq A$ for all $I \in B(0,R)_s$;
(iv) $\operatorname{Im} h(I) = 0$ for all $I \in \mathcal{B}_s(R)$;
(v) $h_{\mathbb{R}}$ is of class C^3 on $\mathcal{B}_s(R)$ (real Fréchet sense; the set is open).

No numerical bound on h is fixed. Clause (v) is redundant by holomorphy ([K:bp:p1:holomorphic-regularity], [K:bp:p1:real-restriction]) and records the regularity that guards every real derivative.

Frozen Lean declaration: `KAM.Nekhoroshev.IsAnalyticIntegrable`

Source: BM11 p. 4 ($A_s(D)$, integrable case, with continuity on the closure).

Definition 1.12 (Analytic perturbation of size ε)**.** `IsAnalyticPerturbation` $R\,s\,\varepsilon\,f$ holds if $\varepsilon \geq 0$ and

(i) f is complex differentiable on $D_s(R)$ and continuous on $\overline{D_s(R)}$;
(ii) $f(\theta + k, I) = f(\theta, I)$ for all $(\theta, I) \in D_s(R)$ and $k \in \mathbb{Z}^n$;
(iii) $\operatorname{Im} f(\theta, I) = 0$ for all $\theta \in \mathbb{R}^n$ and $I \in \mathcal{B}_s(R)$;
(iv) $|f(z)| \leq \varepsilon$ for all $z \in D_s(R)$ (complex absolute value).

The size is measured by $\leq \varepsilon$, not the source's strict bound; the source is applied with 2ε. For $\varepsilon = 0$ this forces $f = 0$ on $D_s(R)$.

Frozen Lean declaration: `KAM.Nekhoroshev.IsAnalyticPerturbation`

Source: BM11 pp. 4–5 ($f \in A_s(D)$, $|f|_s < \varepsilon$); deviation: $\leq \varepsilon$ (ruling Q2, D18).

Definition 1.13 (Derivative bounds $B(M)$)**.** For $U \subset \mathbb{R}^n$, $h : \mathbb{C}^n \to \mathbb{C}$ and $M \in \mathbb{R}$, `HasDerivativeBounds` $U\,h\,M$ holds if $M > 0$, U is open, $h_{\mathbb{R}} \in C^3(U)$, and for all $I \in U$, all $k \in \{1,2,3\}$ and all $(i_1, \ldots, i_k) \in \{1, \ldots, n\}^k$,

$$\left|D^k h_{\mathbb{R}}(I)[e_{i_1}, \ldots, e_{i_k}]\right| \leq M.$$

These are all ordered coordinate partials of orders one to three; for C^3 functions they coincide with the multi-index partials $\partial^k h$ of the source (symmetry, [K:bp:p1:ext-schwarz]). The bound is on coordinate partials, not on operator norms.

Frozen Lean declaration: `KAM.Nekhoroshev.HasDerivativeBounds`

Source: BM11 p. 4, $B(M)$.

Definition 1.14 (Strict quasi-convexity $QC(m)$)**.** For $U \subset \mathbb{R}^n$, $h : \mathbb{C}^n \to \mathbb{C}$, $m \in \mathbb{R}$, `IsStrictlyQuasiConvex` $U\,h\,m$ holds if $m > 0$, U is open, $h_{\mathbb{R}} \in C^2(U)$, and for every $I \in U$: $\omega_h(I) \neq 0$ and

$$\sum_i \omega_h(I)_i v_i = 0 \implies m|v|^2 \leq D^2 h_{\mathbb{R}}(I)[v,v] \qquad (v \in \mathbb{R}^n),$$

with $|v|$ the max norm on the left and the Euclidean dot product in the hypothesis.

Frozen Lean declaration: `KAM.Nekhoroshev.IsStrictlyQuasiConvex`

Source: BM11 p. 4, $QC(m)$ (the source's norm on the right is the max norm, orthogonality is Euclidean).

Definition 1.15 (Complex Hessian bound)**.** `HasComplexHessianBound` $R\,s\,h\,M_{\mathbb{C}}$ holds if $M_{\mathbb{C}} > 0$, h is C^2 in the complex sense on $B(0,R)_s$ and $\|D^2 h(z)\| \leq M_{\mathbb{C}}$ for all $z \in B(0,R)_s$, the norm being that of $D(Dh)(z) \in \mathcal{L}(\mathbb{C}^n, \mathcal{L}(\mathbb{C}^n, \mathbb{C}))$ for the max norm. Every entry $|\partial_i \partial_j h|$ is then $\leq M_{\mathbb{C}}$, and the Euclidean operator norm of the Hessian matrix is $\leq n M_{\mathbb{C}}$. This is not the spectral radius of a complex symmetric matrix.

Frozen Lean declaration: `KAM.Nekhoroshev.HasComplexHessianBound`

Source: Pos93 p. 190, "The Hessian of the integrable hamiltonian h" (conversion by n recorded in `docs/tracks/N/sources.md`, anchor POS190-HESS).

Definition 1.16 (Solutions on a time set)**.** For $H : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$, $U \subset \mathbb{R}^n$, $J \subset \mathbb{R}$ and $z : \mathbb{R} \to \mathbb{R}^n \times \mathbb{R}^n$, `IsSolutionOn` $H\,U\,J\,z$ holds if for every $t \in J$: $z(t)_2 \in U$, H is differentiable at $z(t)$, and z is differentiable at t (two-sided, also at endpoints of J, since z is defined on $\mathbb{R}$) with $z'(t) = X_H(z(t))$.

Frozen Lean declaration: `KAM.Nekhoroshev.IsSolutionOn`
Source: BM11 Theorem 2.1 p. 5 and the maximal-solution argument p. 13.

Definition 1.17 (The integrable orbit)**.** For $h : \mathbb{C}^n \to \mathbb{C}$, $\theta_0, I_0 \in \mathbb{R}^n$ and $t \in \mathbb{R}$: $\gamma_{\theta_0,I_0}(t) = (\theta_0 + t\,\omega_h(I_0),\, I_0)$.
Frozen Lean declaration: `KAM.Nekhoroshev.integrableOrbit`
Source: BM11 p. 1 ($H(\theta, I) = h(I)$).

Remark 1.18 (Lean totality). The frozen definitions and theorems use `fderiv` (which is 0 at a point of non-differentiability), `iteratedFDeriv`, the operator norm, `rpow` ($\varepsilon^{\pm 1/(2n)}$), `Real.exp` and the max norm; there is no matrix inverse, no Bochner integral and no division by a quantity that may vanish. The guards are:

(a) In N-HEAD-C and N-ZERO, ω_h and $D^2 h_{\mathbb{R}}$ are used only on the open set $\mathcal{B}_s(R)$ where $h_{\mathbb{R}}$ is C^3 (Definitions **??**(v), **??**, **??**), so the derivatives are the genuine ones. N-QC uses them on its arbitrary open U, where its hypothesis gives C^2 (item (e)). The definitions `frequencyMap`, `integrableOrbit` themselves are total and keep Lean's default values outside the differentiability locus.
(b) $D^2 h$ for complex h is used only on $B(0,R)_s$, open, where h is C^2 (Definition **??**).
(c) X_H is used only inside `IsSolutionOn`, which requires differentiability of H at every visited point; in N-ZERO and N-HEAD-C this is a proof obligation: $H_{h,f}$ is real-differentiable on $\mathbb{R}^n \times \mathcal{B}_s(R) \supseteq \mathbb{R}^n \times B(0,R)$ (angles first, actions restricted) because h, f are holomorphic on open sets containing the real points ([K:`bp:p1:hamiltonian-real`]).
(d) $\varepsilon^{-1/(2n)}$ and $\varepsilon^{1/(2n)}$ occur only with $\varepsilon > 0$ in N-HEAD-C, and $2n \neq 0$.
(e) In N-QC the map $\Psi(I', \lambda) = (h_{\mathbb{R}}(I'), \lambda\omega_h(I'))$ is differentiable at (I, λ) because $h_{\mathbb{R}} \in C^2$ on the open set $U \ni I$; `fderiv` is then the true derivative.

1.4. **The frozen theorems.**

Theorem 1.19 (N-HEAD-C: exponential stability, complex Hessian bound)**.** *Let $n \geq 3$ and $R, s, M, m, M_{\mathbb{C}}, \omega_{\min} > 0$. Then there are $C, T, c, \varepsilon_0 > 0$ with*

$$C\,\varepsilon_0^{1/(2n)} \leq R/4$$

such that the following holds for every $h : \mathbb{C}^n \to \mathbb{C}$, $f : \mathbb{C}^n \times \mathbb{C}^n \to \mathbb{C}$ and $\varepsilon \in \mathbb{R}$ with

(H1) `IsAnalyticIntegrable` $R\,s\,h$ *(Definition* **??***);*
(H2) `HasComplexHessianBound` $R\,s\,h\,M_{\mathbb{C}}$ *(Definition* **??***);*
(H3) $\omega_{\min} \leq |\omega_h(I)|$ *for every $I \in B(0,R)$ (max norm);*
(H4) `IsStrictlyQuasiConvex` $\mathcal{B}_s(R)\,h\,m$ *(Definition* **??***);*
(H5) `HasDerivativeBounds` $\mathcal{B}_s(R)\,h\,M$ *(Definition* **??***);*
(H6) `IsAnalyticPerturbation` $R\,s\,\varepsilon\,f$ *(Definition* **??***);*
(H7) $0 < \varepsilon \leq \varepsilon_0$.

For every $\theta_0 \in \mathbb{R}^n$ and $I_0 \in B(0, R/2)$, with $\tau := T\exp\bigl(c\,\varepsilon^{-1/(2n)}\bigr)$ and $H = H_{h,f}$, there is $z : \mathbb{R} \to \mathbb{R}^n \times \mathbb{R}^n$ with

(a) $z(0) = (\theta_0, I_0)$;
(b) `IsSolutionOn` $H\,B(0,R)\,[-\tau,\tau]\,z$;
(c) $|z(t)_2 - I_0| \leq C\varepsilon^{1/(2n)}$ *for all t with $|t| \leq \tau$;*
(d) (uniqueness) for every $L \in [0,\tau]$ and every $y : \mathbb{R} \to \mathbb{R}^n \times \mathbb{R}^n$ with $y(0) = (\theta_0, I_0)$ and `IsSolutionOn` $H\,B(0,R)\,[-L,L]\,y$*: $y(t) = z(t)$ for all $t \in [-L, L]$.*

The order of choice is: n; $R, s, M, m, M_{\mathbb{C}}, \omega_{\min}$; then C, T, c, ε_0, depending only on these seven data; then h, f, ε, then θ_0, I_0; the solution z may depend on all of them. There is no dependence on A in Definition **??***(iii). In (H4)–(H5) the set is $\mathcal{B}_s(R) = B(0, R+s)$ (Remark* **??***), in (H3) the smaller ball $B(0,R)$.*

Frozen Lean declaration: `KAM.Nekhoroshev.quasiConvex_nekhoroshev_complexBound`
Source: *Pos93 Theorem 1, pp. 190–191 (with the conversions and anchors of* `docs/tracks/N/sources.md`*); exponent* $1/(2n)$ *as BM11 p. 5 with* $\delta = (2n(n-1))^{-1}$*; ruling D18 Q2, Q4; deviation D20: the extra inputs* $M_{\mathbb{C}}$ *and* $\omega_{\min}$*.*
Source gap: *The printed theorem is for a* 2π*-periodic torus, Euclidean norms and exceptional initial frequencies* $\|\omega(I_0)\| \le mr_0/8$*; the conversion to the frozen statement (rescaling* $H \mapsto 2\pi H(\varphi/2\pi, I)$*, choice of* r_0, s_0 *from the seven data, exclusion of the exceptional branch by* $\omega_{\min}$*, existence and uniqueness of the solution) is given only in* `sources.md` *and is proved in Part 10 of* `PLAN.md`*.*

Theorem 1.20 (N-ZERO: exact integrable dynamics)**.** *Let* $n \in \mathbb{N}$*,* $R, s > 0$*,* h *with* `IsAnalyticIntegrable` $R\,s\,h$ *and* f *with* `IsAnalyticPerturbation` $R\,s\,0\,f$*. For every* $\theta_0 \in \mathbb{R}^n$ *and* $I_0 \in B(0, R/2)$*, with* $\gamma = \gamma_{\theta_0, I_0}$ *(Definition* **??***):*

$$\gamma(0) = (\theta_0, I_0), \qquad \texttt{IsSolutionOn}\, H_{h,f}\, B(0,R)\, \mathbb{R}\, \gamma, \qquad \gamma(t)_2 = I_0 \quad (t \in \mathbb{R}).$$

Frozen Lean declaration: `KAM.Nekhoroshev.zero_perturbation`
Source: *Ours (supplement to BM11 Theorem 2.1, p. 5, whose strict bound excludes* $\varepsilon = 0$*); from* $H = h(I)$*, BM11 p. 1.*

Theorem 1.21 (N-QC: nonsingular isoenergetic derivative)**.** *Let* $n \in \mathbb{N}$*,* $U \subset \mathbb{R}^n$*,* $h : \mathbb{C}^n \to \mathbb{C}$*,* $m \in \mathbb{R}$ *with* `IsStrictlyQuasiConvex` $U\,h\,m$*. For* $I \in U$ *and* $\lambda > 0$*, the derivative at* (I, λ) *of*

$$\Psi : \mathbb{R}^n \times \mathbb{R} \to \mathbb{R} \times \mathbb{R}^n, \qquad \Psi(I', \lambda') = \big(h_{\mathbb{R}}(I'),\ \lambda' \omega_h(I')\big)$$

is a bijective linear map $\mathbb{R}^n \times \mathbb{R} \to \mathbb{R} \times \mathbb{R}^n$*. Explicitly it is* $D\Psi(I,\lambda)[v,u] = \big(\omega_h(I)\cdot v,\ u\,\omega_h(I) + \lambda D^2 h_{\mathbb{R}}(I)v\big)$*. No uniform radius of invertibility is asserted.*
Frozen Lean declaration: `KAM.Nekhoroshev.quasiConvex_isoenergetic_derivative`
Source: *BM11 Lemma 3.3 and its proof, pp. 9–10 (only the derivative assertion).*

2. Interfaces and conversion of hypotheses

begingroup

This part connects the frozen hypotheses of track N (Definitions of Part 0) to the form in which the proofs use them. It proves: the real action domain and the inclusions of tubes (§**??**); that holomorphy gives a smooth real Hamiltonian with $X_H = X_{h+f}$ (§**??**); the translation of the derivative bounds $B(M)$, the complex Hessian bound and $QC(m)$ into the Euclidean, uniform form of Pöschel's Theorem 1 (§**??**); the change of normalisation 2π-periodic↔ 1-periodic and the sup-to-weighted Fourier norm (§**??**); and the dictionary of constants (§**??**). *Convention ruling.* All of Parts 1–10 work in the frozen conventions (period 1, max norms, characters $e^{2\pi i k\cdot\theta}$). Pöschel's normalisation (period 2π, Euclidean norms on actions, weights $e^{|k|_1 s}$) enters only through the labelled conversions Lemmas **??**, **??**, **??**, **??**; quantities in his normalisation carry a tilde or the subscript P. This overrules the suggestion in `PLAN.md` to work in Pöschel's normalisation. The Euclidean norm is written $|\cdot|_2$, the max norm $|\cdot|$ (or $|\cdot|_\infty$), and $|k|_1 = \sum_i |k_i|$; the phase-space norm is $\max(|\theta|_\infty, |I|_\infty)$, Euclidean structure is used only for the dot product and where stated.

2.1. **External facts.**

(E1) *Finite-dimensional linear algebra.* An injective linear map between real vector spaces of equal finite dimension is bijective (rank–nullity; Mathlib `LinearMap.injective_iff_surjective`). Exists.

(E2) *Norm comparison and Cauchy–Schwarz.* For $x \in \mathbb{R}^n$ or $\mathbb{C}^n$: $|x|_\infty \le |x|_2 \le \sqrt{n}|x|_\infty$ and $|x|_1 \le \sqrt{n}|x|_2$, $|x|_2 \le |x|_1$; $|x \cdot y| \le |x|_2 |y|_2$ (for the bilinear, not Hermitian, product too). Exists.

(E3) *Restriction of scalars and chain rule.* A $\mathbb{C}$-differentiable (resp. C^k) map is $\mathbb{R}$-differentiable (resp. C^k) with the same derivative; the chain rule for the $\mathbb{R}$-linear inclusion $\iota : \mathbb{R}^N \to \mathbb{C}^N$ and for Re. (Mathlib `DifferentiableAt.restrictScalars`, `ContDiffOn.restrict_scalars`, `HasFDerivAt.comp`.) Exists.

(E4) *Product rule* for $(x, \lambda) \mapsto \lambda\, v(x)$ with v differentiable, and differentiability of a vector function coordinatewise. Exists.

Track-K results used: [K:`bp:p1:holomorphic-regularity`], [K:`bp:p1:real-restriction`], [K:`bp:p1:hamiltonian-real`], [K:`bp:p1:ext-schwarz`], [K:`bp:p1:decay`], [K:`bp:p1:summation`], [K:`bp:p1:exp-sums`].

2.2. **Domains and norms.**

Lemma 2.1 (Norm and lattice comparisons)**.** *(a) For $x \in \mathbb{R}^n$ (or $\mathbb{C}^n$): $|x|_\infty \le |x|_2 \le \sqrt{n}\,|x|_\infty$, hence $|x|_\infty^2 \ge |x|_2^2/n$ and $|x|_\infty \le |x|_2$. (b) For $k \in \mathbb{Z}^n$: $|k|_2 \le |k|_1 \le \sqrt{n}|k|_2$. Consequently $\{|k|_1 \le K\} \subset \{|k|_2 \le K\}$ and $\{|k|_2 \le K\} \subset \{|k|_1 \le \sqrt{n}K\}$; so a geometric cutoff K_g in the Euclidean norm and an analytic cutoff K_a in $|\cdot|_1$ are compatible when $\sqrt{n}\,K_g \le K_a$. (c) The bilinear bound: if $|Q_{ij}| \le M_0$ for a matrix Q then $|x^\top Q y| \le n M_0 |x|_2 |y|_2$ for $x, y \in \mathbb{C}^n$.*
*Source: Standard (**??**); the lattice inclusions correct the direction stated in `PLAN.md` (review N06).*

Proof. **Step S2.1: Vectors.** (a),(b) are **??**: $|x|_2^2 = \sum x_i^2 \le n \max x_i^2$, $|x|_1 \le \sqrt{n}|x|_2$ is Cauchy–Schwarz with $(1, \dots, 1)$.

Step S2.2: Matrices. $|x^\top Q y| \le M_0 |x|_1 |y|_1 \le M_0\, n |x|_2 |y|_2$ by (b). □

Lemma 2.2 (Domains)**.** *Let $R, s > 0$, $n \ge 0$.*

*(a) $\mathcal{B}_s(R) = B(0, R+s)$; in particular it is open and $B(0,R) \subset \mathcal{B}_s(R)$ (Remark **??**); $W_s := \mathbb{R}^n \times \mathcal{B}_s(R)$ is open in $\mathbb{R}^n \times \mathbb{R}^n$ and $D_s(R) \cap (\mathbb{R}^n \times \mathbb{R}^n) = W_s$ (real points of $D_s(R)$: $\theta \in \mathbb{R}^n \subset T_s$ since $s > 0$).*
(b) $B(0,R)_s \cap \mathbb{R}^n = \mathcal{B}_s(R)$; $D_s(R)$ and $B(0,R)_s$ are open.
(c) (Euclidean tubes) Let $P = \{J \in \mathbb{R}^n : |J| \le R/2\}$ and $0 < r_0 \le \min(s/2, R/8)$. If $I \in \mathbb{C}^n$ and $J \in P$ with $|I - J|_2 < r_0$, then $I \in B(0,R)_s$ and $|I - J|_\infty < s$. If moreover $I \in \mathbb{R}^n$ then $|I|_\infty < 5R/8 \le 3R/4$.
(d) $T_\rho \subset T_s$ for $\rho \le s$; $T_s = \mathbb{C}^0$ is the universal singleton for $n = 0$ (no $n > 0$ hypothesis is needed anywhere in N-ZERO, N-QC).

*Source: Ours (Remark **??**); BM11 p. 4 for the notation.*

Proof. **Step S2.3: Real tube.** ($\subset$) $|I| \le |J| + |I - J| < R + s$. ($\supset$) If $|I| < R + s$ put $J = \frac{R}{R+s} I$; then $|J| < R$ and $|I - J| = \frac{s}{R+s}|I| < s$ (for $I = 0$ take $J = 0$).

Step S2.4: Openness and real points. $B(0,R)_s = \bigcup_{J \in B(0,R)} \{|I - J| < s\}$ is a union of open balls; T_s is open as a preimage of an open set; products of open sets are open; the rest is the definition of ι and $\mathcal{B}_s(R) = B(0,R)_s \cap \mathbb{R}^n$ by Step **??**.

Step S2.5: Euclidean tubes. By **??** $|I - J|_\infty \le |I - J|_2 < r_0 \le s/2 < s$, and $|J| < R$ since $J \in P$; so $I \in B(0,R)_s$. For real I: $|I|_\infty < R/2 + R/8$. □

2.3. **Holomorphy implies real smoothness.**

Throughout this subsection $R, s > 0$, h satisfies `IsAnalyticIntegrable` $R\, s\, h$, f satisfies `IsAnalyticPerturbation` $R\, s\, \varepsilon\, f$ ($\varepsilon \ge 0$), $G(\theta, I) = h(I) + f(\theta, I)$ on $D_s(R)$, $H = H_{h,f}$ and $\iota : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{C}^n \times \mathbb{C}^n$ the componentwise inclusion. Complex partials are $\partial_{\theta_i} G(w) = DG(w)[(e_i, 0)]$, $\partial_{I_i} G(w) = DG(w)[(0, e_i)]$.

Lemma 2.3 (Real smoothness and real Hamiltonian field)**.** *(a) G is holomorphic and C^∞ on $D_s(R)$; h is holomorphic and C^∞ on $B(0,R)_s$.*

(b) *H is C^∞ on the open set $W_s = \mathbb{R}^n \times B(0, R+s)$ (Lemma ??); on W_s it is real and $H = G \circ \iota$, and for all orders $D^k H(z)[\zeta_1, \dots, \zeta_k] = D^k G(\iota z)[\iota\zeta_1, \dots, \iota\zeta_k] \in \mathbb{R}$. Likewise $h_{\mathbb{R}} = h \circ \iota$ is C^∞ on $\mathcal{B}_s(R)$, with $D^k h_{\mathbb{R}}(I)[a_1, \dots, a_k] = D^k h(I)[\iota a_1, \dots]$.*

(c) *(Hamiltonian field) For $z = (\theta, I) \in W_s$: H is differentiable at z and $X_H(z) = \big((\partial_{I_i} G(\iota z))_i, (-\partial_{\theta_i} G(\iota z))_i\big) = X_{h+f}(\iota z)$, the complex Hamiltonian field of G at the real point ιz. The frequency map is $\omega_h(I)_i = \partial_i h(I) \in \mathbb{R}$ and $D^2 h_{\mathbb{R}}(I)[e_i, e_j] = \partial_i \partial_j h(I) = \partial_j \partial_i h(I) \in \mathbb{R}$ for $I \in \mathcal{B}_s(R)$.*

(d) *(Periodicity) $H(\theta + k, I) = H(\theta, I)$ for $(\theta, I) \in W_s$, $k \in \mathbb{Z}^n$; consequently DH, $D^k H$ and X_H are 1-periodic in θ on W_s.*

(e) *If $\varepsilon = 0$ then $f = 0$ on $D_s(R)$ and $H(\theta, I) = h_{\mathbb{R}}(I)$ on W_s.*

Source: BM11 p. 4 (regularity of h on B_s); [K:bp:p1:holomorphic-regularity], [K:bp:p1:hamiltonian-real]*; ours for the packaging. Part 3 uses this as* [N:bpN:p3:holo-input].

Proof. **Step S2.6: Holomorphy.** h is complex differentiable on the open set $B(0, R)_s$ and f on $D_s(R)$ (Definitions ??(i), ??(i); openness Lemma ??(b)); the map $(\theta, I) \mapsto h(I)$ is holomorphic on $D_s(R)$ (composition with the linear projection; $B(0, R)_s$ is the second factor). So G is holomorphic on $D_s(R)$, hence C^∞ with holomorphic derivatives of all orders ([K:bp:p1:holomorphic-regularity]), which is (a); by ?? G is C^∞ in the real sense.

Step S2.7: Reality. $\iota(W_s) \subset D_s(R)$ (Lemma ??(a)) and $W_s = \iota^{-1}(D_s(R))$ is open. For $z = (\theta, I) \in W_s$, $\operatorname{Im} h(I) = 0$ (Definition ??(iv), $I \in \mathcal{B}_s(R)$) and $\operatorname{Im} f(\theta, I) = 0$ (Definition ??(iii)); so $G(\iota z) \in \mathbb{R}$ and $H(z) = \operatorname{Re} G(\iota z) = G(\iota z)$. Likewise $h_{\mathbb{R}} = h \circ \iota$ on $\mathcal{B}_s(R)$.

Step S2.8: Derivatives of all orders. On the open set W_s, $H = G \circ \iota$ with ι $\mathbb{R}$-linear and G real-C^∞ (Step ??): chain rule ?? gives $H \in C^\infty(W_s)$ and the displayed formula for $D^k H$, with values real because H is real-valued (the derivative of a real-valued function along real directions is a limit of real difference quotients). The same for $h_{\mathbb{R}}$.

Step S2.9: The field. By Definition ??, $X_H(z) = ((DH(z)[(0, e_i)])_i, (-DH(z)[(e_i, 0)])_i)$; by Step ?? these are $\partial_{I_i} G(\iota z)$ and $-\partial_{\theta_i} G(\iota z)$. This is [K:bp:p1:hamiltonian-real] (with $V = D_s(R)$, H agreeing with $\operatorname{Re} G$ on the real neighbourhood W_s). Taking $k = 1, 2$ for $h_{\mathbb{R}}$ gives $\omega_h(I)_i = Dh_{\mathbb{R}}(I)[e_i] = \partial_i h(I)$ and $D^2 h_{\mathbb{R}}(I)[e_i, e_j] = \partial_i \partial_j h(I)$, symmetric by [K:bp:p1:holomorphic-regularity] ([K:bp:p1:ext-schwarz]).

Step S2.10: Periodicity. For $(\theta, I) \in W_s$, $k \in \mathbb{Z}^n$ and real θ: $(\theta, I), (\theta + k, I) \in D_s(R)$ and $f(\theta + k, I) = f(\theta, I)$ (Definition ??(ii)), $h(I)$ does not depend on θ. So $H \circ \tau_k = H$ on W_s, $\tau_k(\theta, I) = (\theta + k, I)$, which maps W_s onto itself with $D\tau_k = \mathrm{id}$; differentiating (chain rule) gives $D^j H(\tau_k z) = D^j H(z)$ and hence $X_H(\tau_k z) = X_H(z)$.

Step S2.11: Zero perturbation. $|f(w)| \le 0$ on $D_s(R)$ gives $f = 0$ there, so $H = \operatorname{Re} h(I) = h_{\mathbb{R}}(I)$ on W_s. □

Lemma 2.4 (Ordered partials and first-order bounds)**.** *Let $U \subset \mathbb{R}^n$ be open with* `HasDerivativeBounds` *$U\ h\ M$ (Definition ??). Then for $I \in U$, $a, b \in \mathbb{R}^n$: $|\omega_h(I)_i| \le M$, $|\omega_h(I)|_\infty \le M$, $|\omega_h(I)|_2 \le \sqrt{n}\, M$, and $|D^2 h_{\mathbb{R}}(I)[e_i, e_j]| \le M$. The ordered partial $D^k h_{\mathbb{R}}(I)[e_{i_1}, \dots, e_{i_k}]$ ($k \le 3$) is invariant under permutations of $(i_1, \dots, i_k)$, so it equals the multi-index partial of the source.*

Source: BM11 p. 4 ($B(M)$); [K:bp:p1:ext-schwarz].

Proof. **Step S2.12: First order.** $k = 1$ in the definition gives $|Dh_{\mathbb{R}}(I)[e_i]| = |\omega_h(I)_i| \le M$ (Definition ??); then $|\omega|_2 \le \sqrt{n}|\omega|_\infty$ (Lemma ??).

Step S2.13: Symmetry. $h_{\mathbb{R}} \in C^3$ on the open set U: $D^2 h_{\mathbb{R}}(I)$ and $D^3 h_{\mathbb{R}}(I)$ are symmetric ([K:bp:p1:ext-schwarz] applied to $h_{\mathbb{R}}$, $Dh_{\mathbb{R}}$, $D^2 h_{\mathbb{R}}$), which is the permutation invariance; the entry bound is $k = 2$. □

Lemma 2.5 (Complex Hessian bound in Euclidean form)**.** *Let* `HasComplexHessianBound` $R\,s\,h\,M_{\mathbb{C}}$. *For* $I \in B(0,R)_s$ *let* $Q(I) = (\partial_i\partial_j h(I))_{ij}$ *(complex symmetric). Then* $|Q(I)_{ij}| \le M_{\mathbb{C}}$ *and* $|x^{\top}Q(I)y| \le nM_{\mathbb{C}}|x|_2|y|_2$ *for* $x, y \in \mathbb{C}^n$*; hence the operator norm of* $Q(I) : (\mathbb{C}^n, |\cdot|_2) \to (\mathbb{C}^n, |\cdot|_2)$ *is* $\le nM_{\mathbb{C}}$. *For real* $I \in \mathcal{B}_s(R)$ *this matrix is the real Hessian* $D^2h_{\mathbb{R}}(I)[e_i, e_j]$ *(Lemma* **??***(c)), and for real* x, y*:* $|D^2h_{\mathbb{R}}(I)[x,y]| \le nM_{\mathbb{C}}|x|_2|y|_2$.

Source: Pos93 p. 190 (Hessian bound, anchor POS190-HESS); the conversion factor n *is ours (*`docs/tracks/N/sources.md`*).*

Source gap: Pos93 p. 190 says that for the symmetric Q *the Euclidean operator norm "amounts to the spectral radius norm". For complex symmetric matrices this is false (*$Q = \left(\begin{smallmatrix} 1 & i \\ i & -1 \end{smallmatrix}\right)$ *has* $Q^2 = 0$*). The frozen definition uses the operator norm of* D^2h *for the max norm, and so does this blueprint; no spectral-radius claim is used.*

Proof. **Step S2.14: Entries.** $D^2h(I) = D(Dh)(I) \in \mathcal{L}(\mathbb{C}^n, \mathcal{L}(\mathbb{C}^n, \mathbb{C}))$ has norm $\le M_{\mathbb{C}}$ by hypothesis and $|e_i| = 1$ for the max norm, so $|\partial_i\partial_j h(I)| = |D^2h(I)[e_i, e_j]| \le M_{\mathbb{C}}$ (the order of arguments is immaterial by symmetry, [K:`bp:p1:holomorphic-regularity`]).

Step S2.15: Euclidean bound. Lemma **??**(c). The operator-norm statement: $|Qx|_2 = \sup_{|y|_2=1}|\sum_i \bar{y}_i(Qx)_i| \le nM_{\mathbb{C}}|x|_2$ by the same estimate applied to $\bar{y}$. □

2.4. **Quasi-convexity: the uniform Euclidean form.**

Lemma 2.6 (Transverse coercivity gives conic coercivity (linear algebra core))**.** *Let* $n \ge 1$, $B, q > 0$, $Q : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ *bilinear (not necessarily symmetric) with* $|Q[x,y]| \le B|x|_2|y|_2$, $\omega \in \mathbb{R}^n \setminus \{0\}$, *and*

$$\omega \cdot v = 0 \implies Q[v,v] \ge q|v|_2^2 \qquad (v \in \mathbb{R}^n).$$

If $\ell' \ge 0$ *and* $\ell' \le |\omega|_2\, q/(8(B+q))$, *then for every* $\xi \in \mathbb{R}^n$*:*

$$|\omega \cdot \xi| \le \ell'|\xi|_2 \implies Q[\xi,\xi] \ge \tfrac{5}{8}q|\xi|_2^2 \ \ (\ge \tfrac{q}{2}|\xi|_2^2).$$

Source: Ours; the computation is in `docs/tracks/N/sources.md` *("Uniform conversion"); all inequalities re-checked here.*

Proof. **Step S2.16: Reduction to unit vectors.** Both hypothesis and conclusion are homogeneous (degrees 1 and 2) in ξ, and the case $\xi = 0$ is trivial; so let $|\xi|_2 = 1$.

Step S2.17: Decomposition. Put $e = \omega/|\omega|_2$, $t = \xi \cdot e$, $u = \xi - te$. Then $u \cdot \omega = 0$, $|u|_2^2 = 1 - t^2$, $|t| \le 1$, and $|t| = |\omega \cdot \xi|/|\omega|_2 \le \ell'/|\omega|_2 \le \kappa := q/(8(B+q)) < \frac{1}{8}$.

Step S2.18: Expansion. $Q[\xi,\xi] = Q[u,u] + tQ[u,e] + tQ[e,u] + t^2Q[e,e]$. By hypothesis $Q[u,u] \ge q(1-t^2)$; by the bound on Q and $|u|_2, |e|_2 \le 1$: $|Q[u,e]|, |Q[e,u]|, |Q[e,e]| \le B$. So $Q[\xi,\xi] \ge q - (q+B)t^2 - 2B|t|$.

Step S2.19: Numerics. $(q+B)t^2 \le (q+B)\kappa \cdot \kappa = \frac{q}{8}\kappa \le \frac{q}{8}$ and $2B|t| \le 2B\kappa = \frac{2Bq}{8(B+q)} \le \frac{q}{4}$ (as $B \le B+q$). Hence $Q[\xi,\xi] \ge q - \frac{q}{8} - \frac{q}{4} = \frac{5}{8}q$. □

Lemma 2.7 (Quasi-convexity in Pöschel's quantified form)**.** *Let* $n \ge 1$, $R, s, m, M_{\mathbb{C}}, \omega_{\min} > 0$ *and put*

$$q = \frac{m}{n}, \qquad B_c = nM_{\mathbb{C}} + q, \qquad \mu = \frac{q}{2}, \qquad \ell = \frac{\omega_{\min}\, q}{8(B_c+q)}.$$

Let h *satisfy* `IsAnalyticIntegrable` $R\,s\,h$ *and* `HasComplexHessianBound` $R\,s\,h\,M_{\mathbb{C}}$, `IsStrictlyQuasiConvex` $\mathcal{B}_s(R)\,h\,m$, *and* $\omega_{\min} \le |\omega_h(I)|_\infty$ *for all* $I \in B(0,R)$. *Then for every* $I \in B(0,R)$, *writing* $\omega = \omega_h(I)$, $Q = D^2h_{\mathbb{R}}(I)$*:*

- *(a)* $|Q[x,y]| \le B_c|x|_2|y|_2$; $\ \omega \cdot v = 0 \Rightarrow Q[v,v] \ge q|v|_2^2$;
- *(b) (Pöschel's alternative,* conditional*) for every* $\xi \in \mathbb{R}^n$: $\ |\omega \cdot \xi| \le \ell|\xi|_2 \implies Q[\xi,\xi] \ge \mu|\xi|_2^2$; *equivalently, either* $|\omega \cdot \xi| > \ell|\xi|_2$ *or* $Q[\xi,\xi] \ge \mu|\xi|_2^2$ *(Pos93 p. 190, "*ℓ, m*-quasi-convex" with* $m := \mu$*);*

(c) $|\omega|_2 \ge |\omega|_\infty \ge \omega_{\min}$ *and* $|\omega|_2 \le \sqrt{n}|\omega|_\infty$.

The constants q, B_c, μ, ℓ *depend only on* $(n, m, M_{\mathbb{C}}, \omega_{\min})$. *In Pöschel's normalisation* ($\tilde{h} = 2\pi h$, *Lemma* **??**) *the same holds with* $(\tilde{\ell}, \tilde{\mu}, \tilde{B}_c, \tilde{\omega}, \tilde{Q}) = 2\pi(\ell, \mu, B_c, \omega, Q)$. Warning. *Full positive definiteness* $Q[\xi,\xi] \ge \mu|\xi|_2^2$ *for all* ξ *is false in general and is not claimed: for* $h(I) = I_1 + \frac{1}{2}(I_2^2 + I_3^2)$ *one has* $\omega = e_1 \ne 0$, *QC holds on every bounded set, but* $Q[e_1, e_1] = 0$. *The estimate (b) is only needed (and only true) in the cone* $|\omega \cdot \xi| \le \ell|\xi|_2$.

Source: Pos93 p. 190 (anchors POS190-QC, POS190-HESS, POS191-LOW; I read the scan, p. 190, ℓ, m*-quasi-convexity, hypothesis* $r_0 \le 4\ell/m$*); conversion:* `docs/tracks/N/sources.md`*, "Uniform conversion"; BM11 p. 4 for* $QC(m)$. *Review N05.*

Proof. **Step S2.20: Regularity and Hessian.** Since $B(0,R) \subset \mathcal{B}_s(R)$, $I \in \mathcal{B}_s(R)$; Q is the real Hessian of Lemma **??**, so $|Q[x,y]| \le nM_{\mathbb{C}}|x|_2|y|_2 \le B_c|x|_2|y|_2$, the first half of (a).

Step S2.21: Transverse coercivity in the Euclidean norm. Definition **??** at $I \in \mathcal{B}_s(R)$ gives $\omega \cdot v = 0 \Rightarrow Q[v,v] \ge m|v|_\infty^2 \ge \frac{m}{n}|v|_2^2 = q|v|_2^2$ (Lemma **??**(a)). Also $\omega \ne 0$.

Step S2.22: Frequency size. (c): $|\omega|_\infty \ge \omega_{\min}$ on $B(0,R)$ by hypothesis, $|\omega|_2 \ge |\omega|_\infty$; $|\omega|_2 \le \sqrt{n}|\omega|_\infty$ by Lemma **??**.

Step S2.23: Conic coercivity. Apply Lemma **??** with $B = B_c$, q and $\ell' = \ell$: $\ell = \omega_{\min} q/(8(B_c + q)) \le |\omega|_2 q/(8(B_c+q))$ by (c). The conclusion is $Q[\xi,\xi] \ge \frac{5}{8} q|\xi|_2^2 \ge \mu|\xi|_2^2$.

Step S2.24: Scaling. Multiplying h by 2π multiplies ω, Q, B_c and the three of μ, q, ℓ by 2π, and the implications are homogeneous, so they persist. □

2.5. **Normalisations.**

Lemma 2.8 (Period conversion)**.** *Let* $\Lambda(\theta, I) = (2\pi\theta, I)$ *and, for* $H : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ *and* h, f *complex, define*

$$\tilde{H}(\varphi, I) = 2\pi H(\varphi/2\pi, I), \qquad \tilde{h} = 2\pi h, \qquad \tilde{f}(\varphi, I) = 2\pi f(\varphi/2\pi, I),$$

so $\tilde{H} = H_{\tilde{h},\tilde{f}}$ *when* $H = H_{h,f}$, *and* $\tilde{f}$ *is* $2\pi\mathbb{Z}^n$*-periodic in* φ. *Then:*

- *(a) (Dynamics) For* $U \subset \mathbb{R}^n$, $J \subset \mathbb{R}$: z *satisfies* `IsSolutionOn` $H\,U\,J\,z$ *iff* $\Lambda \circ z$ *satisfies* `IsSolutionOn` $\tilde{H}\,U\,J$ *(same time, same actions). In particular* $|(\Lambda z)_2 - I_0| = |z_2 - I_0|$.
- *(b) (Quantities)* $\omega_{\tilde{h}} = 2\pi\omega_h$, $D^2\tilde{h}_{\mathbb{R}} = 2\pi D^2 h_{\mathbb{R}}$, *and so:* `IsStrictlyQuasiConvex` $U\,h\,m \Rightarrow$ `IsStrictlyQuasiConvex` $U\,\tilde{h}\,2\pi m$; $B(M) \Rightarrow$ *coordinate partials of* $\tilde{h}_{\mathbb{R}}$ *of order* $1,2,3$ *bounded by* $2\pi M$; *the complex Hessian bound* $M_{\mathbb{C}} \to 2\pi M_{\mathbb{C}}$; $\omega_{\min} \to 2\pi\omega_{\min}$; *Euclidean sizes of* ω: $|\tilde{\omega}|_2 = 2\pi|\omega|_2$. *Time is unchanged.*
- *(c) (Domains)* $\tilde{f}$ *is holomorphic on* $\{|\mathrm{Im}\,\varphi| < 2\pi s\} \times B(0,R)_s$, *with* $|\tilde{f}| \le 2\pi\varepsilon$ *if* $|f| \le \varepsilon$ *on* $D_s(R)$; *the strip* T_ρ (ρ *in* θ) *corresponds to the strip of width* $2\pi\rho$ *in* φ.
- *(d) (Fourier)* $\tilde{f}(\varphi, I) = \sum_k \hat{f}_k(I) e^{ik\cdot\varphi}$ *with the same coefficients* $\hat{f}_k(I) = \int_{[0,1]^n} f(\theta, I) e^{-2\pi i k\cdot\theta}\, d\theta$ *times* 2π; *the coefficient radius* s' *of Pöschel's weight* $e^{|k|_1 s'}$ *equals* $2\pi\rho$.

Source: `docs/tracks/N/sources.md` *("For the period conversion"); Pos93 p. 189 (set-up:* $\theta \in \mathbb{R}^n/2\pi\mathbb{Z}^n$, *strip* W_{s_0} *of width* s_0 *in the max norm of* $\mathrm{Im}\,\theta$*).*

Proof. **Step S2.25: Chain rule.** $\tilde{H} = 2\pi\, H \circ \Lambda^{-1}$; Λ is a linear isomorphism, so $\tilde{H}$ is differentiable at Λz iff H is at z, with $\partial_{I_i}\tilde{H}(\Lambda z) = 2\pi\partial_{I_i}H(z)$ and $\partial_{\varphi_i}\tilde{H}(\Lambda z) = \partial_{\theta_i}H(z)$.

Step S2.26: Vector field. Hence $X_{\tilde{H}}(\Lambda z) = (2\pi\partial_I H(z), -\partial_\theta H(z)) = (2\pi\dot{\theta}, \dot{I}) = \frac{d}{dt}\Lambda z$ when $\dot{z} = X_H(z)$, and conversely; the condition $z(t)_2 \in U$ is unchanged. This is (a). (Rescaling the angle without the factor 2π in H would change time.)

Step S2.27: Scalars. (b) is linearity of $h \mapsto \omega_h, D^2 h_{\mathbb{R}}, D^k h_{\mathbb{R}}$ and of the operator norm. (c),(d) are the substitution $\varphi = 2\pi\theta$, $|\mathrm{Im}\,\varphi| = 2\pi|\mathrm{Im}\,\theta|$, and $e^{ik\cdot\varphi} = e^{2\pi i k\cdot\theta}$. □

Lemma 2.9 (From sup norm on a strip to the weighted Fourier norm)**.** *Let $u : T_\rho \to \mathbb{C}$ be holomorphic, bounded and 1-periodic, $\|u\|_\rho = \sup_{T_\rho}|u|$, and $0 < \delta < \rho$. Then*

$$\sum_{k\in\mathbb{Z}^n} |\hat{u}_k|\, e^{2\pi|k|_1(\rho-\delta)} \;\le\; \coth(\pi\delta)^n\, \|u\|_\rho,$$

the left side is Pöschel's weighted norm with weight $e^{|k|_1 s'}$, $s' = 2\pi(\rho-\delta)$, and the factor is $\coth^n\sigma$ with $\sigma = \pi\delta$ (Pos93 p. 190: $|u|_{D,r,s} \le \coth^n\sigma\, |u|_{D,r,s+2\sigma}$, where the right side is the sup norm on the strip of width $s+2\sigma$ in φ, i.e. $\rho = (s'+2\sigma)/2\pi$). Conversely $\|u\|_\rho \le \sum_k |\hat{u}_k| e^{2\pi|k|_1\rho}$. For I-dependent $u(\theta, I)$ the statement holds for each fixed I with the same constant, so the sup over I in the weighted norm is $\le \coth(\pi\delta)^n \sup_{\theta,I}|u|$.
Source: Pos93 p. 190 (the displayed relation between the weighted and sup norms); frozen-convention statement: [K:bp:p1:summation], [K:bp:p1:decay], [K:bp:p1:exp-sums]*(b).*

Proof. **Step S2.28: Decay.** For $\rho' \in (\rho-\delta, \rho)$, [K:bp:p1:decay] gives $|\hat{u}_k| \le e^{-2\pi|k|_1\rho'}\|u\|_\rho$; hence $|\hat{u}_k| e^{2\pi|k|_1(\rho-\delta)} \le e^{-2\pi|k|_1(\rho'-\rho+\delta)}\|u\|_\rho$, and letting $\rho' \uparrow \rho$ (termwise, then summing, or directly [K:bp:p1:summation], converse part) the bound is $e^{-2\pi\delta|k|_1}\|u\|_\rho$ per k.

Step S2.29: Sum. $\sum_{k\in\mathbb{Z}^n} e^{-2\pi\delta|k|_1} = \coth(\pi\delta)^n$ ([K:bp:p1:exp-sums](b) with $x = 2\pi\delta$). The case with parameters is pointwise in I. □

2.6. **The dictionary of constants.**

Lemma 2.10 (Constants dictionary for N-HEAD-C)**.** *Let $n \ge 3$ (only $n \ge 2$ is used by Pöschel) and $R, s, M, m, M_{\mathbb{C}}, \omega_{\min} > 0$, $a = \frac{1}{2n}$. Define successively*

$$q = \tfrac{m}{n},\ B_c = nM_{\mathbb{C}} + q,\ \mu = \tfrac{q}{2},\ \ell = \tfrac{\omega_{\min} q}{8(B_c+q)}, \qquad r_0 = \min\Big(\tfrac{s}{2}, \tfrac{R}{8}, \tfrac{2\ell}{\mu}, \tfrac{4\omega_{\min}}{\mu}\Big), \qquad A = \tfrac{11B_c}{\mu},$$

$$s_0 = \pi s, \quad \sigma = \tfrac{\pi s}{4}, \quad F = 2\pi \coth^n\big(\tfrac{\pi s}{4}\big), \quad \varepsilon_* = \frac{2\pi\mu\, r_0^2}{2^{10} A^{2n}},$$

$$C = \tfrac{r_0}{A}\Big(\tfrac{F}{\varepsilon_*}\Big)^a, \quad c = \tfrac{s_0}{6}\Big(\tfrac{\varepsilon_*}{F}\Big)^a, \quad T = \frac{A^2 s_0}{4\pi n M},$$

$$\varepsilon_0 = \min\Big(\tfrac{\varepsilon_*}{F},\ \big(\tfrac{R}{4C}\big)^{2n}\Big).$$

Pöschel's data (his normalisation, for $\tilde{h} = 2\pi h$): $m_P = 2\pi\mu$, $\ell_P = 2\pi\ell$, $M_P = 2\pi B_c$, $A_P = 11M_P/m_P = A$, $\varepsilon_{0,P} = m_P r_0^2/(2^{10}A_P^{2n}) = \varepsilon_*$, $R_0 = r_0/A$, and the set $P = \{J \in \mathbb{R}^n : |J| \le R/2\}$. Then:

- *(a) All constants are positive and finite and depend only on $(n, R, s, M, m, M_{\mathbb{C}}, \omega_{\min})$ — not on $h, f, \varepsilon, \theta_0, I_0$, nor on the bound A of Definition **??**(iii). Order of choice: $q, B_c, \mu, \ell, r_0, A$; then s_0, σ, F; ε_*; C, c; T; ε_0. And $C\varepsilon_0^a \le R/4$, $0 < \varepsilon_0 \le \varepsilon_*/F$.*
- *(b) $r_0 \le 4\ell_P/m_P$ $(= 4\ell/\mu)$; $A \ge 22$; $R_0 = r_0/A \le r_0/22 \le R/176$.*
- *(c) (Tubes) With $V_{r_0}P = \{I \in \mathbb{C}^n : \exists J \in P,\ |I-J|_2 < r_0\}$ and $U_{r_0}P = V_{r_0}P \cap \mathbb{R}^n$: $V_{r_0}P \subset B(0,R)_s$ and $U_{r_0}P \subset B(0, 5R/8) \subset B(0,R)$.*
- *(d) (Hypotheses of Theorem 1) If h satisfies (H1)–(H5) of Theorem **??**, then for $I \in U_{r_0}P$: $\tilde{h}$ is ℓ_P, μ_P-quasi-convex (Lemma **??**) with $\mu_P = 2\pi\mu$; the complex Hessian operator norm (Euclidean) of $\tilde{h}$ on $V_{r_0}P$ is $\le M_P$ (Lemma **??**); Pöschel's m, M are m_P, M_P and his hypothesis $r_0 \le 4\ell_P/m_P$ holds.*
- *(e) (Perturbation) If f satisfies (H6) with ε, then Pöschel's norm of $\tilde{f}$ on $V_{r_0}P \times$(strip of width s_0) is $\le F\varepsilon$; and $F\varepsilon \le \varepsilon_{0,P}$ if $\varepsilon \le \varepsilon_0$.*
- *(f) (Exceptional branch is empty) For $I_0 \in B(0, R/2)$: $|\tilde{\omega}(I_0)|_2 \ge 2\pi\omega_{\min} > \pi\omega_{\min} \ge m_P r_0/8$.*
- *(g) (Time prefactor) For $I_0 \in B(0, R/2)$, $\Omega_0 = \sup_{|I-I_0|_2 \le R_0} |\tilde{\omega}(I)|_2 \le 2\pi\sqrt{n}M \le 2\pi n M$, so Pöschel's $T_0 = A_P^2 s_0/\Omega_0 \ge 2T$.*

(h) (Conclusion shape) Pöschel's $R_0(\varepsilon_P/\varepsilon_{0,P})^a$ and $T_0\exp\big(\frac{s_0}{6}(\varepsilon_{0,P}/\varepsilon_P)^a\big)$ with $\varepsilon_P = F\varepsilon$ are $C\varepsilon^a$ and $T_0\exp(c\varepsilon^{-a})$, and $T\exp(c\varepsilon^{-a}) \le \frac{1}{2}T_0\exp(c\varepsilon^{-a})$.

Source: `docs/tracks/N/sources.md` "Uniform conversion" (all constants), checked against Pos93 p. 190 (hypotheses, $A = 11M/m$, $\varepsilon_0 = mr_0^2/(2^{10}A^{2n})$, $r_0 \le 4\ell/m$), p. 191 (parameters a, R_0, T_0, exceptional branch) and p. 204 (assembly: $K = (\varepsilon_0/\varepsilon)^a$). Pos93 requires $\|f_\varepsilon\| \le \varepsilon$ (p. 190) with $\varepsilon \le \varepsilon_{0,P}$.
Source gap: Pos93 states no theorem about the $\omega_{\min}$-conversion and the frozen-to-source dictionary: it exists only in `sources.md` and here (Part 0 `bpN:p0:head`). The choice of T with the factor $1/2$ is slack, not needed by the source.

Proof. **Step S2.30: Positivity and dependence.** Each constant is a positive finite expression in earlier ones (denominators: $n, \mu, 8(B_c+q), A, F, \varepsilon_*, 4C, 4\pi nM$, all > 0); $\varepsilon_0 > 0$ as a minimum of positive numbers, and $C\varepsilon_0^a \le C\cdot\frac{R}{4C} = R/4$ since $((R/4C)^{2n})^{1/(2n)} = R/(4C)$. The list of inputs in each formula contains only $n, R, s, M, m, M_{\mathbb{C}}, \omega_{\min}$. This is (a).

Step S2.31: Size relations. $B_c = nM_{\mathbb{C}} + q \ge q = 2\mu$ so $A = 11B_c/\mu \ge 22$. $4\ell_P/m_P = 4\ell/\mu \ge 2\ell/\mu \ge r_0$, and $R_0 = r_0/A \le r_0/22 \le R/176$. This is (b).

Step S2.32: Tubes. $r_0 \le \min(s/2, R/8)$: Lemma **??**(c) gives (c).

Step S2.33: Quasi-convexity and Hessian. For (d): $U_{r_0}P \subset B(0,R)$ by (c), so Lemma **??**(b) applies at every point of $U_{r_0}P$ and, after Lemma **??**(b), gives the (ℓ_P, μ_P)-alternative for $\tilde h$. For $I \in V_{r_0}P \subset B(0,R)_s$ the Euclidean operator norm of $\tilde Q(I)$ is $\le 2\pi nM_{\mathbb{C}} \le 2\pi B_c = M_P$ (Lemma **??**). Pöschel's $A = 11M/m$ with $M = M_P$ and $m = \mu_P = m_P$ is $11B_c/\mu = A$.

Step S2.34: Perturbation. By (H6) $|f| \le \varepsilon$ on $D_s(R) \supset T_{3s/4}\times V_{r_0}P$ ($V_{r_0}P \subset B(0,R)_s$, (c)). For each fixed $I \in V_{r_0}P$ apply Lemma **??** to $u = f(\cdot, I)$ with $\rho = 3s/4$, $\delta = s/4$: $\sum_k |\hat f_k(I)|e^{2\pi|k|_1 s/2} \le \coth(\pi s/4)^n\varepsilon$. Multiplying by 2π (Lemma **??**(d)) and noting $2\pi\cdot s/2 = \pi s = s_0$ gives $\le F\varepsilon$. (In Pöschel's terms $s_0 + 2\sigma = 3\pi s/2 = 2\pi\cdot\frac{3s}{4}$.) $\varepsilon \le \varepsilon_*/F$ gives $F\varepsilon \le \varepsilon_* = \varepsilon_{0,P}$. This is (e).

Step S2.35: Exceptional branch. $I_0 \in B(0,R/2) \subset B(0,R)$ so $|\omega(I_0)|_2 \ge \omega_{\min}$ (Lemma **??**(c)) and $|\tilde\omega|_2 = 2\pi|\omega|_2$. And $m_P r_0/8 = 2\pi\mu r_0/8 \le 2\pi\mu\cdot\frac{4\omega_{\min}}{8\mu} = \pi\omega_{\min}$ by $r_0 \le 4\omega_{\min}/\mu$. This is (f).

Step S2.36: Time. The Euclidean ball $\{|I - I_0|_2 \le R_0\}$ with $R_0 \le r_0$ lies in $\{|I|_\infty < R/2 + R_0\} \subset B(0, 5R/8)$ ($|I - I_0|_\infty \le |I - I_0|_2$), inside $\mathcal{B}_s(R)$ where Lemma **??** gives $|\omega(I)|_2 \le \sqrt{n}M$. So $\Omega_0 \le 2\pi\sqrt{n}M \le 2\pi nM$ and $T_0 = A^2 s_0/\Omega_0 \ge A^2 s_0/(2\pi nM) = 2T$. (g).

Step S2.37: Exponent and constants. $R_0(F\varepsilon/\varepsilon_*)^a = \frac{r_0}{A}(F/\varepsilon_*)^a\varepsilon^a = C\varepsilon^a$ and $\frac{s_0}{6}(\varepsilon_*/(F\varepsilon))^a = c\varepsilon^{-a}$. (h) follows from $T \le T_0/2$. □

Remark 2.11 (Forward use and an unused constant). Part 10 chooses C, T, c, ε_0 as in Lemma **??**, verifies Pos93's hypotheses by (d),(e) and excludes the exceptional branch by (f). The time prefactor T enters only through (g); the remaining slack $\varepsilon_0 \le (R/4C)^{2n}$ gives $C\varepsilon^a \le R/4$ so that the a-priori confinement keeps $I(t) \in B(0, R/2 + R/4) = B(0, 3R/4)$ (continuation, [N:`bpN:p3:continuation`]). The value $M_{\mathbb{C}}$ enters only through B_c; M only through T.

3. N-QC and N-ZERO

begingroup

This part proves two frozen theorems completely: N-QC (Theorem **??**, the nonsingular isoenergetic derivative) and N-ZERO (Theorem **??**, the exact integrable dynamics). Neither uses the normal-form machinery. The linear-algebra core of N-QC is stated in a reusable form (Lemma **??**, cited by track I as the isoenergetic gate, [N:`bpN:p2:qc-kernel`]). Frozen conventions throughout (max norms, period 1); Euclidean structure appears only as the dot product $\omega\cdot v = \sum\omega_i v_i$.

3.1. **N-QC.**

Lemma 3.1 (Kernel core: bordered Hessian is nonsingular)**.** *Let $n \geq 1$, $\omega \in \mathbb{R}^n$, A a real $n \times n$ matrix (not necessarily symmetric, no norm bound), $\lambda \neq 0$, and suppose*

$$\omega \cdot v = 0,\ v \neq 0 \implies v^\top A v \neq 0 \qquad \text{and} \qquad \omega \neq 0.$$

Let $L : \mathbb{R}^n \times \mathbb{R} \to \mathbb{R} \times \mathbb{R}^n$, $L(v,u) = (\omega \cdot v,\ u\,\omega + \lambda A v)$. Then $\ker L = 0$, hence L is bijective. In particular this holds if $\lambda > 0$, $\omega \neq 0$ and $v^\top A v \geq m|v|^2$ for some norm and $m > 0$ whenever $\omega \cdot v = 0$. (For $n = 0$ no $\omega \neq 0$ exists; N-QC is then vacuous, Remark **??**.*)*
Source: BM11 Lemma 3.3 and its proof, pp. 9–10 (kernel computation); linear algebra **??**.

Proof. **Step S3.1: Kernel equations.** Let $L(v,u) = 0$: $\omega \cdot v = 0$ and $u\omega + \lambda A v = 0$.

Step S3.2: Pairing with v. Dot the second equation with v: $u(\omega \cdot v) + \lambda v^\top A v = 0$, so $\lambda v^\top A v = 0$, i.e. $v^\top A v = 0$ as $\lambda \neq 0$. Since also $\omega \cdot v = 0$, the hypothesis forces $v = 0$.

Step S3.3: The scalar. With $v = 0$: $u\omega = 0$; as $\omega \neq 0$, $u = 0$.

Step S3.4: Bijectivity. L is a linear map between spaces of dimension $n+1$ with trivial kernel, hence bijective by **??**. □

Remark 3.2 ($n = 0$). For $n = 0$ the space $\mathbb{R}^0$ is a point, so $\omega = 0$ and Definition **??** requires $\omega_h(I) \neq 0$: no I satisfies it. Hence for $n = 0$ the hypotheses of N-QC are never met with $I \in U$ and the theorem is vacuous; no restriction $n \geq 1$ is added to it (review N03).

Lemma 3.3 (The derivative of Ψ)**.** *Let $U \subset \mathbb{R}^n$ be open, h with $h_{\mathbb{R}} \in C^2(U)$, $I \in U$, $\lambda \in \mathbb{R}$, $\Psi(I', \lambda') = (h_{\mathbb{R}}(I'), \lambda' \omega_h(I'))$. Then Ψ is differentiable at (I, λ) (so* `fderiv` *is the genuine derivative; Remark* **??***(e)) and*

$$D\Psi(I,\lambda)[v,u] = \big(\omega \cdot v,\ u\,\omega + \lambda Q v\big), \qquad \omega = \omega_h(I),\ Q_{ij} = D^2 h_{\mathbb{R}}(I)[e_i, e_j] = Q_{ji}.$$

Source: BM11 p. 10; **??**, [K:`bp:p1:ext-schwarz`].

Proof. **Step S3.5: First component.** $(I', \lambda') \mapsto h_{\mathbb{R}}(I')$ is differentiable at (I, λ) ($h_{\mathbb{R}}$ is C^2 on the open set $U \ni I$) with derivative $[v,u] \mapsto Dh_{\mathbb{R}}(I)[v] = \sum_i v_i Dh_{\mathbb{R}}(I)[e_i] = \omega \cdot v$.

Step S3.6: The frequency map is C^1. $\omega_h(I')_i = Dh_{\mathbb{R}}(I')[e_i]$ is the composition of $I' \mapsto Dh_{\mathbb{R}}(I') \in \mathcal{L}(\mathbb{R}^n, \mathbb{R})$ (which is C^1 on U since $h_{\mathbb{R}} \in C^2(U)$ and U is open) with the continuous linear evaluation $\ell \mapsto \ell(e_i)$. So ω_h is differentiable on U with $D\omega_h(I)[v]_i = D(Dh_{\mathbb{R}})(I)[v][e_i] = D^2 h_{\mathbb{R}}(I)[v, e_i]$.

Step S3.7: Symmetry. $D^2 h_{\mathbb{R}}(I)$ is symmetric ([K:`bp:p1:ext-schwarz`]), so $D\omega_h(I)[v] = Qv$ with Q the symmetric Hessian matrix.

Step S3.8: Product. By the product rule **??**, $(I', \lambda') \mapsto \lambda' \omega_h(I')$ is differentiable at (I, λ) with derivative $[v,u] \mapsto u\,\omega + \lambda Q v$. Both components are differentiable, so Ψ is, with the stated derivative. □

Theorem 3.4 (N-QC)**.** *Theorem* **??** *holds: for* `IsStrictlyQuasiConvex`$U\,h\,m$*, $I \in U$, $\lambda > 0$, the map $D\Psi(I,\lambda)$ is bijective.*
Frozen Lean declaration: `KAM.Nekhoroshev.quasiConvex_isoenergetic_derivative`
Source: BM11 Lemma 3.3 and proof, pp. 9–10.

Proof. **Step S3.9: Unpack the hypothesis.** Definition **??**: $m > 0$, U open, $h_{\mathbb{R}} \in C^2(U)$, $\omega := \omega_h(I) \neq 0$, and $\omega \cdot v = 0 \Rightarrow m|v|^2 \leq D^2 h_{\mathbb{R}}(I)[v,v]$ (max norm on v). Note $n \geq 1$ because $\omega \neq 0$ (Remark **??**).

Step S3.10: The derivative. By Lemma **??**, $D\Psi(I,\lambda) = L$ of Lemma **??** with $A = Q$ the Hessian matrix, $v^\top Q v = D^2 h_{\mathbb{R}}(I)[v,v]$ (bilinearity).

Step S3.11: Positivity on $\omega^\perp$. For $\omega \cdot v = 0$, $v \neq 0$: $v^\top Q v \geq m|v|_\infty^2 > 0$, since $m > 0$ and $|v|_\infty > 0$. So the hypothesis of Lemma **??** holds (with $\lambda > 0$).

Step S3.12: Conclusion. Lemma **??** gives bijectivity of $L = D\Psi(I, \lambda) = \texttt{fderiv}\,\mathbb{R}\,\Psi\,(I, \lambda)$. □

Remark 3.5 (Reusable form for track I). Lemma **??** needs no norm, no symmetry of A and no uniform constant: any quasi-convexity-type hypothesis ("$v^\top A v > 0$ on $\omega^\perp \setminus 0$") together with $\omega \neq 0$ and $\lambda \neq 0$ gives nonsingularity of the bordered matrix $\begin{pmatrix} \omega^\top & 0 \\ \lambda A & \omega \end{pmatrix}$. A quantitative inverse bound is not asserted by N-QC and is not provided here; it would follow, if needed, from the compactness of $\{v : |v|_2 = 1,\ \omega \cdot v = 0\}$ for a fixed (I, λ) but is not uniform in (I, λ).

3.2. **N-ZERO.**

Lemma 3.6 (The field at the integrable orbit)**.** *Let $R, s > 0$, `IsAnalyticIntegrable R s h`, `IsAnalyticPerturbation R s 0 f`, $H = H_{h,f}$, and $W_s = \mathbb{R}^n \times B(0, R + s)$. Then $H(\theta, I) = h_{\mathbb{R}}(I)$ on W_s; H is differentiable at every $z = (\theta, I) \in W_s$ with $DH(z)[(a, b)] = Dh_{\mathbb{R}}(I)[b]$ and*

$$X_H(\theta, I) = (\omega_h(I), 0).$$

Source: BM11 p. 1 ($H = h(I)$ when $f = 0$); ours.

Proof. **Step S3.13: f vanishes.** Definition **??**(iv) with $\varepsilon = 0$: $|f| \leq 0$ on $D_s(R)$, so $f = 0$ there. For $z = (\theta, I) \in W_s$, $(\theta, I) \in D_s(R)$ (Lemma **??**(a)), so $H(z) = \operatorname{Re} h(I) = h_{\mathbb{R}}(I)$ (Lemma **??**(e)).

Step S3.14: Differentiability. W_s is open and on it $H = h_{\mathbb{R}} \circ \mathrm{pr}_2$ with $h_{\mathbb{R}} \in C^3(\mathcal{B}_s(R))$ (Definition **??**(v)), $\mathcal{B}_s(R) = B(0, R + s)$ open. So H is differentiable at each point of W_s with $DH(z)[(a, b)] = Dh_{\mathbb{R}}(I)[b]$ (chain rule with the continuous linear projection). The point-wise hypotheses of the totality remark **??** are met.

Step S3.15: Field. By Definition **??**, $\partial_{I_i} H = Dh_{\mathbb{R}}(I)[e_i] = \omega_h(I)_i$ (Definition **??**) and $\partial_{\theta_i} H = DH(z)[(e_i, 0)] = 0$. □

Theorem 3.7 (N-ZERO)**.** *Theorem **??** holds.*
Frozen Lean declaration: `KAM.Nekhoroshev.zero_perturbation`
Source: Ours (supplement to BM11 Theorem 2.1, p. 5).

Proof. **Step S3.16: Initial value and action.** $\gamma(t) = (\theta_0 + t\omega_h(I_0), I_0)$ (Definition **??**) gives $\gamma(0) = (\theta_0, I_0)$ and $\gamma(t)_2 = I_0$ for all t: the first and third conclusions.

Step S3.17: The action stays in $B(0, R)$. $I_0 \in B(0, R/2) \subset B(0, R)$ since $R > 0$ (max norm: $|I_0| < R/2 < R$).

Step S3.18: Differentiability of H along the orbit. $\gamma(t) \in \mathbb{R}^n \times B(0, R) \subset W_s$ (Lemma **??**(a): $B(0, R) \subset B(0, R + s)$). By Lemma **??** H is differentiable at $\gamma(t)$ for each t.

Step S3.19: Derivative of the orbit. γ is affine in t, so differentiable at every $t \in \mathbb{R}$ (two-sided) with $\gamma'(t) = (\omega_h(I_0), 0)$.

Step S3.20: The equation. Lemma **??** gives $X_H(\gamma(t)) = (\omega_h(I_0), 0) = \gamma'(t)$ for every $t \in \mathbb{R}$. With Steps **??**–**??**, every clause of Definition **??** holds with $U = B(0, R)$ and $J = \mathbb{R}$. □

Remark 3.8 (Scope). N-ZERO uses neither the holomorphy of Part 1 (beyond the stated clauses of the definitions) nor the ODE layer of Part 3; Part 3 supplies uniqueness if ever required for $\varepsilon = 0$ (γ is the unique solution by [N:`bpN:p3:gen-uniqueness`]), which is not asserted by the frozen statement. The case $\varepsilon = 0$ is excluded from N-HEAD-C by (H7).

4. Hamiltonian ODE infrastructure (shared by tracks I, N, P)

This part is the single ODE layer of the three blueprints I, N, P. It has two levels. The *general layer* (§??–§??) is about an autonomous vector field F of class C^1 on an open subset W of a finite-dimensional real normed space E: local existence, uniqueness on *every* interval, maximal solutions, continuation (compact sets, and closed sets with uniform bounds), first exit from an open set. The *Hamiltonian layer* (§??) adds $F = X_H$: smoothness of X_H, energy conservation, translation in the angles. §?? shows that the frozen solution predicates of the three tracks (`IsSolutionOn`, `IsHamiltonianSolutionOn`, `IsRestrictedSolutionOn`, `IsNewtonSolution`, `IsFixedSolution`) are instances of the general predicate, and §?? gives the six track-N results of `PLAN.md`. Tracks I and P cite the general results as [N:`bpN:p3:gen-...`].

4.1. Conventions and external facts.

- E is a finite-dimensional real normed space (hence complete and proper: closed bounded sets are compact), $F : E \to E$ a *total* function, $W \subset E$ open. In the applications $E = \mathbb{R}^n \times \mathbb{R}^n$ with the max norm $|(\theta, I)| = \max(|\theta|, |I|)$ (the product norm of the Lean text, equal to the max norm on the $2n$ coordinates), or $E = (\mathbb{R}^2)^N \times (\mathbb{R}^2)^N$ for the Newtonian problems.
- F is C^1 *on* W means `ContDiffOn` $\mathbb{R}$ 1 F W; as W is open this gives $F \in C^1$ at each point of W, and no information off W. Off W the values of F never enter (Lean totality, below).
- An *interval* is an order-connected subset of $\mathbb{R}$ (Lean `OrdConnected`); open, closed, half-open, bounded or not, possibly a single point. $\overline{B}(x, \delta)$ is the closed ball.
- All derivatives $\dot{z}(t)$ in solution predicates are two-sided (Lean `HasDerivAt`) at every $t \in J$, including endpoints of J, exactly as in the frozen predicates ("z is defined on $\mathbb{R}$").

External facts. Mathlib-level statements, with the declarations that exist (checked in `Mathlib/Analysis/ODE/`); what is *not* in Mathlib is listed after them.

(E1) *Picard–Lindelöf, quantitative form.* Let $F : E \to E$, $x \in E$, $a, M, L \geq 0$ with $\|F(y)\| \leq M$ and F L-Lipschitz on $\overline{B}(x, a)$, $T \geq 0$ with $MT \leq a$. Then there is $\alpha : \mathbb{R} \to E$ with $\alpha(t_0) = x$ and α has derivative $F(\alpha(t))$ within $[t_0 - T, t_0 + T]$ at each t of that interval. (Mathlib: `ODE.IsPicardLindelof` with $r = 0$, time-independent field, and `IsPicardLindelof.exists_eq_forall_mem_Icc_hasDerivWithinAt_0`; the time constraint is the field `mul_max_le`: $M \max(t_{\max} - t_0, t_0 - t_{\min}) \leq a - r$. The contraction on iterates means no smallness of LT is needed.) Exists.

(E2) *Uniqueness with a Lipschitz constant on a set.* If $S \subset E$ and F is L-Lipschitz on S, $f, g : [a, b] \to E$ are continuous, have right derivative $F(f(t))$, $F(g(t))$ on $[a, b)$, take values in S and $f(a) = g(a)$, then $f = g$ on $[a, b]$; and the mirror statement with $f(b) = g(b)$ and left derivatives. (Mathlib: `ODE_solution_unique_of_mem_Icc_right`, `ODE_solution_unique_of_mem_Icc_left`.) Exists.

(E3) C^1 *implies locally Lipschitz.* If F is C^1 at p there are L and a neighbourhood on which F is L-Lipschitz (Mathlib: `ContDiffAt.exists_lipschitzOnWith`). Exists.

(E4) *Mean value inequality*: [K:`bp:p1:ext-mvt`]; for a differentiable map on a convex set with $\|DF\| \leq L$, F is L-Lipschitz there; for a function of one variable with derivative bounded by M on an interval, $\|\alpha(u) - \alpha(s)\| \leq M|u - s|$. Exists.

(E5) *Topology.* Heine–Borel in E (`ProperSpace`); continuous images of compact sets are compact; sequential compactness; $x \mapsto \operatorname{dist}(x, A)$ is continuous and positive on $E \setminus A$ for A closed (`Metric.infDist`). Exists.

(E6) *Calculus.* Chain rule `HasFDerivAt.comp_hasDerivAt`; a function continuous on $[s, t]$ with right derivative 0 on $[s, t)$ is constant (`constant_of_has_deriv_right_zero`); two one-sided derivatives agree ⇒ two-sided one (`HasDerivWithinAt.union` with $(-\infty, b] \cup [b, \infty) = \mathbb{R}$ and `hasDerivWithinAt_univ`); `ContDiffOn.fderiv_of_isOpen`: if H is C^{k+1} on an open set then DH is C^k there. Exists.

Not in Mathlib (checked: no maximal-solution, continuation, first-exit or uniform-existence-time statement in `Analysis/ODE`): everything of §??–§?? beyond (E1)–(E3), notably Lemmas ??, ??, Proposition ??, Theorems ??, ??, ??, and the Hamiltonian layer. They must be proved from the blueprint below.

Remark 4.1 (Lean totality). `KAM.hamiltonianVectorField` H uses `fderiv`, which is 0 at a point of non-differentiability, and the Newtonian fields use $1/\ell^3$ with $\ell =$ `planeLength` or `radius`. In every use below these are guarded: X_H occurs only on an open set where H is C^2 (or inside the frozen predicates, which assert differentiability at every visited point), and $\ell > 0$ at every visited point because the frozen predicates require it. The general layer never uses F off W.

4.2. **The general solution predicate; Lipschitz data.**

Definition 4.2 (Solutions in W)**.** For $F : E \to E$, $W \subset E$, $J \subset \mathbb{R}$ and $z : \mathbb{R} \to E$: `IsSolIn` $F\,W\,J\,z$ holds if for every $t \in J$, $z(t) \in W$ and z has derivative $F(z(t))$ at t (two-sided). Suggested Lean name `KAM.ODE.IsSolutionIn`; it is *not* a frozen declaration.
Source: Standard; shape dictated by the frozen predicates of Definition ?? and §??.

Elementary consequences (each a one-line Lean lemma, ??): (i) restriction: `IsSolIn` $F\,W\,J\,z$ and $J' \subset J$ give `IsSolIn` $F\,W\,J'\,z$; (ii) z is continuous on J (differentiable at each point of J); (iii) *local nature*: if J' is a neighbourhood of each of its points and $z = y$ on it, then `IsSolIn` $F\,W\,J'\,y \Leftrightarrow$ `IsSolIn` $F\,W\,J'\,z$ for open J'.

Lemma 4.3 (Symmetries)**.** *Let* `IsSolIn` $F\,W\,J\,z$.

(a) (time shift) For $c \in \mathbb{R}$, $t \mapsto z(t+c)$ is a solution on $J - c$.
(b) (time reversal) $t \mapsto z(-t)$ is a solution of $-F$ on $-J$, and $-F$ is C^1 on W if F is.
(c) (space translation) If $v \in E$ satisfies $W + v = W$ and $F(x+v) = F(x)$ for $x \in W$, then $t \mapsto z(t) + v$ is a solution of F in W on J.

Source: Standard.

Proof. **Step S4.1: Chain rule.** (a) and (b) are the chain rule for $t \mapsto t+c$, $t \mapsto -t$ (??); (c): $z+v$ has the same derivative $F(z(t)) = F(z(t)+v)$, and $z(t)+v \in W+v = W$. □

Lemma 4.4 (Bounds and Lipschitz constants on compact sets)**.** *Let F be C^1 on the open set W and $K \subset W$ compact. Then F is bounded on K and there is $L \geq 0$ with $\|F(x) - F(y)\| \leq L\|x - y\|$ for $x, y \in K$.*
Source: Standard.

Proof. **Step S4.2: Bounded.** F is continuous on $W \supset K$, so $\|F\| \leq M$ on the compact K (??).

Step S4.3: Contradiction hypothesis. If no L works, for every $k \in \mathbb{N}$ there are $x_k \neq y_k$ in K with $\|F(x_k) - F(y_k)\| > k\|x_k - y_k\|$; then $\|x_k - y_k\| < 2M/k \to 0$.

Step S4.4: Compactness. A subsequence $x_{k_j} \to p \in K$, hence also $y_{k_j} \to p$. By ?? there are L_p and a neighbourhood N_p of p on which F is L_p-Lipschitz. For large j both points lie in N_p and $k_j > L_p$, which contradicts Step ??. □

Definition 4.5 (Uniform data)**.** Let F be C^1 on the open set W. A set $\Phi \subset E$ has *uniform data* (δ, M, L), with $\delta > 0$, $M, L \geq 0$, if for every $x \in \Phi$: $\overline{B}(x, \delta) \subset W$, $\|F\| \leq M$ on $\overline{B}(x, \delta)$ and F is L-Lipschitz on $\overline{B}(x, \delta)$. (Balls are convex, so by ?? a bound $\|DF\| \leq L$ on $\overline{B}(x, \delta) \subset W$ gives the Lipschitz clause.)

Lemma 4.6 (Compact sets have uniform data)**.** *Let F be C^1 on the open set W and $K \subset W$ compact. Then K has uniform data (δ, M, L) for some $\delta > 0$, and $K_\delta = \{y : \operatorname{dist}(y, K) \leq \delta\} \subset W$.*

Proof. **Step S4.5: Choice of δ.** If $W = E$ put $\delta = 1$. Otherwise $x \mapsto \operatorname{dist}(x, E \setminus W)$ is continuous and positive on K, so has a minimum $2\delta > 0$ (**??**); then $K_\delta \subset W$. K_δ is closed and bounded, hence compact.

Step S4.6: Data. Apply Lemma **??** to K_δ: $\|F\| \le M$ and F is L-Lipschitz on $K_\delta \supset \overline{B}(x,\delta)$ for $x \in K$. □

Lemma 4.7 (Uniform data for angle-periodic fields)**.** *Let $E = \mathbb{R}^n \times \mathbb{R}^n$ (max norm), $U \subset \mathbb{R}^n$ open, $W = \mathbb{R}^n \times U$, F C^1 on W with $F(\theta + k, I) = F(\theta, I)$ for $(\theta, I) \in W$, $k \in \mathbb{Z}^n$. If $K \subset U$ is compact, then $\Phi = \mathbb{R}^n \times K$ has uniform data (δ, M, L) for some $\delta \in (0, 1]$.*

Proof. **Step S4.7: A compact cell.** Choose $\delta \in (0, 1]$ with $K_\delta = \{I : \operatorname{dist}(I, K) \le \delta\} \subset U$ (as in Step **??**). $C = [-1, 2]^n \times K_\delta$ is compact and contained in W; let M, L_C be the constants of Lemma **??** for C.

Step S4.8: Translating a ball into the cell. Let $x = (\theta, I) \in \Phi$, $k = (\lfloor \theta_i \rfloor)_i \in \mathbb{Z}^n$ and $x' = x - (k, 0) \in [0, 1)^n \times K$. Since the norm is the max norm and $\delta \le 1$, $\overline{B}(x', \delta) \subset C$, and $\overline{B}(x, \delta) = \overline{B}(x', \delta) + (k, 0) \subset W$. By periodicity $F(y + (k, 0)) = F(y)$ on $\overline{B}(x', \delta)$, so $\|F\| \le M$ and F is L_C-Lipschitz on $\overline{B}(x, \delta)$. □

4.3. **Local existence.**

Lemma 4.8 (A priori short-time confinement)**.** *Let J be an interval, $t_0 \in J$, $\alpha : \mathbb{R} \to E$ with $\alpha(t_0) = x$ and $\alpha'(t) = F(\alpha(t))$ for all $t \in J$ (derivative two-sided; F is only assumed to be a function), and suppose $\|F\| \le M$ on $\overline{B}(x, \delta)$ with $M > 0$. Then $\|\alpha(t) - x\| \le M|t - t_0|$ for all $t \in J$ with $|t - t_0| < \delta/M$; in particular $\alpha(t) \in \overline{B}(x, \delta)$.*
Source: Standard.

Proof. **Step S4.9: Forward bootstrap.** Let $t \in J$, $t_0 < t < t_0 + \delta/M$ (the case $t < t_0$ follows by Lemma **??**(b) and the case $t = t_0$ is trivial). Let $A = \{s \in [t_0, t] : \|\alpha(u) - x\| \le M(u - t_0)\ \forall u \in [t_0, s]\}$ and $s^* = \sup A$. $t_0 \in A$ and A is closed by continuity of α on J, so $s^* \in A$.

Step S4.10: Extension. If $s^* < t$, then $\|\alpha(s^*) - x\| \le M(s^* - t_0) < \delta$, so by continuity there is $\eta > 0$, $s^* + \eta \le t$, with $\alpha([s^*, s^* + \eta]) \subset \overline{B}(x, \delta)$. On this interval $\|\alpha'\| = \|F(\alpha)\| \le M$, so by **??** $\|\alpha(u) - \alpha(s^*)\| \le M(u - s^*)$ and hence $\|\alpha(u) - x\| \le M(u - t_0)$ there, contradicting the definition of s^*. So $s^* = t$. □

Proposition 4.9 (Local existence with uniform time)**.** *Let F be C^1 on the open set W and Φ have uniform data (δ, M, L); put $M' = \max(M, 1)$ and $\varepsilon = \delta/M'$. Then for every $x \in \Phi$ and every $t_0 \in \mathbb{R}$ there is $\alpha : \mathbb{R} \to E$ with $\alpha(t_0) = x$, $\|\alpha(t) - x\| < \delta$ and $\mathtt{IsSolIn}\, F\, W\, (t_0 - \varepsilon, t_0 + \varepsilon)\, \alpha$.*
Source: Standard (Picard–Lindelöf).

Proof. **Step S4.11: Picard–Lindelöf.** Apply **??** with $a = \delta$, bound M', Lipschitz constant L and $T = \varepsilon$ ($M'T = \delta \le a$): there is α with $\alpha(t_0) = x$ and derivative $F(\alpha(t))$ within $[t_0 - \varepsilon, t_0 + \varepsilon]$ at each point of it. At points t of the open interval this is a two-sided derivative, since the closed interval is a neighbourhood of t.

Step S4.12: Staying in W. By Lemma **??** on $J = (t_0 - \varepsilon, t_0 + \varepsilon)$, $\|\alpha(t) - x\| \le M'|t - t_0| < \delta$, so $\alpha(t) \in \overline{B}(x, \delta) \subset W$. □

Theorem 4.10 (Local existence, both time directions)**.** *Let F be C^1 on the open set W, $x_0 \in W$, $t_0 \in \mathbb{R}$. There are $\varepsilon > 0$ and $\alpha : \mathbb{R} \to E$ with $\alpha(t_0) = x_0$ and $\mathtt{IsSolIn}\, F\, W\, (t_0 - \varepsilon, t_0 + \varepsilon)\, \alpha$. Moreover ε may be taken uniform on any compact $K \subset W$ (for all $x_0 \in K$).*
Source: Standard.

Proof. **Step S4.13: Reduction.** $\{x_0\}$ (resp. K) is compact; Lemma **??** gives uniform data and Proposition **??** the claim. □

4.4. **Uniqueness on every interval.**

Theorem 4.11 (Uniqueness on arbitrary intervals)**.** *Let F be C^1 on the open set W, $J \subset \mathbb{R}$ an interval (of any kind: open, closed, half-open, unbounded, or a point), $t_0 \in J$, and $y_1, y_2 : \mathbb{R} \to E$ with* `IsSolIn` $F\,W\,J\,y_i$ *and $y_1(t_0) = y_2(t_0)$. Then $y_1 = y_2$ on J. No confinement, boundedness or smallness of either solution is assumed. In particular, if y_1, y_2 are solutions on intervals J_1, J_2 and agree at one point of $J_1 \cap J_2$, they agree on $J_1 \cap J_2$ (an interval), and two solutions which agree at one time of J agree at all times of J.*
Source: *Standard; used for BM11 Theorem 2.1 (uniqueness) and in the maximal-solution argument of BM11 p. 13.*

Proof. **Step S4.14: A compact piece.** Let $t_1 \in J$, $t_1 > t_0$ ($t_1 < t_0$ is the mirror image, using the left-derivative form of **??**; $t_1 = t_0$ is the hypothesis). Then $[t_0, t_1] \subset J$, each y_i is continuous on it, so $K = y_1([t_0, t_1]) \cup y_2([t_0, t_1])$ is compact (**??**) and $K \subset W$.

Step S4.15: Lipschitz on K. Lemma **??** gives L with F L-Lipschitz on K.

Step S4.16: Apply ??. Both y_i are continuous on $[t_0, t_1]$, have right derivative $F(y_i(t))$ on $[t_0, t_1)$ (a two-sided derivative gives a right derivative) and take values in K; so $y_1(t_1) = y_2(t_1)$. □

4.5. **Maximal solutions and continuation.**

Lemma 4.12 (Extension to an open interval)**.** *Let F be C^1 on the open set W, $J \neq \emptyset$ an interval and* `IsSolIn` $F\,W\,J\,y$*. Then there are an open interval $J' \supset J$ and y' with $y' = y$ on J and* `IsSolIn` $F\,W\,J'\,y'$*.*

Proof. **Step S4.17: Right end.** If J has a largest element b, apply Theorem **??** at $x_0 = y(b) \in W$, $t_0 = b$: α on $(b - \varepsilon, b + \varepsilon)$. Put $y_1(s) = y(s)$ for $s \le b$ and $y_1(s) = \alpha(s)$ for $s > b$. Then y_1 is a solution on $J_1 = J \cup (b, b + \varepsilon)$: at $t \in J$, $t < b$, and at $t \in (b, b + \varepsilon)$ it agrees with y, resp. α, near t; at $t = b$ it has derivative $F(y(b))$ within $(-\infty, b]$ (it equals y there) and within $[b, \infty)$ (it equals α there, as $\alpha(b) = y(b)$), so by **??** it has the two-sided derivative. If J has no largest element, $J_1 = J$, $y_1 = y$.

Step S4.18: Left end. Apply Step **??** to the reversed solution $y_1(-\cdot)$ (Lemma **??**(b)). The result has no endpoint, so is an open interval. □

Theorem 4.13 (Maximal solution)**.** *Let F be C^1 on the open set W and $x_0 \in W$. There are $T_- \in [-\infty, 0)$, $T_+ \in (0, +\infty]$ and $z : \mathbb{R} \to E$ with $z(0) = x_0$ and* `IsSolIn` $F\,W\,(T_-, T_+)\,z$*, such that every solution y with $y(0) = x_0$ on an interval $J \ni 0$ satisfies $J \subset (T_-, T_+)$ and $y = z$ on J. Then (T_-, T_+) is the* maximal interval *and $z|_{(T_-, T_+)}$ the* maximal solution*; both are unique. For initial time t_0 instead of 0 use Lemma **??**(a).*
Source: *BM11 p. 13 ("maximal time of existence"); standard.*

Proof. **Step S4.19: The union.** Let $\mathcal{J}$ be the set of open intervals $J \ni 0$ carrying a solution y_J with $y_J(0) = x_0$; $\mathcal{J} \neq \emptyset$ by Theorem **??**. The union I of $\mathcal{J}$ is an open interval $(T_-, T_+) \ni 0$ (a union of intervals containing 0).

Step S4.20: Gluing. For $t \in I$ choose $J \in \mathcal{J}$ with $t \in J$ and set $z(t) = y_J(t)$ (and $z(t) = x_0$ outside I). This is independent of J: for $J, J' \in \mathcal{J}$, $y_J = y_{J'}$ on $J \cap J' \ni 0$ by Theorem **??**. Each $t \in I$ has an open neighbourhood $J \in \mathcal{J}$ on which $z = y_J$, so z is a solution on I (local nature, (iii) above).

Step S4.21: Maximality. If y is a solution on an interval $J \ni 0$ with $y(0) = x_0$, Lemma **??** gives an open $J' \supset J$ and a solution y' extending y, so $J' \in \mathcal{J}$, $J \subset J' \subset I$, and $y = y' = z$ on J (Theorem **??**). Uniqueness of (T_-, T_+) and z is immediate from the maximality property applied twice. □

Theorem 4.14 (Continuation criterion, uniform form)**.** *Let F be C^1 on the open set W, $x_0 \in W$, (T_-, T_+) and z the maximal interval and solution of Theorem* **??***. Let $\Phi \subset E$ have uniform data (δ, M, L) (Definition* **??***). If $T_+ < \infty$ there is no $t_* \in [0, T_+)$ with $z(t) \in \Phi$ for all $t \in [t_*, T_+)$. Symmetrically, if $T_- > -\infty$ there is no $t_* \in (T_-, 0]$ with $z(t) \in \Phi$ for all $t \in (T_-, t_*]$. Equivalently: if the solution stays in Φ near the right (left) end of its maximal interval, that end is $+\infty$ $(-\infty)$.*
Source: Standard; BM11 p. 13 ("this solution is defined for all time"), Pöschel 1993 p. 198 ("$T = T_*$").

Proof. **Step S4.22: Set-up.** Suppose $T_+ < \infty$ and $z(t) \in \Phi$ on $[t_*, T_+)$. Let $\varepsilon = \delta / \max(M, 1)$ from Proposition **??**. Choose $t_1 \in [t_*, T_+)$ with $t_1 > T_+ - \varepsilon$.

Step S4.23: A solution through $z(t_1)$. Proposition **??** with $x = z(t_1) \in \Phi$ and $t_0 = t_1$ gives α on $(t_1 - \varepsilon, t_1 + \varepsilon)$ with $\alpha(t_1) = z(t_1)$. By Theorem **??** on $(t_1 - \varepsilon, t_1 + \varepsilon) \cap (T_-, T_+)$, $\alpha = z$ there.

Step S4.24: Gluing past T_+. Put $y = z$ on (T_-, T_+) and $y = \alpha$ on $[T_+, t_1 + \varepsilon)$; then $y = \alpha$ on $(t_1 - \varepsilon, t_1 + \varepsilon)$ and $y = z$ on (T_-, T_+), two open sets covering $(T_-, t_1 + \varepsilon)$ on each of which y is a solution. So y is a solution on $(T_-, t_1 + \varepsilon) \ni 0$ with $y(0) = x_0$, and $t_1 + \varepsilon > T_+$ contradicts Theorem **??**. The left end follows by Lemma **??**(b). □

Corollary 4.15 (Compact continuation)**.** *In the setting of Theorem* **??***, if $K \subset W$ is compact and $z(t) \in K$ for all $t \in [t_*, T_+)$ (some $t_* \in [0, T_+)$), then $T_+ = +\infty$; likewise on the left.*

Proof. Lemma **??** gives uniform data for K; apply Theorem **??**. □

Corollary 4.16 (Closed sets, periodic fields)**.** *Let $E = \mathbb{R}^n \times \mathbb{R}^n$, $U \subset \mathbb{R}^n$ open, $W = \mathbb{R}^n \times U$, F C^1 on W with $F(\theta + k, I) = F(\theta, I)$ for $k \in \mathbb{Z}^n$. If the maximal solution through $x_0 \in W$ satisfies $z(t)_2 \in K$ for $t \in [t_*, T_+)$ with $K \subset U$ compact, then $T_+ = +\infty$ (likewise on the left). The angle components $z(t)_1$ may be unbounded.*

Proof. Lemma **??** gives uniform data for $\Phi = \mathbb{R}^n \times K$; apply Theorem **??**. □

Remark 4.17 (What the closed-set variant really needs)*.* Staying in a closed non-compact set $\Phi \subset W$ does *not* suffice in general (the solution of $\dot{x} = x^2$ on $E = \mathbb{R}$, $W = \mathbb{R}$, $x_0 = 1$ stays in the closed set $[1, \infty)$ and blows up at $T_+ = 1$). What suffices is the uniform data (δ, M, L) of Definition **??**: a δ-neighbourhood of Φ inside W on which F is bounded and Lipschitz, uniformly. This holds for compact sets, for angle-periodic fields over a compact action set (Lemma **??**), and for the Newtonian regions bounded away from collisions with bounded velocities (Corollary **??**).

Theorem 4.18 (Confinement glue)**.** *Let F be C^1 on the open set W, $x_0 \in W$, $\tau > 0$, and $\Phi \subset E$ a set with uniform data (e.g. compact $\Phi \subset W$, or as in Lemma* **??***). Assume the* a priori confinement*: every solution y on an open interval $J \ni 0$ with $y(0) = x_0$ satisfies $y(t) \in \Phi$ for all $t \in J$ with $|t| \le \tau$. Then the maximal interval contains $[-\tau, \tau]$ in its interior, so z is a solution on $[-\tau, \tau]$ with $z(t) \in \Phi$ for $|t| \le \tau$, and every solution on $[-L, L]$ $(0 \le L \le \tau)$ with initial value x_0 equals z there.*
Source: BM11 p. 13 (the two-case argument "Therefore this solution is defined for all time"); Pöschel 1993 p. 198.

Proof. **Step S4.25: No exit before τ.** Suppose $T_+ \le \tau$. The maximal solution z on (T_-, T_+) satisfies, for $t \in [0, T_+)$ $(|t| < \tau)$, $z(t) \in \Phi$ by confinement. Theorem **??** gives $T_+ = \infty$, a contradiction. So $T_+ > \tau$, and likewise $T_- > -\tau$.

Step S4.26: Conclusion. $[-\tau, \tau] \subset (T_-, T_+)$ so z is a solution on $[-\tau, \tau]$ and $z(t) \in \Phi$ for $|t| \le \tau$ by confinement applied to z itself. Uniqueness is Theorem **??**. □

4.6. **First exit from an open set; barriers.**

Lemma 4.19 (First exit)**.** *Let* `IsSolIn F W J z` *with J an interval, $0 \in J$, $V \subset E$ open and $z(0) \in V$. Put $A = \{t \in J : t \geq 0,\ z(t) \notin V\}$. Then exactly one of the following holds.*

(i) $A = \emptyset$: $z(t) \in V$ for all $t \in J$, $t \geq 0$.

(ii) $A \neq \emptyset$. Then $t^* = \min A$ exists, $t^* > 0$, $t^* \in J$, $z(t) \in V$ for $t \in [0, t^*)$ and $z(t^*) \in \partial V = \overline{V} \setminus V$. The number t^* is the first exit time *and equals $\sup\{t \in J : t \geq 0,\ z([0,t]) \subset V\}$; in particular t^* is finite and lies in J (t^* is "inside the solution's interval" and the solution is defined, continuous, at t^*).*

The same holds backwards in time (replace $z(t)$ by $z(-t)$, Lemma **??***(b)). Only continuity of z on J is used; F and W play no role.*

Source: Standard; used for BM11 p. 13 ("smallest time t_* such that $\omega(t_*) \in R_K$") and Pöschel 1993 p. 198–199 (stopping-time arguments).

Proof. **Step S4.27: Closedness.** z is continuous on J (consequence (ii)), $E \setminus V$ is closed, so A is closed in $J \cap [0, \infty)$.

Step S4.28: The minimum. If $A \neq \emptyset$, $t^* = \inf A \in [0, \infty)$; $t^* \in J$ since J is an interval containing 0 and a point of A that is $\geq t^*$. By closedness $t^* \in A$, i.e. $z(t^*) \notin V$. As $z(0) \in V$, $t^* > 0$.

Step S4.29: Boundary point. For $t \in [0, t^*)$, $t \notin A$, so $z(t) \in V$. By continuity $z(t^*) = \lim_{t \uparrow t^*} z(t) \in \overline{V}$, and $z(t^*) \notin V$; as V is open, $\overline{V} \setminus V = \partial V$.

Step S4.30: The supremum formula. $z([0,t]) \subset V$ iff $t < t^*$ (for $t \in J$, $t \geq 0$), so the supremum is t^*. □

Definition 4.20 (Invariant sets)**.** $S \subset E$ is *invariant* for (F, W) if for every interval J' and every y with `IsSolIn F W J' y`: if $y(t_1) \in S$ for one $t_1 \in J'$ then $y(t) \in S$ for all $t \in J'$. (Equilibria $\{x : F(x) = 0\}$ are invariant by Theorem **??**; so is any union of such sets and any set $\{G = c\}$ for a first integral G.)

Corollary 4.21 (Barrier)**.** *Let F be C^1 on the open set W, V open,* `IsSolIn F W J z` *with J an interval, $0 \in J$, $z(0) \in V$. Let $G \subset E$ with $z(J) \subset G$, and suppose $\partial V \cap G \subset S$ where S is invariant and $z(0) \notin S$ (e.g. G an energy level, Theorem* **??***, and $\partial V \cap G$ a union of invariant sets not meeting the initial point). Then $z(t) \in V$ for all $t \in J$ (both time directions).*

Proof. **Step S4.31: Forward.** If not, Lemma **??**(ii) gives $t^* \in J$ with $z(t^*) \in \partial V \cap G \subset S$; invariance gives $z(0) \in S$, a contradiction.

Step S4.32: Backward. Apply Step **??** to $t \mapsto z(-t)$, a solution of $-F$ on $-J$; G, ∂V are unchanged and S is invariant for $-F$ as well (reversal maps solutions of F to solutions of $-F$ and back). □

4.7. **Hamiltonian layer.** Phase space $P = \mathbb{R}^n \times \mathbb{R}^n$, points $z = (\theta, I)$, max norm. X_H is the frozen `KAM.hamiltonianVectorField`, Definition **??** (convention of [K:`bp:p0:hamiltonian`]): $X_H(z) = \big((\partial_{I_i} H(z))_i, (-\partial_{\theta_i} H(z))_i\big)$, with $\partial_{I_i} H(z) = DH(z)(0, e_i)$ and $\partial_{\theta_i} H(z) = DH(z)(e_i, 0)$.

Lemma 4.22 (Regularity of X_H)**.** *Let $W \subset P$ be open and $H : P \to \mathbb{R}$ be C^{k+1} on W, $k \geq 0$. Then X_H is C^k on W; for $k \geq 1$ it is C^1 (hence locally Lipschitz) on W, and H is differentiable at every point of W.*

Source: Standard.

Proof. **Step S4.33: Derivative is C^k.** $DH : W \to \mathcal{L}(P, \mathbb{R})$ is C^k on the open W (**??**, `ContDiffOn.fderiv_of_isOpen`).

Step S4.34: Components. Each coordinate of X_H is $\pm$ the composition of DH with the continuous linear evaluation $\ell \mapsto \ell(0, e_i)$ or $\ell \mapsto \ell(e_i, 0)$; hence C^k, and a map into $\mathbb{R}^n \times \mathbb{R}^n$ with C^k coordinates is C^k. □

Theorem 4.23 (Energy conservation)**.** *Let $H : P \to \mathbb{R}$, $J \subset \mathbb{R}$ an interval and $z : \mathbb{R} \to P$ such that for every $t \in J$, H is differentiable at $z(t)$ and z has derivative $X_H(z(t))$ at t. Then $t \mapsto H(z(t))$ is constant on J. (No regularity of H beyond the stated pointwise differentiability, and no confinement.)*
Source: BM11 p. 13 and Pöschel 1993 p. 198 ("energy conservation", "$\Delta H = H(I(t), \theta(t))|_0^T$ vanishes"); standard.

Proof. **Step S4.35: Chain rule.** For $t \in J$, $\frac{d}{dt} H(z(t)) = DH(z(t))\big[X_H(z(t))\big]$ (**??**).

Step S4.36: Cancellation. Write $z = z(t)$, $\ell = DH(z)$ and $X_H(z) = (a, b)$ with $a_i = \ell(0, e_i)$, $b_i = -\ell(e_i, 0)$. By linearity, $\ell(a, b) = \sum_i a_i\, \ell(e_i, 0) + \sum_i b_i\, \ell(0, e_i) = \sum_i \ell(0, e_i)\ell(e_i, 0) - \sum_i \ell(e_i, 0)\ell(0, e_i) = 0$.

Step S4.37: Constancy. Let $s < t$ in J. $H \circ z$ is continuous on $[s, t] \subset J$ and has derivative 0 at every point, so $H(z(t)) = H(z(s))$ (**??**). □

Lemma 4.24 (Periodicity in the angles passes to solutions)**.** *Let $W \subset P$ be open with $W + (k, 0) = W$ for all $k \in \mathbb{Z}^n$, and $H : P \to \mathbb{R}$ with $H(\theta + k, I) = H(\theta, I)$ for $(\theta, I) \in W$, $k \in \mathbb{Z}^n$. Then $X_H(\theta + k, I) = X_H(\theta, I)$ on W. Consequently, if `IsSolIn` $X_H\, W\, J\, z$ then `IsSolIn` $X_H\, W\, J\, (z + (k, 0))$ for every $k \in \mathbb{Z}^n$, and $H \circ (z + (k, 0)) = H \circ z$.*
Source: Standard.

Proof. **Step S4.38: Field.** Let $\tau_k(x) = x + (k, 0)$. On the open set W, $H \circ \tau_k = H$, so $DH = D(H \circ \tau_k) = DH \circ \tau_k$ at each point of W (the derivative of a translate; both sides are 0 if H is not differentiable, by the definition of `fderiv`). Evaluating on $(0, e_i)$ and $(e_i, 0)$ gives $X_H \circ \tau_k = X_H$ on W.

Step S4.39: Solutions. Lemma **??**(c) with $v = (k, 0)$ applies to $F = X_H$. □

4.8. **The frozen solution predicates are instances.**

Proposition 4.25 (Track N and track I Hamiltonian predicates)**.** *Let $H : P \to \mathbb{R}$, $U \subset \mathbb{R}^n$ open, $W \subset P$ open with $\mathbb{R}^n \times U \subset W$, and H be C^2 on W. Put $W_U = \mathbb{R}^n \times U$. For every $J \subset \mathbb{R}$ and $z : \mathbb{R} \to P$,*

$$\texttt{IsSolutionOn}\, H\, U\, J\, z \iff \texttt{IsSolIn}\, X_H\, W_U\, J\, z \iff \texttt{IsHamiltonianSolutionOn}\, H\, U\, J\, z,$$

and X_H is C^1 on W_U. (The three predicates are literally the same formula up to the letter naming the action set, `KAM.Nekhoroshev.IsSolutionOn` and `KAM.Iso.IsHamiltonianSolutionOn`.)
Frozen Lean declaration: `KAM.Nekhoroshev.IsSolutionOn`, `KAM.Iso.IsHamiltonianSolutionOn`
Source: Ours (bookkeeping).

Proof. **Step S4.40: Membership.** $z(t)_2 \in U \iff z(t) \in W_U$ by definition of W_U.

Step S4.41: Differentiability is free. For $z(t) \in W_U \subset W$, H is C^2 on the open W, hence differentiable at $z(t)$ (Lemma **??**); so the clause `DifferentiableAt` $\mathbb{R}\, H\, (z\, t)$ is implied by $z(t) \in W_U$, and the two predicates coincide. Smoothness of X_H: Lemma **??**, $k = 1$, restricted to $W_U \subset W$. □

Proposition 4.26 (Newtonian predicates)**.** *Let $N \geq 2$, $G \in \mathbb{R}$, `mass` $: Fin\, N \to \mathbb{R}$. Let $E_N = (Fin\, N \to \mathbb{R}^2) \times (Fin\, N \to \mathbb{R}^2)$ (max norm), $\ell =$ `planeLength`,*

$$W_N = \{(q, v) : \ell(q_j - q_i) > 0\ \forall i \neq j\}, \qquad F_N(q, v) = \Big(v,\ \big(\textstyle\sum_{j \neq i} \frac{G\, \mathrm{mass}_j}{\ell(q_j - q_i)^3}(q_j - q_i)\big)_i\Big).$$

Then W_N is open, F_N is C^∞ on W_N, and for $Q, V : \mathbb{R} \to (Fin\,N \to \mathbb{R}^2)$ with $x(t) = (Q(t), V(t))$: `IsNewtonSolution` $G\,\mathrm{mass}\,Q\,V \iff$ `IsSolIn` $F_N\,W_N\,\mathbb{R}\,x$. Likewise for `IsFixedSolution` $G\,m_0\,m\,\varepsilon$ with $W_N' = W_N \cap \{\ell(q_i) > 0\ \forall i\}$ and the field $F'(q,v) = (v, (-\frac{Gm_0}{\ell(q_i)^3} q_i + \varepsilon \sum_{j\neq i} \frac{Gm_j}{\ell(q_j-q_i)^3}(q_j - q_i))_i)$, and for `IsRestrictedSolutionOn` $\mu\,J$ with $W_\mu = \{(q,p) :$ `CollisionFree` $\mu\,(q,p)\}$ and the total extension F_μ of `restrictedVectorField` μ by the same formula (with the values off W_μ irrelevant): `IsRestrictedSolutionOn` $\mu\,J\,z \iff$ `IsSolIn` $F_\mu\,W_\mu\,J\,z$. In all three cases F is C^∞ on the open set W and the general layer applies with $J = \mathbb{R}$ (resp. the given J).

Frozen Lean declaration: `KAM.Planetary.IsNewtonSolution`, `KAM.Planetary.IsFixedSolution`, `KAM.Iso.IsRestrictedSolutionOn`

Source: Ours (bookkeeping); Fejoz Section 6.1, Newton equations as in the frozen text.

Proof. **Step S4.42: Open domain and smoothness.** $\ell(w)^2 = \texttt{planeSq}(w) = w_0^2 + w_1^2$ is a polynomial, so $\{\ell > 0\} = \{\texttt{planeSq} > 0\}$ is open, and $\ell = \sqrt{\cdot}$ is C^∞ where $\texttt{planeSq} > 0$. Hence $\ell(q_j - q_i)^{-3}$ is C^∞ on W_N and F_N is a finite sum of products of C^∞ maps. Same for F' and F_μ ($\texttt{radius} = \sqrt{\texttt{radiusSq}}$ with the shifts by $\texttt{primary}_{1,2}\mu$).

Step S4.43: Derivatives of pairs. A map into a finite product has derivative (a,b) at t iff each component has the corresponding derivative (`hasDerivAt_pi`, `HasDerivAt.prodMk` and converse). The first block of the frozen definition (distance positivity) is membership in W_N; the second block says Q has derivative $V(t)$; the third says V has derivative the force. Together they say exactly that x has derivative $F_N(x(t))$ at t. □

Corollary 4.27 (Newtonian continuation)**.** *In the setting of Proposition **??**, let x be the maximal solution through $x_0 \in W_N$ (Theorem **??**), $\rho > 0$, $\bar V \geq 0$ and $\Phi_{\rho,\bar V} = \{(q,v) : \ell(q_j - q_i) \geq \rho\ (i \neq j),\ |v| \leq \bar V\}$. If $T_+ < \infty$ then for every $\rho, \bar V$ and every $t_* \in [0, T_+)$ there is $t \in [t_*, T_+)$ with $x(t) \notin \Phi_{\rho,\bar V}$ (a finite-time singularity needs near-collisions or unbounded velocities; positions may be unbounded). The same for F' with the additional condition $\ell(q_i) \geq \rho$, and for F_μ with $\ell \geq \rho$ from both primaries and $|q|, |p| \leq \bar V$.*

Proof. **Step S4.44: Uniform data.** Let $\delta = \rho/8$. If $|y - x| \leq \delta$ (max norm) with $x \in \Phi_{\rho,\bar V}$ then each $\ell(q_j' - q_i') \geq \rho - 2\sqrt{2}\,\delta \geq \rho/2$, so $y \in W_N$ and $\overline{B}(x,\delta) \subset W_N$. On this ball $\ell \geq \rho/2$ and $|v'| \leq \bar V + \delta$; the entries of F_N and DF_N are polynomials in v', the differences $q_j' - q_i'$ (Euclidean length $\leq \ell(q_j' - q_i')$) and the inverse powers ℓ^{-3}, ℓ^{-5} ($D(w/\ell^3) = \ell^{-3} I - 3\ell^{-5} w w^\top$), hence bounded by a constant $C(N, G, \mathrm{mass}, \rho, \bar V)$ *independent of the position of x*. By **??** on the convex ball, F_N is C-Lipschitz there. So $\Phi_{\rho,\bar V}$ has uniform data.

Step S4.45: Apply. Theorem **??** gives the claim; the other two fields are identical in structure (for F_μ the extra terms are linear in (q,p) and bounded on $|q|, |p| \leq \bar V$). □

4.9. **The track-N statements.** Throughout, $R, s > 0$, $h : \mathbb{C}^n \to \mathbb{C}$ with $\texttt{IsAnalyticIntegrable}\,R\,s\,h$, $\varepsilon \geq 0$ and $f : \mathbb{C}^n \times \mathbb{C}^n \to \mathbb{C}$ with $\texttt{IsAnalyticPerturbation}\,R\,s\,\varepsilon\,f$ (Definitions **??**, **??**), $H = H_{h,f}$ (Definition **??**), $W_s = \mathbb{R}^n \times B(0, R+s) \subset P$ (open; $B(0,R+s) = \mathcal{B}_s(R)$ by Remark **??**), and $U \subset B(0, R+s)$ open (in N-HEAD-C, $U = B(0,R)$).

Lemma 4.28 (Input from Part 1: H is smooth on W_s)**.** *$H_{h,f}$ is C^∞ (in particular C^2) on the open set W_s, $X_H = X_{h+f}$ on $W_s \cap (\mathbb{R}^n \times \mathbb{R}^n)$, and $H(\theta + k, I) = H(\theta, I)$ for $(\theta, I) \in W_s$, $k \in \mathbb{Z}^n$.*

Source: Part 1 forward label [N:bpN:p1:holo-real]*; the two ingredients are* [K:bp:p1:holomorphic-regularity], [K:bp:p1:hamiltonian-real]*; periodicity is Definition **??**(ii).*

Proof. **Step S4.46: Regularity.** The function $G(\theta, I) = h(I) + f(\theta, I)$ is holomorphic on the open set $D_s(R) \supset W_s$ (real points (θ, I) with $I \in \mathcal{B}_s(R) = B(0,R)_s \cap \mathbb{R}^n$ lie in $T_s \times B(0,R)_s$), hence

C^∞ in the real sense ([K:bp:p1:holomorphic-regularity]); it is real on W_s by Definition **??**(iv) and **??**(iii). So $H = \operatorname{Re} G = G$ on W_s is the restriction of C^∞ map to a real subspace: C^∞ on W_s; $X_H = X_G$ at real points is [K:bp:p1:hamiltonian-real]. (Packaged as [N:bpN:p1:holo-real].)

Step S4.47: Periodicity. $f(\theta + k, I) = f(\theta, I)$ for real θ, $I \in \mathcal{B}_s(R)$, and h depends on I only. □

Theorem 4.29 (X_H is C^1, hence locally Lipschitz)**.** *$X_{H_{h,f}}$ is C^1 on $W_s \supset \mathbb{R}^n \times B(0,R)$; consequently it is Lipschitz on every compact subset of W_s, and on every set $\mathbb{R}^n \times K$ with $K \subset B(0, R+s)$ compact there are δ, M, L as in Lemma **??**.*
Source: BM11 p. 13 (implicit in "the solution"); standard.

Proof. **Step S4.48: Smoothness.** Lemma **??** and Lemma **??** ($k = 1$).

Step S4.49: Lipschitz. Lemma **??** on compact sets, Lemma **??** (with $U = B(0, R+s)$, using the periodicity of X_H from Lemma **??** and Lemma **??**) for the cylinders. □

Theorem 4.30 (Local existence and uniqueness for `IsSolutionOn`)**.** *For every $t_0 \in \mathbb{R}$ and $z_0 \in \mathbb{R}^n \times U$ there are $\epsilon > 0$ and $y : \mathbb{R} \to P$ with $y(t_0) = z_0$ and $\mathtt{IsSolutionOn}\, H\, U\, (t_0 - \epsilon, t_0 + \epsilon)\, y$ (both time directions). If J is any interval, $t_0 \in J$ and y_1, y_2 satisfy $\mathtt{IsSolutionOn}\, H\, U\, J\, y_i$ with $y_1(t_0) = y_2(t_0)$, then $y_1 = y_2$ on J.*
*Source: BM11 p. 13 and Theorem 2.1 p. 5; Picard–Lindelöf (external fact **??**).*

Proof. **Step S4.50: Reduction.** Proposition **??** (with $W = W_s$, H C^2 there by Lemma **??**) turns $\mathtt{IsSolutionOn}\, H\, U$ into $\mathtt{IsSolIn}\, X_H\, (\mathbb{R}^n \times U)$ with X_H C^1 on $\mathbb{R}^n \times U$.

Step S4.51: Existence, uniqueness. Theorem **??** and Theorem **??**. □

Theorem 4.31 (Energy conservation for `IsSolutionOn`)**.** *If $\mathtt{IsSolutionOn}\, H\, U\, J\, z$ for an interval J (here H may be any function $P \to \mathbb{R}$ and U any set), then $H(z(t)) = H(z(t'))$ for $t, t' \in J$. In particular, for $H = H_{h,f}$ and $J = [0, T]$ the quantity $\Delta H = H(z(t))|_0^T$ of Pöschel's Lemma 1 vanishes.*
Source: Pöschel 1993 p. 198 ("by energy conservation ... ΔH vanishes"; page read); BM11 p. 13.

Proof. **Step S4.52: Direct.** The definition gives differentiability of H at every $z(t)$ and $z'(t) = X_H(z(t))$; Theorem **??**. □

Theorem 4.32 (Maximal solutions and continuation)**.** *Let $z_0 = (\theta_0, I_0)$ with $I_0 \in U$. There is a unique maximal interval $(T_-, T_+) \ni 0$ and a solution z with $z(0) = z_0$ and $\mathtt{IsSolutionOn}\, H\, U\, (T_-, T_+)\, z$, such that every y with $y(0) = z_0$ and $\mathtt{IsSolutionOn}\, H\, U\, J\, y$ on an interval $J \ni 0$ has $J \subset (T_-, T_+)$ and $y = z$ on J (Theorem **??** through Proposition **??**). If $K \subset U$ is compact and $z(t)_2 \in K$ for all $t \in [t_*, T_+)$ for some $t_* \in [0, T_+)$, then $T_+ = +\infty$; likewise on the left. The angle components $z(t)_1$ may be unbounded: no obstruction arises because X_H is 1-periodic in θ.*
Source: BM11 p. 13 ("From this estimate we can deduce that the solution $I(t)$ belongs to some compact ball ... this solution is defined for all time"); Pöschel 1993 p. 198 ("$T = T_*$").

Proof. **Step S4.53: General statements.** By Proposition **??**, $\mathtt{IsSolutionOn}\, H\, U\, J\, y \iff \mathtt{IsSolIn}\, X_H\, (\mathbb{R}^n \times U)\, J\, y$ and X_H is C^1 on $W = \mathbb{R}^n \times U$; maximality is Theorem **??**.

Step S4.54: Periodic cylinder. X_H is 1-periodic in θ on W (Lemmas **??**, **??**). Corollary **??** yields $T_+ = \infty$. □

Theorem 4.33 (Uniqueness on closed intervals)**.** *For $L \geq 0$ and y_1, y_2 with $y_1(0) = y_2(0)$ and $\mathtt{IsSolutionOn}\, H\, U\, [-L, L]\, y_i$ one has $y_1 = y_2$ on $[-L, L]$. In particular a solution on $[-L, L]$ agrees with every solution on a larger interval containing $[-L, L]$ and the same initial value (clause (d) of Theorem **??**, where $U = B(0, R)$, $L \in [0, \tau]$, y arbitrary and z the exhibited solution). No*

*confinement of y is assumed: Gronwall is replaced by Theorem **??**, whose Lipschitz constant comes from the compact set $y_1([-L,L]) \cup y_2([-L,L])$.*
Source: Standard (Gronwall); clause (d) of the frozen N-HEAD-C.

Proof. **Step S4.55: Apply.** $J = [-L, L]$, $t_0 = 0$ in Theorem **??** (uniqueness part). □

Theorem 4.34 (Confinement glue)**.** *Let $\theta_0 \in \mathbb{R}^n$, $I_0 \in U$, $\tau > 0$ and $\rho > 0$ with $\overline{B}(I_0, \rho) \subset U$ (e.g. $|I_0| < R/2$, $\rho \le R/4$, $U = B(0,R)$). Assume that for every open interval $J \ni 0$ and every y with $y(0) = (\theta_0, I_0)$ and `IsSolutionOn` $H\,U\,J\,y$ one has $|y(t)_2 - I_0| \le \rho$ for all $t \in J$ with $|t| \le \tau$. Then there is z with $z(0) = (\theta_0, I_0)$, `IsSolutionOn` $H\,U\,[-\tau,\tau]\,z$ and $|z(t)_2 - I_0| \le \rho$ for $|t| \le \tau$; and for every $L \in [0,\tau]$ and every y with $y(0) = z(0)$ and `IsSolutionOn` $H\,U\,[-L,L]\,y$, $y = z$ on $[-L,L]$. This is the assembly of clauses (a)–(d) of Theorem **??** from an a priori estimate (Part 10); $\rho = C\varepsilon^{1/(2n)}$ there.*
Source: BM11 p. 13 (two cases, "Therefore this solution is defined for all time"); Pöschel 1993 p. 198.

Proof. **Step S4.56: Compact confinement set.** $K = \overline{B}(I_0, \rho)$ is compact and $\subset U$; $\Phi = \mathbb{R}^n \times K$ has uniform data by Lemma **??** (with $W = \mathbb{R}^n \times U$, $F = X_H$ periodic by Lemma **??**).

Step S4.57: General glue. By Proposition **??** the hypothesis reads: every general solution on an open $J \ni 0$ with this initial value stays in Φ for $|t| \le \tau$. Theorem **??** gives z on $[-\tau,\tau]$ in Φ, i.e. $|z(t)_2 - I_0| \le \rho$, and uniqueness on $[-L,L]$ (it is also Theorem **??**). □

4.10. Label index for tracks I and P.

`bpN:p3:gen-sol`, `gen-symmetry`	predicate; time shift, reversal, space translation
`gen-lip`, `gen-uniform`, `gen-uniform-compact`, `gen-uniform-periodic`	Lipschitz and uniform data
`gen-short`, `gen-uniform-exist`, `gen-local`	local existence, uniform time
`gen-uniqueness`	uniqueness on all intervals, no confinement
`gen-extend`, `gen-maximal`	maximal solutions
`gen-continuation`, `-compact`, `-periodic`, `gen-confinement`	continuation, closed sets with uniform bounds, confinement glue
`gen-exit`, `gen-invariant`, `gen-barrier`	first exit, barrier
`ham-smooth`, `ham-energy`, `ham-periodic`	Hamiltonian layer
`ham-reduction`, `newton-reduction`, `newton-continuation`	frozen predicates of I, N, P
`holo-input`, `lipschitz`, `local-exist`, `energy`, `continuation`, `uniqueness-closed`, `confinement-glue`	track N specializations