

Simply Typed Lambda-calculus

Renato Neves



Universidade do Minho



Table of Contents

The Calculus

Denotational Semantics

Equational System

Disjunctive Types

The essence

Knowledge obtained via assumptions and logical rules

The essence

Knowledge obtained via assumptions and logical rules

Studied since Aristotle ...

... long before the age of artificial computers

What does it have to do with programming ?

A Basic Deductive System

$\mathbb{A}, \mathbb{B} \dots$ denote propositions and
1 a proposition that always holds



If \mathbb{A} and \mathbb{B} are propositions then

- $\mathbb{A} \times \mathbb{B}$ is a proposition – conjunction of \mathbb{A} and \mathbb{B}
- $\mathbb{A} \rightarrow \mathbb{B}$ is a proposition – implication of \mathbb{B} from \mathbb{A}

A Basic Deductive System

Γ denotes a list of propositions (often called context)

$\Gamma \vdash \mathbb{A}$ reads “if the propositions in Γ hold then \mathbb{A} also holds”

$$\frac{\mathbb{A} \in \Gamma}{\Gamma \vdash \mathbb{A}} \text{ (ass)} \quad \frac{}{\Gamma \vdash 1} \text{ (trv)} \quad \frac{\Gamma \vdash \mathbb{A} \times \mathbb{B}}{\Gamma \vdash \mathbb{A}} (\pi_1) \quad \frac{\Gamma \vdash \mathbb{A} \times \mathbb{B}}{\Gamma \vdash \mathbb{B}} (\pi_2)$$

$$\frac{\Gamma \vdash \mathbb{A} \quad \Gamma \vdash \mathbb{B}}{\Gamma \vdash \mathbb{A} \times \mathbb{B}} \text{ (prd)} \quad \frac{\Gamma, \mathbb{A} \vdash \mathbb{B}}{\Gamma \vdash \mathbb{A} \rightarrow \mathbb{B}} \text{ (cry)} \quad \frac{\Gamma \vdash \mathbb{A} \rightarrow \mathbb{B} \quad \Gamma \vdash \mathbb{A}}{\Gamma \vdash \mathbb{B}} \text{ (app)}$$

Exercise

Show that $\mathbb{A} \times \mathbb{B} \vdash \mathbb{B} \times \mathbb{A}$

New Knowledge From Old

The rules below are derivable from the previous system

$$\frac{\Gamma, A, B, \Delta \vdash C}{\Gamma, B, A, \Delta \vdash C} \text{ (exchange)}$$

$$\frac{\Gamma \vdash A}{\Gamma, B \vdash A} \text{ (weakening)}$$

$$\frac{\Gamma, A \vdash B \quad \Gamma \vdash A}{\Gamma \vdash B} \text{ (cut elimination)}$$

Proofs (again) by an appeal to your old friend ... induction :-)

Derive the following judgements

- $A \rightarrow B, B \rightarrow C \vdash A \rightarrow C$
- $A \rightarrow B, A \rightarrow C \vdash A \rightarrow B \times C$

Back to programming ...

The Bare Essentials of Programming

We should think of what are the basic features of programming ...

- variables
- function application and creation
- pairing ...

and base our study on the simplest language with such features ...

Simply-typed λ -calculus

The basis of Haskell, ML, Eff, F#, Agda, Elm and many other programming languages

Simply-typed λ -Calculus

Types are defined by $\mathbb{A} ::= 1 \mid \mathbb{A} \times \mathbb{A} \mid \mathbb{A} \rightarrow \mathbb{A}$

Γ now a non-repetitive list of typed variables ($x_1 : \mathbb{A}_1 \dots x_n : \mathbb{A}_n$)

Programs built according to the following deduction rules

$$\frac{x : \mathbb{A} \in \Gamma}{\Gamma \vdash x : \mathbb{A}} \text{ (ass)} \qquad \frac{}{\Gamma \vdash * : 1} \text{ (triv)} \qquad \frac{\Gamma \vdash t : \mathbb{A} \times \mathbb{B}}{\Gamma \vdash \pi_1 t : \mathbb{A}} \text{ (\pi}_1\text{)}$$

$$\frac{\Gamma \vdash t : \mathbb{A} \quad \Gamma \vdash s : \mathbb{B}}{\Gamma \vdash \langle t, s \rangle : \mathbb{A} \times \mathbb{B}} \text{ (prd)} \qquad \frac{\Gamma, x : \mathbb{A} \vdash t : \mathbb{B}}{\Gamma \vdash \lambda x : \mathbb{A}. t : \mathbb{A} \rightarrow \mathbb{B}} \text{ (cry)}$$

$$\frac{\Gamma \vdash t : \mathbb{A} \rightarrow \mathbb{B} \quad \Gamma \vdash s : \mathbb{A}}{\Gamma \vdash ts : \mathbb{B}} \text{ (app)}$$

Examples of λ -terms

$x : \mathbb{A} \vdash x : \mathbb{A}$ (identity)

$x : \mathbb{A} \vdash \langle x, x \rangle : \mathbb{A} \times \mathbb{A}$ (duplication)

$x : \mathbb{A} \times \mathbb{B} \vdash \langle \pi_2 x, \pi_1 x \rangle : \mathbb{B} \times \mathbb{A}$ (swap)

$f : \mathbb{A} \rightarrow \mathbb{B}, g : \mathbb{B} \rightarrow \mathbb{C} \vdash \lambda x : \mathbb{A}. g(f x) : \mathbb{A} \rightarrow \mathbb{C}$ (composition)

Recall the derivations that lead to the judgement

$$A \rightarrow B, A \rightarrow C \vdash A \rightarrow B \times C$$

Build the corresponding program

Derive as well the judgement

$$A \rightarrow B \vdash A \times C \rightarrow B \times C$$

and subsequently build the corresponding program

Table of Contents

The Calculus

Denotational Semantics

Equational System

Disjunctive Types

A Semantics for Simply Typed λ -calculus

We wish to assign a mathematical meaning to λ -terms

$$\llbracket - \rrbracket : \lambda\text{-terms} \longrightarrow \dots$$

so that we can reason about them rigorously, and take advantage of known mathematical theories

A Semantics for Simply Typed λ -calculus

We wish to assign a mathematical meaning to λ -terms

$$\llbracket - \rrbracket : \lambda\text{-terms} \longrightarrow \dots$$

so that we can reason about them rigorously, and take advantage of known mathematical theories

This is the goal of the next slides. But first ...

Functions: Basic Facts

For every set X there exists a 'trivial' function

$$! : X \longrightarrow \{\star\} = 1 \qquad !(x) = \star$$

We can always pair two functions into $f : X \rightarrow A, g : X \rightarrow B$

$$\langle f, g \rangle : X \rightarrow A \times B \qquad \langle f, g \rangle(x) = (f\ x, g\ x)$$

There exist projection functions

$$\pi_1 : X \times Y \rightarrow X \qquad \pi_1(x, y) = x$$

$$\pi_2 : X \times Y \rightarrow Y \qquad \pi_2(x, y) = y$$

Functions: Basic Facts

We can always ‘curry’ a function $f : X \times Y \rightarrow Z$ into

$$\lambda f : X \rightarrow Z^Y \quad \lambda f(x) = (y \mapsto f(x, y))$$

Consider sets X, Y, Z . There exists an application function

$$\text{app} : Z^Y \times Y \rightarrow Z \quad \text{app}(f, y) = f y$$

Types \mathbb{A} interpreted as sets $\llbracket \mathbb{A} \rrbracket$

$$\llbracket 1 \rrbracket = \{\star\}$$

$$\llbracket \mathbb{A} \times \mathbb{B} \rrbracket = \llbracket \mathbb{A} \rrbracket \times \llbracket \mathbb{B} \rrbracket$$

$$\llbracket \mathbb{A} \rightarrow \mathbb{B} \rrbracket = \llbracket \mathbb{B} \rrbracket^{\llbracket \mathbb{A} \rrbracket}$$

Typing contexts Γ interpreted as Cartesian products

$$\llbracket \Gamma \rrbracket = \llbracket x_1 : \mathbb{A}_1, \dots, x_n : \mathbb{A}_n \rrbracket = \llbracket \mathbb{A}_1 \rrbracket \times \dots \times \llbracket \mathbb{A}_n \rrbracket$$

λ -terms $\Gamma \vdash t : \mathbb{A}$ interpreted as functions

$$\llbracket \Gamma \vdash t : \mathbb{A} \rrbracket : \llbracket \Gamma \rrbracket \longrightarrow \llbracket \mathbb{A} \rrbracket$$

λ -term $\Gamma \vdash t : \mathbb{A}$ interpreted as a function

$$\llbracket \Gamma \vdash t : \mathbb{A} \rrbracket : \llbracket \Gamma \rrbracket \longrightarrow \llbracket \mathbb{A} \rrbracket$$

$$\frac{x_i : \mathbb{A} \in \Gamma}{\llbracket \Gamma \vdash x_i : \mathbb{A} \rrbracket = \pi_i}$$

$$\frac{}{\llbracket \Gamma \vdash * : 1 \rrbracket = !}$$

$$\frac{\llbracket \Gamma \vdash t : \mathbb{A} \times \mathbb{B} \rrbracket = f}{\llbracket \Gamma \vdash \pi_1 t : \mathbb{A} \rrbracket = \pi_1 \cdot f}$$

$$\frac{\llbracket \Gamma \vdash t : \mathbb{A} \rrbracket = f \quad \llbracket \Gamma \vdash s : \mathbb{B} \rrbracket = g}{\llbracket \Gamma \vdash \langle t, s \rangle : \mathbb{A} \times \mathbb{B} \rrbracket = \langle f, g \rangle}$$

$$\frac{\llbracket \Gamma, x : \mathbb{A} \vdash t : \mathbb{B} \rrbracket = f}{\llbracket \Gamma \vdash \lambda x : \mathbb{A}. t : \mathbb{A} \rightarrow \mathbb{B} \rrbracket = \lambda f}$$

$$\frac{\llbracket \Gamma \vdash t : \mathbb{A} \rightarrow \mathbb{B} \rrbracket = f \quad \llbracket \Gamma \vdash s : \mathbb{A} \rrbracket = g}{\llbracket \Gamma \vdash t s : \mathbb{B} \rrbracket = \text{app} \cdot \langle f, g \rangle}$$

The Unravelling

$$\llbracket x \vdash \langle \pi_2 x, \pi_1 x \rangle \rrbracket = \dots$$

$$\llbracket - \vdash \lambda x. \langle \pi_2 x, \pi_1 x \rangle \rrbracket = \dots$$

$$\llbracket f, g, x \vdash g f x \rrbracket = \dots$$

$$\llbracket f, g \vdash \lambda x. g f x \rrbracket = \dots$$

$$\llbracket f, x \vdash \langle f \pi_1 x, \pi_2 x \rangle \rrbracket = \dots$$

$$\llbracket f \vdash \lambda x. \langle f \pi_1 x, \pi_2 x \rangle \rrbracket = \dots$$

$$\llbracket - \vdash \lambda f. \lambda x. \langle f \pi_1 x, \pi_2 x \rangle \rrbracket = \dots$$

(**N.B.** all types omitted for simplicity)

Show that the following equations hold

$$\llbracket x, y \vdash \pi_1 \langle x, y \rangle \rrbracket = \llbracket x, y \vdash x \rrbracket$$

$$\llbracket \Gamma \vdash t \rrbracket = \llbracket \Gamma \vdash \langle \pi_1 t, \pi_2 t \rangle \rrbracket$$

$$\llbracket x \vdash (\lambda y. \langle x, y \rangle) x \rrbracket = \llbracket x \vdash \langle x, x \rangle \rrbracket$$

Denotational Semantics and Equivalence Revisited

Show that the following equations hold

$$\llbracket x, y \vdash \pi_1 \langle x, y \rangle \rrbracket = \llbracket x, y \vdash x \rrbracket$$

$$\llbracket \Gamma \vdash t \rrbracket = \llbracket \Gamma \vdash \langle \pi_1 t, \pi_2 t \rangle \rrbracket$$

$$\llbracket x \vdash (\lambda y. \langle x, y \rangle) x \rrbracket = \llbracket x \vdash \langle x, x \rangle \rrbracket$$

Show that the (complicated) λ -term below is really just the identity

$$z \vdash \lambda x. \langle \pi_2 x, \pi_1 x \rangle \left(\lambda y. \langle \pi_2 y, \pi_1 y \rangle z \right)$$

Denotational Semantics and Equivalence Revisited

Show that the following equations hold

$$\llbracket x, y \vdash \pi_1 \langle x, y \rangle \rrbracket = \llbracket x, y \vdash x \rrbracket$$

$$\llbracket \Gamma \vdash t \rrbracket = \llbracket \Gamma \vdash \langle \pi_1 t, \pi_2 t \rangle \rrbracket$$

$$\llbracket x \vdash (\lambda y. \langle x, y \rangle) x \rrbracket = \llbracket x \vdash \langle x, x \rangle \rrbracket$$

Show that the (complicated) λ -term below is really just the identity

$$z \vdash \lambda x. \langle \pi_2 x, \pi_1 x \rangle \left(\lambda y. \langle \pi_2 y, \pi_1 y \rangle z \right)$$

Hard ?

Table of Contents

The Calculus

Denotational Semantics

Equational System

Disjunctive Types

Logic to the Rescue !

Recall that the rules below are derivable from our logical system

$$\frac{\Gamma, \mathbb{A}, \mathbb{B}, \Delta \vdash \mathbb{C}}{\Gamma, \mathbb{B}, \mathbb{A}, \Delta \vdash \mathbb{C}} \text{ (exchange)}$$

$$\frac{\Gamma \vdash \mathbb{A}}{\Gamma, \mathbb{B} \vdash \mathbb{A}} \text{ (weakening)}$$

$$\frac{\Gamma, \mathbb{A} \vdash \mathbb{B} \quad \Gamma \vdash \mathbb{A}}{\Gamma \vdash \mathbb{B}} \text{ (cut elimination)}$$

$$\frac{\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \vdash t : \mathbb{C}}{\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \vdash t : \mathbb{C}} \text{ (exch)}$$

$$\frac{\Gamma \vdash t : \mathbb{A}}{\Gamma, x : \mathbb{B} \vdash t : \mathbb{A}} \text{ (weak)}$$

$$\frac{\Gamma, x : \mathbb{A} \vdash t : \mathbb{B} \quad \Gamma \vdash s : \mathbb{A}}{\Gamma \vdash \dots : \mathbb{B}} \text{ (cut elimination)}$$

$$\frac{\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \vdash t : \mathbb{C}}{\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \vdash t : \mathbb{C}} \text{ (exch)}$$

$$\frac{\Gamma \vdash t : \mathbb{A}}{\Gamma, x : \mathbb{B} \vdash t : \mathbb{A}} \text{ (weak)}$$

$$\frac{\Gamma, x : \mathbb{A} \vdash t : \mathbb{B} \quad \Gamma \vdash s : \mathbb{A}}{\Gamma \vdash \dots : \mathbb{B}} \text{ (cut elimination)}$$

Filling up the dots will lead us to a fundamental concept

$$\frac{\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \vdash t : \mathbb{C}}{\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \vdash t : \mathbb{C}} \text{ (exch)}$$

$$\frac{\Gamma \vdash t : \mathbb{A}}{\Gamma, x : \mathbb{B} \vdash t : \mathbb{A}} \text{ (weak)}$$

$$\frac{\Gamma, x : \mathbb{A} \vdash t : \mathbb{B} \quad \Gamma \vdash s : \mathbb{A}}{\Gamma \vdash \dots : \mathbb{B}} \text{ (cut elimination)}$$

Filling up the dots will lead us to a fundamental concept

Substitution

Substitution

The essence

Substitution of variables in a λ -term t by another λ -term s

$t[s/x]$ reads *"replace every occurrence of x in t by s "*

Substitution

The essence

Substitution of variables in a λ -term t by another λ -term s

$t[s/x]$ reads *"replace every occurrence of x in t by s "*

Example

$$\langle x, x \rangle[s/x] = \langle s, s \rangle$$

$$\langle x, y \rangle[s/x] = \langle s, y \rangle$$

$$\langle y, z \rangle[s/x] = \langle y, z \rangle$$

Substitution More Formally

We define it by induction

$$x[s/y] = \begin{cases} s & \text{if } x = y \\ x & \text{otherwise} \end{cases}$$

$$*[s/y] = *$$

$$\langle t_1, t_2 \rangle [s/y] = \langle t_1[s/y], t_2[s/y] \rangle$$

$$(t_1 \ t_2)[s/y] = t_1[s/y] \ t_2[s/y]$$

$$(\pi_1 t)[s/y] = \pi_1 t[s/y]$$

$$(\pi_2 t)[s/y] = \pi_2 t[s/y]$$

$$(\lambda x. t)[s/y] = \dots$$

$\lambda x. y$ is a “constant function” (given x return y)

Variable Captures

$\lambda x. y$ is a “constant function” (given x return y)

$(\lambda x. y)[z/y]$ is still a “constant function” (given x return z)

Variable Captures

$\lambda x. y$ is a “constant function” (given x return y)

$(\lambda x. y)[z/y]$ is still a “constant function” (given x return z)

$(\lambda x. y)[x/y]$ is now the identity !?

Variable Captures

$\lambda x. y$ is a “constant function” (given x return y)

$(\lambda x. y)[z/y]$ is still a “constant function” (given x return z)

$(\lambda x. y)[x/y]$ is now the identity !?

The problem: variable x “captured” by the construct “ $\lambda x.$ ”

Somehow similar to variable shadowing in programming

Substitution More Formally

$$x[s/x] = \begin{cases} s & \text{if } x = y \\ x & \text{otherwise} \end{cases}$$

$$*[s/y] = *$$

$$\langle t_1, t_2 \rangle [s/y] = \langle t_1[s/y], t_2[s/y] \rangle$$

$$(t_1 \ t_2)[s/y] = t_1[s/y] \ t_2[s/y]$$

$$(\pi_1 \ t)[s/y] = \pi_1 \ t[s/y]$$

$$(\pi_2 \ t)[s/y] = \pi_2 \ t[s/y]$$

$$(\lambda x. t)[s/y] = \lambda z. t[z/x][s/y]$$

(where z is fresh (i.e. new))

Compute the following substitutions

$$* [t/y][s/z] = \dots$$

$$\langle y, z \rangle [t/y][s/z] = \dots$$

$$(\lambda x. x) [t/x] = \dots$$

$$(\lambda x. \langle x, y \rangle) [z/y] = \dots$$

$$(\lambda x. \langle x, y \rangle) [x/y] = \dots$$

Via the Programming Lens

$$\frac{\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \vdash t : \mathbb{C}}{\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \vdash t : \mathbb{C}} \text{ (exch)}$$

$$\frac{\Gamma \vdash t : \mathbb{A}}{\Gamma, x : \mathbb{B} \vdash t : \mathbb{A}} \text{ (weak)}$$

$$\frac{\Gamma, x : \mathbb{A} \vdash t : \mathbb{B} \quad \Gamma \vdash s : \mathbb{A}}{\Gamma \vdash \dots : \mathbb{B}} \text{ (cut elimination)}$$

$$\frac{\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \vdash t : \mathbb{C}}{\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \vdash t : \mathbb{C}} \text{ (exch)}$$

$$\frac{\Gamma \vdash t : \mathbb{A}}{\Gamma, x : \mathbb{B} \vdash t : \mathbb{A}} \text{ (weak)}$$

$$\frac{\Gamma, x : \mathbb{A} \vdash t : \mathbb{B} \quad \Gamma \vdash s : \mathbb{A}}{\Gamma \vdash t[s/x] : \mathbb{B}} \text{ (cut elimination)}$$

$$\frac{\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \vdash t : \mathbb{C}}{\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \vdash t : \mathbb{C}} \text{ (exch)}$$

$$\frac{\Gamma \vdash t : \mathbb{A}}{\Gamma, x : \mathbb{B} \vdash t : \mathbb{A}} \text{ (weak)}$$

$$\frac{\Gamma, x : \mathbb{A} \vdash t : \mathbb{B} \quad \Gamma \vdash s : \mathbb{A}}{\Gamma \vdash t[s/x] : \mathbb{B}} \text{ (cut elimination)}$$

Substitution also fundamental in the study of equivalence

An Equational System pt. I

$$\begin{array}{lll} \pi_1 \langle t, s \rangle =_{\beta\eta} t & t =_{\beta\eta} * & (\text{if } t : 1) \\ \pi_2 \langle t, s \rangle =_{\beta\eta} s & \lambda x. t \ s =_{\beta\eta} t[s/x] & \\ \langle \pi_1 t, \pi_2 t \rangle =_{\beta\eta} t & \lambda x. (t \ x) =_{\beta\eta} t & \end{array}$$

An Equational System pt. II

$$t =_{\beta\eta} t \qquad \frac{t =_{\beta\eta} s}{s =_{\beta\eta} t} \qquad \frac{t =_{\beta\eta} s \quad s =_{\beta\eta} u}{t =_{\beta\eta} u}$$

$$\frac{\Gamma \vdash t =_{\beta\eta} s}{\pi\Gamma \vdash t =_{\beta\eta} s}$$

$$\frac{t =_{\beta\eta} s}{u[t/x] =_{\beta\eta} u[s/x]}$$

Table of Contents

The Calculus

Denotational Semantics

Equational System

Disjunctive Types