

# ***Computer Graphics***

***by Ruen-Rone Lee***  
***ICL/ITRI***



# *Wrap up from last course*


- ◆ **Geometrical Transformation**
- ◆ **Viewing Transformation**
- ◆ **Projection Transformation**
- ◆ **Viewport Transformation**



# *Part I:*

## *Conventional 3D Graphics Pipeline*

### *Lighting*



Conventional 3D Graphics Pipeline

*Color Model*  
*Illumination Model*  
*Polygon Shading*




# ***Color Model***

- ◆ **A color model is an abstract mathematical model describing the way color can be represented.**
- ◆ **Commonly, three or four values (components) are used to represent a specific color, such as RGB or CMYK**




# Color Space

- ◆ A color space is defined by a mapping function, e.g.,  $f(r, g, b)$ , and the color model




# ***RGB Color Model***

- ◆ **Additive color model**
- ◆ **Used in sensing, representation, and display of images in electronic systems, such as TV and computer**



# ***CMYK Color Model***

- ◆ **Subtractive color model**
- ◆ **Used in color printing**






# Shading

- ◆ Shading is a light-material interaction in determining the color for each point of objects or scenes



**Flat Shading**



**Smooth Shading**



# *Advanced Shading*

## ◆ Global illumination



# ***Factors that affect Shading***

## ◆ **Light sources**

- Ambient light, directional light, positional light, spot light...

## ◆ **Material properties**

- Ambient reflection, diffuse reflection, specular reflection...

## ◆ **Location of the viewer**

- Position of perceiving specular highlight

## ◆ **Surface orientation**


- Surface normal, vertex normal



# *Light Sources*


## ◆ Directional Light

- Light source located at infinite far away. Such as sun



# *Light Sources*


- ◆ **Positional Light (Point Light)**
  - Light source located at a specific position



# Light Sources

## ◆ Spot Light



- Light source located at a specific position with certain cutoff range



# Lighting Equation

◆ Intensity = Ambient + Diffuse + Specular

$$I = I_a k_a + \sum_{p=1}^m f_p I_p (k_d (N \cdot L_p) + k_s (R_p \cdot V)^n)$$



# Ambient Light


- ◆ Illumination surrounding a scene without providing any specific light source

$$I = I_a k_a$$

$I$ : resulting intensity

$I_a$ : ambient light intensity

$k_a$ : ambient reflection coefficient






# Diffuse Reflection

$I_p$ : point light source intensity  
 $k_d$ : diffuse reflection coefficient  
 $N$ : normalized normal vector  
 $L$ : normalized light direction vector


## ◆ Lambert's Cosine Law




# Diffuse Reflection

## ◆ Example


- Fixed point light source at (1.0, 1.0, 1.0)




$k_d = 0.4$



$k_d = 0.55$



$k_d = 0.7$




$k_d = 1.0$

# Ambient + Diffuse Reflection


## ◆ Example

■  $k_d = 0.4$


$$I = I_a k_a + I_p k_d (N \cdot L)$$




$k_a = 0.0$



$k_a = 0.3$



$k_a = 0.6$




$k_a = 0.9$

# Light Source Attenuation


- ◆ Light source intensity will attenuate with respect to the distance between the light source and the object

$$I = I_a k_a + f_{att} I_p k_d (N \cdot L) \quad f_{att} = \min\left(\frac{1}{c_1 + c_2 d_L + c_3 d_L^2}, 1\right)$$




# Specular Highlight

- ◆ Specular highlight is the bright spot of light on object being illuminated



Edit = Specular Material  
Ka = (0.247, 0.199, 0.075)  
Kd = (0.752, 0.606, 0.226)  
Ks = (0.628, 0.556, 0.366) Ke = 200.00  
Ambient ON Diffuse ON Specular ON



Edit = Specular Material  
Ka = (0.247, 0.199, 0.075)  
Kd = (0.752, 0.606, 0.226)  
Ks = (0.628, 0.556, 0.366) Ke = 200.00  
Ambient ON Diffuse ON Specular ON

# Specular Highlight

$$I_s = I_p k_s \cos^n \alpha = I_p k_s (R_p \cdot V)^n$$

$$I = I_a k_a + f_p I_p (k_d (N \cdot L_p) + k_s (R_p \cdot V)^n)$$

$I$ : Intensity of final illumination

$I_a$ : Intensity of ambient light

$k_a$ : Ambient reflection coefficient

$f_p$ : Attenuation function of point light source  $p$

$k_d$ : Diffuse reflection coefficient

$N$ : Normalized normal vector


$L_p$ : Normalized light source direction of point light source  $p$

$k_s$ : Specular reflection coefficient

$R_p$ : Normalized light source reflection vector of point light source  $p$

$V$ : Normalized viewpoint direction vector


$n$ : Material's specular reflection exponent



# Specular Highlight

## ◆ Phong Reflection Model

$$I = I_a k_a + f_p I_p (k_d (N \cdot L_p) + k_s (R_p \cdot V)^n)$$





# ***Modified Phong Reflection Model***

- ◆ **Also called Blinn-Phong Reflection Model**
  - **Original Phong model requires to calculate the reflection vector and view vector for each point**
  - **Blinn suggested to use an approximated way to calculate the specular reflection term by introducing the halfway vector**


# The Halfway Vector

- ◆ The halfway vector is the normalized vector halfway between the viewpoint and the light vector

$$H = \frac{L+V}{|L+V|}$$

$$\theta + \beta = \theta - \beta + \alpha$$

$$\beta = \frac{1}{2}\alpha$$



# Multiple Light Sources

$$I = I_a k_a + \sum_{p=1}^m f_p I_p (k_d (N \cdot L_p) + k_s (N \cdot H_p)^{n'})$$

$I$ : Intensity of final illumination

$I_a$ : Intensity of ambient light

$k_a$ : Ambient reflection coefficient

$f_p$ : Attenuation function of point light source  $p$

$k_d$ : Diffuse reflection coefficient

$N$ : Normalized normal vector

$L_p$ : Normalized light source direction of point light source  $p$

$k_s$ : Specular reflection coefficient


$H_p$ : Normalized half vector between viewpoint and the point light source  $p$

$n'$ : Material's specular reflection exponent (Blinn-Phong Reflection Model)



# *Flat Shading*

- ◆ Using face normal to derive polygon color




# *Smooth Shading*

- ◆ The colors for the interior of the polygon are interpolated between vertex colors



# *Smooth Shading*


## ◆ Gouraud Shading



# Smooth Shading

## ◆ Gouraud Shading

- Compute colors for each vertices respectively
- Interpolate pixel colors through vertex colors



$$P_a = P_0 \left(1 - \frac{y_a - y_0}{y_1 - y_0}\right) + P_1 \frac{y_a - y_0}{y_1 - y_0}$$


$$P_b = P_0 \left(1 - \frac{y_b - y_0}{y_2 - y_0}\right) + P_2 \frac{y_b - y_0}{y_2 - y_0}$$

$$P_c = P_a \left(1 - \frac{x_c - x_a}{x_b - x_a}\right) + P_b \frac{x_c - x_a}{x_b - x_a}$$




# Smooth Shading


## ◆ Phong Shading




Ambient



Diffuse



Specular




= Phong Reflection

# Smooth Shading

## ◆ Phong Shading

- Interpolate pixel normal through vertex normals
- Compute pixel color with derived pixel normal



$$P_a = P_0 \left(1 - \frac{y_a - y_0}{y_1 - y_0}\right) + P_1 \frac{y_a - y_0}{y_1 - y_0}$$


$$P_b = P_0 \left(1 - \frac{y_b - y_0}{y_2 - y_0}\right) + P_2 \frac{y_b - y_0}{y_2 - y_0}$$

$$P_c = P_a \left(1 - \frac{x_c - x_a}{x_b - x_a}\right) + P_b \frac{x_c - x_a}{x_b - x_a}$$

# Smooth Shading


## ◆ Phong Shading

- Interpolate pixel normal through vertex normals




# Smooth Shading

- ◆ **Difference between Gouraud shading and Phong shading**
  - If specular highlight is located inside the triangle only, then Gouraud shading cannot reveal such kind of highlight





# *Polygon Size Mattered*

- ◆ Large polygon size might degrade the lighting quality
  - Darker if the light source is closer to the polygon
  - No specular highlight can be perceived in the middle of a polygon



# *Polygon Size Mattered*

- ◆ **Subdivide large polygon into smaller polygons to gain better shading result**




# Vertex Normal Derivation

## ◆ Flat shading

- Vertex normal is equal to polygon face normal

$$N = (P_2 - P_1) \times (P_3 - P_1)$$






# Vertex Normal Derivation

## ◆ Smooth shading

- Vertex normal is equal to the sum of polygon face normals of adjacent polygons

$$N_{sum} = (N_1 + N_2 + \dots + N_m)$$

$$N = N_{sum} / |N_{sum}|$$




# *Normal Transformation*

- ◆ **Model transform and viewing transform together can transform a vertex from object space to eye space**
- ◆ **Normal is related to lighting process and the lighting calculation is performed in eye space**
- ◆ **So, normal has to transform to eye space as well**
- ◆ **But, what will happen if we transform normal using the same model-view matrix?**



# Normal Transformation

- ◆ If we transform normal using model-view matrix  $M$ , then...



$$T = P_2 - P_1$$

$$M \cdot T = M \cdot (P_2 - P_1)$$

$$T' = M \cdot P_2 - M \cdot P_1 = P'_2 - P'_1$$

$$N = Q_2 - Q_1$$

$$M \cdot N = M \cdot (Q_2 - Q_1)$$

$$N' = M \cdot Q_2 - M \cdot Q_1 = Q'_2 - Q'_1$$

$$N \cdot T = 0$$

But, after normal transformed by model-view matrix  $M$

$$N' \cdot T' \neq 0$$

# Normal Transformation

- ◆ **Normal transformation should be taking care if you have done any model and viewing transformations to the geometric data**

A vector in homogeneous coordinate is represented as

$$T = (x, y, z, 0) = (x_2, y_2, z_2, 1) - (x_1, y_1, z_1, 1)$$

A normal in homogeneous coordinate is represented as  $N = (n_x, n_y, n_z, 0)$

Since  $T$  and  $N$  are orthogonal, thus  $N \cdot T = 0$

We also know that after model-view transformation,  $T'$  and  $N'$  should remain orthogonal.

That is,  $N' \cdot T' = 0$ .



# Normal Transformation

- ◆ Represent the dot product by matrix multiplication and let  $M$  be the model-view matrix, we have

**Normal Transformation**

$$N \cdot T = \begin{pmatrix} n_x & n_y & n_z & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ 0 \end{pmatrix} = 0$$

**Eye space normal**

**Object space normal**

$$N' \cdot T' = \begin{pmatrix} n_x & n_y & n_z & 0 \end{pmatrix} M^{-1} M \begin{pmatrix} x \\ y \\ z \\ 0 \end{pmatrix} = 0 \Rightarrow \begin{pmatrix} n'_x \\ n'_y \\ n'_z \\ 0 \end{pmatrix} = (M^{-1})^T \begin{pmatrix} n_x \\ n_y \\ n_z \\ 0 \end{pmatrix}$$

**Model-view Transformation**

# *Lighting Procedure*

- ◆ **Define the vertex normals**
  - Lighting is the interaction between vertex normals and the light source
- ◆ **Define light sources**
  - Light source properties
- ◆ **Select lighting model**
  - Determine which lighting equation is used
- ◆ **Define material properties**
  - Define the percentage of reflectance to the light source




# Complete OpenGL Lighting Formula

$$\begin{aligned}
 \text{vertex color} = & \boxed{\text{emission}_{\text{material}}} + \text{Object can emit light itself} \\
 & \boxed{\text{ambient}_{\text{light model}} * \text{ambient}_{\text{material}}} + \text{Global ambient light} \\
 & \boxed{\sum_{i=0}^{n-1} \left( \frac{1}{k_c + k_l d + k_q d^2} \right)_i * (\text{spotlight effect})_i *} \\
 & \boxed{[\text{ambient}_{\text{light}} * \text{ambient}_{\text{material}} +} \\
 & \boxed{(\max \{ \mathbf{L} \cdot \mathbf{n}, 0 \}) * \text{diffuse}_{\text{light}} * \text{diffuse}_{\text{material}} +} \\
 & \boxed{(\max \{ \mathbf{s} \cdot \mathbf{n}, 0 \})^{\text{shininess}} * \text{specular}_{\text{light}} * \text{specular}_{\text{material}} ]_i} + \text{Light source contribution}
 \end{aligned}$$



# Spotlight Effect



# Spotlight Effect

## ◆ Spotlight Effect =

- 1, if the light source is not a spotlight
- 0, if the light source is a spotlight but the vertex lies outside the cone of illumination produced by the spotlight
- Otherwise, spotlight effect =  $(\max\{v \cdot d, 0\})^{\text{spot\_exp}}$ 
  - $v$  is the unit vector from the spotlight to the vertex
  - $d$  is the spotlight direction



# Q&A

