

Identification of Undocumented Meat Samples Seized at Heathrow Airport

1935263

Abstract

Introduction

Rarity status attributed to a species often increases its price and demand, with critically endangered species fetching the highest prices [1, 2]. The fallout of this is a positive feedback loop between rarity and demand which may drive an endangered species towards extinction and threaten whole ecosystems [3]. Species-level identification of undocumented meat samples is key in public health for validating the meat being delivered to consumers, but may also uncover networks carrying out illegal imports of endangered species, thus aiding conservation efforts by providing routes to intervene in the trade of these species.

If the morphology of a sample is compromised, DNA barcoding - comparing genetic sequences in a sample against known references - can aid in resolving the species of the sample [4, 5]. Mitochondrial DNA (mtDNA) genes are commonly used for DNA barcoding because they occur in high copy numbers relative to nuclear DNA and typically lack recombination, keeping within-species diversity low which enables species-level identification [6]. Cytochrome c oxidase I (COI), found within mtDNA, is the primary barcode marker in the animal kingdom, and reference sequences for COI are abundant in databases such as BOLD and NCBI, permitting barcoding against a diverse set of organisms [7, 8].

This analysis aims to resolve the species of four meat samples intercepted by UK Border Force officers at Heathrow Airport. The sequenced samples provide the starting point of this analysis pipeline.

Methods

Overview

A modular pipeline was constructed, with reproducibility in mind, to take the provided sequence data (samples.zip) through to the visualisation of a phylogenetic tree for sample identification. The four key pipeline stages are described here.

QC and Pre-Processing

FASTQ sequence files were first extracted from samples.zip before undergoing quality control to enforce a consistent FASTQ structure - including headers and line counts. Every part file

(1-3) belonging to the four samples (A-D) were then concatenated into multi-read FASTQ files by sample. seqtk [9] was used to convert each to FASTA while masking any bases with quality scores < 20 to 'N', ensuring bases were called at a confidence $\geq 99\%$ during sequencing to reduce the chances of false positives and increase confidence in downstream BLAST results.

BLAST and FASTA Concatenation

BLAST [10] searches were carried out using the FASTA files as query sequences against the NCBI Nucleotide database using blastn [11]. As the samples were labeled parts 1-3 without information on genomic region, each part was treated as a separate fragment and BLAST results for each sample were split into part-specific files, enabling downstream alignment per-part across all samples. These results were then sorted by ascending e-value, descending bitscore, and descending alignment length before collapsing by species with taxonkit [12]. The top four for each part and sample were selected. sampleD_part3 returned *Homo sapiens*, consistent with human contamination, and was therefore removed.

efetch [13] was used to download the FASTA sequences for each BLAST hit accession, and the sstart and ssend values were used to trim these sequences to match the query region using seqtk, reverse complementing if an opposite strand is matched in BLAST. COI sequences of three *Selachii* species were retrieved using efetch as an outgroup before all FASTA sequences were concatenated by part into three complete FASTA files (parts 1-3) using seqkit [14].

Translation and Alignment

Translation of the FASTA sequences using the vertebrate mitochondrial code was carried out using Biopython [15]. To counteract potential frame shift errors during translation, ORFs were identified and selected as the protein sequence for each read.

MUSCLE5 [16] was used to carry out multiple sequence alignments (MSA) on the protein sequences, and trimAl [17] removed poorly aligned regions while retaining internal sites for a more stable tree build. Inspection of BLAST results indicated that each part derived from COI, so the trimmed alignments were concatenated into a supermatrix alignment with catfasta2phym1 [18], retaining partitions for each part.

Tree Building and Viewing

Phylogenetic inference was performed using maximum likelihood in IQ-TREE3 [19] with ModelFinder Plus selecting the best model to improve accuracy, and included 1000 ultrafast bootstrap replicates to evaluate branch support. The Python library ete3 [20] was used to visually display the phylogenetic tree as well as rooting to the outgroup *Selachii* clade.

Results

BLAST searches suggested that the samples corresponded to COI, and provided preliminary identifications for each sample. Sample A consistently matched Right and Bowhead Whales, with sample B returning Beluga and Narwhal. Sample C matched true tunas, and sample D matched sea turtle species and *Homo sapiens* (Table 1).

Table 1: Top BLAST hit per part across all samples

Sample	Part	Species	% Identity	e-value
A	1	<i>Eubalaena japonica</i>	97.5	0.0
A	2	<i>Eubalaena japonica</i>	99.6	0.0
A	3	<i>Eubalaena japonica</i>	93.4	7.0e-160
B	1	<i>Delphinapterus leucas</i> ; <i>Monodon monoceros</i> (tie)	92.3	0.0
B	2	<i>Delphinapterus leucas</i>	92.8	0.0
B	3	<i>Monodon monoceros</i>	89.2	0.0
C	1	<i>Thunnus thynnus</i> ; <i>Thunnus orientalis</i> (tie)	81.7	3.7e-103
C	2	—	—	—
C	3	<i>Thunnus thynnus</i>	89.2	1.2e-157
D	1	<i>Dermochelys coriacea</i>	98.3	0.0
D	2	<i>Dermochelys coriacea</i>	96.2	0.0
D	3	<i>Homo Sapiens</i>	100	6.3e-46




Figure 1: Window of supermatrix alignment with potential frame-shift error in sampleD near the highly-conserved NFITTIIN region.

The phylogenetic tree (Fig. 2) recovered four well-resolved vertebrate clades: a Cetacean

clade (whales); a Testudine clade (turtle and tortoises); a Scombridae clade (tuna); and the outgroup Selachii clade (sharks).




Figure 2: Maximum likelihood phylogenetic tree with placement of unknown meat samples and top BLAST hits. Scale bar corresponds to 0.01 substitutions per site. Bootstrap support (%) shown at nodes.

Sample A placement was strongly supported within Balaenidae (95%), but species-level resolution within was poor. However, BLAST consistently returned *Eubalaena japonica* as the top hit with a high percentage identity across all parts, suggesting North Pacific right whale as the leading candidate species.

Sample B placement within Monodontidae was fully supported (100%), but internal node support and species resolution was weak (55%). Only *Delphinapterus leucas* and *Monodon monoceros* exist in this family, therefore sample B must represent one of these species.

Sample C placement was strongly supported within *Thunnus* (97%), with low-moderate internal node support. BLAST suggested *Thunnus thynnus*, but this is not confirmed by the phylogeny. As such, sample C can only be assigned to the *Thunnus* genus.

Sample D placement within Testudines was strongly supported (92%), with very low support within this clade (11-59%). Sample D is placed with *Dermochelys coriacea*, consistent

with BLAST, suggesting Leatherback sea turtle as the leading candidate species.

Discussion

Across all samples, order-level placement was strongly supported with confident assignment to major vertebrate groups. However, species confidence was generally low and relied on BLAST compliance with phylogeny where candidate species were identified.

The COI fragment for barcoding is 650bp long which makes species inference sensitive to sequencing errors, with low quality reads providing limited phylogenetic signal [5]. FastQC [21] reports showed all samples displayed regions of low quality calls, reducing usable bases in already short sequences, and likely contributed to weakened inter-specific discriminatory power with inflated branch lengths and reduced bootstrap support. An exploratory analysis investigated the inflated branch length of sample D: a Needleman-Wunsch alignment (Fig. 3) indicated a potential sequencing mis-read leading to the apparent frame-shift in Fig. 1. Adding an ambiguous base and re-running the analysis reduced branch length and increased bootstrap support within the Testudine clade (Figs. 4, 5), illustrating the downstream effects a single base can have on phylogenetic confidence. Furthermore, references from databases such as GenBank (which BLAST searches) which do not require strict standards or quality control must be interpreted cautiously for species identification [6, 22].

The candidate species of sample A, *Eubalaena japonica*, is currently listed as ‘Endangered’ on the IUCN Red List [23]. The presence of this sample raises serious conservation concerns if confirmed, and could reflect trade of a vulnerable whale species. The sample B candidate monotonid species are both listed under ‘Least Concern’ on the IUCN Red List, with regulated international trade [24, 25]. Their presence may indicate by-catch from fisheries in or near conservation sites, but without sample origin the conservation implications cannot be assessed fully. IUCN Red List assessments for the *Thunnus* genus range from ‘Least Concern’ (*Thunnus albacares*) to ‘Endangered’ (*Thunnus maccoyii*) [26, 27]. Without species level identification the conservation implications of sample C cannot be assessed fully. The sample D candidate, *Dermochelys coriacea*, is listed on the IUCN Red List as ‘Vulnerable’. This species inhabits conservation sites across its range as well as protected areas. The presence of this sample raises conservation concerns and may reflect illegal trade, depending on the origin of the sample.

Overall, this analysis demonstrated the use of COI in order and family level identification, but displays a lack of resolution in identifying the unknown samples to the species level. Subsequent sequencing of samples using longer mitochondrial barcodes may improve resolution among more recently diverged species. Re-sequencing to obtain multiple reads of each sequence may reduce mis-reads and base ambiguity, while the use of curated reference databases, such as BOLD, is also suggested to increase confidence in reference sequences [22]. These suggestions should improve species level resolution and provide more confident conservation claims for the meat samples. Nevertheless, the detection of candidate species from undocumented meat illustrates the value of DNA barcoding in the context of the illegal wildlife trade.

References

- [1] Yik-Hei Sung and Jonathan J Fong. Assessing consumer trends and illegal activity by monitoring the online wildlife trade. *Biological conservation*, 227:219–225, 2018.
- [2] Franck Courchamp, Elena Angulo, Philippe Rivalan, Richard J Hall, Laetitia Signoret, Leigh Bull, and Yves Meinard. Rarity value and species extinction: the anthropogenic allee effect. *PLoS biology*, 4(12):e415, 2006.
- [3] Liam J Hughes, Oscar Morton, Brett R Scheffers, and David P Edwards. The ecological drivers and consequences of wildlife trade. *Biological Reviews*, 98(3):775–791, 2023.
- [4] Pilar A Haye, Nicolás I Segovia, Raúl Vera, María de los Ángeles Gallardo, and Cristian Gallardo-Escárate. Authentication of commercialized crab-meat in chile using dna barcoding. *Food Control*, 25(1):239–244, 2012.
- [5] Sarfraz Ahmed, Muhammad Ibrahim, Chanin Nantasenamat, Muhammad Farrukh Nisar, Aijaz Ahmad Malik, Rashem Waheed, Muhammad Z Ahmed, Suvash Chandra Ojha, and Mohammad Khurshed Alam. Pragmatic applications and universality of dna barcoding for substantial organisms at species level: a review to explore a way forward. *BioMed Research International*, 2022(1), 2022.
- [6] Nick Dawnay, Rob Ogden, Ross McEwing, Gary R Carvalho, and Roger S Thorpe. Validation of the barcoding gene coi for use in forensic genetic species identification. *Forensic science international*, 173(1):1–6, 2007.
- [7] Teresita M Porter and Mehrdad Hajibabaei. Over 2.5 million coi sequences in genbank and growing. *PloS one*, 13(9):e0200177, 2018.
- [8] Sujeevan Ratnasingham, Catherine Wei, Dean Chan, Jireh Agda, Josh Agda, Liliana Ballesteros-Mejia, Hamza Ait Boutou, Zak Mohammad El Bastami, Eddie Ma, Ramya Manjunath, et al. Bold v4: A centralized bioinformatics platform for dna-based biodiversity data. In *DNA barcoding: Methods and protocols*, pages 403–441. Springer, 2024.
- [9] Heng Li. seqtk: Toolkit for processing sequences in FASTA/Q formats . <https://github.com/lh3/seqtk>, 2012.
- [10] Stephen F Altschul, Warren Gish, Webb Miller, Eugene W Myers, and David J Lipman. Basic local alignment search tool. *Journal of molecular biology*, 215(3):403–410, 1990.
- [11] Christiam Camacho, George Coulouris, Vahram Avagyan, Ning Ma, Jason Papadopoulos, Kevin Bealer, and Thomas L Madden. Blast+: architecture and applications. *BMC bioinformatics*, 10(1):421, 2009.
- [12] Wei Shen and Hong Ren. Taxonkit: A practical and efficient ncbi taxonomy toolkit. *Journal of genetics and genomics*, 48(9):844–850, 2021.
- [13] Jonathan Kans. Entrez direct: E-utilities on the unix command line. In *Entrez programming utilities help [Internet]*. National Center for Biotechnology Information (US), 2024.
- [14] Wei Shen, Shuai Le, Yan Li, and Fuquan Hu. Seqkit: a cross-platform and ultrafast toolkit for fasta/q file manipulation. *PloS one*, 11(10):e0163962, 2016.
- [15] Peter JA Cock, Tiago Antao, Jeffrey T Chang, Brad A Chapman, Cymon J Cox, Andrew Dalke, Iddo Friedberg, Thomas Hamelryck, Frank Kauff, Bartek Wilczynski, et al. Biopython: freely available python tools for computational molecular biology and bioinformatics. *Bioinformatics*, 25(11):1422, 2009.

- [16] Robert C Edgar. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology and phylogeny. *Nature communications*, 13(1):6968, 2022.
- [17] Salvador Capella-Gutiérrez, José M Silla-Martínez, and Toni Gabaldón. trimal: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, 25(15):1972–1973, 2009.
- [18] Johan Nylander. catfasta2phym1 – Concatenate FASTA alignments to PHYML, PHYLIP, or FASTA format. <https://github.com/nylander/catfasta2phym1>, 2010.
- [19] Thomas KF Wong, Nhan Ly-Trong, Huaiyan Ren, Hector Baños, Andrew J Roger, Edward Susko, Chris Bielow, Nicola De Maio, Nick Goldman, Matthew W Hahn, et al. Iq-tree 3: Phylogenomic inference software using complex evolutionary models. 2025.
- [20] Jaime Huerta-Cepas, François Serra, and Peer Bork. Ete 3: reconstruction, analysis, and visualization of phylogenomic data. *Molecular biology and evolution*, 33(6):1635–1638, 2016.
- [21] Simon Andrews. FastQC: A quality control analysis tool for high throughput sequencing data. <https://github.com/s-andrews/FastQC>, 2007.
- [22] Kelly A Meiklejohn, Natalie Damaso, and James M Robertson. Assessment of bold and genbank—their accuracy and reliability for the identification of biological materials. *PloS one*, 14(6):e0217084, 2019.
- [23] J. G. Cooke and P. J. Clapham. *Eubalaena japonica*. <https://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T41711A50380694.en>, 2018. Accessed 07 December 2025.
- [24] L. Lowry, K. Laidre, and R Reeves. *Monodon monoceros*. <https://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T13704A50367651.en>, 2017. Accessed 07 December 2025.
- [25] L. Lowry, K. Laidre, and R Reeves. *Delphinapterus leucas*. <https://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T6335A50352346.en>, 2017. Accessed 07 December 2025.
- [26] B.B. Collette, A. Boustany, W. Fox, J. Graves, M. Juan Jorda, and V Restrepo. *Thunnus albacares*. <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T21857A46624561.en>, 2021. Accessed 07 December 2025.
- [27] B.B. Collette, A. Boustany, W. Fox, J. Graves, M. Juan Jorda, and V Restrepo. *Thunnus maccoyii*. <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T21858A170082633.en>, 2021. Accessed 07 December 2025.

Supplementary Data

Supplementary data and all scripts can be found in the GitHub repository for this project: https://github.com/archiebenn/BIOLM0051_analysis.git

Appendix: Exploratory Figures



Figure 3: Needleman-Wunsch alignment of sampleD_part1 against closest reference. Gap indicates a possible mis-read during sequencing.




Figure 4: Supermatrix alignment after inserting one ambiguous base (N) to sampleD_part1. This base resolves the apparent frame-shift, resulting in a protein sequence which aligns much more closely to highly-conserved region




Figure 5: Phylogenetic tree resulting from insertion of an ambiguous base (N) into sampleD_part1. Demonstrates the impact of sequencing error on branch lengths and bootstraps compared to Fig. 2.