



The Effect of Forest Age on Machine Learning Evapotranspiration Generalisation

Author

Archie BENN

Supervisor

Professor Martin DE KAUWE

ABSTRACT (200 words)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum. Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo.

Key words: Evapotranspiration, Forest Age, Machine Learning, FLUXNET.

Word count: 4000 (Literature Review: 3000; Project Plan: 1000)

A dissertation submitted to the University of Bristol in accordance with the requirements of the degree of Master of Science by advanced study in Bioinformatics in the Faculty of Health and Life Sciences.

DECLARATION

I declare that the work in this dissertation was carried out in accordance with the requirements of the University's Regulations and Code of Practice for Taught Programmes and that it has not been submitted for any other academic award. Except where indicated by specific reference in the text, this work is my own work. Work done in collaboration with, or with the assistance of others, is indicated as such. I have identified all material in this dissertation which is not my own work through appropriate referencing and acknowledgement. Where I have quoted or otherwise incorporated material which is the work of others, I have included the source in the references. Any views expressed in the dissertation, other than referenced material, are those of the author.

Archie BENN

May 6, 2026

CONTENTS

LITERATURE REVIEW	4
Introduction	4
Evapotranspiration	4
Forest Complexity	4
Forest Age	5
Evapotranspiration Prediction ~1400 Words	6
Why predict ET?	6
Contemporary ET Prediction Methods (in. +s and -s) and Feature Selection .	6
Machine Learning Models for ET Prediction	6
Machine Learning and Applicability of Incorporating Age ~800 Words	
<i>(What is the case for using ML and forest age here?)</i>	7
Why use Machine Learning?	7
Why Incorporate Age?	7
Effect on Generalisation	7
Quick section closing off the review and what the project will aim to investigate	7
PROJECT AIMS AND OBJECTIVES (500 WORDS)	8
Aims	8
PROJECT PLAN (500 WORDS)	9
Data Acquisition	9
Site Selection	9
Analysis	9
Feature Engineering	9
ML Model Setup and Training	9
ML Model Evaluation	9
ML Model Predictions	9
Tools	9
Interpretation	9
GANTT CHART	10
RISK REGISTER (~ 250 WORDS)	11
REFERENCES	12
FIGURES AND TABLES	15

LITERATURE REVIEW

Introduction

Evapotranspiration

Terrestrial evapotranspiration (ET) is a key component of the global hydrological cycle describing the transition of water from liquid to gas at Earth's surface and further transfer to the atmosphere, and is a crucial component of the energy and water exchange between terrestrial ecosystems and the climate system. The term emphasises the combination of water vapourisation flux from (1) abiotic evaporation from soil and plant canopy surfaces, and (2) biotic transpiration through stomata to the atmosphere (Katul et al. 2012). Approximately 60% of precipitation on the land surface is returned to the atmosphere through ET (Oki and Kanae 2006), and the associated latent heat flux, λET (where λ is the latent heat of vaporisation), accounts for 50% of the solar radiation absorbed by the land surface (Trenberth et al. 2009). ET can be generally described through:

$$ET = T + E_{Soil} + E_{Canopy} \quad (1)$$

where T refers to transpiration through plant stomata, E_{soil} to evaporation from the soil surface, and E_{Canopy} to evaporation from plant canopies (via rainfall interception) (Lawrence et al. 2007). The contribution of these variables to overall ET is not equal, with Lian et al. (2018) estimating a global T/ET ratio of 0.62 ± 0.06 , indicating plant transpiration as the leading component of ET worldwide. The partitioning of ET into E ($E_{soil} + E_{Canopy}$) and T within ecosystems is associated with "water use efficiency" and has use in irrigation management, where T is linked with plant productivity, while E does not directly contribute to production (Kool et al. 2014). Accurate partitioning estimates are also important in aiding carbon cycle projections given the coupled effect of transpiration and land-atmosphere carbon dioxide exchanges through stomata.

Forest Complexity

The magnitude of ET is governed by many environmental factors such as solar radiation, leaf area index (LAI), air temperature, vapour pressure deficit (VPD), soil moisture, and wind speed (Yang et al. 2023). Of these, Lian et al. (2018) identified LAI as the primary driver of T/ET spatial patterns, with Kool et al. (2014) suggesting that in systems with full canopy cover (and therefore high LAI) ET is often assumed to be "similar enough to T to permit correlation of ET with biomass". However, this assumption may break down in densely vegetated systems with multi-layer canopies such as forests, despite high LAI. In these systems, the 2-dimensional remote measure of LAI will not capture the vertical canopy complexity which drives rainfall interception and subsequent E_{Canopy} , implying that canopy cover indices may struggle to explain T/ET ratios in more complex vertical environments. This is reflected by S. Zhou et al. (2016) who report higher coefficients of

determination between enhanced vegetation index (EVI) and T/ET in corn and soybean sites (0.84 and 0.82) with simple canopies, compared to evergreen needle leaf forests (0.37), a result possibly due to increased E_{Canopy} driving this ratio down. Hardiman et al. (2013) also demonstrate that while LAI may be stable in stands older than 50 years, canopy complexity continues to increase with age through >160 years of stand development and had a greater impact on annual net primary productivity (NPP) and overall resource use efficiency than LAI across all stands. Furthermore, LAI has shown to only explain around 20% of mean annual T/ET (X. Li et al. 2019), suggesting simple vegetation indices which don't take into account 3-dimensional complexity may be insufficient for predicting T/ET in structurally complex systems.

Oishi et al. (2020), in a chronosequence study on broadleaf forests in the US, report that interception of incoming rainfall strongly increases with age from ~2% (12 year old stand) to 20% (200 year old stand) despite only small differences in LAI, an effect in part due to branch architecture which varies with age. This study also reported that annual canopy transpiration (T) was over twice as high in the 35 year old stand compared to the 12 and 200 year old stands, with increasing canopy interception in the oldest stand largely offsetting the reduction in T such that total ET was similar between the two older stands, but remained lower in the youngest stand (Fig. 1). This demonstrates how age-driven effects in canopy architecture introduce complexity and can affect overall ET, as well as the relative contributions of T and E_{Canopy} which may vary significantly over time within stands.

Alongside canopy architecture, stomatal conductance (g_s) declines as trees grow taller due to gravitational reductions in leaf water potential and decreasing hydraulic conductance with increasing path length from soil to leaf (Ryan and Yoder 1997). For example, Schäfer et al. (2000) demonstrated that the mean stomatal conductance of tree crowns decreased by approximately 60% over a 30m increase in tree height, and Hubbard et al. (1999) established a 44% drop in hydraulic conductance in old trees compared to young trees in a mixed age stand. As g_s directly influences T (Monteith 1965), these age related changes introduce further complexity into ET dynamics and suggest that age could be a key source of variability in ET in forest ecosystems.

Forest Age

Forest age distributions are increasingly being affected by climate change-driven disturbances - such as fire, drought, insect attack, and pathogens - which are expected to continue in the coming decades (Solomon et al. 2009; Seidl et al. 2017). Some of these disturbances are able to completely change forest demography by causing widespread mortality (Dale et al. 2001), effectively resetting the age of the forest landscape. Land use changes also directly affect forest age distributions, for example the recovery of deforested areas in southern China following the implementation of forestation policies around the year 2000 has resulted in a complex pattern of forest age structure in recent years, and an increase in average tree cover from 21% in 1987 to 38% in 2016, leading to large areas now being dominated by young forests (Tong et al. 2020; Leng et al. 2024), with similar patterns of

young forests due to recent disturbance found worldwide (Besnard et al. 2021).

As forests are exhibiting greater age heterogeneity, and with climate change increasing the severity and frequency of potential forest disturbances worldwide, stand age may represent an increasingly important variable for ET estimation in forests, capturing age specific effects, such as canopy complexity and alterations to g_s , and linking these to ET beyond what simple vegetation indices such as LAI are capable of. Yet whether these age-driven changes in forests are a large enough source of ET variability to affect the predictions made by current ET models which do not take age into account remains largely unexplored.

Evapotranspiration Prediction ~1400 Words

Why predict ET?

Estimating ET accurately is important in understanding hydrological processes, management of irrigation systems for effective water use in agriculture, prediction and mitigation of droughts, determining the adaptive mechanisms of vegetation, and in assessing the overarching patterns of global climatic shifts (Amani and Shafizadeh-Moghadam 2023; Xue et al. 2025). These assessments can, in turn, steer regulations and policies in climate science and water resource management, representing a critical link between the quantification of terrestrial water fluxes and land systems governance (Haiyang Shi et al. 2024).

The direct measurement of ET uses lysimeters or eddy covariance (EC) methods, but these are costly to use and only measure at the plot scale, resulting in spatially constrained measurements and making them unsuitable for ET predictions at larger ecosystem scales (Liyew et al. 2025). The uneven distribution and low sampling rate of EC data in ecoregions of Africa, Oceania (excluding Australia), and South America may be attributed to the costs associated with these methods, despite efforts to mitigate this (Hill et al. 2017; Markwitz and Siebicke 2019). As a result of these issues, various methods to estimate ET indirectly have been derived, including: (1) traditional physics-based models which rely on pre-defined equations such as the Penman-Monteith (PM) and Hargreaves-Samani (HS) equations (Monteith 1965; Hargreaves and Samani 1985); (2) remote sensing methods, for example the SEBS model (Su 2002); and more recently, (3) machine learning (ML) methods such as artificial neural networks and random forest models trained on historical data (T. Xu et al. 2018). The development of these methods has permitted global ET estimates to be produced (S. Pan et al. 2020), yet significant uncertainties remain, particularly in regions with sparse EC coverage and in complex or heterogeneous landscapes where the assumptions underlying these models may break down.

Contemporary ET Prediction Methods (in. +s and -s) and Feature Selection

The prediction of ET has

Machine Learning Models for ET Prediction

Machine Learning and Applicability of Incorporating Age ~800 Words

(What is the case for using ML and forest age here?)

Why use Machine Learning?

Why Incorporate Age?

Effect on Generalisation

Quick section closing off the review and what the project will aim to investigate

PROJECT AIMS AND OBJECTIVES (500 WORDS)

Aims

- Overview of how this project will answer the question/area proposed at end of introduction.

PROJECT PLAN (500 WORDS)

Data Acquisition

Site Selection

Analysis

Feature Engineering

ML Model Setup and Training


ML Model Evaluation

ML Model Predictions

Tools

Interpretation

GANTT CHART



RISK REGISTER (~ 250 WORDS)

Table 1: Risk register

Nature of the risk	Likelihood and potential impact on the project	Mitigation strategy
ahhh	oh	lipsum
yeh	risk	register

REFERENCES

- Amani, Shima and Hossein Shafizadeh-Moghadam (2023). "A review of machine learning models and influential factors for estimating evapotranspiration using remote sensing and ground-based data". In: *Agricultural water management* 284, p. 108324.
- Besnard, S et al. (2021). *Mapping global forest age from forest inventories, biomass and climate data*, *Earth Syst. Sci. Data*, 13, 4881–4896.
- Dale, Virginia H et al. (2001). "Climate change and forest disturbances: climate change can affect forests by altering the frequency, intensity, duration, and timing of fire, drought, introduced species, insect and pathogen outbreaks, hurricanes, windstorms, ice storms, or landslides". In: *BioScience* 51.9, pp. 723–734.
- Hardiman, Brady S et al. (2013). "Maintaining high rates of carbon storage in old forests: A mechanism linking canopy structure to forest function". In: *Forest Ecology and Management* 298, pp. 111–119.
- Hargreaves, George H and Zohrab A Samani (1985). "Reference crop evapotranspiration from temperature". In: *Applied engineering in agriculture* 1.2, pp. 96–99.
- Hill, Timothy, Melanie Chocholek, and Robert Clement (2017). "The case for increasing the statistical power of eddy covariance ecosystem studies: why, where and how?" In: *Global Change Biology* 23.6, pp. 2154–2165.
- Hubbard, Robert M, Barbara J Bond, and Michael G Ryan (1999). "Evidence that hydraulic conductance limits photosynthesis in old *Pinus ponderosa* trees". In: *Tree physiology* 19.3, pp. 165–172.
- Katul, Gabriel G et al. (2012). "Evapotranspiration: A process driving mass transport and energy exchange in the soil-plant-atmosphere-climate system". In: *Reviews of Geophysics* 50.3.
- Kool, Dilia et al. (2014). "A review of approaches for evapotranspiration partitioning". In: *Agricultural and forest meteorology* 184, pp. 56–70.
- Lawrence, David M et al. (2007). "The partitioning of evapotranspiration into transpiration, soil evaporation, and canopy evaporation in a GCM: Impacts on land-atmosphere interaction". In: *Journal of Hydrometeorology* 8.4, pp. 862–880.
- Leng, Yi et al. (2024). "Forest aging limits future carbon sink in China". In: *One Earth* 7.5, pp. 822–834.
- Li, Xi et al. (2019). "A simple and objective method to partition evapotranspiration into transpiration and evaporation at eddy-covariance sites". In: *Agricultural and Forest Meteorology* 265, pp. 171–182.

- Lian, Xu et al. (2018). "Partitioning global land evapotranspiration using CMIP5 models constrained by observations". In: *Nature Climate Change* 8.7, pp. 640–646.
- Liyew, Chalachew Muluken et al. (2025). "A review of feature selection methods for actual evapotranspiration prediction". In: *Artificial Intelligence Review* 58.10, p. 292.
- Markwitz, Christian and Lukas Siebicke (2019). "Low-cost eddy covariance: a case study of evapotranspiration over agroforestry in Germany". In: *Atmospheric measurement techniques* 12.9, pp. 4677–4696.
- Monteith, John Lennox (1965). "Evaporation and environment". In.
- Oishi, AC et al. (2020). "Quantifying the effects of stand age on components of forest evapotranspiration". In: *Acta Horticulturae* 1300, pp. 89–96.
- Oki, Taikan and Shinjiro Kanae (2006). "Global hydrological cycles and world water resources". In: *science* 313.5790, pp. 1068–1072.
- Pan, Shufen et al. (2020). "Evaluation of global terrestrial evapotranspiration using state-of-the-art approaches in remote sensing, machine learning and land surface modeling". In: *Hydrology and earth system sciences* 24.3, pp. 1485–1509.
- Ryan, Michael G and Barbara J Yoder (1997). "Hydraulic limits to tree height and tree growth". In: *Bioscience* 47.4, pp. 235–242.
- Schäfer, KVR, Ram Oren, and JD Tenhunen (2000). "The effect of tree height on crown level stomatal conductance". In: *Plant, Cell & Environment* 23.4, pp. 365–375.
- Seidl, Rupert et al. (2017). "Forest disturbances under climate change". In: *Nature climate change* 7.6, pp. 395–402.
- Shi, Haiyang et al. (2024). "Machine learning-based investigation of forest evapotranspiration, net ecosystem productivity, water use efficiency and their climate controls at meteorological station level". In: *Journal of Hydrology* 641, p. 131811.
- Solomon, Susan et al. (2009). "Irreversible climate change due to carbon dioxide emissions". In: *Proceedings of the national academy of sciences* 106.6, pp. 1704–1709.
- Su, Zhongbo (2002). "The Surface Energy Balance System (SEBS) for estimation of turbulent heat fluxes". In: *Hydrology and earth system sciences* 6.1, pp. 85–100.
- Tong, Xiaowei et al. (2020). "Forest management in southern China generates short term extensive carbon sequestration". In: *Nature communications* 11.1, p. 129.
- Trenberth, Kevin E, John T Fasullo, and Jeffrey Kiehl (2009). "Earth's global energy budget". In: *Bulletin of the american meteorological society* 90.3, pp. 311–324.
- Xu, Tongren et al. (2018). "Evaluating different machine learning methods for upscaling evapotranspiration from flux towers to the regional scale". In: *Journal of Geophysical Research: Atmospheres* 123.16, pp. 8674–8690.

- Xue, Yayong et al. (2025). “A review of evapotranspiration estimation models: advances and future development”. In: *Water Resources Management* 39.8, pp. 3641–3657.
- Yan, Xiaohui et al. (2023). “Deep learning for daily potential evapotranspiration using a HS-LSTM approach”. In: *Atmospheric Research* 292, p. 106856.
- Yang, Yuting et al. (2023). “Evapotranspiration on a greening Earth”. In: *Nature Reviews Earth & Environment* 4.9, pp. 626–641.
- Zhou, Sha et al. (2016). “Partitioning evapotranspiration based on the concept of underlying water use efficiency”. In: *Water Resources Research* 52.2, pp. 1160–1175.

FIGURES AND TABLES




Figure 1: Components of annual evapotranspiration by stand age, redrawn from Oishi et al. (2020). Sub canopy evapotranspiration (E_{sub}), canopy transpiration (E_c), and canopy interception (E_i) correspond approximately to E_{Soil} , T , and E_{Canopy} in Equation 1, although E_{sub} additionally captures under canopy transpiration.

Table 2: Advantages and limitations of ML models in the literature.

Method	Incorporated Features	Limitations & Unresolved
Random Forest (T. Xu et al. 2018)	Outperformed other models (ANN, cubist, SVM, DBN) based on R^2, MAPE, RMSE, and relative uncertainties from three-cornered hat. RF has an ability to handle complex nonlinear relationships between flux and climate and environmental predictors.	Model performs better over areas of dense vegetation compared to barren land. More spatiotemporal continuous land surface variables at fine resolution should be added to potentially enhance predictability.
HS-LSTM coupled model (Yan et al. 2023)	Shown to outperform the standalone ML and classical ET_0 models (RMSE reduced ~20%). Can be used to estimate future ET_0 with only air temperature forecasts which can be easily obtained.	Slight underestimates of extreme ET_0 values. Each station required a differently-tuned model, so further studies are required to evaluate potential for a generally valid model and assess whether transfer learning may be applied across different regions.