

# The Effect of Some Plant Growth Regulators on Cell Biomass in the Cell Suspension Culture of *Calendula officinalis* L. and *Calendula arvensis* L. Species

Nergis Kaya<sup>1\*</sup>

<sup>1</sup>Canakkale Onsekiz Mart University, Department of Food Processing, Food Technology Program, 17200, Canakkale, Turkey, nergisskkaya[at]gmail.com

**Abstract:** The cell suspension culture *C. officinalis* and *C. arvensis* plants having medicinal and economical important was carried out in the four different MS medium which supplemented with different concentrations of NAA:BAP (1:1, 0.5:5 mg/l) and IAA:BAP (1:1, 0.5:5 mg/l) under sterile conditions. For this purpose, four months old calli which grown in the callus cultures were passed into the cell suspension culture. In this cell suspension cultures, cell counting and measurement of fresh/dry weight process were realized for two *Calendula* species during 40 days. This process repeated once in a five days. All of the research results were evaluated with Tukey Multiple Comparison Test.

**Keywords:** *Calendula officinalis*, *Calendula arvensis*, cell suspension culture, plant growth regulators

## 1. Introduction

*Calendula officinalis* (pot marigold) L., known for its ornamental plant characteristics, is a medicinal plant which is belonging to Asteraceae (Compositae) family. The species grows to 20 up to 40 cm height and has 20 varieties. Its flower appears yellow [1]. Its chemical constituents include triterpene glycosides, triterpene alcohols, flavonol glycosides, essential oil, polysaccharides and fatty oil [2]. Many studies have reported that the plant have pharmacological effects such as anti-cancer [3; 4; 5; 6], anti-microbial [7; 8; 9; 10; 11], anti-leishmanial [12; 13], anti-HIV [14], antioxidants [15;16;17], cytotoxic, anti-tumor [18; 3; 19], anti-viral [20], anti-inflamatur [21; 19], oedema diuretic [22], hypoglycemic [23], uterotonic [24], lymphocyte activator effect [3], in venous ulcer treatment [25] and for biligenic function [26].

It was known that the in vitro cultures has some advantages. Some of this advantages is secondary metabolite production, the cell proliferation, etc. under sterile and controlled conditions during culture period in the laboratory [27].

The callus was defined the accumulation containing undifferentiated/unorganized parenchymatic cells that was occurred at the injured areas of undifferentiated cells and tissues [28]. The callus culture was state that the beginning of cell suspension culture [29]. It was clarified that the auxin and cytokinin amounts was important in the in vitro cultures [30] and the equal ratio of auxin/cytokinin was caused the callus formation [31]. It was explained for the in vitro culture that the different biotic (elicitation, etc.) and abiotic (plant growth regulators, metal ions, etc.) stimulating compounds added in the nutrient medium [32; 33; 34; 35].

The cell suspension culture can be prepared from explant from differentiated tissue or callus with undifferentiated cell mass [36]. Transition from undifferentiated structures to cell suspension cultures requires a shorter time compared to the

transition from differentiated plant parts. Therefore, it is indicated that callus is preferred as starting material in cell suspension cultures [37]. This process is achieved by varying concentrations two major plant growth regulators (auxin and cytokinin) in the nutrient medium [38]. Under appropriate conditions, callus cells can continue to grow in the suspension cultures without differentiation [39].

The objective of myresearch was to evaluate the effect of different auxin:cytokinin combinations which added to MS medium to enhance cell biomass induction capability of *C. officinalis* and *C. arvensis* species in the cell suspension culture. Therefore, the cell suspension culture would be an important step to obtain medically important secondary metabolites containing of *C. officinalis* and *C. arvensis* species. In this research I focused on the establishment of the cell suspension cultures and development of optimal conditions in suspension cultures for cell biomass growth.

## 2. Experimental Work

### 2.1. Plant Material

In this research two species of *Calendula* were used as a plant material. Certificated seeds of *C. officinalis* and *C. arvensis* were bought from Ceylan Agricultural Company in Turkey.

### 2.2. MS Medium Preparing

The MS mediums was prepared as ¼ strength and supplied with different combinations and concentration of auxin and cytokinin, pH adjusted to 5.80 and 8 g/l agar, 15g of sucrose were added to the MS medium [40]. The materials used in the laboratory and the MS nutrient mediums were autoclaved for 15 min.

In order to begin the cell suspension cultures, it was prepared the four different MS mediums which supplemented with the

different concentrations of auxin (NAA, IAA) and cytokinin (BAP)[41].

### 2.3. Calli Growth and Cell Suspension Culture

Calli growth were established as described from [41]. After the 4<sup>th</sup> callus subculture 120 days old calli of *C. officinalis* and *C. arvensis* species were transferred to the cell suspension culture correlated with MS1, MS3, MS4, MS6 nutrient medium. At the 0. culture initiation of cell suspension, the transfer of calli was performed as 1 g. into the 100 ml of erlenmeyer flask. The cell suspension cultures of two *Calendula* species were placed in the orbital shaker adjusted to 110 rpm (Table 1).

**Table 1:** The MS nutrient mediums used the cell suspension cultures

| Medium | Auxin (mg/l) | Cytokinin (mg/l) |
|--------|--------------|------------------|
|        | NAA (mg/l)   | BAP (mg/l)       |
| MS1    | 1            | 1                |
| MS3    | 0,5          | 5                |
|        | IAA (mg/l)   | BAP (mg/l)       |
| MS4    | 1            | 1                |
| MS6    | 0,5          | 5                |

### 2.4. The Growth Parameters of Cell Suspension Cultures

It was determined the culture period reached the maximum cell viability and the maximum fresh/dry weight in the 0. culture of cell suspension. For this purpose, the cell counting and the measurement of fresh/dry weight was carried out. So, the cell suspension cultures were taken into the 1<sup>th</sup> subculture before the cell deaths began.

The cell counting was carried out to determine the cell viability in every 5 days during the 40 days from the 0. day of the cell suspension cultures of *C. officinalis* and *C. arvensis* species. 1 ml sample which taken from the cell suspension culture was colored with 1 ml of methylene blue and so the sample volume was completed to 2 ml. Cell counting were realized with Thoma chamber for the viability test. Thus, the average cell viability (%) was detected.


The measurements of fresh and dry weight was performed to determine the cell growth of the suspension cultures. For this purpose, the vacuum filtration system be composed of filtration pump (Lab 312, DrVAC-300), filtration flask and sieve with a pore size of 100  $\mu$ m (Sartorius) was utilized. The once in every 5 days from 0. day during 40 days, it was filtered by the vacuum filtration system via milipore filter by taking 5 ml of sample between the 0. culture and 1<sup>th</sup> culture of cell suspension cultures. With this way, the fresh weight and the liquid medium was separated from each other. The fresh weight (g/l) was determined by weighting the cell

weight remaining onto the filter on a precision scale. The fresh weight by isolating from the cell suspension cultures was dried by standing in the room temperature until the stable callus biomass was obtained (2 days). The results of measurements was recorded by weighting the drying cells on a precision scale. The measurement of fresh and dry weight was practiced in three repetition. The results of cell viability, fresh and dry weight measurement was evaluated with Tukey Multiple Comparison Test.

## 3. Results and Discussion

### 3.1. Results of The Cell Suspension Culture

In the 0. culture of cell suspension culture, it was observed that the nutrient medium was clear, become blurred, turned to brown blurring at the 0., 15. and 30. day, respectively. Besides it was observed that 0., 15. and 30. day of the cells was not separated, separated partially and separated from each other almost completely in the cell suspension 0. culture, respectively. At the end of cell suspension 4. culture (120 days), photography of the cell biomass in the MS1, MS3, MS4, MS6 nutrient mediums was showed at the Figure 1.



**Figure 1:** At the end of cell suspension 4<sup>th</sup> culture (120 days)  
a) MS1, b) MS3, c) MS4, d) MS6 nutrient mediums

Table 2. Descriptive statistics and Tukey Multiple Comparison Test results according to day, nutrient medium with regard to percent cell viability (%)

| Day     | Nutrient medium           |                           |                           |                           |
|---------|---------------------------|---------------------------|---------------------------|---------------------------|
|         | MS6                       | MS4                       | MS3                       | MS1                       |
|         | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ |
| 0. day  | 77.170 $\pm$ 1.320Cg      | 81.460 $\pm$ 1.340Bf      | 82.220 $\pm$ 0.290Bf      | 85.250 $\pm$ 1.080Af      |
| 5. day  | 79.770 $\pm$ 1.180Cf      | 83.570 $\pm$ 1.190Be      | 84.730 $\pm$ 1.110Be      | 87.910 $\pm$ 1.040Ae      |
| 10. day | 82.440 $\pm$ 1.030De      | 84.615 $\pm$ 0.000Cde     | 86.616 $\pm$ 0.982Bd      | 88.709 $\pm$ 0.834Ae      |
| 15. day | 84.921 $\pm$ 0.892Dd      | 86.453 $\pm$ 0.806Cc      | 89.775 $\pm$ 0.758Bc      | 91.399 $\pm$ 0.641Ad      |
| 20. day | 89.412 $\pm$ 0.636Dbc     | 91.017 $\pm$ 0.543Cb      | 94.742 $\pm$ 0.516Bb      | 96.906 $\pm$ 0.438Ab      |
| 25. day | 90.863 $\pm$ 0.552Dab     | 93.843 $\pm$ 0.463Ca      | 96.906 $\pm$ 0.438Ba      | 98.589 $\pm$ 0.352Aa      |
| 30. day | 91.166 $\pm$ 0.534Da      | 94.095 $\pm$ 0.445Ca      | 97.099 $\pm$ 0.411Ba      | 98.686 $\pm$ 0.327Aa      |
| 35. day | 88.974 $\pm$ 0.662Dc      | 91.964 $\pm$ 0.487Cb      | 95.579 $\pm$ 0.435Bab     | 97.629 $\pm$ 0.336Aab     |
| 40. day | 82.090 $\pm$ 0.883De      | 85.960 $\pm$ 0.533Ced     | 89.724 $\pm$ 0.449Be      | 93.428 $\pm$ 0.337Ae      |

Note 1. Differences between nutrient medium averages shown with different capital letters in the same day is important ( $p \leq 0.05$ ).

Note 2. Differences between day averages shown with different small letters in the same nutrient medium is important ( $p \leq 0.05$ ).

### 3.2. Results of The Growth Parameters of Cell Suspension Cultures

The cell counting, the measurement of fresh and dry weight was carried out during 40 days in the cell suspension 0. culture (40 days) of *C. officinalis* and *C. arvensis* species and determined the average cell viability (%) and the average fresh and dry weight (g/l) was specified in the cell suspension 0. culture.

Because of the alkaline feature of methylene blue, methylene blue was colored the asidic compartment of cell (the nucleic acid in the cell nucleus and the organelle of cell). According to this method, while the cells colored the light blue compared to the nutrient medium was counted as the alive cell, the cells colored the dark blue was counted as the dead cell.

Day X nutrient medium interaction of average cell viability (%) in the cell suspension culture (MS1, MS3, MS4, MS6 nutrient mediums) was indicated in Table 2.

Day X plant species of average cell viability (%) Tukey Multiple Comparison Test results in the cell suspension culture (MS1, MS3, MS4, MS6 nutrient mediums) was indicated in Table 3.

**Table 3:** Descriptive statistics and Tukey Multiple Comparison Test results according to day and plant species with regard to percent cell viability


| Day     | Plant Species         | $\bar{X} \pm S_{\bar{x}}$ |
|---------|-----------------------|---------------------------|
| 0. day  | <i>C. officinalis</i> | 82.890 $\pm$ 1.230Af      |
|         | <i>C. arvensis</i>    | 80.170 $\pm$ 1.050Bf      |
| 5. day  | <i>C. officinalis</i> | 84.960 $\pm$ 1.110Ae      |
|         | <i>C. arvensis</i>    | 83.030 $\pm$ 1.140Be      |
| 10. day | <i>C. officinalis</i> | 86.352 $\pm$ 0.938Ad      |
|         | <i>C. arvensis</i>    | 84.837 $\pm$ 0.787Bd      |
| 15. day | <i>C. officinalis</i> | 88.954 $\pm$ 0.937Ac      |
|         | <i>C. arvensis</i>    | 87.320 $\pm$ 0.871Bc      |
| 20. day | <i>C. officinalis</i> | 93.571 $\pm$ 0.955Ab      |
|         | <i>C. arvensis</i>    | 92.468 $\pm$ 0.944Bb      |
| 25. day | <i>C. officinalis</i> | 95.513 $\pm$ 0.931Aa      |
|         | <i>C. arvensis</i>    | 94.588 $\pm$ 0.935Ba      |
| 30. day | <i>C. officinalis</i> | 95.701 $\pm$ 0.904Aa      |
|         | <i>C. arvensis</i>    | 94.822 $\pm$ 0.912Ba      |
| 35. day | <i>C. officinalis</i> | 94.030 $\pm$ 1.030Ab      |
|         | <i>C. arvensis</i>    | 93.040 $\pm$ 1.060Bb      |
| 40. day | <i>C. officinalis</i> | 88.380 $\pm$ 1.280Ac      |
|         | <i>C. arvensis</i>    | 87.220 $\pm$ 1.360Bc      |

Note 1. Differences between plant species averages shown with different capital letters in the same day is important ( $p \leq 0,05$ ).


Note 2. Differences between day averages shown with different small letters in the same day is important ( $p \leq 0,05$ ).

Results of the cell counting made, the average percentage cell viability was increased till the 25<sup>th</sup> day, reached a constant value between 25<sup>th</sup> and 30<sup>th</sup> day and was begun to decrease after 30<sup>th</sup> day in the cell suspension culture of *C. officinalis* and *C. arvensis*. It was revealed that the maximum of average percentage cell viability was acquired in the 30<sup>th</sup> day and from the MS1, MS4, MS3, MS6 nutrient medium, respectively. It was found that day X plant species X nutrient medium triple interaction p value is 0.9784 and it is not statistically significant.

The cell viability (%) graphs composed as regards to the cell counting in the cell suspension culture was indicated for *C. officinalis* and *C. arvensis* species (Figure 2-3).



**Figure 2:** The cell viability (%) in the cell suspension culture of *C. officinalis* species



**Figure 3:** The cell viability (%) in the cell suspension culture of *C. arvensis* species

The fresh and dry weight was increased from the 0. day to the 30. day, reached to the maximum value on the 30. day and decreased till on the 40. day from this point. The maximum fresh and dry weight was gotten from the MS1, MS4, MS3 and MS6 nutrient medium, respectively.

It was decided that the fresh and dry weight of *C. officinalis* species in the MS1 nutrient medium of the cell suspension culture was to be 115.07 g/l (fresh weight) and 56.176 g/l (dry weight) on the 30<sup>th</sup> day, while decreased to 109.06g/l (fresh weight) and 49.349 g/l (dry weight) on the 35. day. Otherwise, it was confirmed that the fresh and dry weight of *C. arvensis* species in the MS1 nutrient medium of cell suspension culture was to be 112.07 g/l (fresh weight) and 56.286 g/l (dry weight) on the 30. day, while decreased to 107.21 g/l (fresh weight) and 54.244 g/l (dry weight) on the 35. day. It was found that plant species X nutrient medium X day triple interaction of fresh/dry weight (g/l) is significant statistically (Table 4-5). In terms of fresh weight, difference between all day averages in all plant species and all nutrient medium is statistically important ( $p \leq 0,05$ ). That is, all of the average fresh weight in all days is different from each other. This wasn't lettered in the table (Tablo 4). In terms of dry weight, differences between day averages except for shown

romen numeral is statistically significant in the same nutrient medium and same plant species ( $p \leq 0.05$ ) (Table 5).

Plant species X nutrient medium X day triple interaction of fresh weight (g/l) during the cell suspension 0. culture of *C. officinalis* and *C. arvensis* species was indicated in Table 4.

Table 4. Descriptive statistics and Tukey Multiple Comparison Test results according to day, plant species, nutrient medium with regard to fresh weight

| Plant species   | <i>C. officinalis</i>     |                           |                           |                           |                           |                           | <i>C. arvensis</i>        |                           |                           |                           |                           |                           |
|-----------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
|                 | MS6                       | MS4                       | MS3                       | MS1                       | MS6                       | MS4                       | MS3                       | MS4                       | MS3                       | MS1                       | MS6                       | MS4                       |
| Nutrient medium | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ |
| 0. day          | 50.051 $\pm$ 0.001Aa      | 50.040 $\pm$ 0.004Aa      | 50.055 $\pm$ 0.002Aa      | 50.054 $\pm$ 0.001Aa      | 50.048 $\pm$ 0.003Aa      | 50.052 $\pm$ 0.007Aa      | 50.064 $\pm$ 0.010Aa      | 50.068 $\pm$ 0.019Aa      |                           |                           |                           |                           |
| 5. day          | 53.051 $\pm$ 0.001Ca      | 54.046 $\pm$ 0.002Bb      | 54.058 $\pm$ 0.003Ba      | 55.057 $\pm$ 0.003Aa      | 53.056 $\pm$ 0.003Da      | 54.207 $\pm$ 0.161Ba      | 54.063 $\pm$ 0.003Ca      | 55.102 $\pm$ 0.003Aa      |                           |                           |                           |                           |
| 10. day         | 57.064 $\pm$ 0.006Da      | 61.055 $\pm$ 0.006Ca      | 62.077 $\pm$ 0.003Ba      | 65.055 $\pm$ 0.005Ab      | 57.094 $\pm$ 0.007Da      | 61.058 $\pm$ 0.003Ca      | 62.048 $\pm$ 0.002Ba      | 65.168 $\pm$ 0.003Aa      |                           |                           |                           |                           |
| 15. day         | 65.065 $\pm$ 0.006Da      | 73.055 $\pm$ 0.007Ca      | 74.070 $\pm$ 0.003Bb      | 80.052 $\pm$ 0.002Aa      | 64.058 $\pm$ 0.003Da      | 73.090 $\pm$ 0.003Ca      | 75.167 $\pm$ 0.006Ba      | 79.099 $\pm$ 0.004Ab      |                           |                           |                           |                           |
| 20. day         | 78.050 $\pm$ 0.001Db      | 91.054 $\pm$ 0.003Ca      | 92.069 $\pm$ 0.003Ba      | 100.06 $\pm$ 0.001Aa      | 80.073 $\pm$ 0.002Da      | 90.034 $\pm$ 0.002Cb      | 91.054 $\pm$ 0.002Bb      | 100.06 $\pm$ 0.006Aa      |                           |                           |                           |                           |
| 25. day         | 84.063 $\pm$ 0.005Db      | 98.053 $\pm$ 0.006Ca      | 102.07 $\pm$ 0.004Ba      | 110.06 $\pm$ 0.007Ab      | 88.059 $\pm$ 0.003Da      | 97.024 $\pm$ 0.003Cb      | 101.06 $\pm$ 0.002Bb      | 110.53 $\pm$ 0.001Aa      |                           |                           |                           |                           |
| 30. day         | 86.064 $\pm$ 0.004Db      | 101.05 $\pm$ 0.005Ca      | 106.07 $\pm$ 0.004Ba      | 115.07 $\pm$ 0.006Aa      | 90.034 $\pm$ 0.003Da      | 99.056 $\pm$ 0.004Cb      | 104.04 $\pm$ 0.004Bb      | 112.07 $\pm$ 0.003Ab      |                           |                           |                           |                           |
| 35. day         | 84.044 $\pm$ 0.004Db      | 95.043 $\pm$ 0.003Ca      | 99.085 $\pm$ 0.006Ba      | 109.06 $\pm$ 0.005Aa      | 85.272 $\pm$ 0.034Da      | 94.216 $\pm$ 0.001Cb      | 99.108 $\pm$ 0.002Ba      | 107.21 $\pm$ 0.003Ab      |                           |                           |                           |                           |
| 40. day         | 72.039 $\pm$ 0.006Db      | 80.040 $\pm$ 0.004Ca      | 82.077 $\pm$ 0.003Ba      | 89.070 $\pm$ 0.005Aa      | 72.416 $\pm$ 0.002Da      | 79.206 $\pm$ 0.002Cb      | 82.098 $\pm$ 0.003Ba      | 87.108 $\pm$ 0.002Ab      |                           |                           |                           |                           |

Note 1. Differences between nutrient medium averages shown with different capital letters in the same species and same day is important ( $p \leq 0.05$ ).

Note 2. Differences between plant species averages shown with different small letters in the same day and same nutrient medium is important ( $p \leq 0.05$ ).




Table 5. Descriptive statistics and Tukey Multiple Comparison Test results according to day, plant species, nutrient medium with regard to dry weight

| Plant species   | <i>C. officinalis</i>     |                           |                           |                           |                           |                           | <i>C. arvensis</i>        |                           |                           |                           |                           |                           |
|-----------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
|                 | MS6                       | MS4                       | MS3                       | MS1                       | MS6                       | MS4                       | MS3                       | MS4                       | MS3                       | MS1                       | MS6                       | MS4                       |
| Nutrient medium | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ | $\bar{X} \pm S_{\bar{X}}$ |
| 0. day          | 10.009 $\pm$ 0.003Aa      | 10.021 $\pm$ 0.004Aa      | 10.042 $\pm$ 0.003Aa      | 10.052 $\pm$ 0.003Aa      | 10.119 $\pm$ 0.003Aa      | 10.108 $\pm$ 0.004Aa      | 10.108 $\pm$ 0.003Aa      | 10.108 $\pm$ 0.004Aa      | 10.108 $\pm$ 0.003Aa      | 10.327 $\pm$ 0.003Aa      | 10.327 $\pm$ 0.003Aa      | 10.327 $\pm$ 0.003Aa      |
| 5. day          | 11.079 $\pm$ 0.001Ca      | 12.021 $\pm$ 0.003Ba      | 12.071 $\pm$ 0.004Ba      | 13.040 $\pm$ 0.003Aa      | 11.111 $\pm$ 0.005Ca      | 12.052 $\pm$ 0.002Ba      | 12.175 $\pm$ 0.003Ba      | 12.052 $\pm$ 0.002Ba      | 12.175 $\pm$ 0.003Ba      | 13.269 $\pm$ 0.002Aa      | 13.269 $\pm$ 0.002Aa      | 13.269 $\pm$ 0.002Aa      |
| 10. day         | 14.010 $\pm$ 0.002Ca      | 15.024 $\pm$ 0.003Ba      | 15.142 $\pm$ 0.004Ba      | 16.189 $\pm$ 0.002Aa      | 14.008 $\pm$ 0.002Ca      | 15.011 $\pm$ 0.003Ba      | 15.057 $\pm$ 0.002Ba      | 15.011 $\pm$ 0.003Ba      | 15.057 $\pm$ 0.002Ba      | 16.093 $\pm$ 0.003Aa      | 16.093 $\pm$ 0.003Aa      | 16.093 $\pm$ 0.003Aa      |
| 15. day         | 22.220 $\pm$ 0.002Da      | 26.199 $\pm$ 0.002Ca      | 27.092 $\pm$ 0.002Bb      | 30.159 $\pm$ 0.001Aa      | 21.060 $\pm$ 0.002Db      | 26.287 $\pm$ 0.003Ca      | 28.178 $\pm$ 0.003Ba      | 26.287 $\pm$ 0.003Ca      | 28.178 $\pm$ 0.003Ba      | 29.840 $\pm$ 0.006Aa      | 29.840 $\pm$ 0.006Aa      | 29.840 $\pm$ 0.006Aa      |
| 20. day         | 37.791 $\pm$ 0.005Db      | 45.727 $\pm$ 0.003Ca      | 47.537 $\pm$ 0.003Ba      | 50.193 $\pm$ 0.002Ab      | 39.029 $\pm$ 0.003Da      | 45.033 $\pm$ 0.002Cb      | 45.928 $\pm$ 0.004Bb      | 45.033 $\pm$ 0.002Cb      | 45.928 $\pm$ 0.004Bb      | 51.091 $\pm$ 0.004Aa      | 51.091 $\pm$ 0.004Aa      | 51.091 $\pm$ 0.004Aa      |
| 25. day         | 42.428 $\pm$ 0.004Db      | 48.007 $\pm$ 0.003Cb      | 50.262 $\pm$ 0.002Bb      | 55.018 $\pm$ 0.003Ab      | 43.400 $\pm$ 0.003Da      | 49.218 $\pm$ 0.002Ca      | 51.053 $\pm$ 0.002Ba      | 49.218 $\pm$ 0.002Ca      | 51.053 $\pm$ 0.002Ba      | 56.208 $\pm$ 0.001Aa      | 56.208 $\pm$ 0.001Aa      | 56.208 $\pm$ 0.001Aa      |
| 30. day         | 43.085 $\pm$ 0.002Db      | 49.068 $\pm$ 0.002Cb      | 51.324 $\pm$ 0.002Bb      | 56.176 $\pm$ 0.002Aa      | 43.785 $\pm$ 0.083Da      | 50.102 $\pm$ 0.041Ca      | 52.101 $\pm$ 0.011Ba      | 50.102 $\pm$ 0.041Ca      | 52.101 $\pm$ 0.011Ba      | 56.286 $\pm$ 0.045Aa      | 56.286 $\pm$ 0.045Aa      | 56.286 $\pm$ 0.045Aa      |
| 35. day         | 34.182 $\pm$ 0.002Db      | 41.734 $\pm$ 0.003Cb      | 44.269 $\pm$ 0.005Bb      | 49.349 $\pm$ 0.004Ab      | 41.240 $\pm$ 1.330Da      | 49.197 $\pm$ 0.018Ca      | 50.328 $\pm$ 0.212Ba      | 49.197 $\pm$ 0.018Ca      | 50.328 $\pm$ 0.212Ba      | 54.244 $\pm$ 0.036Aa      | 54.244 $\pm$ 0.036Aa      | 54.244 $\pm$ 0.036Aa      |
| 40. day         | 27.256 $\pm$ 0.016Db      | 33.528 $\pm$ 0.004Ca      | 36.569 $\pm$ 0.003Ba      | 43.210 $\pm$ 0.003Aa      | 27.837 $\pm$ 0.004Da      | 32.968 $\pm$ 0.004Cb      | 33.427 $\pm$ 0.004Bb      | 32.968 $\pm$ 0.004Cb      | 33.427 $\pm$ 0.004Bb      | 39.235 $\pm$ 0.015Ab      | 39.235 $\pm$ 0.015Ab      | 39.235 $\pm$ 0.015Ab      |


Note 1. Differences between nutrient medium averages shown with different capital letters in the same species and same day is important ( $p \leq 0.05$ ).Note 2. Differences between plant species averages shown with different small letters in the same day and same nutrient medium is important ( $p \leq 0.05$ ).

Plant species X nutrient medium X day triple interaction of dry weight (g/l) during the cell suspension culture of *C. officinalis* and *C. arvensis* species was indicated in Table 5.


The changing amount of fresh and dry weight of *C. officinalis* species was indicated in the Figure 4-5.



**Figure 4:** The fresh weight (g/l) in the cell suspension culture of *C. officinalis*




**Figure 7:** The dry weight (g/l) in the cell suspension culture of *C. Arvensis*



**Figure 5:** The dry weight (g/l) in the cell suspension culture of *C. officinalis*

The changing amount of fresh and dry weight of *C. arvensis* species was indicated in the Figure 6-7.



**Figure 6:** The fresh weight (g/l) in the cell suspension culture of *C. arvensis*

According to the measurement of cell viability (%), fresh and dry weight, it was proven that the cell suspension culture of *C. officinalis* and *C. arvensis* species (MS1, MS3, MS4, MS6 nutrient medium) was to be lag phase between on the 0.-5.day, log phase between on the 5.-25. day and the death phase on the 25.-30. day (the fresh and dry weight increased very little on the 25.-30. day). In reference to cell viability counting and the measurement of fresh/dry weight, it was determined that the cell viability and fresh/dry weight was begun to decrease after the 30<sup>th</sup> day. As a result of this, the subcultures was carried out by repeating in the every 30 days.

The cells in suspension culture can exhibit greater rates of cell division than cells in callus culture[42; 43; 44; 45; 46]. To optimize biomass production of an important medicinal plant, *C. officinalis* and *C. arvensis*, I studied the impact of adding factors such as the plant growth regulators (i.e., NAA, BAP) on the growth of biomass of *C. officinalis* and *C. arvensis* in cell suspension culture at 5-day intervals during 120 days. Nguyen and Paek found that MS media containing 2,4-D is optimal for culturing *P. ginseng* in cell suspension culture. Thus, the results obtained in the present study are interesting because most cell suspension cultures of ginseng cells previously documented required 2,4-D, which is unsuitable for pharmaceutical and food industrial use due to its potency as an herbicide and a carcinogen. Generally, cytokinins and auxins promote cell division and cell expansion in plant cell suspension cultures[47; 48; 49; 50]. As general information, [51] reported that auxin and cytokinin are key regulators of plant secondary growth. A study by [52] also demonstrated the essential role of cytokinins in the cell cycle and primary metabolite formation in crop plant cell suspension cultures.

In contrast to [53], I established that the fresh and dry weight was increased and decreased with parallel to each other. I identified that the fresh and dry weight wasn't increased during the cell suspension culture (40 days). Otherwise, [53] was indicated that the fresh weight was increased during the cell suspension culture (45 days). Consistent with my research, [53] was stated that the cells were died in the cell suspension culture after a while. Otherwise, [53] was defined that the fresh and dry weight was increased in the logarithmic phase up to the 20. day in the cell suspension culture done with *C. officinalis* cotyledon and hypocotyl. In contrast

to[53], I specified that the fresh and dry weight wasn't increased until the end of cell suspension culture (40 days).

Myresearch results were parallel with [54]in respect to the color of cell suspension cultures differentiated according to the culture days.Consistent with myresearch results, [55] reported that light conditions increase cell clustering in *C. officinalis* cell suspension cultures. It was determined that the cell amounts dispersed in the nutrient mediums and the size of aggregates increased during three months.

#### 4. Conclusion

In conclusion, cell proliferation has been achieved from callus by administration of NAA, BAP and IAA to the cell suspension cultures.The growth parameters of the cultures were detected by determining the period in which cell suspension cultures of *C. officinalis* and *C. arvensis* species had the highest cell viability and the maximum cell weight.The most appropriate period for the production of excess secondary metabolite, which may be the next stage of my study, could be determined.Plant growth regulators, which may be a stress factor in the nutrient medium, are also associated with the amount of secondary metabolite.

#### Acknowledgements

This research is a part of Miss. Nergis KAYA's Doctorate Thesis in Biological Science which is successfully completed in Graduate School of Natural and Applied Sciences in Canakkale Onsekiz Mart University under the supervisory of Prof. Dr. Cüneyt AKI. This research project which numbered as FDK 2015-401 was supported by Canakkale Onsekiz Mart University Commission of Scientific Research Projects.

#### References

- [1] Davis P.H., Flora of Turkey and the East Aegean Islands, Edinburg Universty, **1982**, 8.
- [2] Gantait S., Chattopadhyay T.K., Agrotechniques to maximize productivity of hydroalcoholic extract from medicinal garden herb calendula. J. Nat. Prod. Rad., **2005**, 4(2) 113-116.
- [3] Jimenez-Medina E., Garcia-Lora A., Paco L., Collada A., Garrido F. A New Extract of The Plant *Calendula officinalis* Produces A Dual In vitro Effect: Cytotoxic Anti-tumor Activity and Lymphocyte Activation. BMC Cancer,**2006**,6: 119.
- [4] Mazzio EA, Soliman KFA., In vitro screening for the tumoricidal properties of international medicinal herbs, Phytother Res., **2009**, 23: 385–398.
- [5] Matić IZ, Juranic Z, Šavikin K, Zdunić G, Nađvinski N, Gođevac D., Chamomile and marigold tea: Chemical characterization and evaluation of anticancer activity, Phytother. Res., **2013**, 27: 852–858.
- [6] Teiten MH, Gaascht F, Dicato M, Diederich M., Anticancer bioactivity of compounds from medicinal plants used in European medieval traditions, Biochem. Pharmacol., **2013**, 86: 1239–1247.
- [7] Dumenil G, Chemli R, Balansard C, Guiraud H, Lallemand M., Evaluation of antibacterial properties of marigold flowers (*Calendula officinalis* L.) and other homeopathic tinctures of *C. officinalis* L. and *C. arvensis* L., Ann. Pharm. Fr., **1980**, 38(6): 493-499.
- [8] Modesto A, Lima KC, Uzeda M., Effects of three different infant dentifrices on biofilms and oral microorganisms, J. Clin. Pediatr. Dent., **2000**, 24(3): 237-243.
- [9] Efstratiou E, Hussain AI, Nigam PS, Moore JE, Ayub MA, Rao JR., Antimicrobial activity of *Calendula officinalis* petal extracts against fungi, as well as Gram-negative and Gram-positive clinical pathogens, **2012**, Complement. Ther. Clin. Pract., 18: 173- 176.
- [10] Farjana A, Zerín N, Kabir MS. 2014. Antimicrobial activity of medicinal plant leaf extracts against pathogenic bacteria, Asian Pac. J. Trop. Dis., **2012**, 4:2, pp. 920-923.
- [11] Vieira O., Laranjinha J., Madeira V., Almeida L. Cholesteryl Ester Hydroperoxide Formation in Myoglobin-Catalyzed Low Density Lipoprotein Oxidation. Concerted Antioxidant Activity of Caffeic and p-Coumaric Acids with Ascorbate. Biochemical Pharmacology 1998, 55 (3): 333–340.
- [12] Nabi S., Ahmed N, Khan MJ, Bazai Z, Yasinzi M, Al-Kahraman Y.M.S.A., In vitro Antileishmanial, Antitumor Activities and Phytochemical Studies of Methanolic Extract and its Fractions of Juniperus Excelsa Berries, World Appl. Sci. J., **2012**, 19 (10): 1495- 1500.
- [13] Nikmehr B., Ghaznavi H, Rahbar A, Sadr S, Mehrzadi S, In vitro anti leishmanial activity of methanolic extracts of *Calendula officinalis* flowers, Chinese J. Nat. Med., **2014**, 12(6):423–427.
- [14] Kalvatchev Z., Walder R., Garzaro D. Anti-HIV Activity of Extracts From *Calendula officinalis* Flowers, Biomed Pharmacother., 1997, 51 (4): 176-80.
- [15] Çetkovic GS, Djilas SM, Canadanovic-Brunet JM, Tumbas VT., Antioxidant properties of marigold extracts, Food Res. Int., **2004**, 37: 643–650.
- [16] Erçetin T, Senol F, Orhan I, Toker G., Comparative assessment of antioxidant and cholinesterase inhibitory properties of the marigold extracts from *Calendula arvensis* L. and *Calendula officinalis* L., Ind. Crop. Prod., **2012**, 36: 203–208.
- [17] Babaee N, Moslemi D, Khalilpour M, Vejdani F, Moghadamnia Y, Bijani A, Baradaran M, Kazemi MT, Khalilpour A, Pouramir M, Moghadamnia AA., Antioxidant capacity of *Calendula officinalis* flowers extract and prevention of radiation induced oropharyngeal mucositis in patient s with head and neck cancers: a randomized controlled clinical study, J. Pharm. Sci., **2013**, 7: 21(1)-18.
- [18] Boucaud-Maitre Y, Algernon O, Raynaud J., Cytotoxic and antitumoral activity of *Calendula officinalis* extracts, Pharmazie, **1988**, 43(3):220-1
- [19] Ukiya M., Akihisa T., Yasukawa K., Tokuda H., Suzuki T., Kimura Y. Anti-Inflammatory, Anti-Tumor-Promoting, and Cytotoxic Activities of Constituents of Marigold (*Calendula officinalis*) Flowers. J. Nat. Prod., 2006,69 (12): 1692-6.
- [20] De Tommasi N., Conti C., Stein M.L., Pizza C., Structure and in vitro antiviral activity of triterpenoid saponins from *Calendula arvensis*, Planta Med., **1991**, 57(3): 250- 253.



- [21] Hamburger M, Adler S, Baumann D, Förg A., Weinreich B., Preparative purification of the major anti-inflammatory triterpenoid esters from Marigold (*Calendula officinalis*), *Fitoterapia*, **2003**, 74: 328–338.
- [22] Eglseer-Zitterl K, Sosa S, Jurenitsch J, SchubertZsilavec M, Della Loggia R, Tubaro A, Bertoldi M, Franz C., Anti-oedematous activities of the main triterpendiol esters of marigold (*Calendula officinalis* L.), *J. Ethnopharmacol.*, **1997**, 57: 139 – 144.
- [23] Marukami T, Kishi A, Yoshikawa M., Medicinal flowers. IV. Marigold. (2). Structures of new ionone and sesquiterpene glycosides from egyptian *Calendula officinalis*, *Chem. Pharm. Bull.*, **2001**, 49(8): 974-978 (Tokyo).
- [24] Shipochliev T., Uterotonic action of extracts from a group of medicinal plants, *Vet. Med. Nauki.*, **1981**, 18(4): 94-98.
- [25] Duran V, Matic M, Jovanovic M, Mimica N, Gajinov Z, Poljacki M, Boza P., Results of the clinical examination of an ointment with marigold (*Calendula officinalis*) extract in the treatment of venous leg ulcers, *Int. J. Tissue React.*, **2005**, 27(3): 101-106.
- [26] Uğulu İ, Başlar S, Yörek N, Doğan Y., The investigation and quantitative ethnobotanical evaluation of medicinal plants used around Izmir province. Turkey, *J. Med. Plant Res.*, **2009**, 3(5): 345-367.
- [27] İşlek C., Serbest ve Tutuklanmış *Capsicum annuum* L. Hücre Süspansiyon Kültürlerinde Kapsaisin Üretimi Üzerine Bazı Uyarıcıların Etkisi. Doktora Tezi. Ankara Üniversitesi (Ankara, Türkiye, **2009**).
- [28] Kaya N., *Catharanthus roseus* (L.) G. Don. Türünde Bitki Büyüme Düzenleyicilerinin In vitro Somaklonal Varyasyon üzerine Etkilerinin Belirlenmesi. Yüksek Lisans Tezi. Çanakkale Onsekiz Mart Üniversitesi (Çanakkale, Türkiye, **2012**).
- [29] Türkmen O.S., Kazdağı'nda Yetişen Oğulotu, Adaçayı ve Kekik Türlerinin Doku Kültürü Yöntemiyle Muhafazası ve Çoğaltılması, Yüksek Lisans Tezi, Çanakkale Onsekiz Mart Üniversitesi (Çanakkale, Türkiye, **2009**).
- [30] Molnár Z., Ördög V., 2005. The effect of cyanobacterial compounds on the organogenesis of pea cultured in vitro. *Proceedings of the 8<sup>th</sup> Hungarian Congress on Plant Physiology and the 6<sup>th</sup> Hungarian Conference on Photosynthesis*, 49(1-2):37-38.
- [31] Yamaguchi M., Kato H., Yoshida S., Yamamura S., Uchimiya H. ve Umeda M., Control of In Vitro Organogenesis by Cyclin-Dependent Kinase Activities in Plants, *PNAS*, **2003**, 100 (13): 8019–8023.
- [32] Di Cosmo F., Misawa M., *Plant Cell Cultures and Microbial Insult: Interactions*, *Trends Biotechnol.*, **1985**, 3: 318-322.
- [33] Zhao J., Davis L.C., Verpoorte R., Elicitor Signal Transduction Leading to Production of Plant Secondary Metabolites, *Biotechnol. Adv.*, **2005**, 23: 283–333.
- [34] Savitha B.C., Thimmaraju R., Bhagyalakshmi N., Ravinshankar A.G., Different Biotic and Abiotic Elicitors Influence Betalain Production in Hairy Root Cultures of *Beta vulgaris* in Shake-Flask and Bioreactor, *Process Biochem.*, **2006**, 41 (1): 50-60.
- [35] Namdeo A.G., Plant Cell Elicitation For Production of Secondary Metabolites: A Review, *Pharmacognosy. Rev.*, **2007**, 1: 69-79.
- [36] Butcher D.N., Ingram D.S., 1976. *Plant Tissue Culture*. London, Arnold.
- [37] Dodds J.H., Roberts L.W., 1985. *Experiments in Plant Tissue Culture* sec. Edi. Cambridge Univ. Press Presented by Britain.
- [38] Skoog F, Miller CO. Chemical regulation of growth and organ formation in plant tissues cultured in vitro. *Symp Soc Exp Biol*, 1957, 54: 118– 130.
- [39] Lee M, Phillips RL. The chromosomal basis of somaclonal variation. *Annu Rev Plant Physiol Plant Mol Biol*, 1998, 39: 413–437.
- [40] Murashige T., Skoog F., A Revised Medium for Rapid Growth and Bioassays with Tobacco Tissue Culture, *Physiol. Plant.*, **1962**, 15: 473-497.
- [41] Kaya N., Akı C. In vitro Effects of Plant Growth Regulators on Callus Formation in *Calendula officinalis* L. and *Calendula arvensis* L. Species. *Annals of Biological Research*, 2017, 8 (3): 1-7.
- [42] Mustafa NR, de Winter W, van Iren F, Verpoorte R. Initiation, growth and cryopreservation of plant cell suspension cultures. *Nat Protoc* 2011; 6(6): 715-742.
- [43] Kikowska M, Kedziora I, Krawczyk A, Thiem B. Methyl jasmonate, yeast extract and sucrose stimulate phenolic acids accumulation in *Eryngium planum* L. shoot cultures. *Acta Biochim Pol* 2015; 62(2): 197-200.
- [44] Abyari M, Nasr N, Soorni J, Sadhu D. Enhanced accumulation of scopoletin in cell suspension culture of *Spilanthes acmella* Murr. using precursor feeding. *Braz Arch Biol Technol* 2016; 59. e16150533.
- [45] Te-chato S, Hilae A, Komgrit IP. Effects of cytokinin types and concentrations on growth and development of cell suspension culture of oil palm. *J Agric Technol* 2008; 4: 157-163.
- [46] Baskaran P, Kumari A, Van Staden J. Embryogenesis and synthetic seed production in *Mondia Whitei*. *Plant Cell Tissue Organ Cult* 2015; 121(1): 205-214.
- [47] Lian ML, Chakrabarty D, Paek KY. Effect of plant growth regulators and medium composition on cell growth and saponin production during cell-suspension culture of mountain ginseng (*Panax ginseng* C. A. Mayer). *J Plant Biol* 2002; 45(4): 201-206.
- [48] Ascough GD, Fennell CW, van Staden J. The regulation of plant growth and development in liquid culture. *South Afr J Bot* 2004; 70(2): 181-190.
- [49] Shi L, Wang C, Zhou X, Zhang Y, Liu Y, Ma C. Production of salidroside and tyrosol in cell suspension cultures of *Rhodiola crenulata*. *Plant Cell Tissue Organ Cult* 2013; 114(3): 295-303.
- [50] Bienaim'e C, Melin A, Bensaddek L, Attoumbr'e J, Nava-Saucedo E, Baltora-Rosset S. Effects of plant growth regulators on cell growth and alkaloids production by cell cultures of *Lycopodiella inundata*. *Plant Cell Tissue Organ Cult* 2015; 123(3): 523-533.
- [51] Jang G, Lee JH, Rastogi K, Park S, Oh SH, Lee JY. Cytokinin-independent secondary growth determines root biomass in radish (*Raphanus sativus* L.). *J Exp Bot* 2015; 66(15): 4607-4619.
- [52] Schm'ulling T, Werner T, Riefler M, Krupkov'a E, Bartrina I, Manns Y, et al. Cytokinin as a regulatory factor for yield and biomass distribution in crop plants, in phytohormones in plant biotechnology and agriculture. In: Mach'a'ckov'a I, Romanov GA, editors. *Proceedings of the NATO-Russia Workshop*; 12–16 May

2002. Moscow; Dordrecht: Springer Netherlands; 2003, p. 97-108.
- [53] Długosz M., Wiktorowska E., Wiśniewska A., Pączkowski C. Production of Oleanolic Acid Glycosides by Hairy Root Established Cultures of *Calendula officinalis* L. Acta Biochimica Polonica 2013, 60 (3): 467–473.
- [54] Dwivedi S., Alam A., Shekhawat G.Y. Antioxidant Response of *Stevia rebaudiana* (Bertoni) Bertoni (Angiosperms; Asteraceae) During Developing Phase of Suspension Cell Culture. Plant Science Today 2016,3 (2): 115-123.
- [55] Grzelak A, Janiszowska W. Initiation and growth characteristic of suspension culture of *Calendula officinalis* cells. Plant Cell Tiss Org Cult 2002; 71: 29–40.

## Author Profile

**Nergis Kaya** graduated in 2009 from Uludag University, Department of Biology. In 2010, she graduated from Uludag University, Institute of Education and Biology Teaching. In 2012, she graduated from Canakkale Onsekiz Mart University, Graduate School of Natural and Applied Sciences, Department of Biology. She received her PhD in 2016 from the Onsekiz Mart University. She has the title of Asst. Prof. Dr. at Canakkale Onsekiz Mart University since 2017.