



1. Schrodinger Equation - Solutions

Practice Set-1

1. Let v, p and E denote the speed, the magnitude of the momentum, and the energy of a free particle of rest mass m . Then

[NET DEC 2012]

A. $dE/dp = \text{constant}$

B. $p = mv$

C. $v = cp/\sqrt{p^2 + m^2c^2}$

D. $E = mc^2$

Solution:

$$p = m'v = \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}} \Rightarrow p^2 = \frac{m^2v^2}{1 - \frac{v^2}{c^2}} \Rightarrow m^2v^2 = p^2 - \frac{p^2v^2}{c^2}, m \rightarrow \text{rest mass energy}$$

$$\Rightarrow v^2 \left(m^2 + \frac{p^2}{c^2} \right) = p^2 \Rightarrow v^2 = \frac{p^2}{\frac{m^2c^2 + p^2}{c^2}} \Rightarrow v = \frac{pc}{\sqrt{p^2 + m^2c^2}}$$

The correct option is (c)

2. If a particle is represented by the normalized wave function

$$\psi(x) = \begin{cases} \frac{\sqrt{15}(a^2 - x^2)}{4a^{5/2}}, & \text{or } -a < x < a \\ 0 & , \text{ otherwise} \end{cases}$$

the uncertainty Δp in its momentum is

[NET DEC 2012]

A. $2\hbar/5a$

B. $5\hbar/2a$

C. $\sqrt{10}\hbar/a$

D. $\sqrt{5}\hbar/\sqrt{2}a$

Solution:

$$\Delta p = \sqrt{\langle p^2 \rangle - \langle p \rangle^2} \text{ and } \langle p \rangle = \frac{\langle \psi | -i\hbar \frac{\partial}{\partial x} | \psi \rangle}{\langle \psi | \psi \rangle}$$

$$\Rightarrow \langle p \rangle = \int_{-a}^a \frac{\sqrt{15}(a^2 - x^2)}{4a^{5/2}} (-i\hbar) \frac{\sqrt{15}}{4a^{5/2}} \frac{\partial}{\partial x} (a^2 - x^2) dx$$

$$= \int_{-a}^a \frac{15}{16a^5} (-i\hbar) (a^2 - x^2) (-2x) dx = +i\hbar \frac{2 \times 15}{16 \times a^5} \int_{-a}^a (a^2 x - x^3) dx = 0, \quad (\because \text{odd function})$$

$$\langle p^2 \rangle = -\hbar^2 \times \frac{15}{16a^5} \int_{-a}^a (a^2 - x^2) \frac{\partial^2}{\partial x^2} (a^2 - x^2) dx$$

$$= -\hbar^2 \times \frac{15}{16a^5} \times (-2) \int_{-a}^a (a^2 - x^2) dx = \hbar^2 \times \frac{15}{16a^5} \times 2 \left\{ a^2 \cdot x - \frac{x^3}{3} \right\}_{-a}^a$$

$$= \hbar^2 \times \frac{15}{16a^5} \times 2 \left[2a^3 - \frac{2a^3}{3} \right] = \hbar^2 \times \frac{15}{16} \times \frac{2}{a^5} \times 2a^3 \left[1 - \frac{1}{3} \right] = \frac{15\hbar^2}{4a^2} \times \frac{2}{3} = \frac{5\hbar^2}{2a^2}$$

$$\text{Now, } \Delta p = \sqrt{\langle p^2 \rangle - \langle p \rangle^2} = \sqrt{\frac{5\hbar^2}{2a^2} - 0} = \frac{\sqrt{5}\hbar}{\sqrt{2}a}$$

The correct option is (d)

3. The energies in the ground state and first excited state of a particle of mass $m = \frac{1}{2}$ in a potential $V(x)$ are -4 and -1 , respectively, (in units in which $\hbar = 1$). If the corresponding wavefunctions are related by $\psi_1(x) = \psi_0(x) \sinh x$, then the ground state eigenfunction is

[NET DEC 2012]

A. $\psi_0(x) = \sqrt{\sec hx}$

B. $\psi_0(x) = \sec hx$

C. $\psi_0(x) = \text{sech}^2 x$

D. $\psi_0(x) = \sec h^3 x$

Solution: Given that ground state energy $E_0 = -4$, first excited state energy $E_1 = -1$ and ψ_0, ψ_1 are corresponding wave functions.

Solving Schrödinger equation (use $m = \frac{1}{2}$ and $\hbar = 1$)

Solving Schrödinger equation (use $m = \frac{1}{2}$ and $\hbar = 1$)

$$\frac{-\hbar^2}{2m} \frac{\partial^2 \psi_0}{\partial x^2} + V \psi_0 = E_0 \psi_0 \Rightarrow -\frac{\partial^2 \psi_0}{\partial x^2} + V \psi_0 = -4 \psi_0 \dots (1)$$

$$\frac{-\hbar^2}{2m} \frac{\partial^2 \psi_1}{\partial x^2} + V \psi_1 = E_1 \psi_1 \Rightarrow -\frac{\partial^2 \psi_1}{\partial x^2} + V \psi_1 = -1 \psi_1 \dots (2)$$

Put $\psi_1 = \psi_0 \sinh x$ in equation (2) one will get

$$-\left[\frac{\partial^2 \psi_0}{\partial x^2} \cdot \sinh x + 2 \frac{\partial \psi_0}{\partial x} \cosh x + \psi_0 \sinh x \right] + V \psi_0 \sinh x = -\psi_0 \sinh x$$

$$-\left[\frac{\partial^2 \psi_0}{\partial x^2} + 2 \frac{\partial \psi_0}{\partial x} \coth x + \psi_0 \right] + V \psi_0 = -\psi_0$$

$$\left[-\frac{\partial^2 \psi_0}{\partial x^2} + V \psi_0 \right] - 2 \frac{\partial \psi_0}{\partial x} \coth x - \psi_0 = -\psi_0$$

$$\text{using relation } -\frac{\partial^2 \psi_0}{\partial x^2} + V \psi_0 = -4 \psi_0$$

$$-4 \psi_0 - 2 \frac{\partial \psi_0}{\partial x} \coth x - \psi_0 = -\psi_0 \Rightarrow \frac{d\psi_0}{\psi_0} = -2 \tanh x dx \Rightarrow \psi_0 = \sec h^2 x$$

The correct option is (c)

4. If $\psi(x) = A \exp(-x^4)$ is the eigenfunction of a one dimensional Hamiltonian with eigen value $E = 0$, the potential $V(x)$ (in units where $\hbar = 2m = 1$) is

[NET DEC 2013]

A. $12x^2$

B. $16x^6$

C. $16x^6 + 12x^2$

D. $16x^6 - 12x^2$

Solution: Schrodinger equation $-\nabla^2\psi + V\psi = 0$ (where $\hbar = 2m = 1$ and $E = 0$)

$$-\frac{\partial^2}{\partial x^2} (Ae^{-x^4}) + VAe^{-x^4} = 0 \Rightarrow -\frac{\partial}{\partial x} [e^{-x^4} \times -4x^3] + Ve^{-x^4} = 0$$

$$4 \left[\left\{ 3x^2 e^{-x^4} + x^3 (-4x^3 e^{-x^4}) \right\} \right] + Ve^{-x^4} = 0 \Rightarrow 12x^2 e^{-x^4} - 16x^6 e^{-x^4} + Ve^{-x^4} = 0$$

$$\Rightarrow V = 16x^6 - 12x^2$$

The correct option is (d)

5. A particle of mass m moves in one dimension under the influence of the potential $V(x) = -\alpha\delta(x)$, where α is a positive constant. The uncertainty in the product $(\Delta x)(\Delta p)$ in its ground state is

[NET JUNE 2016]

A. $2\hbar$

B. $\frac{\hbar}{2}$

C. $\frac{\hbar}{\sqrt{2}}$

D. $\sqrt{2}\hbar$

Solution: $V(x) = -\alpha\delta(x)$

For this potential wavefunction

$$\psi(x) = \begin{cases} \sqrt{\alpha}e^{\alpha x}, & x < 0 \\ \sqrt{\alpha}e^{-\alpha x}, & x > 0 \end{cases}$$

which evenfunction about $x = 0$ so $\langle x \rangle = 0, \langle p \rangle = 0$

$$\langle p^2 \rangle = -\hbar^2 \int_{-\infty}^{\infty} \psi^* \frac{d^2}{dx^2} \psi dx = -\hbar^2 \int_{-\infty}^0 \sqrt{\alpha}e^{\alpha x} \frac{d^2}{dx^2} \sqrt{\alpha}e^{\alpha x} dx - \hbar^2 \int_0^{\infty} \sqrt{\alpha}e^{-\alpha x} \frac{d^2}{dx^2} \sqrt{\alpha}e^{-\alpha x} dx$$

$$= -\hbar^2 \alpha^3 \int_{-\infty}^0 e^{2\alpha x} dx - \hbar^2 \alpha^3 \int_0^{\infty} e^{-2\alpha x} dx = -\frac{\hbar^2 \alpha^3}{2\alpha} - \frac{\hbar^2 \alpha^3}{2\alpha} = -\hbar^2 \alpha^2, \text{ which is not possible}$$

so, we will use the formula $\langle p \rangle^2 = \hbar^2 \int_{-\infty}^{\infty} \left| \frac{d\psi}{dx} \right|^2 dx = \hbar^2 \alpha^2, \Delta p = \sqrt{\langle p^2 \rangle - \langle p \rangle^2} = \hbar\alpha$ now,

$$\Delta x \cdot \Delta p = \frac{1}{\sqrt{2}\alpha} \cdot \hbar\alpha = \frac{\hbar}{\sqrt{2}}$$

The correct option is (c)

6. Consider the two lowest normalized energy eigenfunctions $\psi_0(x)$ and $\psi_1(x)$ of a one dimensional system. They satisfy $\psi_0(x) = \psi_0^*(x)$ and $\psi_1(x) = \alpha \frac{d\psi_0}{dx}$, where α is a real constant. The expectation value of the momentum operator in the state ψ_1 is

[NET DEC 2016]

A. $-\frac{\hbar}{\alpha^2}$

B. 0

C. $\frac{\hbar}{\alpha^2}$

D. $\frac{2\hbar}{\alpha^2}$

Solution: $\psi_1(x) = \alpha \frac{d\psi_0}{dx}$

$$\begin{aligned}\langle p_x \rangle &= \int_{-\infty}^{\infty} \psi_1^* p_x \psi dx = \int_{-\infty}^{\infty} \psi_1^* \left(-i\hbar \frac{\partial \psi_1}{\partial x} \right) dx = \int_{-\infty}^{\infty} \alpha^* \frac{d\psi_0}{dx} (-i\hbar \alpha) \frac{d^2 \psi_0}{dx^2} dx \\ &= -i\hbar |\alpha|^2 \int_{-\infty}^{\infty} \frac{d\psi_0}{dx} \frac{d^2 \psi_0}{dx^2} dx\end{aligned}$$

Integrate by parts

$$I = -i\hbar |\alpha|^2 \left(\left. \frac{d\psi_0}{dx} \frac{d\psi_0}{dx} \right|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} \frac{d^2 \psi_0}{dx^2} \frac{d\psi_0}{dx} dx \right) = 0 - (-i\hbar) |\alpha|^2 \int_{-\infty}^{\infty} \frac{d\psi_0}{dx} \frac{d^2 \psi_0}{dx^2} dx$$

$$I = 0 - (-i\hbar) |\alpha|^2 \int_{-\infty}^{\infty} \frac{d\psi_0}{dx} \frac{d^2 \psi_0}{dx^2} dx$$

$$\frac{d\psi_0}{dx} \rightarrow \frac{\psi_0}{\alpha}, \quad \psi_0 = 0, x \rightarrow \infty$$

$$I = 0 - I \Rightarrow 2I = 0 \Rightarrow I = 0 \Rightarrow \langle p_x \rangle = 0$$

The correct option is (b)

7. A particle in one dimension is in a potential $V(x) = A\delta(x-a)$. Its wavefunction $\psi(x)$ is continuous everywhere. The discontinuity in $\frac{d\psi}{dx}$ at $x = a$ is

[NET DEC 2016]

A. $\frac{2m}{\hbar^2} A\psi(a)$

B. $A(\psi(a) - \psi(-a))$

C. $\frac{\hbar^2}{2m} A$

D. 0

Solution: $-\frac{\hbar^2}{2m} \frac{d^2 \psi(x)}{dx^2} + A\delta(x-a)\psi(x) = E\psi(x)$
Integrates both side within limit

$$\begin{aligned}& a - \epsilon \text{ to } a + \epsilon \\ & -\frac{\hbar^2}{2m} \int_{a-\epsilon}^{a+\epsilon} \frac{d^2 \psi}{dx^2} dx + \int_{a-\epsilon}^{a+\epsilon} A\delta(x-a)\psi dx = E \int_{a-\epsilon}^{a+\epsilon} \psi(x) dx \\ & -\frac{\hbar^2}{2m} \left(\frac{d\psi_{II}}{dx} - \frac{d\psi_I}{dx} \right) + A\psi(a) = 0 \\ & \frac{d\psi_{II}}{dx} - \frac{d\psi_I}{dx} = \frac{2mA}{\hbar^2} \psi(a) \\ & \text{so discontinues in } \frac{d\psi}{dx} \text{ at } x = a \text{ is } \frac{2mA}{\hbar^2} \psi(a).\end{aligned}$$

The correct option is (a)

8. The normalized wavefunction of a particle in three dimensions is given by $\psi(r, \theta, \phi) = \frac{1}{\sqrt{8\pi a^3}} e^{-r/2a}$ where $a > 0$ is a constant. The ratio of the most probable distance from the origin to the mean distance from the origin, is [You may use $\int_0^\infty dx x^n e^{-x} = n!$]

[NET DEC 2017]

A. $\frac{1}{3}$

B. $\frac{1}{2}$

C. $\frac{3}{2}$

D. $\frac{2}{3}$

Solution: $\psi(r, \theta, \phi) = \frac{1}{\sqrt{8\pi a^3}} e^{-r/2a}$

$$\langle r \rangle = \iiint r \psi^* \psi r^2 dr \sin \theta d\theta d\phi = \frac{3}{2}(2a) = 3a$$

one can compare the wave function at hydrogen atom with Bohr radius $a_0 = 2a$ most probable distance,

$$\frac{d}{dr} r^2 e^{-r/a} = 0$$

$$r_P = 2a$$

$$\frac{r_P}{\langle r \rangle} = \frac{2a}{3a} = \frac{2}{3}$$

The correct option is (d)

9. The normalized wavefunction in the momentum space of a particle in one dimension is $\phi(p) = \frac{\alpha}{p^2 + \beta^2}$, where α and β are real constants. The uncertainty Δx in measuring its position is

[NET DEC 2017]

A. $\sqrt{\pi} \frac{\hbar \alpha}{\beta^2}$

B. $\sqrt{\pi} \frac{\hbar \alpha}{\beta^3}$

C. $\frac{\hbar}{\sqrt{2}\beta}$

D. $\sqrt{\frac{\pi}{\beta}} \frac{\hbar \alpha}{\beta}$

Solution: $\phi(p) = \frac{\alpha}{p^2 + \beta^2}$

From inverse Fourier transformation Normalize, $\psi(x) = \sqrt{\frac{\beta}{\hbar}} e^{-\frac{\beta|x|}{\hbar}} \langle x \rangle = 0$

$$\langle x^2 \rangle = \frac{\beta}{\hbar} \int_{-\infty}^{\infty} x^2 e^{-2\frac{\beta|x|}{\hbar}} dx = \frac{\hbar^2}{2\beta^2}$$

$$\Delta x = \sqrt{\langle x^2 \rangle - \langle x \rangle^2} = \frac{\hbar}{\sqrt{2}\beta}$$

The correct option is (c)

10. The product $\Delta x \Delta p$ of uncertainties in the position and momentum of a simple harmonic oscillator of mass m and angular frequency ω in the ground state $|0\rangle$, is $\frac{\hbar}{2}$. The value of the product $\Delta x \Delta p$ in the state, $e^{-i\hat{p}\ell/\hbar}|0\rangle$ (where ℓ is a constant and \hat{p} is the momentum operator) is

[NET DEC 2018]

A. $\frac{\hbar}{2} \sqrt{\frac{m\omega\ell^2}{\hbar}}$

B. \hbar

C. $\frac{\hbar}{2}$

D. $\frac{\hbar^2}{m\omega\ell^2}$

Solution: The correct option is (c)

Practice Set-2

1. Which of the following is an allowed wavefunction for a particle in a bound state? N is a constant and $\alpha, \beta > 0$. [GATE 2010]

A. $\psi = N \frac{e^{-\alpha r}}{r^3}$

B. $\psi = N(1 - e^{-\alpha r})$

C. $\psi = N e^{-\alpha x} e^{-\beta(x^2+y^2+z^2)}$

D. $\psi = \begin{cases} \text{non - zero constant} & \text{if } r < R \\ 0 & \text{if } r > R \end{cases}$

Solution: The correct option is (c)

Common data questions for 2 and 3

The wavefunction of particle moving in free space is given by, $\psi = (e^{ikx} + 2e^{-ikx})$

2. The energy of the particle is [GATE 2012]

A. $\frac{5\hbar^2 k^2}{2m}$

B. $\frac{3\hbar^2 k^2}{4m}$

C. $\frac{\hbar^2 k^2}{2m}$

D. $\frac{\hbar^2 k^2}{m}$

Solution: $H\psi = E\psi$, $H\psi = \frac{-\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} = \frac{-\hbar^2}{2m} [(ik)(ik)e^{ikx} + 2(-ik)(-ik)e^{-ikx}]$
 $\Rightarrow H\psi = \frac{\hbar^2 k^2}{2m} (e^{ikx} + 2e^{-2ikx}) = \frac{\hbar^2 k^2}{2m} \psi$
 The correct option is (c)

3. The probability current density for the real part of the wavefunction is

A. 1

B. $\frac{\hbar k}{m}$

C. $\frac{\hbar k}{2m}$

D. 0

Solution: The real part of the wave function $\psi_{\text{real}} = \cos kx + 2 \cos kx$
 Current density for real part of wave function = 0
 The correct option is (d)

4. Consider the wave function $Ae^{ikr}(r_0/r)$, where A is the normalization constant. For $r = 2r_0$, the magnitude of probability current density up to two decimal places, in units of $(A^2 \hbar k/m)$ is [GATE 2013]

Solution: $\vec{J} = |\psi|^2 \frac{\hbar k}{m} = |A|^2 \left| \frac{r_0}{r} \right|^2 \frac{\hbar k}{m} \Rightarrow J = |A|^2 \left| \frac{r_0}{2r_0} \right|^2 \frac{\hbar k}{m} \Rightarrow J = |A|^2 \frac{\hbar k}{4m} = (0.25) |A|^2 \frac{\hbar k}{m}$

5. The recoil momentum of an atom is p_A when it emits an infrared photon of wavelength 1500 nm, and it is p_B when it emits a photon of visible wavelength 500 nm. The ratio $\frac{p_A}{p_B}$ is [GATE 2014]

A. 1 : 1

B. 1 : $\sqrt{3}$

C. 1 : 3

D. 3 : 2

Solution: $p = \frac{h}{\lambda}$, $\frac{p_A}{p_B} = \frac{\lambda_B}{\lambda_A}$, $\frac{\lambda_B}{\lambda_A} = \frac{500}{1500} = 1 : 3$
 The correct option is (c)

6. The state of a system is given by $|\psi\rangle = |\phi_1\rangle + 2|\phi_2\rangle + 3|\phi_3\rangle$, where $|\phi_1\rangle$, $|\phi_2\rangle$ and $|\phi_3\rangle$ form an orthonormal set. The probability of finding the system in the state $|\phi_2\rangle$ is

[GATE 2016]

Solution: Probability that ψ in state $|\phi_2\rangle = \frac{2^2}{1^2+2^2+3^2} = \frac{4}{1+4+9} = \frac{4}{14} = \frac{2}{7} = 0.28$

7. The Compton wavelength of a proton is fm. (up to two decimal places).

[GATE 2017]

Solution: ($m_p = 1.67 \times 10^{-27}$ kg, $h = 6.626 \times 10^{-34}$ Js, $e = 1.602 \times 10^{-19}$ C, $c = 3 \times 10^8$ ms⁻¹)

8. Consider a one-dimensional potential well of width 3 nm. Using the uncertainty principle ($\Delta x \cdot \Delta p \geq \frac{\hbar}{2}$), an estimate of the minimum depth of the well such that it has at least one bound state for an electron is ($m_e = 9.31 \times 10^{-31}$ kg, $h = 6.626 \times 10^{-34}$ Js, $e = 1.602 \times 10^{-19}$ C)

[GATE 2017]

A. 1μeV

B. 1meV

C. 1eV

D. 1MeV

Solution: $E = \frac{p^2}{2m}$, $\Delta p = \frac{\hbar}{2\Delta x} \Rightarrow \Delta p = \frac{\hbar}{2a}$
 So, $E = \frac{\hbar^2}{8ma^2} = \frac{\hbar^2}{32\pi^2ma^2} = \frac{(6.6 \times 10^{-34})^2}{32 \times 10 \times 9.31 \times 10^{-31} \times 9 \times 10^{-18}} = .001 \times 10^{-19} \text{ J} \approx 1\text{meV}$
 The correct option is (b)

9. The Hamiltonian for a quantum harmonic oscillator of mass m in three dimensions is

$$H = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 r^2$$

where ω is the angular frequency. The expectation value of r^2 in the first excited state of the oscillator in units of $\frac{\hbar}{m\omega}$ (rounded off to one decimal place) is

[GATE 2018]

Solution: $\langle r^2 \rangle = \langle x^2 \rangle + \langle y^2 \rangle + \langle z^2 \rangle$

$$= \frac{\hbar}{2m\omega} [(2n_x + 1) + (2n_y + 1) + (2n_z + 1)]$$

For first excited state $n_x = 1, n_y = 0, n_z = 0$ Hence it is triply degenerate one can take

$$n_x = 0, n_y = 1, n_z = 0 \text{ or } n_x = 0, n_y = 0, n_z = 1$$

putting any one combination, expectation value of $r^2 = \frac{5}{2} \frac{\hbar}{m\omega} = 2.5 \frac{\hbar}{m\omega}$

10. Consider the motion of a particle along the x - axis in a potential $V(x) = F|x|$. Its ground state energy E_0 is estimated using the uncertainty principle. Then E_0 is proportional to

[GATE 2019]

A. $F^{1/3}$ B. $F^{1/2}$ C. $F^{2/5}$ D. $F^{2/3}$

Solution: $E = \frac{p^2}{2m} + F|x|$ $E = \frac{p^2}{2m} + Fx$ for $x > 0$ $E = \frac{p^2}{2m} - Fx < 0$ from uncertainty theory

$$\Delta x \cdot \Delta p = \hbar \Rightarrow \Delta p = \frac{\hbar}{\Delta x}$$

$$E = \frac{(\Delta p)^2}{2m} + F(\Delta x) \Rightarrow E = \frac{\hbar^2}{2m(\Delta x)^2} + F\Delta x$$

For minimum energy,

$$\frac{dE}{d\Delta x} = -\frac{\hbar^2}{m(\Delta x)^3} + F = 0 \Rightarrow \Delta x = \left(\frac{\hbar^2}{mF}\right)^{1/3} \frac{\hbar^2}{2m} \left(\frac{mF}{\hbar^2}\right)^{2/3} + F \left(\frac{\hbar^2}{mF}\right)^{1/3} \Rightarrow E \propto F^{2/3}$$

The correct option is **(d)**

