

Advanced Calculus

Alec Zabel-Mena

Text

An Introduction to Analysis (3^{rd} edition)

William R. Wade

October 16, 2020

Chapter 1

Chapter 1: The Real Number System.

1.1 The Ordered Field Axioms

We want to study the algebraic structure of the real numbers; the so called set \mathbb{R} . We assume standard knowledge of set theory. We display the field axioms below:

Postulate 1. *There are binary operations $+$ and \cdot defined on the set $\mathbb{R} \times \mathbb{R}$, satisfying the following properties $\forall a, b, c \in \mathbb{R}$:*

- (1) $a + b \in \mathbb{R}$.
- (2) $a + (b + c) = (a + b) + c$ and $a \cdot (b \cdot c) = (a \cdot b) \cdot c$.
- (3) $a + b = b + a$ and $a \cdot b = b \cdot a$.
- (4) $a \cdot (b + c) = (ab) + (a \cdot c)$.
- (5) *There exists a unique element $0 \in \mathbb{R}$ such that $\forall a \in \mathbb{R}$, $a + 0 = a$.*
- (6) *There exists a unique element $1 \neq 0 \in \mathbb{R}$ such that $\forall a \in \mathbb{R}$, $a \cdot 1 = a$.*
- (7) *For each $a \in \mathbb{R}$, there exists a unique element $-a \in \mathbb{R}$ such that $a + (-a) = 0$.*
- (8) *For each $a \in \mathbb{R} \setminus \{0\}$, there exists a unique element $a^{-1} \in \mathbb{R}$ such that $a \cdot a^{-1} = 1$.*

We call postulate 1 the **field axioms** for real numbers; in essence what it says is that the pairs $(\mathbb{R}, +)$ and (\mathbb{R}, \cdot) form abelian groups, and that \cdot distributes over $+$. We now get the following properties from the field axioms:

- (1) $(-1)(-1) = 1$.
- (2) $0a = 0$.
- (3) $-(a - b) = b - a$.
- (4) For $a, b \in \mathbb{R}$ and $ab = 0$, then either $a = 0$ or $b = 0$.

The proofs for these are relatively easy, and can be done at anytime as an exercise. We now introduce the second postulate called the **order axioms**:

Postulate 2. *There is a relation $<$ on $\mathbb{R} \times \mathbb{R}$ satisfying the following properties:*

- (1) *Given $a, b \in \mathbb{R}$, one and only one of the following hold: either $a < b$, $b < a$, or $a = b$.*
- (2) *If $a < b$ and $b < c$, then $a < c$.*
- (3) *If $a < b$ and $c \in \mathbb{R}$, then $a + c < b + c$.*
- (4) *If $a < b$ and $c > 0$, then $ac < bc$; if $c < 0$, then $bc < ac$.*

We may make the following remarks: by $b > a$ we mean that $a < b$, by $a \leq b$, we mean that $a < b$ or $a = b$, and by $a < b < c$, we mean that $a < b$ and $b < c$.

Definition. For $a \in \mathbb{R}$, a is called **nonnegative** if $a \geq 0$, a is called **nonpositive** if $a \leq 0$. We call a **positive** if $a > 0$ and we call a **negative** if $a < 0$.

Example 1.1. If $a \in \mathbb{R}$, show that $a \neq 0$ implies $a^2 > 0$; in particular that $-1 < 0 < 1$.

Proof. If $a \neq 0$, then either $a > 0$ or $a < 0$. If $a > 0$, then clearly $a^2 > 0$ and we are done. If $a < 0$, then $-a > 0$, hence $(-a)(-a) > (-a)0$, and so $a^2 > 0$.

Now we have that $1 \neq 0$, and that $1^2 = 1 > 0$, subtracting we get $1 - 1 > 0 - 1$ so $0 > -1$. ■

Example 1.2. If $a \in \mathbb{R}$, show that $0 < a < 1$ implies that $0 < a^2 < a$, and $a > 1$ implies that $a^2 > a$.

Proof. We have already that if $a > 0$, then $a^2 > 0$. Now suppose that $a < 1$ then multiplying by a we get $(a)(a) < 1(a)$, hence $a^2 < a$. Likewise, by the same reasoning, if $a > 1$ we get $a^2 > a$. ■

Example 1.3. Prove that:

- (1) $0 \leq a < b$ and $0 \leq c < d$ imply that $ac < bd$.
- (2) $0 \leq a < b$ imply that $0 \leq a^2 < b^2$ and $0 \leq \sqrt{a} < \sqrt{b}$.
- (3) $0 < a < b$ implies $0 < \frac{1}{b} < \frac{1}{a}$.

Proof. (1) Let $0 \leq a < b$ and $0 \leq c < d$. Then $0 \leq ac < bc$, and since $c < d$, then $bc < bd$, hence we have that $0 \leq ac < bd$.

- (2) Let $0 \leq a < b$. Then $0 \leq aa < ab$, and notice that $ab < bb$, so $0 \leq aa < bb$, i.e. $0 \leq a^2 < b^2$. Now notice that $a = (\sqrt{a})(\sqrt{a})$ and $b = (\sqrt{b})(\sqrt{b})$, by the previous result, we have $0 \leq (\sqrt{a})(\sqrt{a}) < (\sqrt{b})(\sqrt{b})$, hence we have that $0 \leq \sqrt{a} < \sqrt{b}$.

- (3) Let $0 < a < b$, multiplying by $\frac{1}{b}$, we get that $0 < a\frac{1}{b} < b\frac{1}{b}$ and get $0 < a\frac{1}{b} < 1$, now multiplying again by $\frac{1}{a}$, we get $0 < \frac{1}{a}a\frac{1}{b} < \frac{1}{a}$, thus we have that $0 < \frac{1}{b} < \frac{1}{a}$. ■

Definition. The **absolute value** of an element $a \in \mathbb{R}$ is a real number $|a|$ defined such that:

$$|a| = \begin{cases} a, & \text{if } a \geq 0 \\ -a, & \text{if } a < 0 \end{cases} \quad (1.1)$$

Remark. The absolute value is multiplicative, i.e. $|ab| = |a||b|$ for all $a, b \in \mathbb{R}$.

Proof. We do a casewise evaluation.

- (1) If $a = 0$, or $b = 0$, then $|ab| = 0 = |a||b|$ (since $|a| = a = 0$ or $|b| = b = 0$).
- (2) Let $a, b > 0$. then $|a| = a$ and $|b| = b$. Then $|ab| = ab$ as $ab > 0$. Hence $|ab| = ab = |a||b|$.
- (3) Let $a > 0$ and $b < 0$, then $|a| = a$ and $|b| = -b$ si $|a||b| = a(-b) = -ab = |ab|$ as $ab < 0$.
- (4) Let $a, b < 0$, then $|a| = -a$ and $|b| = -b$ and $|a||b| = (-a)(-b) = ab = |ab|$, since $ab > 0$.

■

Theorem 1.1.1. Let $a \in \mathbb{R}$ and let M be nonnegative. Then $|a| \leq M$ if and only if $-M \leq a \leq M$.

Proof. Notice that if $|a| \leq M$, then $-|a| \geq -M$. Suppose then that $a \geq 0$, then $|a| = a \leq M$, and since $-M \leq 0$, we have $-M \leq a \leq M$. Now if $a < 0$, $-|a| = a$, then $a \geq -M$, and since $a < 0$, $a \leq M$.

Conversely suppose that $-M \leq a \leq M$. Then $-M \leq a$ and $a \leq M$. For $-M \leq a$, multiplying by -1 we have $M \geq -a$. If $a \leq 0$ then $|a| = -a$ and $|a| \leq M$. If $a < 0$ then $|a| = -a \leq M$. Hence we have in both cases that $|a| \leq M$. ■

Theorem 1.1.2. The absolute value satisfies the following three properties For all $a, b \in \mathbb{R}$:

- (1) $|a| \geq 0$ with $|a| = 0$ if and only if $a = 0$.
- (2) $|a - b| = |b - a|$.
- (3) $|a + b| \leq |a| + |b|$.

Proof. If $a \leq 0$, then clearly $|a| \geq 0$, if $a < 0$, then $|a| = -a \geq 0$. Now if $a = 0$ then $|0| = 0$, and if $|a| = 0$, then $\pm a = 0$; then $a = 1a = (\pm 1)^2 a = (\pm 1)(\pm 1)a = (\pm 1)(\pm a)$, since $\pm a = 0$, $(\pm 1)0 = 0$ hence $a = 0$.

Now we have $(a - b) = -(b - a)$, so $|a - b| = |-1(b - a)| = |-1||b - a| = 1|b - a| = |b - a|$.

Notice for all $x \in \mathbb{R}$, $|x| \leq |x|$. So $-|x| \leq x \leq |x|$, then we have for $a, b \in \mathbb{R}$, $-|a| \leq a \leq |a|$ and $-|b| \leq b \leq |b|$, adding we get $-(|a| + |b|) \leq a + b \leq |a| + |b|$, thus $|a + b| \leq |a| + |b|$. ■

Corollary. For all $a, b \in \mathbb{R}$, $|a - b| \geq |a| - |b|$ and $||a| - |b|| \leq |a - b|$.

Proof. Now $|a| - |b| = |a + b - b| - |b|$, by the above theorem, $|a| - |b| \leq |a - b| + |b| - |b| = |a - b|$.

Now $||a| - |b|| \leq |a - b|$ implies $-|a - b| \leq |a| - |b| \leq |a - b|$, we need to show that $-|a - b| \leq |a| - |b|$. Then $|a - b| \geq |b| - |a|$ hence $|b - a| \geq |b| - |a|$ which reduces to the second equality. ■

Example 1.4. Show that if $-2 < x < 1$ then $|x^2 - x| < 6$.

Proof. We have that $-2 < x < 2$, so $|x| < 2$. Then $|x^2 + x| \leq |x^2| + |x| = |x||x| + |x| < (2)(2) + 2 = 6$. ■

Theorem 1.1.3. Let $x, y, a \in \mathbb{R}$.

(1) $x < y + \epsilon$ for all $\epsilon > 0$ if and only if $x \leq y$.

(2) $x > y - \epsilon$ for all $\epsilon > 0$ if and only if $x \geq y$.

(3) $|a| < \epsilon$ for all $\epsilon > 0$ if and only if $a = 0$.

Proof. (1) Suppose that $x < y + \epsilon$ for all $\epsilon > 0$, but that $x > y$. Let $\epsilon = \epsilon_0 = x - y$, then $x = y + \epsilon_0$. Then it is not true that $x < y + \epsilon_0$. A contradiction.

Conversely, suppose that $x \leq y$ and that $\epsilon > 0$. Then either $x < y$ or $x = y$. For $x < y$, $x + 0 < y + 0 < y + \epsilon$, hence $x < y + \epsilon$. Similarly, for $x = y$, $x < y + \epsilon$.

(2) This proof is analogous to the previous. Suppose that $x > y - \epsilon$ for $\epsilon > 0$. Then let $\epsilon = \epsilon_0 = y - x$. Then $x = y - \epsilon_0$, which contradicts our assumption that $x > y - \epsilon$.

Conversely, let $x \geq y$ and let $\epsilon > 0$. Then either $x > y$ or $x = y$. For $x > y$, we have $x - 0 > y - 0 > y - \epsilon$. For $x = y$, we see clearly that $x > y - \epsilon$.

(3) Notice first that if $\epsilon > 0$ then $0 > -\epsilon$. Now suppose that for $a \in \mathbb{R}$, $|a| \leq \epsilon$. Then $-\epsilon < a < \epsilon$, thus by the transitivity of $<$ we have that $0 < a$ and $a < 0$, which cannot happen, so it must be that $a = 0$.

Now let $a = 0$, then clearly, by our assumptions, we see that $a < \epsilon$ and $-\epsilon < a$, thus $|a| \leq \epsilon$. ■

Definition. For all $a, b \in \mathbb{R}$, we define the **closed interval** to be the set $[a, b] = \{x : a \leq x \leq b\}$. We define the **open interval** to be the set $(a, b) = \{x : a < x < b\}$. We denote a **half open interval** to be a set of the form $[a, b) = \{x : a \leq x < b\}$ or $(a, b] = \{a < x \leq b\}$.

We denote $(a, \infty) = \{x : a < x\}$, $(-\infty, a) = \{x : x < a\}$ and $(-\infty, \infty) = \{x : x \in \mathbb{R}\}$.

Definition. An interval I is **bounded** if and only if it has the form: $[a, b]$, $[a, b)$, $(a, b]$, or (a, b) . We call a and b the **endpoints**, or **bounds** of the interval. We call all other intervals **unbounded**.

Now if $a = b$, the two bounds coincide and we call the interval **degenerate**, and if $a < b$, it is called **nondegenerate**. Also notice that an interval $(1, 1) = \emptyset$, but $[1, 1] = \{1\}$. So we notice that a degenerate open interval is the empty set, and a degenerate closed interval is a point. Now for bounded intervals, we call $|a - b|$ is called the **length** of the interval, and sometimes denoted $|I|$.

Homework. Do exercises 1, 5, 7, 9, and 10.

1.2 The Well Ordering Principle.

So far we have the set of all natural numbers \mathbb{N} , the set of all integers \mathbb{Z} , the rationals \mathbb{Q} and the reals \mathbb{R} . Now \mathbb{N} is special from these, as it has a “minimal” element. We clarify this below.

Definition. An element $x \in \mathbb{R}$ is a **least element** in a set $E \subseteq \mathbb{R}$ if and only if $x \in E$ and $x \leq \alpha$ for all $\alpha \in E$.

Postulate 3 (The Well Ordering Principle.). *Every nonempty subset of \mathbb{N} has a least element.*

Now this property does not apply to \mathbb{Z} , \mathbb{Q} , and \mathbb{R} ; if one takes the subset \mathbb{Q} , one sees immediately that \mathbb{Q} has no least element. Another thing is that the well ordering principle implies the principle of mathematical induction.

Theorem 1.2.1. *Suppose that for each $n \in \mathbb{N}$, that $A(n)$ is a proposition such that:*

(1) $A(1)$ is true.

(2) For every $k \in \mathbb{N}$ for which $A(k)$ is true, then $A(k+1)$ is also true.

Then $A(n)$ is true for all $n \in \mathbb{N}$.

Proof. Suppose there is some n for which $A(n)$ is false. Then the set $E = \{n \in \mathbb{N} : A(n) \text{ is false}\} \neq \emptyset$. Then by the well ordering principle, E has a least element x . Then $A(x)$ is false. However, since $A(1)$ is true, $x \neq 1$, then $x-1 \in \mathbb{N}$ and $x-1 < x$, since x is the least element, then $A(x-1)$ is true. By the second condition, we get that $A(x)$ is true, contradicting that $x \in E$. Therefore, E must be empty, and $A(n)$ is true for all $n \in \mathbb{N}$. ■

Example 1.5. Show that $\sum_{k=1}^n (3k-1)(3k+2) = 3n^3 + 6n^2 + n$.

Proof. For $k=1$ we see that $(3(1)-1)(3(1)+2) = 10 = 2(1)^3 + 6(1)^2 + 1$.

Now suppose that the proposition is true for $n \geq 1$. Then:

$$\begin{aligned} \sum_{k=1}^{n+1} ((3k-1)(3k+2)) &= \sum_{k=1}^n ((3k-1)(3k+2)) + (3(n+1)-1)(3(n+1)+2) \\ &= 3n^3 + 6n^2 + n + (3(n+1)-1)(3(n+1)+2) \\ &= 3(n+1)^3 + 6(n+1)^2 + (n+1) \end{aligned}$$

■

Remark. If $m, n \in \mathbb{N}$, then $m+n \in \mathbb{N}$ and $mn \in \mathbb{N}$. This also implies that $m+n, mn \in \mathbb{Z}$.

Definition. Let $a, b \in \mathbb{Z}$. We call a **binomial** an expression of the form $(a+b)^n$ for some $n \in \mathbb{N}$.

We wish to study binomials further.

Definition. We define, for $n, k \in \mathbb{N}$ the **binomial coefficient** $\binom{n}{k}$ is defined such that:

$$(1) \binom{0}{0} = 1$$

$$(2) \binom{n}{k} = \frac{n!}{(n-k)!k!}.$$

The binomial coefficient is itself a natural number, and can be visualized via pascal's triangle.

Lemma 1.2.2. $\binom{n}{k} + \binom{n}{k-1} = \binom{n+1}{k}$ For $n, k \in \mathbb{N}$ and $1 \leq k \leq n$.

Proof.

$$\binom{n}{k} + \binom{n}{k-1} = \frac{n!}{(n-k)!k!} + \frac{n!}{(n-k+1)!(k-1)!} \quad (1.2)$$

$$= n! \left(\frac{1}{(n-k)!k!} + \frac{1}{(n-k+1)!(k-1)!} \right) \quad (1.3)$$

$$= n! \left(\frac{1}{(k-1)!} \left(\frac{1}{(n-k)!k} + \frac{1}{(n-k+1)!} \right) \right) \quad (1.4)$$

$$= \frac{n!}{(k-1)!(n-k)!} \left(\frac{1}{k} + \frac{1}{n-k+1} \right) \quad (1.5)$$

$$= \frac{(n+1)!}{(n-k+1)!k!} \quad (1.6)$$

$$= \binom{n+1}{k} \quad (1.7)$$

■

Theorem 1.2.3 (The Binomial Theorem). If $a, b \in \mathbb{R}$, and $n \in \mathbb{N}$, then:

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \quad (1.8)$$

Proof. We use induction in the proof. We see that for $n = 1$, $(a+b)^1 = a+b = \binom{1}{0}a^{1-0} + \binom{1}{1}b^1$. Now suppose that the theorem is true for all $n \geq 1$, that is $(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$; and now consider $n+1$. Then:

$$(a+b)^{n+1} = (a+b)(a+b)^n \quad (1.9)$$

$$= (a+b) \left(\sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \right) \quad (1.10)$$

$$(1.11)$$

by the distributive law:

$$= a \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k + b \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k = \sum_{k=0}^n \binom{n}{k} a^{n-k+1} b^k + \sum_{k=0}^n \binom{n}{k} a^{n-k} b^{k+1}. \quad (1.12)$$

Listing the terms we get $\binom{n}{0}a^{n+1} + \binom{n}{1}a^n b + \binom{n}{2}a^{n-1}b^2 + \dots + \binom{n}{n+1}ab^n + \binom{n+1}{n+1}b^n$; adding like terms, and by lemma 1.2.2 we get: $\binom{n+1}{0}a^{n+1} + \binom{n+1}{1}a^n b + \binom{n+1}{2}a^{n-1}b^2 + \dots + \binom{n+1}{n}ab^n + \binom{n+1}{n+1}b^n = \sum_{k=0}^{n+1} \binom{n+1}{k} a^{n-k+1} b^k$. ■

Homework. Exercises: 1,2,3,5,6 on page 17. These exercises practice the principle of mathematical induction.

1.3 The Axiom of Completeness.

Definition. Let $E \subseteq \mathbb{R}$ be nonempty. The set E is **bounded above** if and only if there is an $M \in \mathbb{R}$ such that $a \leq M$ for all $a \in E$; we call M the **upperbound**. A number s is called a **supremum** or **least upperbound** of E if s is an upper bound, and for all other upper bounds M , $s \leq M$. We denote the least upper bound as $\sup E$.

Example 1.6. If $E = [0, 1]$ prove that $\sup E = 1$.

Proof. By definition of $[0, 1]$, 1 is an upperbound. Now let $M \in \mathbb{R}$ be an upper bound, then $x \leq M$ for all $x \in E$. Now $1 \in E$, so $1 \leq M$, so by definition, we get that $1 = \sup E$. ■

Remark. If a set has one upperbound, then it has infinitely many upper bounds.

Proof. If M_0 is an upperbound, then for any $M > M_0$, M is an upper bound. ■

Remark. If a set has a least upperbound, then that least upperbound is unique.

Proof. Assume that $s_1, s_2 \in \mathbb{R}$ are least upperbounds. Then s_1 is an upperbound and so $s_2 \leq s_1$. Likewise, s_2 is an upperbound, so $s_1 \leq s_2$, hence it must be that $s_1 = s_2$. ■

Theorem 1.3.1 (Approximation Property for Least Upperbounds). *If E has a least upperbound, and $\epsilon > 0$ is a positive number, then there is an element $a \in E$ such that $\sup E - \epsilon < a \leq \sup E$.*

Proof. Suppose not, that is there is some $\epsilon_0 > 0$ for all $a \in E$ for which a does not lie between $\sup E - \epsilon_0$ and $\sup E$. Now $a \leq \sup E$, hence $a \leq \sup E - \epsilon_0$, so $\sup E - \epsilon_0$ is an upperbound of E . So $\sup E \leq \sup E - \epsilon_0$, implying $\epsilon_0 \leq 0$, a contradiction. ■

Now it is not always true that for some set E , that $\sup E \in E$.

Remark. If $E \subseteq \mathbb{N}$ has a least upperbound, then $\sup E \in E$.

Proof. Let $s = \sup E$. By the approximation property, for $\epsilon = 1$, there is an x_0 in E such that $s - 1 < x_0 \leq s$. Now if $x_0 = s \in E$, then we are done.

Otherwise, we have $s - 1 < x_0 < s$. Then applying the approximation property again for $\epsilon = s - x_0$, $s - \epsilon = s - (s - x_0) = x_0$, then there is an $x_1 \in E$ for which $x_0 < x_1 \leq s$. Again if $x_1 = s$ we are done. Now if $x_1 < s$ we get $0 < x_1 - x_0 < s - x_0 < s - (s - 1) = 1$. We also know that $x_1 - x_0 \geq 1$ for different integers $x_1 > x_0$, a contradiction. ■

This brings us to the axiom of completeness.

Postulate 4 (The Axiom of Completeness). *If E is a nonempty subset of \mathbb{R} that is bounded above, then E has a finite least upperbound.*

This does not apply to \mathbb{Q} , for example, take π . We can take the set E to be $E = \{3.14, 3.141, 43.1415, \dots\}$. We see that π is an upperbound, so is 4; however, π is the least upperbound. If we only consider $E \in \mathbb{Q}$, then E has upperbounds, but no least upperbound (as $\pi \notin \mathbb{Q}$). In essence, \mathbb{Q} has a “hole”; infact it has many “holes” that is \mathbb{Q} is not complete. But if $E \in \mathbb{R}$, we see that $\sup E = \pi$. In essence, \mathbb{R} is complete as \mathbb{R} has no “holes”.

Theorem 1.3.2 (The Archimedean Principle). *Given positive real numbers a , and b ; there is an integer n such that $b < na$.*

Proof. We want to build a nonempty set of integers such that it is bounded above with a largest integer. The set $E = \{k \in \mathbb{Z} : ka \leq b\}$ has an upper bound; hence it has a least upperbound k_0 . Then $k_0 + 1 \notin E$, hence $(k_0 + 1)a > b$. ■

Example 1.7. Let $A = \{1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\}$ and $B = \{\frac{1}{2}, \frac{3}{4}, \frac{7}{8}, \dots\}$. Show that $\sup A = \sup B = 1$.

Proof. First, 1 is an upperbound for both A and B . Now for A it is clear that 1 is the least upperbound. Now let M be an upperbound for B and $M < 1$, so $1 - M > 0$ and $\frac{1}{1-M} > 0$. Then by the Archimedean principle, for $\frac{1}{1-M}$ and 1, there is an $n \in \mathbb{Z}$ such that $\frac{1}{1-M} < n$. Now we have that $n < 2^n$; hence it follows that $1 - \frac{1}{2^n} > M$. So M is not an upperbound; a contradiction. Hence $\sup B = 1$. ■

Theorem 1.3.3. *If $a, b \in \mathbb{R}$ such that $a < b$, then there exists a rational $q \in \mathbb{Q}$ such that $a < q < b$.*

Proof. If $b - a > 0$, we have a pair of positive numbers 1 and $b - a$; so by the Archimedean principle, there is an $n \in \mathbb{N}$ such that $n(b - a) > 1$.

Now suppose that $b > 0$. Consider the set $E = \{k \in \mathbb{N} : b \leq \frac{k}{n}\}$. Clearly $E \neq \emptyset$ as $1 \in E$. Then by the well ordering principle, E has a least element k_0 . Now let $m = k_0 - 1$ and let $q = \frac{m}{n}$. Now $m < k_0$, so $m \notin E$; now there are two possible cases: either $m \leq 0$ or $b > \frac{m}{n} = q$, either way we have that $b > q$. And $k_0 \in E$, so $b \leq \frac{k_0}{n}$ so $a = b - (b - a) < \frac{k_0 - 1}{n} = q$.

Now suppose that $b \leq 0$, then $-b > 0$; applying the Archimedean principle for 1 and b , there is a $k \in \mathbb{N}$ such that $-b < (k)(1)$, thus $k + b > 0$. Now if $b = 0$, clearly $1 + 0 > 0$. By the first case, there is a rational $q \in \mathbb{Q}$ such that $a + k < q < b + k$. Subtracting k , we get $a < q - k < b$, which finishes the proof as $q - k \in \mathbb{Q}$. ■

Remark. If $x > 1$ and $x \notin \mathbb{N}$, then there is a natural number $n \in \mathbb{N}$ such that $n < x < n + 1$.

Proof. For the pair $(x, 1)$ by the Archimedean principle, there is a $k \in \mathbb{N}$ such that $E = \{m \in \mathbb{N} : x < m\}$ is nonempty. Then by the well ordering principle, there is a least element m_0 , hence $x < m_0$. Let $n = m_0 - 1$, then $n \notin E$. Now either $n \leq 0$, or $n < x$. As $x \notin \mathbb{N}$, $x \neq n$. Now if $n \leq 0$, then $n < x < m_0 = n + 1$; and if $n < x$, then $n < x < m_0 = n + 1$. ■

Remark. If $n \in \mathbb{N}$ is not a perfect square, then \sqrt{n} is irrational.

Proof. Assume that \sqrt{n} is rational for a nonperfect square $n \in \mathbb{N}$. Then $\sqrt{n} = \frac{p}{q}$ for $p, q \in \mathbb{Z}$. Now since 1 is a perfect square, let $n \geq 2$. Then $\sqrt{n} > 1$. Then there is an $m_0 \in \mathbb{N}$ such that $m_0 < \sqrt{n} < m_0 + 1$. Consider the set $E = \{k \in \mathbb{N} : k\sqrt{n} \in \mathbb{N}\}$. Now $q\sqrt{n} = p \in \mathbb{N}$, so $E \neq \emptyset$. By the well ordering principle, E has a least element n_0 . Then $n_0\sqrt{n} \in \mathbb{N}$ and $n_0m_0 \in \mathbb{N}$. Now $n_0(\sqrt{n} - m_0) = x \in \mathbb{N}$. Now $0 < \sqrt{n} - m_0 < 1$, then $0 < x < n_0$. Now $x \notin E$, on the other hand, $x\sqrt{n} = n_0(\sqrt{n} - m_0)\sqrt{n}$, so $x \in E$; a contradiction. ■

Definition. Let $E \subseteq \mathbb{R}$ be nonempty. The set E is **bounded below** if and only if there is an $m \in \mathbb{R}$ such that $a \geq m$ for all $a \in E$. We call m a **lowerbound** of E . A number t is called an **infimum** or a **greatest lowerbound** if and only if t is a lowerbound of E and $t \geq m$ for all lowerbounds m of E . We denote the greatest lowerbound as $t = \inf E$.

It should be worthwhile to say that the least element of a set, and its greatest lowerbound are not the same.

Definition. A nonempty set $E \subseteq \mathbb{R}$ is said to be **bounded** if it is bounded above and bounded below. That is there exist $m, M \in \mathbb{R}$ such that $m \leq a \leq M$ for all $a \in E$. If $\sup E \in E$ then we write $\sup E = \max E$ and call it the **maximum** of E . If $\inf E \in E$, then we write $\inf E = \min E$ and call it the **minimum** of E .

Definition. The **reflection** of a set $E \subseteq \mathbb{R}$ is defined to be the set $-E = \{-a : a \in E\}$.

Theorem 1.3.4. *Let $E \subseteq \mathbb{R}$ be a nonempty set. Then E has a greatest upperbound if and only if $-E$ has a greatest lowerbound; in which case we have that $\inf(-E) = -\sup E$. Likewise, E has a greatest lowerbound if and only if $-E$ has a least upperbound, in which case $\sup(-E) = -\inf E$.*

Proof. Suppose that E has a least upperbound s , and let $t = -s$. Now we have that $-a \geq -s = t$, for all $a \in E$, so t is a lowerbound of $-E$. Now suppose m is any lowerbound of $-E$. Then $m \leq -a$ for all $a \in E$. Now, $-m \geq a$ for all $a \in E$, so $-m$ is an upperbound of E . Since s is the least upperbound, $s \leq -m$, hence $t = -s \geq m$.

Conversely, suppose that $-E$ has a greatest lowerbound t ; then $t \leq -a$ for all $a \in E$, hence $-t \geq a$ for all $a \in E$, so $-t$ is an upperbound of E . Let M be any upperbound of E , then $M \geq a$ for all $a \in E$. Then $-M \leq -a$ and $-M$ is a lowerbound of $-E$. Then $-M \leq t$, hence $M \geq -t = s$, so s is the least upperbound of E . ■

Homework. Exercises 2,3,4,5(a), and 6 on page 23.

Theorem 1.3.5 (Monotone property). *Let $A \subseteq B$ be nonempty subsets of \mathbb{R} . Then:*

- (1) *If B has a least upper bound, then $\sup A \leq \sup B$.*
- (2) *If B has a greatest lower bound then $\inf A \geq \inf B$*

Proof. 1. We have that $A \subseteq B$, so any upper bound of B is an upper bound of A . Therefore $\sup A$ is an upper bound of A ; By completeness, $\sup A$ exists, and moreover, $\sup A \leq \sup B$.

- 2. Consider $-A \subseteq -B$, by part (1), we have $\sup -A \leq \sup -B$, and we also have that $\sup -A = -\inf A$ and $\sup -B = -\inf B$; thus $-\inf A \leq -\inf B$ therefore, $\inf A \geq \inf B$. ■

To consider $\sup A$ and $\inf A$, we need nonempty and bounded sets of \mathbb{R} . Now we would like to talk about the least upper and greatest lower bounds for any subset of \mathbb{R} ; not just the bounded ones.

Definition. An **extended real number** x is real number such that either $x = \infty$ or $x = -\infty$.

Now if E is not bounded above, then we define the least upper bound of E to be $\sup E = \infty$. If E is not bounded below we define $\inf E = -\infty$. Now for the empty set \emptyset , we define $\sup \emptyset = -\infty$ and $\inf \emptyset = \infty$. It is worth considering why this phenomena occurs with the empty set.

1.4 Functions, Countability, and The Algebra of Sets.

Definition. Given a mapping $f : E \rightarrow \mathbb{R}$, we call f a **real valued function** of a **real variable**. If we are given $f : \mathbb{R}^n \rightarrow \mathbb{R}$ we call the mapping a real valued **multivariable function**. If we are given $f : \mathbb{R}^m \rightarrow \mathbb{R}^m$, we call it a **vector valued** multivariable function.

We are interested in a whole slew of functions; the trigonometric functions: \sin , \cos , \tan , \cot , \sec , and \csc . Other functions of interest are the natural logarithmic function, and exponential function: \log and e^x , as well as arbitrary power functions x^α . We can define the power function to be $x^\alpha = e^{\alpha \log x}$; where $x > 0$ and $\alpha \in \mathbb{R}$.

The derivatives of these functions carry over, and will be examined with more scrutiny later on.

Definition. A mapping $f : X \rightarrow Y$ is called **1-1** if and only if for $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies that $x_1 = x_2$. f is called **onto** if for every $y \in Y$, there exists an $x \in X$ such that $y = f(x)$.

Example 1.8. $f(x) = x^2$ is 1-1 on the interval $[0, \infty)$, but it is not 1-1 on $(0, \infty)$.

Theorem 1.4.1. Let X , and Y and let $f : X \rightarrow Y$ be a mapping. The f is 1-1 from X onto Y if there is a unique mapping $g : Y \rightarrow X$ that satisfies: $f(g(y)) = y$ and $g(f(x)) = x$.

Proof. Suppose that f is 1-1 and onto. Then for each $y \in Y$, choose the unique $x \in X$ such that $f(x) = y$. We define $g(y) = x$. Then it is clear to see that g takes Y onto X , and g is also 1-1. Then $g(y) = g(f(x)) = x$ and $f(x) = f(g(y)) = y$.

Conversely, suppose that there exists a mapping $g : Y \rightarrow X$ satisfying the $g(y) = g(f(x)) = x$ and $f(x) = f(g(y)) = y$. Now let $x_1, x_2 \in X$ and let $f(x_1) = f(x_2)$. Then $x_1 = g(f(x_1)) = g(f(x_2)) = x_2$; so f is 1-1. For any $y \in Y$, let $x = g(y)$, then $f(x) = f(g(y)) = y$; thus f is onto.

Now suppose there is another mapping $h : Y \rightarrow X$ satisfying the conditions that g satisfies. Let $y \in Y$ and choose $x \in X$ such that $f(x) = y$. Then $h(y) = h(f(x)) = x = g(f(x)) = g(y)$; then it must be that $g = h$. ■

Definition. if $f : X \rightarrow Y$ is 1-1 and onto, we say that f has a **inverse mapping** and denote it f^{-1} .

If $f : E \rightarrow \mathbb{R}$ is a real valued function, then the pair $(x, f(x))$ is on the graph of f ; $(f^{-1}(y), y)$ is also on the graph of f . However the pair $(y, f^{-1}(y))$ is on the graph of f^{-1} . The graphs of f and f^{-1} are symmetric with respect to the line $y = x$.

Remark. If f is differentiable on an open interval I and $f'(x) \neq 0$ for all $x \in I$.

This remark is a sufficient, but not a necessary condition. The concept of “differentiability” must also be discussed. We can prove this remark with the “mean value theorem” which will be discussed later.

Example 1.9. Prove that $f(x) = e^x - e^{-x}$ is 1-1 and find f^{-1} .

Proof. By the chain rule, we have $f'(x) = e^x + e^{-x} > 0$ for all $x \in \mathbb{R}$. Now let $y = e^x - e^{-x}$, then $e^x y = e^{2x} - 1$, by the quadratic formula we get:

$$e^x = \frac{y}{2} + \frac{\sqrt{y^2 + 4}}{2}$$

So taking the natural logarithm:

$$y = \frac{x}{2} + \frac{\sqrt{x^2 + 4}}{2} = f^{-1}(x)$$

■

Definition. A set E is **finite** if and only if $e =$ or there is a 1-1 mapping f between $\{1, \dots, n\}$ onto E . E is said to be **countable** if and only if there is a 1-1 mapping from \mathbb{N} onto E . E is said to be **at most countable** if and only if E is either finite or countable. E is said to be **uncountable** if and only if E is neither finite, nor countable.

To show a set to be countable, we need a 1-1 mapping from \mathbb{N} onto E . For example, let $2\mathbb{N} = \{2, 4, 6, 8, \dots\}$, and defining $f(n) = 2n$, then $f : \mathbb{N} \rightarrow 2\mathbb{N}$ is 1-1 and onto. Therefore the set $2\mathbb{N}$ is countable. We know that if a set B is finite, then any proper subset A of B has strictly less number of elements.

Definition. A set B is an **infinite** set if and only if there is a proper subset A so that A and B have the same number of elements.

Now all countable sets are infinite, but not all infinite sets are countable.

Theorem 1.4.2 (Cantor’s Diagonalization Argument). *The open interval $(0, 1)$ is uncountable.*

Proof. Suppose that the interval $(0, 1)$ is countable; then by definition, there exists a mapping $f : \mathbb{N} \rightarrow (0, 1)$ that is 1-1 and onto. Then this mapping exhausts all the elements of $(0, 1)$. We wish to construct an element $x \in (0, 1)$ such that $x \neq f(n)$ for all $n \in \mathbb{N}$. For each $f(n)$ we have:

$$f(n) = 0.\alpha_{n1}\alpha_{n2}\dots$$

This expansion may not be unique, consider $0.1 = 0.0999\dots$. We require that the decimal expansion does not terminate in 9’s. Then $f(n)$ and $0.\alpha_{n1}\alpha_{n2}\dots$ are 1-1:

$$f(1) = 0.\alpha_{11}\alpha_{12}\alpha_{13}\alpha_{14}\dots \quad (1.13)$$

$$f(2) = 0.\alpha_{21}\alpha_{22}\alpha_{23}\alpha_{24}\dots \quad (1.14)$$

$$f(3) = 0.\alpha_{31}\alpha_{32}\alpha_{33}\alpha_{34}\dots \quad (1.15)$$

$$f(4) = 0.\alpha_{41}\alpha_{42}\alpha_{43}\alpha_{44}\dots \quad (1.16)$$

$$\vdots \quad (1.17)$$

So we get an infinite matrix; an all α_{ij} are digits in $\{0, 1, \dots, 9\}$, and none of them terminate in 9's. Now let $x = 0.\beta_1\beta_2\beta_3\dots$ with:

$$\beta_k = \begin{cases} \alpha_{kk} + 1 & \text{if } \alpha_{kk} \leq 5 \\ \alpha_{kk} - 1 & \text{if } \alpha_{kk} > 5 \end{cases}$$

By this construction, we have that $\beta_1 \neq \alpha_{11}$, $\beta_2 \neq \alpha_{22}$, *dots*, $\beta_n \neq \alpha_{nn}$; so $x \neq f(n)$ for all $n \in \mathbb{N}$ and where $x \in (0, 1)$. This is a contradiction of the fact that we assumed f to be onto. Therefore $(0, 1)$ is uncountable. ■

Now we want to study the countability of the sets \mathbb{Z} , \mathbb{Q} , and \mathbb{R} . We study some general properties first:

Lemma 1.4.3. *A nonempty set E is at most countable if and only if there is a mapping $g : \mathbb{N} \rightarrow E$ that is onto.*

Proof. Assume that E is at most countable, then by definition, we are done. If E is finite, by definition there is an $n \in \mathbb{N}$ and a 1-1 mapping f taking $\{1, \dots, n\}$ onto E . Now define:

$$g(j) = \begin{cases} f(j) & \text{if } j \leq n \\ f(1) & \text{if } j > n \end{cases}$$

Then g takes \mathbb{N} onto E .

Conversely, suppose that g takes \mathbb{N} onto E , we need to construct a 1-1 f from \mathbb{N} or a subset of \mathbb{N} onto E . Let $k_1 = 1$ and $E_1 = \{k \in \mathbb{N} : g(k) \neq g(k_1)\}$. Now if $E = \emptyset$, we are done. Else, by the well ordering principle, there is a least element k_2 and define $E_2 = \{k \in \mathbb{N} : g(k) \neq g(k_1), \text{ and } g(k) \neq g(k_2)\}$. Now we have that $k_2 > k_1$ and $k_2 \geq 2$. If $E_2 = \emptyset$ then $E = \{g(k_1), g(k_2)\}$ is finite and we are done. Now if $E_2 \neq \emptyset$, then we continue along both our method until we exhaust all possible set until we reach $E_n = \emptyset$ for some n .

If this process ever terminates, then $E = \{g(k_1), \dots, g(k_n)\}$ and is finite; now if it never terminates, then we have a sequence $k_1 < k_2 < k_3 < \dots$, and so k_{j+1} is the least element of E_j and $k_j \geq j$. We define $f(j) = g(k_j)$ and we show that f is 1-1. For $j < l$ we have $k_j < k_l$, so $k_j \leq k_{l-1}$; by construction, $g(k_l) \in E_{l-1}$, so $g(k_l) \neq g(k_j)$, f is 1-1. Now for any $x \in E$ there is an $n \in \mathbb{N}$ such that $x = g(n)$, and there is an l such that $x = g(n) = g(k_l)$; and so $E = \bigcup_{l=1}^{\infty}$, and so f is onto. Therefore, E is countable. ■

Theorem 1.4.4. *Suppose that A and B are sets and that $A \subseteq B$ and B is at most countable. Then A is at most countable. Likewise if $A \subseteq B$ and A is uncountable, then B is uncountable; in particular, \mathbb{R} is uncountable.*

Proof. If B is at most countable, then there is a function $g : \mathbb{N} \rightarrow B$. If A is empty, we are done. If A is not empty, then choose an element $a_0 \in A$ and define $f : \mathbb{N} \rightarrow A$ such that $f(n) = g(n)$ if $g(n) \in A$, and $f(n) = a_0$ if $g(n) \notin A$. Therefore, f is onto, and A is at most countable.

Assume that B is at most countable, then by the above, since $A \subseteq B$, A is also at most countable. This is a contradiction by assumption, thus, B must also be uncountable.

Now, since the set $(0, 1)$ is uncountable, and $(0, 1) \subseteq \mathbb{R}$, then by the above \mathbb{R} is uncountable. ■

Theorem 1.4.5. *Let A_1, A_2, \dots be atmost countable sets, then:*

(1)

$$A_1 A_2$$

is atmost countable.

(2) *If $E = \bigcup_{j \in \mathbb{N}} A_j$, then E is atmost countable.*

Proof. A_1 and A_2 are atmost countable, then there exist onto functions $\phi_1 : \mathbb{N} \rightarrow A_1$, and $\phi : \mathbb{N} \rightarrow A_2$, (more over we notice that since A_n is also atmost countable, then there exists an onto function $\phi_n : \mathbb{N} \rightarrow A_n$).

Now define $f : \mathbb{N} \times \mathbb{N} \rightarrow A_1 \times A_2$ by $f(n, m) = (\phi_1(m), \phi_2(n))$, clearly f is onto, now if we can define another onto function $g : \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$, then the composition $f \circ g : \mathbb{N} \rightarrow A_1 \times A_2$ is also onto; now $g(1) = (1, 1)$, $g(2) = (2, 1)$, $g(3) = (1, 2)$, $g(4) = (3, 1)$, \dots , by observing the behaviour of g as g moves through \mathbb{N} , we can deduce a formula. Assume that $g(j)$ lies on the n -th line, so $g(1) = (1, n+1)$, $g(2) = (2, n-1)$, $g(3) = (3, n-2)$, and so on. Thus we deduce that for $l \in \mathbb{N}$ that $g(j) = (l, n+1-l)$. Now what is the relation between j and n ? We have $j > 1 + 2 + 3 + \dots + (n-1) = \frac{n(n-1)}{2}$. Now $j \geq 1$, so $j+1 \geq 2$, hence $\frac{j+1}{2} \geq 1$, to $\frac{f(j+1)}{2} \geq j$. Now let $E = \{k \in \mathbb{N} : j \leq \frac{k(k+1)}{2}\}$ which is nonempty, hence E has a least element n . so $j \leq \frac{n(n+1)}{2}$. Thus $l = j - \frac{n(n-1)}{2}$, therefore $j = l + \frac{n(n+1)}{2}$.

Now we have for arbitrary $j \in \mathbb{N}$ that A_j is countable, thus there is an onto function $\phi_j : \mathbb{N} \rightarrow A_j$. Define $f : \mathbb{N} \times \mathbb{N} \rightarrow \bigcup_{j \in \mathbb{N}} A_j$ by $f(m, n) = \phi_n(m)$, then f is also onto. Therefore $\bigcup_{j \in \mathbb{N}} A_j$ is atmost countable. ■

Remark. We have $\mathbb{Z} = \mathbb{N} \cup -\mathbb{N}$ is countable, and $\mathbb{Q} = \bigcup_{n \in \mathbb{N}} \{\frac{p}{n} : p \in \mathbb{Z}\}$ is also countable. Now \mathbb{R} is uncountable, if \mathbb{Q}^* is also countable, then we get $\mathbb{Q}^* \cup \mathbb{Q} = \mathbb{R}$ is countable, contradicting the uncountability of \mathbb{R} , hence \mathbb{Q}^* must also be uncountable.

Definition. A collection of sets \mathcal{E} is said to be indexed by a set A if and only if there is a function F from A onto \mathcal{E} . We call A the **index set** of \mathcal{E} . We may write $\mathcal{E} = \{E_\alpha\}_{\alpha \in A}$.

Definition. Let $\{E_\alpha\}_{\alpha \in A} = \mathcal{E}$ be a collection of sets. Then:

$$(1) \bigcup_{\alpha \in A} E_\alpha = \{x : x \in E_\alpha \text{ for some } \alpha \in A\}$$

$$(2) \bigcap_{\alpha \in A} E_\alpha = \{x : x \in E_\alpha \text{ for all } \alpha \in A\}$$

Theorem 1.4.6 (DeMorgan's Law). *Let \mathcal{X} be a set and let $\{E_\alpha\}_{\alpha \in A}$ be a collection of subsets of \mathcal{X} . For each $E \subseteq \mathcal{X}$ then:*

$$(\bigcap E_\alpha)^C = \bigcup E_\alpha^C$$

where $E^C = \mathcal{X} \setminus E$.

Definition. Let X and Y be sets and let $f : X \rightarrow Y$ be function. Then the **image** of X under f is the set $f(X) = \{f(x) : x \in X\}$. The **inverse image** is the set $f^{-1}(E) = \{x \in X : y = f(x) \text{ for some } y \in E\}$.

For the inverse image, we do not require that the mapping f be 1-1 as we require it for the existence of f^{-1} . Now let X, Y be sets and let $f : X \rightarrow Y$. If $\{E_\alpha\}_{\alpha \in A}$ for some index set A then:

$$f\left(\bigcup_{\alpha \in A} E_\alpha\right) = \bigcup_{\alpha} f(E_\alpha) \quad (1.18)$$

and

$$f\left(\bigcap_{\alpha \in A} E_\alpha\right) = \bigcap_{\alpha} f(E_\alpha) \quad (1.19)$$

Now if B and C are subsets of X , then:

$$f(B) \setminus f(C) \subseteq f(B \setminus C) \quad (1.20)$$

and if $B, C \subseteq Y$ then:

$$f^{-1}(B \setminus C) = f^{-1}(B) \setminus f^{-1}(C) \quad (1.21)$$

and if $\{E_\alpha\}_{\alpha \in A}$ then:

$$f^{-1}\left(\bigcup_{\alpha \in A} E_\alpha\right) = \bigcup_{\alpha} f^{-1}(E_\alpha) \quad (1.22)$$

and

$$f^{-1}\left(\bigcap_{\alpha \in A} E_\alpha\right) = \bigcap_{\alpha} f^{-1}(E_\alpha) \quad (1.23)$$

Finally, if $E \subseteq f(X)$, then $f(f^{-1}(E)) = E$, but if $E \subseteq X$, then $E \subseteq f^{-1}(f(E))$. We accept these properties without proof as they can be demonstrated through elementary set theory.

Now to illustrate an example of these properties, let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $x \rightarrow x^2$, let $E_1 = \{1\}$ and $E_2 = \{-1\}$. Then $E_1 \cap E_2 = \emptyset$, then $f(E_1 \cap E_2) = \emptyset$; but $f(E_1) = \{1\}$ and $f(E_2) = \{1\}$, hence $f(E_1) \cap f(E_2) = \{1\}$. It is interesting to note that if f is 1-1, then all the relations established above are equalities.

Homework. Exercises 6, 9, 10, and 11 on page 33 of the book.

Chapter 2

Sequences in \mathbb{R} .

2.1 Limits of Sequences.

Definition. A **sequence** is a mapping $f : \mathbb{N} \rightarrow X$ for some nonempty set X , whose terms are $x_n = f(n)$. We denote a sequence as $\{x_n\}_n \in \mathbb{N}$ or just simply $\{x_n\}$. We say that a sequence is **real valued** if $X = \mathbb{R}$.

Definition. A sequence of real numbers $\{x_n\}$ is said to **converge** to a real number $a \in \mathbb{R}$ if and only if for every $\epsilon > 0$, there is an $N \in \mathbb{N}$ (in general depending on ϵ) such that for all $n \geq N$, $|x_n - a| < \epsilon$. We say that $\{x_n\}$ (or x_n) **converges** to a , and we write:

$$\lim_{n \rightarrow \infty} x_n = a \quad (2.1)$$

or simply, $x_n \rightarrow a$ as $n \rightarrow \infty$. We call a the **limit** of the sequence. A sequence which does not converge is said to **diverge**.

We can consider x_n as a sequence of “approximations” to a and ϵ as an upperbound for the “error” of those approximations. $N \rightarrow \infty$ as $\epsilon \rightarrow 0$, that is N gets larger as ϵ gets smaller.

Prove that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$

Proof. Let $\epsilon > 0$, by the Archimedean principle, there is an $N \in \mathbb{N}$ such that $\frac{1}{\epsilon} < N$. Then $\frac{1}{N} < \epsilon$. For all $n \geq N$, $\frac{1}{n} < \frac{1}{N} < \epsilon$, hence, $|\frac{1}{n} - 0| < \epsilon$. ■

The sequence $\{(-1)^n\}_{n \in \mathbb{N}}$ does not converge

Proof. Assume that $(-1)^n \rightarrow a$ as $n \rightarrow \infty$. By definition, we have that for any $\epsilon > 0$, there is an $N \in \mathbb{N}$ such that $|(-1)^n - a| < \epsilon$ for $n \geq N$. Choose $\epsilon = \frac{1}{2}$. then for n odd we have $(-1)^n = -1$, and for n even, $(-1)^n = 1$. Then $|(-1)^n - (-1)^{n+1}| = |((-1)^n - a) - ((-1)^{n+1} - a)|$; by the triangle inequality, we have $|(-1)^n - (-1)^{n+1}| \leq |(-1)^n - a| + |(-1)^{n+1} - a| < \frac{1}{2} + \frac{1}{2} = 1$, when $n \geq N$. But $|(-1)^n - (-1)^{n+1}| = 2$, a contradiction. Thus $\{(-1)^n\}$ does not converge. ■

Remark. A sequence can have at most one limit.

Proof. Suppose it has at least two limits, that is $x_n \rightarrow a$ and $x_n \rightarrow b$ (with $a \neq b$) as $n \rightarrow \infty$. For $\epsilon > 0$ there is an N_1 such that for all $n \geq N_1$ $|x_n - a| \leq \epsilon$, and there is an N_2 such that $|x_n - b| \leq \epsilon$. Choose $N = \max(N_1, N_2)$, then for $n \geq N$ we have:

$$|x_n - a| < \epsilon \quad \text{and} \quad |x_n - b| < \epsilon$$

Then $|a - b| = |(a - x_n) - (b - x_n)| \leq |a - x_n| + |b - x_n| < 2\epsilon$ for every ϵ . Now choose $\epsilon = |a - b|/4 > 0$, then $2\epsilon = |a - b|/2$ hence $|a - b| < |a - b|/2$, which is a contradiction. Then x_n converges to at most one limit. ■

Definition. A **subsequence** of a sequence $\{x_n\}_{n \in \mathbb{N}}$ is a sequence of the form $\{x_{n_k}\}_{k \in \mathbb{N}}$ where $n_k \in \mathbb{N}$ and $n_1 < n_2 < \dots$.

Recall that the sequence $\{(-1)^n\}$ does not converge, however, suppose we only select even terms and form $\{(-1)^{2n}\}$, then this latter sequence converges. Like wise the sequence $\{\frac{1}{n}\} \rightarrow 0$ rather slowly, but forming the subsequence $\{\frac{1}{2^n}\}$, the latter converges rather quickly. So the immediate use of subsequences is in the correction of sequences that behave badly (i.e they don't converge), or to make them converge quickly. Now if $n_k = k$, then the subsequence is the original sequence, if $n_k > k$, then the subsequence is a **proper** subsequence.

Remark. If $\{x_n\}$ converges to a and $\{x_{n_k}\}$ is any subsequence of $\{x_n\}$, then $\{x_{n_k}\}$ also converges to a .

Proof. For any $\epsilon > 0$, there is an $N > 0$ such that for all $n \geq N$, $|x_n - a| < \epsilon$. Now if $n_k = k$, we are done, so suppose that $n_k > k$, so for $k = N$, $n_k > N$ then $|x_{n_k} - a| < \epsilon$. ■

Definition. Let $\{x_n\}$ be a sequence of real numbers. Then we say that $\{x_n\}$ is **bounded above** if there is an $M \in \mathbb{R}$ such that $x_n \leq M$ for all $n \in \mathbb{N}$. Similarly, $\{x_n\}$ is **bounded below** if there is an $m \in \mathbb{R}$ such that $m \leq x_n$ for all $n \in \mathbb{N}$. We say $\{x_n\}$ is **bounded** if it is both bounded above and bounded below

Now $\{x_n\}$ is bounded if and only if there is some $C \in \mathbb{R}$ with $C > 0$ such that $|x_n| \leq C$ for all $n \in \mathbb{N}$.

Theorem 2.1.1. *Every convergent sequence is bounded.*

Proof. Let $\{x_n\}$ converge to a . Then for every $\epsilon > 0$, there is an $N > 0$ such that for every $n \geq N$, $|x_n - a| < \epsilon$. Choosing $\epsilon = 1$, we have $|x_n - a| < 1$, so $|x_n| = |x_n - a + a| \leq |x_n - a| + |a| < 1 + |a|$. Now let $C = |x_1| + |x_2| + \dots + |x_{N-1}| + |a| + 1$. Then we see that $|x_n| \leq C$ for all $n \in \mathbb{N}$. Thus $\{x_n\}$ is bounded. ■

The converse of this theorem is not true; take $\{(-1)^n\}$ which is bounded by 1, but $\{(-1)^n\}$ diverges.

Homework. Exercises 1, 4, 5, and 6 on page 38.

2.2 Limit Theorems

There are two problems that are worth discussing, the first is that if we have a sequence, how do we know if it converges? The second problem is determining to where it converges, that is to say, what is the limit; if we cannot find the concrete value, how can we approximate it?

Theorem 2.2.1 (Sandwich Theorem). *Consider real valued sequences $\{x_n\}$, $\{y_n\}$, and $\{w_n\}$. Suppose that $\lim x_n = \lim y_n = a$ and that there is an $N_0 \in \mathbb{N}$ such that $x_n \leq w_n \leq y_n$ for all $n \geq N_0$. Then $\lim_{n \rightarrow \infty} w_n = a$.*

Proof. Let $\epsilon > 0$ and let $\{x_n\}$ and $\{y_n\}$ both converge to a . Then by definition there are $N_1, N_2 \in \mathbb{N}$ such that $|x_n - a| < \epsilon$ and $|y_n - a| < \epsilon$ for $n \geq N_1, n_2$. Now choose $N = \max N_0, N_1, N_2$, if $n \geq N$, we have $-\epsilon < x_n - a < \epsilon$, and we also have $x_n - a < w_n - a < y_n - a$, thus we have that:

$$-\epsilon < x_n - a < w_n - a < y_n - a < \epsilon$$

Thus we have that $|w_n - a| < \epsilon$. ■

Corollary. *If $x_n \rightarrow 0$ as $n \rightarrow \infty$, and $\{y_n\}$ is bounded, then $x_n y_n \rightarrow 0$ as $n \rightarrow \infty$.*

Proof. We have that $\{y_n\}$ is bounded, hence, there is $M > 0$ such that $|y_n| < M$ for all $n \in \mathbb{N}$. And since $\{x_n\}$ converges to 0 we have that for any ϵ there is an $N \in \mathbb{N}$ such that for $n \geq N$, $|x_n - 0| < \frac{\epsilon}{M}$. For $|x_n y_n - 0| = |x_n y_n| < M|x_n| < M \frac{\epsilon}{M} = \epsilon$. Therefore, $x_n y_n \rightarrow 0$ as $n \rightarrow \infty$. ■

Example 2.1. Find $\lim_{n \rightarrow \infty} 2^{-n} \cos(n^3 - n^2 + n - 13)$.

solution. We have that \cos is bounded by 1 for any value it takes, and we also know that $2^n > n$ hence $0 < 2^{-n} < \frac{1}{n}$. We know that $\frac{1}{n} \rightarrow 0$ as $n \rightarrow \infty$, thus by the sandwich theorem, $2^{-n} \rightarrow 0$. Then by the corollary, since \cos is bounded, we get that $\lim_{n \rightarrow \infty} 2^{-n} \cos(n^3 - n^2 + n - 13) = 0$.

Theorem 2.2.2. *Let $E \subseteq \mathbb{R}$. If E has a finite least upper bound (respectively a finite greatest lower bound), then there is a sequence $\{x_n\} \subseteq E$ such that $x_n \rightarrow \sup E$ as $n \rightarrow \infty$.*

Proof. If the least upper bound of E is finite, then for every $\epsilon > 0$, there is an $x \in E$ such that $\sup E - \epsilon < x \leq \sup E$. For all $n \geq N$, choose $\epsilon = \frac{1}{n}$. Then there is one element $x_n \in E$ such that $\sup E - \frac{1}{n} < x_n \leq \sup E$. Now construct sequences $\{w_n\}$, and $\{y_n\}$ such that $w_n = \sup E - \frac{1}{n}$ and $y_n = \sup E$. Then $\lim w_n = \lim y_n = \sup E$ and $w_n < x_n \leq y_n$; thus by the sandwich theorem, $\lim x_n = \sup E$. Analogously, we get the same result for $\inf E$. ■

Theorem 2.2.3. *Suppose that $\{x_n\}$ and $\{y_n\}$ are real valued sequence, and let $\alpha \in \mathbb{R}$. If $\{x_n\}$ and $\{y_n\}$ converge, and suppose that $n \rightarrow \infty$. Then:*

$$(1) \lim x_n + y_n = \lim x_n + \lim y_n.$$

$$(2) \lim \alpha x_n = \alpha \lim x_n.$$

$$(3) \lim x_n y_n = \lim x_n \lim y_n.$$

(4) $\lim \frac{x_n}{y_n} = \frac{\lim x_n}{\lim y_n}$, provided that $y_n \neq 0$ does not converge to 0.

Proof. Suppose that $\{x_n\} \rightarrow x$ and $\{y_n\} \rightarrow y$ as $n \rightarrow \infty$, for $x, y \in \mathbb{R}$. Then:

- (1) For any $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for $n \geq N$ we have $|x_n - x| < \frac{\epsilon}{2}$ and $|y_n - y| < \frac{\epsilon}{2}$. Then $|(x_n - x) + (y_n - y)| = |(x_n + y_n) - (x + y)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$.
- (2) For $\epsilon > 0$, let $N \in \mathbb{N}$ such that for $n \geq N$, $|x_n - x| < \frac{\epsilon}{|\alpha|}$. Then multiplying the inequality by $|\alpha|$, we get $|\alpha||x_n - x| = |\alpha x_n - \alpha x| < \epsilon$.
- (3) Notice that $y_n - y \rightarrow 0$, and x_n is bounded (by theorem 2.1.1), so $x_n(y_n - y) \rightarrow 0$. Similarly, $y_n(x_n - x) \rightarrow 0$. Also notice that $|x_n y_n - xy| = |x_n(y_n - y) + y_n(x_n - x)| \leq |x_n(y_n - y)| + |y_n(x_n - x)|$.

Now for $\epsilon > 0$, there is some $N \in \mathbb{N}$ such that for $n \geq N$, $|x_n - y| < \frac{\epsilon}{2}$ and $|y_n - y| < \frac{\epsilon}{2}$. Then $|x_n(y_n - y)| + |y_n(x_n - x)| < \epsilon$. Therefore, $|x_n y_n - xy| < \epsilon$.

- (4) Provided that $y_n \neq 0$ does not converge to 0, notice that:

$$\left| \frac{x_n}{y_n} - \frac{x}{y} \right| = \left| \frac{x_n y - x y_n}{y y_n} \right| = \frac{1}{|y y_n|} |x_n(y - y_n) + y_n(x_n - x)|$$

By the triangle inequality we have:

$$\frac{1}{|y y_n|} (|x_n(y - y_n)| + |y_n(x_n - x)|) = \left| \frac{x_n}{y y_n} (y - y_n) \right| + \left| \frac{1}{y} (x_n - x) \right| \quad (2.2)$$

Now we see that $\frac{1}{y_n}(x_n - x) \rightarrow 0$ we also have that $y - y_n \rightarrow 0$, and x_n converges, so x_n is bounded. Now since $y_n, y \neq 0$, we must show that $\frac{1}{y_n}$ is bounded. We have that $\lim |y_n| = |y|$, then $|y| > 0$, for $\epsilon = \frac{1}{2}|y|$, there is an $N \in \mathbb{N}$ such that for all $n \geq N$

$$||y_n| - |y|| < \frac{1}{2}|y| = \epsilon$$

then we get that $\frac{1}{|y_n|} < \frac{2}{|y|}$; hence $\frac{1}{y_n}$. For $k = 1, \dots, N-1$ let $M = \frac{2}{|y|} + \max \left\{ \frac{1}{|y_1|}, \dots, \frac{1}{|y_{N-1}|} \right\}$, thus $\frac{1}{|y_n|}$ is bounded by M .

Now since $\frac{1}{|y_n|}$ is bounded, so is $\frac{x_n}{|y y_n|}$, thus we get that $\frac{x_n}{y y_n}(y - y_n) \rightarrow 0$, thus we get that

$$\frac{x_n y - x y_n}{y y_n} = \frac{x_n}{y_n} - \frac{x}{y} \rightarrow 0$$

This completes the proof. ■

Example 2.2. Find the limit: $\lim_{n \rightarrow \infty} \frac{n^3 + n^2 - 1}{1 - 3n^3}$.

solution. We divide the numerator and the denominator by the highest power of n to get:

$$\frac{1 + \frac{1}{n} - \frac{1}{n^3}}{\frac{1}{n^3} - 3} \quad (2.3)$$

Then we get that as $n \rightarrow \infty$ then

$$\frac{1 + \frac{1}{n} - \frac{1}{n^3}}{\frac{1}{n^3} - 3} \rightarrow \frac{1 + 0 - 0}{0 - 3} = -\frac{1}{3} \quad (2.4)$$

Hence, $\lim_{n \rightarrow \infty} \frac{n^3 + n^2 - 1}{1 - 3n^3} = -\frac{1}{3}$

A sequence $\{n\}$ diverges, so does the sequence $\{(-1)^n\}$, however, they diverge differently; the sequence $\{n\}$ just gets bigger and bigger as $n \rightarrow \infty$, however, $\{(-1)^n\}$ oscillates as $n \rightarrow \infty$.

Definition. Let $\{x_n\}$ be a real valued sequence. Then

- (1) $\{x_n\}$ is said to **diverge** to $+\infty$ if and only if for every $M \in \mathbb{R}$, there is an $N \in \mathbb{N}$ such that for $n \geq N$, $x_n > M$.
- (2) $\{x_n\}$ is said to **diverge** to $-\infty$ if and only if for each $M \in \mathbb{R}$, there is an $N \in \mathbb{N}$ such that for $n \geq N$, $x_n < M$.

Theorem 2.2.4. Suppose that $\{x_n\}$ and $\{y_n\}$ are real valued sequence such that $x_n \rightarrow +\infty$ as $n \rightarrow \infty$.

- (1) If $\{y_n\}$ is bounded below, then $x_n + y_n \rightarrow +\infty$ as $n \rightarrow \infty$. Similarly if $\{y_n\}$ is bounded above, then $x_n + y_n \rightarrow -\infty$.
- (2) If $\alpha > 0$, then $\alpha x_n \rightarrow +\infty$ as $n \rightarrow \infty$. Similarly, if $\alpha < 0$, then $\alpha x_n \rightarrow -\infty$.
- (3) If $y_n > M_0$ for some $M_0 > 0$ and for all $n \in \mathbb{N}$. Then $x_n y_n \rightarrow +\infty$ as $n \rightarrow \infty$.
- (4) If $\{y_n\}$ is bounded, and $x_n \neq 0$, then $\frac{y_n}{x_n} \rightarrow 0$ as $n \rightarrow \infty$.

We defer the proof.

Corollary. Let $\{x_n\}$ and $\{y_n\}$ be real valued sequences and α, x, y be extended real numbers. If $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$, then $x_n + y_n \rightarrow x + y$ as $n \rightarrow \infty$. and $\alpha x_n \rightarrow \alpha x$ and $x_n y_n \rightarrow xy$; We disclude of course the indeterminate forms for the extended real numbers.

Theorem 2.2.5 (The Comparison Theorem). Suppose that $\{x_n\}$ and $\{y_n\}$ are convergent sequences, if there is an $N_0 \in \mathbb{N}$ such that $x_n \leq y_n$ for $n \geq N_0$, then $\lim x_n \leq \lim y_n$ as $n \rightarrow \infty$. In particular, if $x_n \in [a, b]$ converges to c , then $c \in [a, b]$.

Proof. Suppose that the statement is false, then $\lim x_n > \lim y_n$ as $n \rightarrow \infty$. Let $\lim x_n = x$ and let $\lim y_n = y$, and let $\epsilon = \frac{x-y}{4}$, then there is an $N \in \mathbb{N}$ such that for $n \geq N$, $|x_n - x| < \epsilon$, and $|y_n - y| < \epsilon$. Now we have that $x_n > x - \epsilon > y + \epsilon > y_n$, a contradiction. Therefore $\lim x_n \leq \lim y_n$.

Now in particular, we have that if $x_n \in [a, b]$, then $a \leq x_n \leq b$, letting $\lim_{n \rightarrow \infty} x_n = c$, we get from above that $a \leq c \leq b$, hence $c \in [a, b]$. ■

Homework. Exercises 3, 5, and 6 on page 44.

2.3 The Bolzano-Weierstrass Theorem.

The sequence $\{(-1)^n\} = (-1, 1, -1, 1, \dots)$, has the properties that it is a bounded divergent sequence that contains convergent subsequences. This turns out to be a special case of the “Bolzano-Weierstrass theorem”. The rest of this section is devoted to stating and proving the theorem.

Definition. Let $\{x_n\}$ be a real valued sequence. Then:

- (1) $\{x_n\}$ is **increasing** if and only if $x_i \leq x_{i+1}$; and $\{x_n\}$ is **strictly increasing** if $x_i < x_{i+1}$ for all $i \in \mathbb{N}$
- (2) $\{x_n\}$ is **decreasing** if and only if $x_{i+1} \leq x_i$; and $\{x_n\}$ is **strictly decreasing** if and only if $x_{i+1} < x_i$ for all $ui \in \mathbb{N}$.
- (3) $\{x_n\}$ is **monotone** if and only if it is either increasing or decreasing.

If $\{x_n\}$ is increasing, and converges to a , we may write $x_n \uparrow a$, if $\{x_n\}$ is decreasing and convergent to a , we may write $x_n \downarrow a$.

Claim. (1) If a sequence is strictly increasing, then it is increasing.

(2) If a sequence is strictly decreasing, then it is decreasing.

(3) If $\{x_n\}$ is a real valued sequence, then $\{x_n\}$ is increasing if and only if $\{-x_n\}$ is decreasing. Similarly $\{x_n\}$ is decreasing if and only if $\{-x_n\}$ is increasing.

We know that all convergent sequences are bounded, however, the converse is not true. However, what if we only consider monotone sequences?

Theorem 2.3.1 (The Monotone Convergence Theorem). *If $\{x_n\}$ is increasing and bounded above, or decreasing and bounded below, then $\{x_n\}$ converges to a finite limit.*

Proof. We show that an increasing sequence bounded above will converge to its least upperbound. Suppose $\{x_n\}$ is increasing and bounded above. By the axiom of completeness, $\{x_n\}$ has a finite least upperbound a . Now for any $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that $a - \epsilon < x_N \leq a$. Then for any $n \geq N$, $\{x_n\}$ is increasing, hence $x_n \geq x_N$, so $a - \epsilon < x_N \leq x_n \leq a$. Hence $|x_n - a| < \epsilon$. Therefore, $x_n \rightarrow a$. By a similar argument, we get that if $\{x_n\}$ is and bounded below, then $x_n \inf \{x_n\}$.

We can also by direct application of the first part, note that by the axiom of completeness, $\{-x_n\} \rightarrow -b$, and by the limit laws, we get that $x_n \rightarrow b$, hence we get that $\{-x_n\}$ is increasing and bounded above, hence $-b = \sup \{-x_n\}$, therefore $b = \inf \{x_n\}$. ■

Example 2.3. If $|a| < 1$, then $a^n \rightarrow 0$ as $n \rightarrow \infty$.

solution. If $\{x_n\}$ is a sequence with the limit $\lim |x_n| = 0$, then $\lim x_n = 0$. Then it suffices to show that $\lim |a|^n = 0$. We have that $|a|^n < |a|^{n-1}$, so $\{|a^n|\}$ is decreasing, and we see that 0 is a lowerbound, hence by theorem 2.3.1, $\lim |a^n| = L$, we must show that $L = 0$. Suppose it is not 0, then $|a|^{n+1} = |a||a|^n$, so we consider the sequences $\{|a|^{n+1}\} = |a|\{|a|^n\}$, then $\{|a|^{n+1}\}$ is a subsequence of $\{|a|^n\}$. Hence we get $\lim |a||a|^n = |a| \lim |a|^n = |a|L$, hence we get that $L = |a|L$, implying that $|a| = 1$. A contradiction, so $L = 0$.

Example 2.4. If $a > 0$, then $\lim a^{\frac{1}{n}} = 1$

solution. We consider 3 cases. Suppose first that $a = 1$, then $a^{\frac{1}{n}} = 1$ for all n , and hence we have the limit of a constant sequence. Now assume that $a > 1$. Then it suffices to show that $\{a^{\frac{1}{n}}\}$ is decreasing. For any $n \in \mathbb{N}$, $a^{n+1} > a^n > 1$, taking the n th root, we have $(a^n)^{\frac{1}{n}} > a$, hence we have $(a^{\frac{1}{n}})^n > a$, so we get that $a^{\frac{1}{n}} > a^{\frac{1}{n+1}}$ which shows that $\{a^{\frac{1}{n}}\}$ is decreasing. Therefore $a^{\frac{1}{n}} \geq 1$. Therefore, by theorem 2.3.1, the limit exists and $\lim a^{\frac{1}{n}} = L$.

Consider then $a^{\frac{1}{2n}}$ and produce the appropriate subsequence. Notice then that $a^{\frac{1}{n}} = (a^{\frac{1}{n}})^{\frac{1}{2}}$ and the limit is L . We also see that the limit of $(a^{\frac{1}{n}})^{\frac{1}{2}} = L^{\frac{1}{2}}$. So we have that $L = L^{\frac{1}{2}}$. Therefore $L^2 - L = 0$, and so $L = 0$ or $L = 1$. However, since $a^{\frac{1}{n}} \geq 1$, $L \neq 0$, so $L = 1$.

Now suppose that $0 < a < 1$, then we know that $\frac{1}{a} > 1$. Thus by the previous case, we get that $\lim \frac{1}{a^{\frac{1}{n}}} = 1$. Using some algebra, we get that

$$\lim a^{\frac{1}{n}} = \frac{1}{\frac{1}{a^{\frac{1}{n}}}} = \frac{1}{1} = 1$$

Definition. A sequence of sets $\{I_n\}_{n \in \mathbb{N}}$ is said to be **nested** if and only if $I_{i+1} \subseteq I_i$ for all $i \in \mathbb{N}$.

Theorem 2.3.2. If $\{I_n\}$ is a nested sequence of nonempty, closed, and bounded intervals then:

$$E = \bigcap_{n \in \mathbb{N}} I_n$$

contains at least one point. Moreover, if the length of these intervals satisfy $|I_n| \rightarrow 0$ as $n \rightarrow \infty$, then E contains exactly one point.

It is interesting to observe some things before we begin the proof. First note that if $\{I_n\}$ is a sequence of nested sets, and $E = \bigcap I_n$ for all n , then $E = [a, b]$ for some $a, b \in \mathbb{R}$. Moreover, we see that $E \subseteq \dots I_n \subseteq \dots \subseteq I_2 \subseteq I_1$, hence E is a nonempty, closed and bounded interval, and $E \in \{I_n\}$. What this illustrates, is that we can make E as arbitrarily small an interval we need it to be, and it will always contain at least one element. Moreover, if we construct $\{I_n\}$ right, then we can make E have exactly one element.

Proof. Let $I_n = [a_n, b_n]$, and that $\{I_n\}$ is nested. Then $a_n \leq a_{n+1}$ and $b_{n+1} \leq b_n$. Thus $\{a_n\}$ is increasing, and $\{b_n\}$ is decreasing, with an upperbound b_1 and a lowerbound a_1 , respectively. Then we have a and b with $a_n \rightarrow a$ and $b_n \rightarrow b$ as $n \rightarrow \infty$. Now since $a_n < b_n$ for all n , we get $a \leq b$. Therefore, $E = [a, b]$ contains at least one point. Moreover, if the length of $|I_n| \rightarrow 0$ as $n \rightarrow \infty$, then $b_n - a_n \rightarrow 0$. So $b - a = 0$. Therefore $b = a$ and we get that E contains exactly one point. ■

Remark. The nested interval property does not hold if “closure” is omitted. The interval $I_n = (0, \frac{1}{n})$ is nested, nonempty, and bounded, however $\bigcap I_n = \emptyset$.

Remark. The nested interval property may not hold if the interval is not bounded. Consider the interval $I_n = [n, \infty)$, which is nested and closed, but not bounded. Then $\bigcap I_n = \emptyset$.

Theorem 2.3.3 (The Bolzano-Weierstrass Theorem). *Every bounded sequence of real numbers has a convergent subsequence.*

Proof. We begin with a general observation. Let $\{x_n\}$ be a sequence, and let E be a set. Then $E = A \cup B$ for sets A and B , and E contains x_n for infinitely many values of n ; then at least one of the sets A or B contains $\{x_n\}$ for infinitely many values of n .

Now let $\{x_n\}$ be a bounded sequence, choose $a, b \in \mathbb{R}$ such that all $x_n \in [a, b]$. Set $I_0 = [a, b]$; now divide I_0 into two subintervals $[a, \frac{1+b}{2}]$ and $[\frac{1+b}{2}, b]$. Now I_0 contains x_n for infinitely many values of n , and $I_0 = [a, \frac{1+b}{2}] \cup [\frac{1+b}{2}, b]$. Then at least one of the aforementioned intervals contains x_n for infinitely many values of n . Call that set I_1 . and we know that $|I_1| = \frac{1}{2}|I_0| = \frac{b-a}{2}$. Now divide I_1 equally into two sub intervals I_1' and I_1'' , then as before, at least one of these intervals contains x_n for infinitely many values of n . Call that interval I_2 , and so $|I_2| = \frac{1}{2}|I_1| = \frac{b-a}{4}$. Repeating the process, we get a nested sequence of intervals $\{I_n\}$ where $|I_n| = \frac{b-a}{2^n}$ and I_n is nonempty, closed, and bounded; so the intersection $E = \bigcap I_n = \{x\}$ by theorem 2.3.2. Now we have that I_n is nonempty, so we will have $x_{n_k} \in I_n$ because I_n contains x_n for infinitely many values of n . Choose $n_k < n_{k+1}$. So $\{x_{n_k}\}$ is a subsequence of $\{x_n\}$ with $x_{n_k} \in I_n$.

Now observe that $x_{n_k}, x \in I_n$, so $0 \leq |x_{n_k} - x| \leq |I_n| \leq \frac{b-a}{2^k}$ for $n_k \geq k$. Hence, by definition of the limit, when $k \rightarrow \infty$, $\lim x_{n_k} = x$, choosing $\epsilon > \frac{b-a}{2^k}$. ■

Homework. Problems 1, 2, 4, 5, and 6 on page 48.

2.4 Cauchy Sequences.

Whenever we say a sequence converges, we would like to find the limit. However, the limit may lie outside of the sequence. It is here that building up the notion of a “Cauchy sequence” which establishes a property that is independent from the limit, but that guarantees that a sequence will be convergent to said limit (which of course we may or may not know).

Definition. A realvalued sequence $\{x_n\}$ is said to be a **Cauchy sequence** if and only if for every $\epsilon > 0$, there is an $N \in \mathbb{N}$ such that $n, m \geq N$ implies that $|x_n - x_m| < \epsilon$.

Remark. If $\{x_n\}$ is a convergent sequence, then $\{x_n\}$ is Cauchy.

Proof. Suppose that $\{x_n\} \rightarrow x$ as $n \rightarrow \infty$. For any $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that for all $n \geq N$, $|x_n - x| < \frac{\epsilon}{2}$. Now then, for any $m \geq N$ we get $|x_m - x| < \frac{\epsilon}{2}$. Then by the triangle inequality: $|x_n - x_m| = |(x_n - x) - (x_m - x)| \leq |x_n - x| + |x_m - x| < \epsilon$. we see that $\{x_n\}$ is a Cauchy sequence. ■

Theorem 2.4.1 (Cauchy’s Theorem). *Let $\{x_n\}$ be a realvalued sequence. Then $\{x_n\}$ is a Cauchy sequence if and only if $\{x_n\}$ converges to some point in \mathbb{R} .*

Proof. We need only to show that Cauchy sequences converge (the remark before hand proves the converse implication). Let $\{x_n\}$ be a Cauchy sequence, then $\{x_n\}$ is bounded. Let $\epsilon = 1$, then there is some $K \in \mathbb{N}$ such that for all $m, n \geq K$, $|x_n - x_m| < 1$. We have that $|x_1|, |x_2|, \dots, |x_K|$, and $|x_{K+1} - x_K| < 1$. Now $|x_{K+1}| = |x_{K+1} - x_K + x_K| \leq |x_{K+1} - x_K| + |x_K| < 1 + |x_K|$. Hence choose $M = \max\{|x_1|, |x_2|, \dots, |x_K|, 1 + |x_K|\}$. Then $|x_n| \leq M$ for all $n \in \mathbb{N}$. Hence every Cauchy sequence is bounded, hence by the Bolzano-Weierstrass theorem, $\{x_n\}$ has a convergent subsequence $\{x_{n_k}\}$. So there is an $x \in \mathbb{R}$, such that for all $\epsilon > 0$, there exists $K \in \mathbb{N}$ such that for all $k \geq K$, $|x_{n_k} - x| < \frac{\epsilon}{2}$.

$\{x_n\}$ is a Cauchy sequence, hence for this same $\epsilon > 0$, there is an $N_1 \in \mathbb{N}$ such that for $m, n \geq N_1$, $|x_n - x_m| < \frac{\epsilon}{2}$. Choosing $K \geq N_1$, we have that $k \geq K$, hence $n_k \geq k \geq K \geq N_1$. For all $n \geq K$, For all $n \geq K$, $|x_n - x| = |x_n - x_{n_k} + x_{n_k} - x| \leq |x_n - x_{n_k}| + |x_{n_k} - x| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$. Thus $x_n \rightarrow x$ as $n \rightarrow \infty$, hence all Cauchy sequences are convergent. ■

Example 2.5. Prove that any real valued sequence $\{x_n\}$ satisfying $|x_n - x_{n+1}| < \frac{1}{2^n}$ is convergent.

solution. To prove that it is convergent, first prove that it is a Cauchy sequence. For $n < m$, we have $|x_n - x_m| = |x_n - x_{n+1} + x_{n+1} - x_{n+2} + x_{n+2} + \cdots + x_{m-1} - x_m| \leq |x_n - x_{n+1}| + |x_{n+1} - x_{n+2}| + \cdots + |x_{m-1} - x_m| < \frac{1}{2^n} + \frac{1}{2^{n+1}} + \cdots + \frac{1}{2^{m-1}} < \frac{1}{2^{n-1}}$. Then for $\epsilon > 0$, there exists $N \in \mathbb{N}$ with $\frac{1}{2^{N-1}} < \epsilon$. Hence, for $m, n \geq N$ we have that $|x_n - x_m| < \frac{1}{2^{n-1}} < \frac{1}{2^{N-1}} < \epsilon$. Therefore, $\{x_n\}$ is a Cauchy sequence, and hence converges to a point in \mathbb{R} .

Remark. If $|x_{n+1} - x_n| \rightarrow 0$, then it is not sufficient to deduce that it is a Cauchy sequence. For example, consider $x_n = \log n$. Then $\{x_n\}$ is unbounded and divergent. However, $\log n + 1 - n = \frac{\log n + 1}{\log n} = \log 1 = 0$.

Homework. Exercises 1, 2, 3, and 6 on page 51.

2.5 Limits, Least Upperbounds and Greatest Lowerbounds.

We know that a sequence converges to a point if and only if it is a Cauchy sequence. We would like to talk about related concepts of sequences in general. Consider the sequence x_1, x_2, x_3, \dots . And construct the following sequence: $S_1 = \sup\{x_1, x_2, x_3, \dots\}$, $T_1 = \inf\{x_1, x_2, x_3, \dots\}$. Continuing, constructs $S_2 = \sup\{x_2, x_3, x_4, \dots\}$ and $T_2 = \inf\{x_2, x_3, x_4, \dots\}$. Continuing along this line, we get the terms $S_n = \sup\{x_n, x_{n+1}, \dots\}$ and $T_n = \inf\{x_n, x_{n+1}, \dots\}$. We see that $\{S_n\}$ is a decreasing sequence of extended real numbers, now it could turn out that $\{S_n\}$ is a complete sequence of ∞ ; if at a certain point, the term $s_n \neq \infty$, then the rest of the sequence is not ∞ .

Similarly, $\{T_n\}$ is an increasing sequence of extended real numbers. It could happen that $\{T_n\}$ is $-\infty$ everywhere, however, if at a certain point $t_n \neq -\infty$, then the whole of the sequence is not $-\infty$.

Definition. Let $\{x_n\}$ be a real valued sequence. Then the **limit supremum** is the extended real number, $\limsup x_n = \lim(\sup_{k \geq n}\{x_k\})$ as $n \rightarrow \infty$. And the **limit infimum** is the extended real number $\liminf x_n = \lim(\inf_{k \geq n}\{x_k\})$, as $n \rightarrow \infty$.

We discuss some cases. If $s_n = \infty$ for all $n \in \mathbb{N}$, then $\limsup x_n = \infty$. Now if $s_n \neq \infty$, and if $\{S_n\}$ is bounded below, then $\limsup x_n$ is a real number. Now if $S_n \neq \infty$ for all $n \in \mathbb{N}$, and $\{S_n\}$ is not bounded below, then $\limsup x_n = -\infty$.

Similarly, if $t_n = -\infty$ for all $n \in \mathbb{N}$, then $\liminf x_n = -\infty$. Now if $t_n \neq -\infty$ for all $n \in \mathbb{N}$, and if $\{T_n\}$ is bounded above, then $\liminf x_n$ is a real number. Now if $t_n \neq -\infty$ for all $n \in \mathbb{N}$ and $\{T_n\}$ is not bounded above, then $\liminf x_n = \infty$.

Theorem 2.5.1. *Let $\{x_n\}$ be a real valued sequence, and let $s = \limsup x_n$ and let $t = \liminf x_n$. Then there are subsequences $\{x_{n_k}\}$ and $\{x_{l_j}\}$ such that $x_{n_k} \rightarrow s$ as $k \rightarrow \infty$, and $x_{l_j} \rightarrow t$ as $j \rightarrow \infty$.*

Proof. We prove it only for the \liminf . We know that $t_n = \inf x_k$, and that $t_n \rightarrow t$ as $n \rightarrow \infty$ (for t_n increasing). Now if $t = -\infty$, then $t_n = -\infty$ for all $n \in \mathbb{N}$. Then $t_1 = -\infty$, so there is an n_1 such that $x_{n_1} < -1$. Now $t_{n_1+1} = -\infty$, then there is an n_2 such that $x_{n_2} < -2$ and $n_2 > n_1$. Continuing along this process, we have $n_1 < n_2 < n_3 < \dots$ and $x_{n_k} < -k$. Then we have that $\lim x_{n_k} = -\infty$.

Now if $t = \infty$, then $t_n \neq -\infty$ for all $n \in \mathbb{N}$, and $\{T_n\}$ is not bounded above. Now without loss of generality, assume that $t_n \neq -\infty$ for all $n \in \mathbb{N}$. For $M = 1$, there is an n_1 such that $t_{n_1} > 1$. So $x_{n_1} \geq t_{n_1} > 1$. Now for $M = 2$, there is an n_2 such that $n_2 \geq n_1$, and $t_{n_2} > 2$. Continuing this process, we get that $n_1 \leq n_2 \leq \dots$ and $x_{n_k} > k$.

Suppose that $-\infty < t < \infty$, then t is not $t_n \neq -\infty$ for all $n \in \mathbb{N}$, and $\lim t_n = t$. So $t_n \rightarrow t$ (where t_n is increasing). Then for every $\epsilon > 0$, there is an $N \in \mathbb{N}$ such that for $n \geq N$, we have $|t - t_n| = t - t_n < \frac{\epsilon}{2}$. Now if $t_n = \inf\{x_k\}$, for the same ϵ , there is an $n_k \geq n$ such that $t_n \leq x_{n_k} < t_n + \frac{\epsilon}{2}$. Then by the triangle inequality, we have $|t - x_{n_k}| = |t - t_n| + |t_n - x_{n_k}| < \epsilon$. Now consider $\epsilon = \frac{1}{n}$, if $n = 1$, then there is an $N_1 \in \mathbb{N}$, such that for $n \geq N_1$, $|t - t_n| = t - t_n < \frac{1}{2}$. Then for $t_{N_1} = \inf x_k$, there is an $n_1 \geq N_1$, with $t_{N_1} \leq x_{n_1} < t_{N_1} + \frac{1}{2}$. Combining these two relations we have $|t - x_{n_1}| < \frac{1}{2}$. Now for $n = 2$, there is an $N_2 \in \mathbb{N}$ with $N_2 > n_1$ such that for $n \geq N_2$, we have $|t - x_{n_2}| < \frac{1}{4}$. Continuing this process, we find a sequence $\{n_k\}$ such that $n_1 < n_2 < \dots$, and $|t - x_{n_k}| < \frac{1}{k}$. Thus we have that $\lim_{k \rightarrow \infty} x_{n_k} = t$. ■

Theorem 2.5.2. *Let $\{x_n\}$ be a real valued sequence and let x be an extended real number. Then $\{x_n\} \rightarrow x$ as $n \rightarrow \infty$ if and only if:*

$$\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n = x$$

Proof. Suppose that $x_n \rightarrow x$ as $n \rightarrow \infty$. Then all subsequences $\{x_{n_k}\}$ will converge to x as $k \rightarrow \infty$. Then by theorem 2.5.1, we know that $\limsup x_n = \liminf x_n = x$. Conversely, suppose that $\limsup x_n = \liminf x_n = x$, then we have three cases. If $x = \infty$, then $s_n = \infty$, and $\{T_n\}$ is not bounded above. Thus for all $M \in \mathbb{R}$, there is an $N \in \mathbb{N}$ such that for $n \geq N$, $t_n = M$, so $x_n \geq t_n \geq M$. Thus $\lim x_n = \infty$.

Suppose now that $x = -\infty$, then $t_n = -\infty$ and $\{S_n\}$ is not bounded below. So for all $M \in \mathbb{R}$, there is an $N \in \mathbb{N}$ such that for $n \geq N$, $s_n \leq M$. So $\lim x_n = -\infty$. Now suppose that $-\infty < x < \infty$. Then for all $\epsilon > 0$, there is an $N \in \mathbb{N}$, such that $0 \leq s_n - x < \frac{\epsilon}{2}$ and $0 \leq x - t_n < \frac{\epsilon}{2}$. Then for $m, n \geq N$, assume that $|x_n - x_m| = x_m - x_n \leq s_m - t_n \leq \epsilon$. Hence, $\{x_n\}$ is a Cauchy sequence, so $x_n \rightarrow x'$ in \mathbb{R} . So all its subsequences converge to x' . But we know that $\limsup x_n = \liminf x_n = x$, thus $x' = x$ and we are done. ■

Theorem 2.5.3. *Let $\{x_n\}$ be a real valued sequence. Then $\limsup x_n$ is the largest value to which some subsequence of $\{x_n\}$ converges, as $n \rightarrow \infty$. Similarly, $\liminf x_n$ is the smallest value for which some subsequence of $\{x_n\}$ converges. In particular, if $x_{n_k} \rightarrow x$ as $k \rightarrow \infty$, then:*

$$\liminf_{n \rightarrow \infty} x_n \leq x \leq \limsup_{n \rightarrow \infty} x_n.$$

Proof. Assume that a subsequence $\{x_{n_k}\}$ converges to x as $k \rightarrow \infty$. Then fix $N \in \mathbb{N}$, and choose K so that for $k \geq K$, $n_k \geq N$. Now under this condition, $\inf x_j \leq x_{n_k} \leq \sup x_j$ for $j \geq N$ for all $k \geq K$. Now we let $k \rightarrow \infty$, then $x_{n_k} \rightarrow x$, then by the comparison of the limit, we have $\inf x_j \leq x \leq \sup x_j$ for $j \geq N$, again, by the comparison of limits, letting $N \rightarrow \infty$, we have that $\inf x_j$ is increasing, and $\sup x_j$ is decreasing, so we get that $\liminf x_n \leq x \leq \limsup x_n$. ■

Corollary. *If $\{x_n\}$ is any real valued sequence, then :*

$$\liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n.$$

Proof. The proof follows from theorem 2.5.3. ■

Remark. A real valued sequence $\{x_n\}$ is bounded above if and only if $\limsup x_n < \infty$ as $n \rightarrow \infty$; and $\{x_n\}$ is bounded below if and only if $\liminf x_n > -\infty$.

Proof. If $\{x_n\}$ is bounded above, there is an M such that $x_n \leq M$ for all n . Then $\sup x_j \leq M$ for $j \geq N$. Then we get that $\limsup x_n \leq M$ as $n \rightarrow \infty$. Conversely, if $\limsup x_n = M$ then for every $\epsilon > 0$, there exists an $N \in \mathbb{N}$ such that:

$$\left| \sup_{j \geq N} x_j - M \right| < \epsilon$$

Thus $M - \epsilon < \sup x_j < M + \epsilon$. Then choose $M' = |x_1| + |x_2| + \cdots + |x_{N-1}| + M + \epsilon$. Then $x_j \leq M'$ for all $j \in \mathbb{N}$. The proof is similar for $\liminf x_n$. ■

Theorem 2.5.4. *If $x_n \leq y_n$ for sufficiently large N , then $\limsup x_n \leq \limsup y_n$, and $\liminf x_n \leq \liminf y_n$ as $n \rightarrow \infty$.*

Proof. We have $\sup x_j \leq \sup y_j$ for $j \leq n$ for n sufficiently large. Then by the comparison theorem of the limit, we will have $x_n \leq \limsup y_n$ as $n \rightarrow \infty$. The same holds for \liminf . ■

Homework. Problems 2, 3, 4, 5, and 6 on page 35.

Chapter 3

Continuity on \mathbb{R} .

3.1 Two Sided Limits.

Definition. Let $f : A \rightarrow B$ be a function. We call f a **real valued function** if $B \subseteq \mathbb{R}$.

It need not be that $A \subseteq \mathbb{R}$ for f to be a real valued function, as long as B is a subset of \mathbb{R} , then the definition is satisfied. However, it is also common to take $A \subseteq \mathbb{R}$ (either an interval, or one of the sets $\mathbb{N}, \mathbb{Z}, \mathbb{Q}$ or \mathbb{Q}^*). \mathbb{B} necessarily, is any of those sets.

Definition. Let $a \in \mathbb{R}$ and let I be an open interval containing a , and let f be a real valued function defined everywhere on I except possibly at a . Then $f(x)$ is said to **converge** to L as $x \rightarrow a$ if and only if for every $\epsilon > 0$, there is a $\delta > 0$ (in general depending on ϵ , f , I , and a) such that: Whenever $0 < |x - a| < \delta$, $|f(x) - L| < \epsilon$. We write $\lim f(x) = L$ as $x \rightarrow a$, and we call L the **limit** of f as x approaches a .

Example 3.1. Suppose that $f(x) = mx + b$ where $m, b \in \mathbb{R}$, then $\lim f(x) = f(a)$ as $x \rightarrow a$ for all $a \in \mathbb{R}$.

Example 3.2. If $f(x) = x^2 + x - 3$, show that $f(x) \rightarrow -1$ as $x \rightarrow 1$.

solution. We check $f(x) - L$, to be $x^2 + x - 3 - (-1) = x^2 + x - 2 = (x + 2)(x - 1)$, we choose δ sufficiently small such that $0 < \delta \leq 1$. Then $|x - 1| < \delta$, then $-1 < x - 1 < 1$, hence $0 < x < 2$. Then by the triangle inequality $|x + 2| \leq |x| + 2 < 2 + 2 = 4$. In order to have $|f(x) - L| = |x + 2||x - 1| < 4\delta$, if $4\delta < \epsilon$, we require $\delta = \frac{\epsilon}{4}$, hence for $\delta < \frac{1}{4}$ and $\delta < 1$, we choose $\delta = \min\{\frac{\epsilon}{4}, 1\}$. Then we have for $|x - 1| < \delta$, we get $|f(x) - L| < 4\delta < \epsilon$. Hence by definition $\lim f(x) = -1$ as $x \rightarrow 1$.

Remark. Let $a \in \mathbb{R}$ and let I be an open interval containing a . Let f and g be real valued functions defined everywhere on I except possibly at a . If $f(x) = g(x)$ for all $x \in I \setminus \{a\}$ and $f(x) \rightarrow a$ as $x \rightarrow a$, then g also has a limit as $x \rightarrow a$, and

$$\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} f(x)$$

Proof. For every $\epsilon > 0$, there is a $\delta > 0$ such that $0 < |x - a| < \delta$ implies that $|f(x) - L| < \epsilon$. Now since $|x - a| > 0$, $x \neq a$, then $f(x) = g(x)$. Thus we get that for $0 < |x - a| < \delta$, that $|g(x) - L| < \epsilon$, so by definition of the limit, we see that g has a limit, and moreover, $\lim g = \lim f$. ■

What this remark says is that those two functions are equal to each other, except (possibly) at a , however if f has a limit, then g must have the same limit.

Example 3.3. Prove that $g(x) = \frac{x^3+x^2-x-1}{x^2-1}$ has a limit as x approaches 1.

solution. First, $g(1)$ is undefined, simplifying $g(x)$ to $(x+1)$ for $x \neq \pm 1$. Letting $f(x) = x+1 = g(x)$, we have $f(1) \neq g(1)$, so by the remark, we have $\lim f = \lim g = 2$ as $x \rightarrow 1$.

We now set up the connection between the limit of a function and the limit of a sequence. This will allow us to translate results from limits of sequences, to those of functions and vice versa.

Theorem 3.1.1 (The Sequential Characterization of Limits.). *Let $a \in \mathbb{R}$, and let I be an open interval containing a . Let f be a real valued function defined everywhere on I except possibly at a . Then:*

$$\lim_{x \rightarrow a} f(x) = L$$

exists if and only if $f(x_n) \rightarrow L$ as $n \rightarrow \infty$ for every sequence $\{x_n\} \subseteq I \setminus \{a\}$ convergent to a as $n \rightarrow \infty$.

Proof. Suppose that f has limit L , then by definition, for every $\epsilon > 0$ there is a $\delta > 0$ such that $0 < |x - a| < \delta$ implies $|f(x) - L| < \epsilon$. For the same δ , there is an $N \in \mathbb{N}$ such that for $n \geq N$ implies $|x_n - a| < \delta$. Hence, for every $\epsilon > 0$ there is an N such that for $n \geq N$, $|x_n - a| < \delta$ implies $|f(x_n) - L| < \epsilon$. So $\lim f(x_n) = L$ as $n \rightarrow \infty$.

Conversely suppose that $f(x_n) \rightarrow L$ as $n \rightarrow \infty$ for all sequences $\{x_n\}$ in I converging to a . We show that $\lim f = L$ as $x \rightarrow a$; for suppose to the contrary. Then there is some $\epsilon > 0$ such that for any $\delta > 0$, there is some x_δ for which $0 < |x_\delta - a| < \delta$ and $|f(x_\delta) - L| \geq \epsilon$. Choosing $\delta = \frac{1}{n}$, denote x_δ as x_n . Then $0 < |x_n - a| < \frac{1}{n}$ implies that $\lim x_n = a$, but $|f(x_n) - L| \geq \epsilon$ for all $n \in \mathbb{N}$, thus $\lim f(x_n) \neq L$ as $n \rightarrow \infty$. A contradiction. Hence $\lim f = L$ as $x \rightarrow a$. ■

What this theorem means if x_n is a point of a sequence in I convergent to a , then when f is evaluated at x_n , then f approaches L . This theorem gives us a way to show that some functions have no limit at certain points. We can construct 2 sequences converging to a , but the sequence of f values do not equal a .

Example 3.4. Prove that $f(x) = \sin \frac{1}{x}$ for $x \neq 0$ and $f(x) = 0$ otherwise has no limit as $x \rightarrow 0$.

solution. We know that $\sin 2k\pi + \frac{\pi}{2} = 1$, so let $x_n = \frac{1}{2n\pi + \frac{\pi}{2}}$, then $f(x_n) = \sin \frac{1}{x_n} = 1$, and $\lim x_n = 0$ as $n \rightarrow \infty$. Similarly, we have that $\sin 2k\pi - \frac{\pi}{2} = -1$, so constructing $y_n = \frac{1}{2n\pi - \frac{\pi}{2}}$, then $f(y_n) = \sin y_n = -1$ and $\lim y_n = 0$ as $n \rightarrow \infty$. So, f has no limit as $x \rightarrow 0$.

We define the algebra of functions.

Definition. A vector space with **vector multiplication** defined over it is called an **algebra**.

Theorem 3.1.2. Suppose that f and g are real valued functions. Define $+$ of functions to be $f + g(x) = f(x) + g(x)$, define αf by $(\alpha f)(x) = \alpha f(x)$. Now define \cdot by $fg(x) = f(x)g(x)$, and finally define the inverse of \cdot to be $f/g(x) = f(x)/g(x)$ for $g(x) \neq 0$ for all x . Then the space of all realvalued functions, together with these operations is an algebra.

Theorem 3.1.3. Suppose that $a \in \mathbb{R}$, and I is an open interval containing a , and that f and g are realvalued functions defined everywhere on I , except possibly at a . If f and g converge as $x \rightarrow a$, then so do $f + g$, αf for $\alpha \in \mathbb{R}$, fg , and f/g .

Proof. We apply theorem 3.1.1, and then repeat the proof analogous to that for the sequences. ■

Theorem 3.1.4 (The Squeeze Theorem for Limits of Functions). Let $a \in \mathbb{R}$, and let I be an open interval containing a , and let f , g , and h be realvalued functions defined every where on I , except possibly at a . If $g(x) \leq h(x) \leq f(x)$ for all $x \neq a \in I$, and $\lim f = \lim g = L$ as $x \rightarrow a$, then the limit of h exists, and $\lim h = L$ as $x \rightarrow a$.

Corollary. If $|g| \leq M$ for all $x \neq a \in I$, and $f \rightarrow 0$ as $x \rightarrow a$, then $\lim fg = 0$ as $x \rightarrow a$.

The proof of theorem 3.1.4 and its corollary are identical to the analogous proofs for sequences, with the added caveat that we are applying theorem 3.1.1.

Theorem 3.1.5 (Comparison Theorem for Limits of Functions). Suppose that $a \in \mathbb{R}$ and that I is an open interval containing a . Let f , and g be realvalued functions defined everywhere on I , except possibly at a . If f and g have limits as $x \rightarrow a$ and $f \leq g$ for all $x \in I \setminus \{a\}$, then $\lim f \leq \lim g$ as $x \rightarrow a$.

This theorem is also a direct application of theorem 3.1.1. We can now easily evaluate limits of functions with these theorems out of the way.

Example 3.5.

$$\lim_{x \rightarrow 1} \frac{x-1}{3x-1} = \frac{1-1}{3+1} = \frac{0}{4} = 0.$$

Homework. Exercises 1, 2, 3, and 4 on page 63.

3.2 One Sided Limits.

If we have a realvalued function $f(x) = \sqrt{x-1}$, we know that $x \geq 1$, and we would like to talk about $\lim f$ at $x = 1$. Then we can only talk about $x \rightarrow 1$ from the righthand side of the function since because of our restriction on x .

Definition. Let $a \in \mathbb{R}$. A realvalued function is said to **converge** to L as x **approaches** a **from the right** if and only if f is efied on some open interval I containing a as a left endpoint and for every $\epsilon > 0$, there is a $\delta(\epsilon) > 0$ such taht $a + \delta \in I$, and $a < x < a + \delta$ implies $|f(x) - L| < \epsilon$. We then call L the **righthanded limit** of f as x approaches a and denote it $\lim f$ as $x \rightarrow a^+$.

Definition. Let $a \in \mathbb{R}$. A realvalued function is said to **converge** to L as x **approaches** a **from the left** if and only if f is defined on some open interval I containing a as a right endpoint and for every $\epsilon > 0$, there is a $\delta(\epsilon) > 0$ such that $a + \delta \in I$, and $a - \delta < x < a$ implies $|f(x) - L| < \epsilon$. We then call L the **lefthanded limit** of f as x approaches a and denote it $\lim_{x \rightarrow a^-} f$.

In general, we may also write $\lim_{x \rightarrow a^+} f$ and $\lim_{x \rightarrow a^-} f$ respectively.

Example 3.6. Let $f(x) = x + 1$ for $x \geq 0$ and $f(x) = x - 1$ for $x < 0$. We see that $\lim_{x \rightarrow 0^+} f = \lim_{x \rightarrow 0^+} x + 1 = 1$ and $\lim_{x \rightarrow 0^-} f = \lim_{x \rightarrow 0^-} x - 1 = -1$. Does the limit of f in general exist? Since the limit of f must be unique, then it may be either of these two limits, however, since the righthanded and left handed limits converge at two different values, that would imply two different limits at the same x , which is impossible. So in general we cannot say if $\lim f$ exists as $x \rightarrow 0$.

Theorem 3.2.1. Let f be a realvalued function. Then the limit $\lim f$ as $x \rightarrow a$ exists and equals L if and only if:

$$\lim_{x \rightarrow a^+} f = \lim_{x \rightarrow a^-} f = L$$

Proof. Suppose that f has limit L as $x \rightarrow a$. Then for any $\epsilon > 0$, there is a $\delta > 0$ such that $0 < |x - a| < \delta$ implies $|f(x) - L| < \epsilon$. Then $a - \delta < x < a + \delta$ implies $|f(x) - L| < \epsilon$. Thus we have that $x - \delta < x < a$ and $a < x < a + \delta$ both imply $|f(x) - L| < \epsilon$. Hence we get that $\lim_{x \rightarrow a^-} f = \lim_{x \rightarrow a^+} f = L$.

Conversely Suppose that $\lim_{x \rightarrow a^-} f = \lim_{x \rightarrow a^+} f = L$. then by definition for the left and righthanded limits we get that $a - \delta_1 < x < a$ and $a < x < a + \delta_2$ imply $|f(x) - L| < \epsilon$, thus we have that $a - \delta_1 < x < a - \delta_2$, then choosing $\delta = \min\{\delta_1, \delta_2\}$, and noting that $x \neq a$, we get that $0 < |x - a| < \delta$ implies $|f(x) - L| < \epsilon$. Thus $\lim f = L$ as $x \rightarrow a$. ■

Another thing we would like to study is the limit of a function f (with domain A) as x gets arbitrarily large, and as x gets arbitrarily small, i.e. as $x \rightarrow \infty$ and $x \rightarrow -\infty$. We say that $f \rightarrow L$ as $x \rightarrow \infty$ if and only if there exists a $C > 0$ such that $(C, \infty) \subseteq A$ and $\epsilon > 0$ such that there is an $M \in \mathbb{R}$ such that $x > M$ implies $|f(x) - L| < \epsilon$; likewise, $f \rightarrow L$ as $x \rightarrow -\infty$ if and only if there is a $C > 0$ such that $(-\infty, C) \subseteq A$ and $\epsilon > 0$ such that there is an $M \in \mathbb{R}$ such that $x < M$ implies $|f(x) - L| < \epsilon$.

Similarly, we can go on to define “infinite limits” of a function f . We say that $f \rightarrow \infty$ as $x \rightarrow a^+$ if and only if there is an open interval I with left endpoint a such that for any $M > 0$, there is a $\delta > 0$ such that $a < x < a + \delta$ implies $f(x) > M$. Other infinite limits can be defined similarly. We also have infinite limits at infinity, for example $\lim f = \infty$ as $x \rightarrow \infty$ (take $f(x) = x$). What we mean by “infinity” is just to say that for x sufficiently large (or small), f tends to L (in the case for limits at infinity), and that f grows without bound in either direction as x approaches some value a (for infinite limits). In the case of infinite limits at infinity, then we are saying that f grows without bound, (in either direction), for x sufficiently large (or small). The symbol ∞ should not be taken as a literal number, but as a concept denoting something as arbitrarily big, or without bound.

Example 3.7. Show that $\lim_{x \rightarrow 0^+} \frac{1}{x} = \infty$ as $x \rightarrow 0^+$ and that $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$ as $x \rightarrow \infty$.

solution. For any $M > 0$, we need $\frac{1}{x} > M$, then $x < \frac{1}{M}$, so choose $\delta = \frac{1}{M}$, then for $0 < x < \delta$, $\frac{1}{x} > M$, so by definition, $\lim_{x \rightarrow 0^+} \frac{1}{x} = \infty$ as $x \rightarrow 0^+$.

Similarly, for any $\epsilon > 0$, we need $|\frac{1}{x}| < \epsilon$, hence $x > \frac{1}{\epsilon}$, so choose $M = \frac{1}{\epsilon}$. Then by definition, the limit follows.

3.3 Continuity.

In general, we can say that realvalued function f is **continuous** at a if f is defined at a , and $\lim f(x) = f(a)$ as $x \rightarrow a$. We give the formal definition below.

Definition. Let $E \subseteq \mathbb{R}$ be nonempty, and let $f : E \rightarrow \mathbb{R}$ be a realvalued function. f is said to be **continuous** at a point $a \in E$, if given $\epsilon > 0$, there is a $\delta(\epsilon) > 0$ such that $|x - a| < \delta$, and $x \in E$ imply $|f(x) - f(a)| < \epsilon$. f is said to be **continuous**, on E if and only if f is continuous at x for all $x \in E$.

Remark. we have that a realvalued function f is continuous at a point a on its domain if and only if $\lim f = f(a)$ as $x \rightarrow a$. We see that $\lim f(x) = f(\lim x) = f(a)$ as $x \rightarrow a$. That is to say, if you can commute f and \lim , then f is a continuous function.

Theorem 3.3.1. Suppose that $E \subseteq \mathbb{R}$ is nonempty, and that $a \in E$, and $f : E \rightarrow \mathbb{R}$ is a realvalued function. Then the following are equivalent:

(1) f is continuous at a .

(2) If $x_n \rightarrow a$, and $x_n \in E$, then $f(x_n) \rightarrow f(a)$ as $n \rightarrow \infty$.

Proof. We apply the sequential characterization of limits of functions. ■

Theorem 3.3.2. Let $E \subseteq \mathbb{R}$ be nonempty, and let $f : E \rightarrow \mathbb{R}$ and $g : E \rightarrow \mathbb{R}$ be realvalued functions. If f and g are continuous at a point $a \in \mathbb{R}$, then so is $f + g$, fg , αf for $\alpha \in \mathbb{R}$. Moreover, f/g is continuous at $a \in E$ when $g(a) \neq 0$ (specifically, when $g(x) \neq 0$ for all $a \in E$).

We omit the proof. What this theorem says algebraically, is the the set of all continuous functions over a domain E form an algebra. Now consider for realvalued functions continuous at a point $a \in E$, then

$$f^+(x) = \frac{f(x) + |f(x)|}{2} = \max\{f(x), 0\}$$

$$f^-(x) = \frac{|f(x)| - f(x)}{2} = \max\{-f(x), 0\}$$

Then notice that $f = f^+ - f^-$, and so any function is the difference of two non negative functions, moreover $|f| = f^+ + f^-$. Moreover, all polynomial functions are continuous over \mathbb{R} .

Definition. We define the composition of functions as the binary operation \circ such that if $f : A \rightarrow B$ are functions and $g : B \rightarrow C$, then $f \circ g : A \rightarrow C$ is the function defined by $f \circ g(x) = f(g(x))$.

It is well known that \circ is not commutative, i.e. $f \circ g \neq g \circ f$.

Theorem 3.3.3. Suppose that $A, B \subseteq \mathbb{R}$ and that $f : A \rightarrow \mathbb{R}$ and $g : B \rightarrow \mathbb{R}$ are realvalued functions, with $f(A) \subseteq B$. Then if $A = I \setminus \{a\}$ where I is a nondegenerate interval, that either contains a , or has a as one of its endpoints. If $L = \lim f$ as $x \rightarrow a$ for $x \in I$ exists and belongs to B , and if g is continuous at $L \in B$, then $\lim g \circ f = g(\lim f)$ as $x \rightarrow a$ for $x \in I$.

Proof. Let $\{x_n\} \subseteq I \setminus \{a\}$, such that $x_n \rightarrow a$ as $n \rightarrow \infty$. Then if $f(A) \subseteq B$ with $f(x_n) \in B$, then $L = \lim f$ as $x \rightarrow a$, then by the sequential characterization of limits of functions, for $x \in I$, so $f(x_n) \rightarrow L$ as $n \rightarrow \infty$. This implies that $g(f(x_n)) \rightarrow g(L)$ as $n \rightarrow \infty$, and we are done. ■

Corollary. If f is continuous at $a \in A$, and g is continuous at $f(a) \in B$, then $g \circ f$ is continuous at $a \in A$.

Definition. Let $E \subseteq \mathbb{R}$ be a nonempty subset of \mathbb{R} . A realvalued function $f : E \rightarrow \mathbb{R}$ is said to be **bounded** if and only if there is an $M \in \mathbb{R}$ such that $|f| \leq M$ for all $x \in E$.

Theorem 3.3.4 (The Extreme Value Theorem). If I is closed, bounded interval, and $f : I \rightarrow \mathbb{R}$ is continuous on I , then f is bounded on I ; moreover, if $M = \sup f(I)$ and $m = \inf f(I)$, then there exists points x_m and x_M such that $f(x_m) = m$ and $f(x_M) = M$.

Proof. Suppose that f is not bounded on I , then there is some $x_n \in I$ such that $|f(x_n)| > n$ for every $n \in \mathbb{N}$. Now I is bounded, hence so is the sequence $\{x_n\}$ of points of I . Thus, by the Bolzano Weierstrass theorem, $\{x_n\}$ has a convergent subsequence, $x_{n_k} \rightarrow a$ as $k \rightarrow \infty$. Now since I is closed, it contains all the limit points, so $a \in I$. Since $f(a) \in \mathbb{R}$, and since f is continuous, then $\lim |f| = \lim |f(x_{n_k})| = \infty$ as $k \rightarrow \infty$, which is a contradiction. Hence f is bounded on I .

Now because f is bounded on I , then M, m are finite. We show that there is an $x_M \in I$ with $f(x_M) = M$. Suppose that $f < M$ for all $x \in I$, define $g : I \rightarrow \mathbb{R}$ by

$$g(x) = \frac{1}{M - f(x)}$$

We have that g is continuous, and bounded on I . Then there is a $C > 0$ such that $|g| \leq C$, so $g(x) \leq C$ for all $x \in I$. Then by definition of g we have that $f \leq M - \frac{1}{C} < M$ for all $x \in I$, so $M - \frac{1}{C}$ is an upperbound of f over I which is less than $\sup f(I)$, a contradiction. Thus $f(x_M) = M$ for some $x_M \in I$. Similarly, we get that $f(x_m) = m$. ■

We can prove the second statement alternatively by noting that $M = \sup f(I)$, so for every $n \in \mathbb{N}$, there is an $x_n \in I$ such that $M - \frac{1}{n} < f(x_n) \leq M$, then the sequence $\{x_n\} \subseteq I$ has a convergent subsequence $x_{n_k} \rightarrow a$ by the Bolzano Weierstrass theorem. Since I is closed, then we get that $a \in I$, and since f is continuous, then by the comparison theorem

$$M - \frac{1}{n_k} < f(x_{n_k}) \leq M$$

Then $M \leq f(a) \leq M$, then $f(a) = M$. The proof is also analogous for $m = \inf f(I)$.

The downside of this proof for the second part of the extreme value theorem is that it repeats the proof of the first part; that aside, the formal proof given also illustrates the relation between the extreme values of f , and the extreme values of g .

For the extreme value theorem to work, the interval I must be closed, and bounded. That is it does not work for intervals of the type (a, b) , (a, ∞) and $[a, \infty)$, or $(-\infty, b)$ and $(-\infty, b]$. f must also be a continuous function. Remove any one of those properties, and the theorem falls apart.

Chapter 4

Solutions to Chapter 1.

4.1 The Ordered Field Axioms

- (1) i. We have $a^+ + a^- = \frac{1|a|+a-a}{2} = |a|$, and $a^+ - a^- = \frac{|a|-|a|+a+a}{2} = a$.
- ii. Now Let $a \geq 0$, then $a^+ = \frac{a+a}{2} = a$ and $a^- = \frac{a-a}{2}$, and for $a < 0$, then $a^+ = \frac{a-a}{2} = 0$ and $a^- = \frac{-a-a}{2} = -a$.
- (5) (a) If $a < a < 1$, then $0 < 1-a < 1$ so $(1-a)^2 < 1-a$, then $1-a = -(a-1) < \sqrt{q-a}$, hence $-\sqrt{1-a} < a-1$, adding 1 to both sides we have $1 - \sqrt{1-a} = b < a$.
- (b) If $2 < a$, then $1 < a-1$, so $a-1 < (a-1)^2$, thus $\sqrt{a-1} < a-1$, adding 1 again to both sides, we again have $1 + \sqrt{a-1} = b < a$.
- (c) Let $a \leq b$, then $0 \leq ab \leq b^2$, so $G(a, b) = \sqrt{ab} \leq b$. Likewise, $0 \leq a+b \leq 2b$ so $A(a, b) = \frac{a+b}{2}$. Now if $A(a, b) < G(a, b)$, then $a+b \leq 2\sqrt{ab}$, so $a^2 + 2ab + b^2 < 4ab$, and so $a^2 + b^2 < 2ab$, which is absurd; thus $G(a, b) \leq A(a, b)$. Moreover, if $G(a, b) = A(a, b)$, then $2\sqrt{ab} = a+b$ implying $a = b$, and now if $a = b$, then $G(a, b) = \sqrt{b^2} = b$ and $\frac{b+b}{2} = b$, hence $G(a, b) = A(a, b)$.
- (7) (a) $|x| \leq 1$, then $|x|+1 = |x+1| \leq 2$, multiplying by $|x-1|$, then we get $|x+1||x-1| = |(x+1)(x-1)| = |x^2 - 1| \leq 2|x-1|$.
- (b) Let $-1 \leq x \leq 2$, then $1 \leq x+2 \leq 4$, hence $|x+2| \leq 4$, then again, $|x^2 + x - 2| \leq 4|x-1|$.
- (c) We have $|x| - 2 = |x-2| \leq -1 < 3$, so $|(x-2)(x+1)| = |x^2 - x - 2| \leq 3|x+1|$.
- (d) If $0 < |x-1| < 1$, adding 1, then $0 < |x| < 2$, by part (b), $|x^2 + x - 2| \leq 4|x-1| < 8|x-1|$. Now if $|x-1| = 0$, then $x = 1$ and so $1^2 + 1 - 2 = 0 < 0$ which is absurd, so this result is not true for $0 \leq |x-1|$.
- (9) We have $(a_1b_1 + a_2b_2)^2 = (a_1b_1)^2 + 2a_1b_1a_2b_2 + (a_2b_2)^2$, and $(a_1^2 + a_2^2)(b_1^2 + b_2^2) = (a_1b_1)^2 + a_1^2b_2^2 + a_2b_1^2 + (a_2b_2)^2$. Now $a_1 \leq a_1^2$ and $a_2 \leq a_2^2$, so $a_1b_1 \leq a_1^2b_1^2$ and $a_2b_2 \leq a_2^2b_2^2$. So $a_1b_1a_2b_2 \leq (a_1b_1a_2b_2)^2 = (a_1b_2a_2b_1)^2$, by problem (5c) $a_1b_1a_2b_2 \leq \frac{a_1b_2+a_2b_1}{2} \leq \frac{a_1^2b_2^2+a_2^2b_1^2}{2}$, therefore $2a_1b_1a_2b_2 \leq a_1^2b_2^2 + a_2^2b_1^2$; and so $(a_1b_1 + a_2b_2)^2 \leq (a_1^2 + a_2^2)(b_1^2 + b_2^2)$.

- (10) (a) Observe that $|x| < |a| + \epsilon$ and $|y| < |b| + \epsilon$. Also observe that $|(xy - xb) + (xb - ab)| \leq |xy - xb| + |xb - ab| < (|ab| - b\epsilon) - (|a| - \epsilon)b + (|a| - \epsilon) - ab = |(|ab| - a\epsilon - b\epsilon + \epsilon^2)| + ||ab| - b\epsilon - ab| = |-a\epsilon + \epsilon^2| + ||ab| - b\epsilon - ab| = |-a\epsilon| + |-b\epsilon| = |-a\epsilon| + |-b\epsilon| + \epsilon^2 = (|a| + |b|)\epsilon + \epsilon^2$.
- (b) The method is the same, noting that $x^2 = (|a| + \epsilon)$ and $y^2 = (|b| + \epsilon)$.

4.2 The Well Ordering Principle.

- (2) (a) $2^n = (1 + 1)^n = \sum \binom{n}{k} 1^{n-k} 1^k = \sum \binom{n}{k}$.
- (b) $(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k = a + na^{n-1}b + A$, hence $a + na^{n-1}b \leq (a + b)^n$
- (c) $(1 + \frac{1}{n})^n = \sum_{k=0}^n \binom{n}{k} \frac{1}{n^k} = 1 + \frac{1}{n} + A = \frac{n+1}{n} + A$. Now since $n \neq 0$, and $n \in \mathbb{N}$, then $n \geq 1$, so $\frac{n+1}{n} \geq 2$, therefore so is $(1 + \frac{1}{n})^n$

(3)

$$\begin{aligned} \frac{(x+h)^n - x^n}{h} &= \frac{x^n - x^n + \sum_{k=0}^n \binom{n}{k} x^{n-k} h^k}{h} \\ &= \frac{\sum \binom{n}{k} x^{n-k} h^k}{h} \\ &= \sum_{k=2}^n \binom{n}{k} x^{n-k} h^{k-1} \end{aligned}$$

- (5) Since $0 < x_1 < 2$, $x_1^2 - x_1 - 2 < 0$, hence $x_1 < \sqrt{2 + x_1} = x_2$. Now suppose that for $n \geq 1$ that $0 < x_n < x_{n+1} < 2$, and let $k = n + 2$. Then $x_k = \sqrt{2 + x_{n+1}}$; and we have that $x_{n+1}^2 - x_{n+1} - 2 < 0$, hence $x_{n+1} < \sqrt{2 + x_{n+1}} = x_k$. Therefore $0 < x_{n+1} < x_{n+2} < 2$.

- (6) (a) For $n = 1$, $1 < 2^1$. Now suppose for $n \geq 1$ that $n < 2^n$, then $n + 1 < 2^n + 1 < 2^n + 2^n = 2^{n+1}$.
- (b) For $n = 1$, $1 \leq 2^1 + 1 = 3$. Now suppose that for $n \geq 1$ that $n^2 \leq 2^n + 1$. Notice that for $n = 2$ that $2n + 1 > 2^n$, so for $n \geq 3$ let $2n + 1 < 2^n$, then $2n + 1 + 2 < 2^n + 2 < 2^n + 2^n < 2^{n+1}$ so

.Noticethen

$$(n+1)^2 = n^2 + 2n + 1 \leq 2^n + 1 + 2n + 1 \leq 2^n + 2^n + 1 \leq 2^{n+1} + 1.$$

- (c) For $n = 1$, then $13 < 3^1$; now suppose for $n \geq 1$ that $(n + 1)^3 < 3^n$. Then $(n + 1)^3 = n^3 + 3n^2 + 3n + 1 < 3^n + 3n^2 + 3n + 1 < 3^n + 3^n + 3^n = 3^{n+1}$.

4.3 The Axiom of Completeness.

- 2 Suppose $E \subseteq \mathbb{Z}$ and $E \neq \emptyset$ and that E is bounded. Then E is bounded above, so by the axiom of completeness, $\sup E$ exists. Now by the approximation property, for $\epsilon > 0$ and $a \in \mathbb{Z}$ $\sup E - \epsilon \leq a \leq \sup E$; making ϵ negligible enough we see that $\sup E \in \mathbb{Z}$. Since E has infinitely many upperbounds, choose $M \in \mathbb{Z}$ to be the least upperbound of E not in E and $M \neq \sup E$, then $n \leq \sup E \leq M$, which makes $\sup E \in E$.

Again, by the above argument, using the approximation property for greatest lower bounds (which we prove later) and letting $m \in \mathbb{Z}$ the largest lowerbound not in E with $m \neq \inf E$, we see that $\inf E \in E$.

- 3 Let $a, b \in \mathbb{R}$ with $a < b$, then notice that $a - \sqrt{2} < b - \sqrt{2}$. By the density of \mathbb{Q} in \mathbb{R} , there is a $\frac{p}{q}$ such that $a - \sqrt{2} < \frac{p}{q} < b - \sqrt{2}$. Then $a < \frac{p}{q} + \sqrt{2} < b$. Now $\frac{p}{q} + \sqrt{2} \in \mathbb{Q}$, but $\sqrt{2} \notin \mathbb{Q}$; thus $\frac{p}{q} + \sqrt{2} \notin \mathbb{Q}$, therefore \mathbb{Q}^* is dense in \mathbb{R} .
- 4 Let $a \in \mathbb{R}$ and $n \in \mathbb{N} \subseteq \mathbb{R}$, Now either $a < n$, $n < a$ or $a = n$. Suppose that $a < n$, then $\frac{1}{n} < \frac{1}{a} < a$, so $\frac{1}{n} < a$. By the density of \mathbb{Q} in \mathbb{R} , there is an $r_n \in \mathbb{Q}$ such that $\frac{1}{n} < r_n < a$. Now $r_n - \frac{1}{n} < a - \frac{1}{n} < a$, so $r_n - \frac{1}{n} < a$, it remains to prove that $a < r_n + \frac{1}{n}$. We have that $nr_n < na$, then

$$na - 1 < nr_n$$

, then $na < nr_n + 1$, thus $a < r_n + \frac{1}{n}$. Therefore we have $r_n - \frac{1}{n} < a < r_n + \frac{1}{n}$, so $|a - r_n| < \frac{1}{n}$. The proof for the case of $n < a$ is analogous; and if $a = n$, then the case reduces to the one we proved, with $\frac{1}{a} < a$.

5(a)

Theorem 4.3.1 (Approximation Property for Greatest Lowerbounds). *If E has a least upperbound, and $\epsilon > 0$ is a positive number, then there is an element $a \in E$ such that $\inf E < a \leq \inf E + \epsilon$.*

Proof. Suppose for some $\epsilon > 0$ that there is no element of E lying between $\inf E$ and $\inf E + \epsilon$. Now $\inf E + \epsilon \leq a$ for all $a \in E$, so $\inf E + \epsilon$ is a lower bound, thus $\inf E + \epsilon \leq \inf E$, implying that $\epsilon \leq 0$, a contradiction. ■

- 6 (a) Suppose that E has a lower bound m . Then for some $m_0 < m$, m_0 is also a lower bound for E . Now let m_0, m_1 be greatest lowerbounds of E , then $m \leq m_0$ for all other lowerbounds m , likewise $m \leq m_1$. Then we get that m_1 is a lowerbound, so $m_1 \leq m_0$, likewise $m_0 \leq m_1$, therefore $m_0 = m_1$ and the greatest lowerbound is unique.
- (b) Let $E \subseteq \mathbb{R}$ be nonempty and bounded below, then $-E \neq \emptyset$ and bounded above, therefore by the axiom of completeness, $\sup -E$ exists, and by theorem 1.3.4, $\sup -E = -\inf E$, which implies that $\inf E$ exists (and is finite).

Solutions to Chapter 2.

4.4 Limits of Sequences

- 1 (a) Consider the sequence $\{3 + \frac{1}{n}\}$ and choose $N > \frac{1}{\epsilon}$, then for $n \geq N$, $\frac{1}{n} \leq \frac{1}{N} < \epsilon$, then $|\frac{1}{n}| = |(3 + \frac{1}{n}) - 3| < \epsilon$, thus $\lim (3 + \frac{1}{n}) = 3$ as $n \rightarrow \infty$.
(b) Choose $N > \frac{2}{\epsilon}$, then $\frac{1}{n} \leq \frac{1}{N} < \frac{\epsilon}{2}$, implies $|\frac{1}{n}| = 2|(1 - \frac{1}{n}) - 1| < 2\frac{\epsilon}{2} = \epsilon$.
(c) Choose $N > \frac{1}{\sqrt{\epsilon}}$.
(d) Choose $N > \frac{1}{\epsilon^2}$.
- 4 (a) Let $\{b_n\} \rightarrow 0$ and let $\{x_n\}$ be a realvalued sequence such that $|x_n - a| \leq b_n$ for n sufficiently large. Then, for any $\epsilon > 0$, there is an $N \in \mathbb{N}$ such that whenever $n \geq N$, $|b_n| < \epsilon$, that is for the same N , $||x_n - a|| \leq |b_n| < \epsilon$. Hence, $x_n \rightarrow a$ as $n \rightarrow \infty$.
(b) Now suppose that $\{b_n\}$ and $\{x_n\}$ are the same sequences as before such that $|x_n - a| \leq Cb_n$ for some $C \in \mathbb{R}$. Then for any $\epsilon > 0$, choose $N = \frac{\epsilon}{C}$ so that $C|b_n| < \epsilon$. Then for the same N , we have $|x_n - a| \leq C|b_n| < \epsilon$, and we get the same result.
- 5 (a) If $\{x_n\}$ is a bounded realvalued sequence, then by definition, there is a $C > 0$ such that $|x_n| \leq C$. Conversely, suppose that there is a $C > 0$ for which $|x_n| \leq C$, then $-C \leq x_n \leq C$. So we see that $\{x_n\}$ is bounded below (by $-C$) and bounded above (by C), thus $\{x_n\}$ is bounded.
(b) If $\{x_n\}$ is a bounded realvalued sequence, then, $|x_n| \leq C$ for $C > 0$. Thus $|\frac{x_n}{n^k}| \leq \frac{C}{|n^k|}$. Thus, for any $\epsilon > 0$, choose $N = \frac{1}{\sqrt[k]{\epsilon}}$, thus we get that for $n \geq N$, $|\frac{x_n}{n^k}| \leq \frac{C}{|n^k|} < \epsilon$. Hence, $\frac{x_n}{n^k} \rightarrow 0$ as $n \rightarrow \infty$.
- 6 (a) Let $x_n \rightarrow a$ and let $y_n \rightarrow at$, then for any $\epsilon > 0$, there is are $N_1, N_2 \in \mathbb{N}$ such that $|x_n - a| < \epsilon$ and $|y_n - a| < \epsilon$. Choosing $N = \max\{N_1, N_2\}$, we get $n \geq N$ $|x - n - y_n| = |((x_n - y_n) - (a - a))| \leq |x_n - a| + |y_n - a| < \epsilon$. Therefore $x_y - y_n \rightarrow 0$ as $n \rightarrow \infty$.
(b) Consider $\{n\}$ for all $n \in \mathbb{N}$. Then n is either even or odd, so choose subsequences $\{2k\}$ and $\{2k + 1\}$ for $k \in \mathbb{N}$. Now suppose that $\lim 2k = \lim (2k + 1) = a$ as $k \rightarrow \infty$. Then $(2k + 1) - (2k) = 1 \rightarrow 0$, a contradiction ($\{1\}$ is a constant sequence), thus $\{n\}$ is divergent.

- (c) Consider the constant sequence $\{1\}$ and the sequence $\{\frac{1}{n}\}$ with $n \in \mathbb{N}$, then $1 - \frac{1}{n} = \frac{n-1}{n} \rightarrow 0$, but $1 \rightarrow 1$ and $\frac{1}{n} \rightarrow 0$ as $n \rightarrow \infty$. Hence the converse of (a) is not necessarily true.

4.5 Limit Theorems.

- (3) refer to theorem 2.2.3(4)
- (5) Let $x \in \mathbb{R}$, and let $\{x_n\}$ and $\{y_n\}$ be realvalued sequences such that $x_n \rightarrow x$ and $y_n \rightarrow x$ as $n \rightarrow \infty$ and $x_n \leq y_n$ for all $n \in \mathbb{N}$. Then by the density of \mathbb{Q} in \mathbb{R} , there are rationals $r_n \in \mathbb{Q}$ such that $x_n \leq r_n \leq y_n$, then constructing $\{r_n\}$ out of these r_n , by the sandwich theorem, $r_n \rightarrow x$ as $n \rightarrow \infty$.
- (6) Let $x, y \in \mathbb{R}^\infty$, and suppose that $\{x_n\}$, $\{y_n\}$, and $\{w_n\}$ are realvalued sequences then:
- (a) If $x_n \rightarrow x$ and $y_n \rightarrow x$ as $n \rightarrow \infty$, and $x_n \leq w_n \leq y_n$, for all $n \in \mathbb{N}$. Then suppose that $x = \infty$, then for $N_1, N_2 \in \mathbb{N}$, there is an $M \in \mathbb{R}$ such that $n \geq N_1, N_2$ implies $x_n > M$ and $y_n > M$, then by supposition, $w_n > M$, so choose $N = \max\{N_1, N_2\}$ and we have $n \geq N$ implies that $w_n > M$, thus $\lim w_n = x$. Similarly, we get the same result for $x = -\infty$.
- (b) Let $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$. There are four cases, but we only examine two (for the other two follow very similar reasoning). Suppose that $x, y = \infty$. Then For all $M_1, M_2 \in \mathbb{R}$, there are $N_1, N_2 \in \mathbb{N}$ such that $n \geq N_1, N_2$ implies $x_n > M_1$ and $n > M_2$. Thus since $x_n \leq y_n$, we have that $M_1 \leq M_2$, so $x \leq y$.
Now suppose that $x = \infty$, and $y = -\infty$. Then we have $n \geq N_1, N_2$ implies that $x > M_1$, and $y < M_2$, so we have $M_1 < x_n \leq y < M_2$, so $M_1 \leq M_2$, and we get $x \leq y$.

4.6 The Bolzano-Weierstrass Theorem.

- (1) Notice that $\sin n^3$ is bounded by 1 ($|\sin x| \leq 1$ for all $x \in \mathbb{R}$), then the sequence $\{x_n\}$ defined by:

$$x_n = \frac{(n^2 + 20n + 35) \sin n^3}{n^2 + n + 1}$$

is bounded as well. Thus by the Bolzano-Weierstrass theorem, $\{x_n\}$ has a convergent subsequence.

- (2) Let $E \subseteq \mathbb{R}$ be nonempty and bounded, and suppose that $\sup E \notin \mathbb{R}$. Let $x = \sup E$, and let $\{x_n\} \subseteq E$ be a realvalued sequence such that $x_n < x_{n+1}$. Then $\{x_n\}$ is strictly increasing, and bounded (since E is bounded), hence, by the monotone convergence theorem, $\lim x_n$ exists and is finite. All that is left is to show that $\lim x_n = x$; for suppose not. First suppose that $\lim x_n > x$, then let $x_k = x$ for $k \geq n$, then $x_n < x_k$, and so $x \in \{x_n\}$, which contradicts the supposition that $x \notin E$. Now suppose that $\lim x_n < x$, then there is some $k \geq n$ for which $x_n \leq x_k$, which violates the supposition

that $\{x_n\}$ is strictly increasing. Thus we must have $\lim x_n = x$ as $n \rightarrow \infty$. Similarly, if we suppose that $\inf E < E$, we get that there is a strictly decreasing sequence $\{y_n\}$ converging to $\inf E$, where $y_n \in E$ for all n .

- (4) By exercise 1.1.5(a), we have $x_{n+1} < x_n$ for all n , so $\{x_n\}$ is strictly decreasing. Then choose $I = (0, 1)$, and let $\{I_n\} \subseteq I$ be a sequence of nested intervals. Notice that $|I_n| \rightarrow 0$, as $n \rightarrow \infty$. So, by the nested interval theorem, choose

$$E = \bigcap I_n = (0, x_k)$$

for some $k \geq n$, we see that $x_k = 0$, and since we suppose that $x_k < x_n$, we see that $x_n \rightarrow 0$.

Now consider the sequence $\{\frac{x_{n+1}}{x_n}\}$, rationalizing $\frac{x_{n+1}}{x_n}$, we get $\frac{x_{n+1}}{x_n} = \frac{1}{2} + A$ for some term A . Then the sequence $A \rightarrow 0$, so $\lim \frac{x_{n+1}}{x_n} = \lim (\frac{1}{2}) + A = \lim \frac{1}{2} + \lim A = \frac{1}{2}$.

- (1) We see that of $x_{n+1} = \sqrt{2x_n + 3}$, then $x_n = \frac{x_{n+1}^2 - 3}{2}$, so $x_n < x_{n+1}$, so $\{x_n\}$ is strictly increasing. Again, by the nested interval property for $I = (0, 3]$, we see that $\lim x_n = 3$.
- (2) By exercise 1.1.5(b), we have that $x_{n+1} < x_n$, so $\{x_n\}$ is strictly decreasing, moreover by the nested interval theorem, choosing $I = (2, \infty)$, we see that $\lim x_n = 2$. Now what happens when we suppose $1 \leq x_1 < 2$? If we let $x_1 = 1$, then $x_2 = 1$. If we let $x_1 = \frac{3}{2}$, then we see that $x_2 = 1 + \frac{\sqrt{2}}{2}$, hence $x_{n+1} \leq x_n$, so $\{x_n\}$ is monotone increasing. By the same reasoning as before, we also see $\lim x_n = 2$.

4.7 Cauchy Sequences.

- (1) Let $\{x_n\}$ and $\{y_n\}$ be Cauchy sequences, we prove that $\{x_n + y_n\}$

By theorem 2.4.1, both $\{x_n\}$ and $\{y_n\}$ converge to points in \mathbb{R} , then by the limit laws, so does $\{x_n + y_n\}$, hence again by theorem 2.4.1, $\{x_n + y_n\}$ is Cauchy.

Alternatively, we have by definition that for all $\epsilon > 0$, there are $N_1, N_2 \in \mathbb{N}$, such that $m, n \geq N_1, N_2$ implies that $|x_n - x_m| < \frac{\epsilon}{2}$, and $|y_n - y_m| < \frac{\epsilon}{2}$. Then choosing $N = \max\{N_1, N_2\}$, we have that $|(x_n + y_n) - (x_m + y_m)| = |x_n - x_m| + |y_n - y_m| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$. Thus we get that $\{x_n + y_n\}$ is a Cauchy sequence.

- (2) Let $\{x_n\}$ be a Cauchy sequence for $x_n \in \mathbb{N}$. Then for all $\epsilon > 0$, there is an $N \in \mathbb{N}$ such that for $m, n \geq N$, $|x_n - x_m| < \epsilon$. Notice that $|x_n - x_m| = |x_m - x_n|$, and that $x_n \rightarrow a$ for some $a \in \mathbb{R}$ (by theorem 2.4.1). Then $\lim x_n = a$, and by definition of the limit, we also have that $\lim x_n = x_n$; thus by the uniqueness of limits, $x_n = a$.
- (6) Let $S_n = \sum_{k=1}^n \frac{(-1)^k}{k}$, then notice that for some $m \leq n$, that $S_n = S_{2m} + S_{2m+1}$, We see that $|S_{2m}| < 1$ (for all values of k), and we observe that $|S_{2m} - S_{2l}| = |S_{2m-2l}|$, hence $\{S_{2m}\}$ is Cauchy, likewise we get that $\{S_{2m+1}\}$ is Cauchy, thus so is S_n , thus by theorem 2.4.1, $\lim S_n$ exists and is finite.

4.8 Limits of Least Upperbounds and Greatest Lowerbounds.

- (2) Let $\{x_n\}$ be a real valued sequence. By theorem 1.3.4 we have that $\sup x_n = -\inf x_n$, so by definition, we have $\limsup x_n = \lim(\sup x_k) = \lim(-\inf x_k) = -\lim(\inf x_k) = -\liminf x_n$ as $n \rightarrow \infty$ and $\text{fork} \geq n$.
- (3) Suppose that $\limsup x_n < r$, then $\limsup x_n$ is finite, so we get that $\{x_n\}$ is bounded above, moreover, notice that $\sup x_k < r$ for all $k \geq n$, thus we get that r is an upperbound for $\{x_n\}$.

Similarly, if $\limsup x_n > r$, then we get that $\limsup x_n$ is not bounded above, but that $\sup x_k$ is bounded below by r for all $k \geq n$; so we get that $\{x_n\}$ is bounded below by r .

- (4) We first prove a lemma.

Lemma 4.8.1. *let $\{x_n\}$ and $\{y_n\}$ be real valued sequences such that $\sup x_n, \sup y_n$, and $\inf x_n, \inf y_n$ exist. Then $\sup x_n + y_n \leq \sup x_n + \sup y_n$ and $\inf x_n + \inf y_n \leq \inf x_n + y_n$.*

Proof. Let $x = \sup x_n$ and $y = \sup y_n$. Then we have that $x \geq x_n$ and $y \geq y_n$ for all n , hence, $x + y \geq x_n + y_n$, and we get that $\sup x_n + y_n \leq x + y$. By similar reasoning, we get that $\inf x_n + \inf y_n \leq \inf x_n + y_n$. ■

- (a) $\liminf x_n + \liminf y_n = \lim(\inf x_k + \inf y_k) \leq \lim(\inf x_k + y_k) = \liminf x_n + y_n \leq \limsup x_n + \liminf y_n \leq \limsup x_n + y_n \leq \limsup x_n + \limsup y_n$.
- (b) Let $x = \lim x_n$ as $n \rightarrow \infty$. Then by theorem 2.5.3, we see that $\liminf x_n \leq x \leq \limsup x_n$, thus we have by part (a) that $\liminf x_n + \liminf y_n \leq x + \liminf y_n \leq \liminf x_n + y_n \leq x + \liminf y_n \leq \limsup x_n + \liminf y_n$, thus $\liminf x_n + y_n = x + \liminf y_n$.

Similarly, we get that $\limsup x_n + y_n = x + \limsup y_n$.

- (5) Similar to part (4), we have the following lemma:

Lemma 4.8.2. *Let $\{x_n\}$ and $\{y_n\}$ be realvalued sequences, such that $\sup x_n, \sup y_n$ and $\inf x_n$ and $\inf y_n$ exist. Then $\sup x_n y_n \leq \sup x_n \sup y_n$ and $\inf x_n \inf y_n \leq \inf x_n y_n$.*

Proof. Let $x = \sup x_n$ and $y = \sup y_n$. Then $x \geq x_n$ and $y \geq y_n$, for all n , thus $xy \geq x_n y_n$, hence we get that $\sup x_n y_n \leq \sup x_n \sup y_n$. By similar reasoning, we get that $\inf x_n \inf y_n \leq \inf x_n y_n$. ■

- (a) $\limsup x_n y_n = \lim(\sup x_k y_k) \leq \lim(\sup x_k \sup y_k) = \lim(\sup x_k) \lim(\sup y_k) = \limsup x_n \limsup y_n$.
- (b) $\liminf x_n \liminf y_n = \lim(\inf x_k) \lim(\inf y_k) = \lim(\inf x_k \inf y_k) \leq \lim(\inf x_k y_k) = \liminf x_n y_n$.
- (6) Let $x = \lim x_n$ as $n \rightarrow \infty$. By theorem 2.5.3m, $x \leq \limsup x_n$, thus $x \limsup x_n \leq \limsup x_n y_n \leq x \limsup y_n$, thus $\limsup x_n y_n = x \limsup y_n$, for $x_n \geq 0$ and $y_n \geq 0$.

Solutions to Chapter 3.

4.9 Two sided Limits

- (1) Let $\epsilon > 0$, we have that $x^2 - x + 1 - 3 = (x - 2)(x + 2)$. If $0 \leq \delta \leq 1$, then $|x - 2| < \delta$ implies $0 < x < 4$. Now by the triangle inequality.