

Topology

Alec Zabel-Mena

Text

Topology (2rd edition)

James Munkres.

November 9, 2020

Chapter 1

Topological Spaces and Continuous Functions.

1.1 Topological Spaces.

Definition. A **topology** on a set X is a collection \mathcal{T} of subsets of X such that:

- (1) $\emptyset, X \in \mathcal{T}$.
- (2) For any subcollection $\{U_\alpha\}$ of subsets of X , $\bigcup_\alpha U_\alpha \in \mathcal{T}$.
- (3) For any finite subcollection $\{U_i\}_{i=1}^n$ of subsets of X , $\bigcap_{i=1}^n U_i \in \mathcal{T}$.

We call the pair (X, \mathcal{T}) a **topological space**, and we call the elements of \mathcal{T} **open sets**.

Example 1.1. (1) Let X be any set, the collection of all subsets of X , 2^X is a topology on X , which we call the **discrete topology**. We call the topology $\mathcal{T} = \{\emptyset, X\}$ the **indiscrete topology**.

- (2) The set of three points $\{a, b, c\}$ has the 9 following topologies in figure 1.1.




Figure 1.1: The Topologies on $\{a, b, c\}$.

- (3) Let X be any set, and let $\mathcal{T}_f = \{U \subseteq X : X \setminus U \text{ is finite, or } X \setminus U = X\}$. Then \mathcal{T}_f is a topology and called the **finite complement topology**.
- (4) Let X be any set, and let $\mathcal{T}_c = \{U \subseteq X : X \setminus U \text{ is countable, or } X \setminus U = X\}$. Then \mathcal{T}_c is a topology on X .

Definition. Let X be a set, and let \mathcal{T} and \mathcal{T}' be topologies on X . We say that \mathcal{T} is **coarser** than \mathcal{T}' , and \mathcal{T}' **finer** than \mathcal{T} if $\mathcal{T} \subseteq \mathcal{T}'$. If two topologies are either coarser, or finer than each other, we call them **comparable**.

Example 1.2. The topologies \mathcal{T}_f and \mathcal{T}_c are comparable, and we see that $\mathcal{T}_c \subseteq \mathcal{T}_f$, so \mathcal{T}_f is coarser than \mathcal{T}_c , and \mathcal{T}_c is finer than \mathcal{T}_f .

1.2 The Basis and Subbasis for a Topology.

Definition. If X is a set, the **basis** for a topology on X is a collection \mathcal{B} of subsets of X , called **basis elements**, such that:

- (1) For every $x \in X$, there is a $B \in \mathcal{B}$ such that $x \in B$.
- (2) For $B_1, B_2 \in \mathcal{B}$, if $x \in B_1 \cap B_2$, then there is a $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.

We define the topology \mathcal{T} **generated** by \mathcal{B} to be collection of open sets: $\mathcal{T} = \{U \subseteq X : x \in B \text{ for some } B \in \mathcal{B}\}$.

Theorem 1.2.1. Let X be a set, and \mathcal{B} a basis of X , then the collection of subsets of X , $\mathcal{T} = \{U \subseteq X : x \in B \text{ for some } B \in \mathcal{B}\}$ is a topology on X .

Proof. Let \mathcal{B} be a basis for a topology in X , and consider \mathcal{T} as defined above. Clearly, $\emptyset \in \mathcal{T}$ and so is X .

Now let $\{U_\alpha\}$ be a subcollection of subsets of X , and let $U = \bigcup U_\alpha$. Then if $x \in U$ for some α , there is a B_α such that $x \in B_\alpha \subseteq U_\alpha$, thus $x \in B_\alpha \subseteq U$.

Now let $x \in U_1 \cap U_2$, and choose $B_1, B_2 \in \mathcal{B}$ such that $x \in B_1 \subseteq U_1$ and $x \in B_2 \subseteq U_2$. Then by definition, there is a B_3 for which $x \in B_3 \subseteq B_1 \cap B_2$. Now suppose for arbitrary n , that $U = \bigcap_{i=1}^n U_i \in \mathcal{T}$, for some finite subcollection $\{U_i\}$ of subsets of X . Then by let $B_n, B_{n+1} \in \mathcal{B}$ such that $x \in B_n \subseteq U$ and $x \in B_{n+1} \subseteq U_{n+1}$. Then by our hypothesis, there is a B for which $x \in B \subseteq B_n \cap B_{n+1}$, thus $U \cap U_{n+1} = \bigcap_{i=1}^{n+1} U_i \in \mathcal{T}$. This make \mathcal{T} a topology on X . ■

Example 1.3. (1) Let \mathcal{B} be the set of all circular regions in the plane $\mathbb{R} \times \mathbb{R}$, then \mathcal{B} satisfies the conditions needed for a basis.

- (2) The collection \mathcal{B}' in $\mathbb{R} \times \mathbb{R}$ of all rectangular region also forms a basis for a topology on $\mathbb{R} \times \mathbb{R}$.
- (3) For any set X , the set of all 1-point elements of X forms a basis for a topology on X .




Figure 1.2: The basis for \mathcal{B} and \mathcal{B}' in $\mathbb{R} \times \mathbb{R}$ (see example (2)).

Lemma 1.2.2. *Let X be a set, and \mathcal{B} be a basis for a topology \mathcal{T} on X . Then $\mathcal{T} = \{\bigcup B : B \in \mathcal{B}\}$.*

Proof. Given a collection $\{B\}$ of basis elements in \mathcal{B} , since they are all in \mathcal{T} , their unions are also in \mathcal{T} . Conversely, given $U \in \mathcal{T}$, then for every point $x \in U$, choose a $B_x \in \mathcal{B}$ such that $x \in B_x \subseteq U$, then $U = \bigcup_{x \in U} B_x$. ■