

A Retained Mixed Gravity–Maxwell Bracket under Cyclic Homological Reduction through ℓ_3 Exact Transfer and Causal Cartan Compatibility

Asger Alstrup Palm*

Working draft, July 2026

Abstract

We transfer the classical gravity–Maxwell BV interaction on a compact positive Berger-clock background from a complete 64-row presentation to a typed cyclic 36-row retained complex. The contraction carries the exact odd pairing and the unary causal Green data. In the repository’s suspended graded-symmetric convention the retained mixed ternary operation is nonzero. Two zero-derivative witnesses have exact coefficients $+1$ and -1 in the gravity-equation and Maxwell-equation sectors, respectively. Thus nonvanishing is witnessed by explicit evaluations rather than inferred from the total sparse-coefficient count.

The homological exchange channel is computed rather than omitted. It contains 342 canonical full-complex coefficients in the only nonempty channel, all supported on full row 38, and the certified retained projection annihilates that row. A one-entry projection mutation exposes all 342 terms. Thus the retained exchange contribution vanishes for a structural reason while the direct q_3 contact survives for this frozen contraction. Standard cyclic homotopy transfer supplies the abstract L_∞ and cyclic identities; an independent sparse implementation audit replays them on all 36 rows. Lowering ℓ_3 with the retained typed pairing independently reproduces all 25,662 degree-zero lowered quartic coefficients with zero cyclicity defects; changing the Maxwell pairing weight from two to one produces 17,108 exact defects on 14 rows. The remaining 288 ghost/antifield coefficients also replay exactly under the suspended-Darboux transpose, closing cyclicity of the full retained BV tensor.

For the explicitly specified cyclic strong deformation retract these calculations establish

The retained mixed gravity–Maxwell bracket survives this
cyclic homological reduction through ℓ_3 .

The separately certified causal theorem gives compatible advanced and retarded unary homotopies and two-sided-causal-hull Cartan primitives for the background-fixing helical generator K_{Berger} . It is a parallel compatibility theorem: ℓ_3 is transferred with the support-local algebraic homotopy S , not by integrating out fields with a causal propagator. The statement is classical and carries separate LOCAL-ALGEBRAIC and LORENTZIAN-CAUSAL dependencies. It does not prove a nonzero operation on cohomology, a nontrivial cyclic deformation class, invariance under changing the cyclic retract, or nonremovability by a cyclic L_∞ field redefinition. Residual mixing, Hadamard products, QME restoration, and every quantum or particle interpretation remain open. A separate exact audit obstructs the canonical local same-bundle branch projector on the 36-row carrier. The resulting rank-46 spatial-STF2 graph prolongation and its cyclic dual are now constructed as an exact local cyclic carrier, but no branch projector or nonlinear lift has yet been certified on it.

*Corresponding author: asger@area9.dk.

AI-use disclosure and computational accountability

OpenAI GPT-5 was used substantively for mathematical synthesis, verification code, claim auditing, and manuscript drafting. The human author directed the programme and remains responsible for the claims, citations, and final text. Every numerical count in the theorem is an exact sparse-coefficient count over $\mathbb{Q}(\sqrt{10})$ and is tied to a deterministic certificate. No floating point calculation enters an identity, rank, cyclicity test, or mutation witness. Submission remains conditional on final human review.

Contents

1	Question and answer	2
2	Complexes, conventions, and claim boundary	3
3	Ternary homological transfer	5
4	Frozen-SDR nonvanishing and its audit	6
5	Independent cyclic implementation audits	7
6	Causal compatibility through ternary order	8
7	Residual descent and the projector obstruction	9
8	Interpretation and nonclaims	10
9	Reproducibility and certificate map	11
10	Outlook	12
A	Exact inclusion columns for the displayed witnesses	12

1 Question and answer

The free BV complex identifies which classes survive gauge reduction, but it does not determine whether nonlinear gravity–matter vertices survive the same reduction. A transferred interaction can disappear in three distinct ways: its direct contact term can project to zero; homological exchange terms can cancel it; or cyclicity and the L_∞ identities can fail after compression. Conversely, a nonzero tensor on a gauge-fixed carrier need not descend to any physical branch until its residual projection is computed.

This paper answers the algebraic question through ternary order for one explicit gravity–Maxwell extension and one frozen cyclic contraction of the positive Berger-clock complex. Write

The retained mixed gravity–Maxwell bracket survives this
cyclic homological reduction through ℓ_3 .

(1)

Here “survives” means that the explicitly transferred mixed ternary bracket is nonzero in the displayed retained coordinates and obeys the transferred arity-three identity. “Cyclic” means

that the contraction is typed cyclic and that the complete lowered BV quartic form has been independently replayed under cyclic transposition. Causal Green and Cartan compatibility are proved separately in Section 6; they are not part of the definition of the algebraic transfer.

Nonvanishing here is deliberately representative-level. We have not proved

$$[\ell_3^{\text{mixed}}] \neq 0 \quad (2)$$

in a cyclic coderivation or Chevalley–Eilenberg deformation complex. Nor have we proved a nonzero induced operation on $H(\mathcal{R}_{36}, \ell_1)$, survival under all cyclic L_∞ equivalences or further minimal transfer, or an obstruction to removing the selected component by a cyclic quadratic change of retained coordinates. Those are the nearest invariant strengthenings of the present theorem.

The theorem stops one rung before physical branch interpretation. A subsequent exact audit now obstructs the canonical support-local same-bundle Einstein-like/extra-Weyl projector on the 36-row carrier. Residual descent therefore requires an enlarged or differently organized carrier. The natural rank-46 STF2 graph carrier has now passed its exact unary cyclic-SDR gate, but its branch projector and nonlinear lift remain open, so a branch mixing table is still unauthorized. Deformation/vertex classes belong to a separate complex and are not used in the theorem below.

2 Complexes, conventions, and claim boundary

Let $(\mathcal{C}_{64}, q_1, \omega_{64})$ be the complete typed classical gravity–clock–Maxwell BV complex at the frozen rational Berger fixture. Its degree multiplicities are

$$6, 26, 26, 6 \quad \text{in degrees } -1, 0, 1, 2. \quad (3)$$

Let $(\mathcal{R}_{36}, \ell_1, \omega_{36})$ denote the retained typed carrier, with degree multiplicities

$$4, 14, 14, 4. \quad (4)$$

The exact contraction is

$$(\mathcal{C}_{64}, q_1) \begin{matrix} \xrightarrow{\pi} \\ \xleftarrow{\iota} \end{matrix} (\mathcal{R}_{36}, \ell_1), \quad \pi\iota = \text{id}, \quad q_1 S + S q_1 = \text{id} - \iota\pi, \quad (5)$$

with the exact side conditions

$$S^2 = 0, \quad S\iota = 0, \quad \pi S = 0, \quad (6)$$

and typed cyclic adjoint relations fixed by the carrier certificate. The coefficients lie in $\mathbb{Q}(\sqrt{10})$. The Maxwell field–equation pairing has absolute weight two, while the gravity weight is one. The signed odd-pairing ledger contains ten gravity entries of sign -1 and four Maxwell entries of sign $+2$ on the degree-zero field–equation pairs.

For completeness, let $\dagger$ denote the formal PBW adjoint, including integration by parts, and let Ω_{64} and Ω_{36} be the signed typed pairing matrices. All suspension and Koszul signs are already included in these matrices. The exact convention-specific cyclic SDR relations are

$$\iota^\dagger \Omega_{64} = \Omega_{36} \pi, \quad \iota^\dagger \Omega_{64} \iota = \Omega_{36}, \quad S^\dagger \Omega_{64} + \Omega_{64} S = 0. \quad (7)$$

Equivalently, for homogeneous retained x and full-carrier y ,

$$\omega_{64}(\iota x, y) = \omega_{36}(x, \pi y), \quad \omega_{64}(Sx, y) = -\omega_{64}(x, Sy). \quad (8)$$

The last displayed sign is therefore the suspended-coordinate sign actually verified by the certificate, not an unsuspended sign imported from another convention.

Table 1 gives the carrier map needed to interpret the row indices. The retained gravity representatives are dressed combinations, so the table states their image by bundle type rather than pretending that the gravity part of ι is a coordinate inclusion. The Maxwell summand is retained by the identity.

Table 1: Human-readable carrier map. A zero retained image means that the displayed full block belongs to the contractible complement.

Full rows	Degree	Full carrier content	Retained image
0–4	−1	three spatial diffeomorphism ghosts, clock ghost τ , Weyl ghost σ	spatial ghosts 0–2
5–16	0	ten $\hat{h}_{\mu\nu}$ fields, R , Θ	dressed $\hat{h}$ rows 3–12
17–26	0	nonminimal multipliers and antighost-duals	0
27–37	1	ten $\hat{h}_{\mu\nu}^*$ equations and R^*	dressed $\hat{h}^*$ rows 13–22
38	1	Θ^* , the clock/Stueckelberg equation dual	0 under π
39–48	1	nonminimal dual and antighost rows	0
49–53	2	three spatial ghost antifields, τ^* , σ^*	spatial ghost-antifields 23–25
54	−1	Maxwell ghost c_M	c_M (row 26)
55–58	0	Maxwell fields $A_0, \dots, A_3$	rows 27–30
59–62	1	Maxwell equations $A_0^+, \dots, A_3^+$	rows 31–34
63	2	Maxwell ghost antifield c_M^+	row 35

The row-38 support statement is therefore geometric. Rows 3, 16, 38, 52 of the full carrier are $\tau, \Theta, \Theta^*, \tau^*$, and the certified blocks include

$$\text{pr}_\Theta q_1(\tau) = \Theta, \quad q_1(\Theta^*) = -\tau^*, \quad S(\Theta) = \tau, \quad S(\tau^*) = -\Theta^*. \quad (9)$$

Moreover

$$\pi|_{\langle \Theta^*, \tau^* \rangle} = 0, \quad \text{pr}_{\langle \Theta^*, \tau^* \rangle} \iota = 0, \quad \omega_{64}(\Theta, \Theta^*) = 1, \quad \omega_{64}(\tau, \tau^*) = 1. \quad (10)$$

Thus row 38 is one half of the cyclic dual contractible clock doublet, not an unnamed discarded equation. At block level, ι , π , and S have respectively 86, 86, and 19 nonzero row-pair blocks; the vanishing in (10) is checked against those complete support ledgers.

The based classical BV vector field is known through cubic Taylor order,

$$q = q_1 + q_2 + q_3 + O(4), \quad (11)$$

in the repository’s suspended graded-symmetric factorial convention. In this convention every q_n and ℓ_n has cohomological degree +1. The full tensors are action-derived, support-local, typed cyclic, and equivariant under the background-fixing helical generator

$$K_{\text{Berger}} = D - \omega R. \quad (12)$$

Raw cylinder translation D is affine at the rotating background and is not substituted for K_{Berger} .

The result combines two dependency layers:

Layer	Certified content	Explicit limitation
LOCAL-ALGEBRAIC	exact q_2, q_3, ℓ_2, ℓ_3 , identities and cyclicity	no residual branch interpretation
LORENTZIAN-CAUSAL	unary Green homotopies and K -Cartan support through arity three	no Hadamard products or QME

Neither layer is silently promoted to RESIDUAL-TRANSFERRED or LORENTZIAN-CERTIFIED quantum status.

3 Ternary homological transfer

Proposition 3.1 (Formal cyclic transfer through arity three). *Let a cyclic L_∞ algebra be given through q_3 , and let (π, ι, S) be a cyclic SDR satisfying (5)–(6) and (7). The rooted-tree formulas (14)–(17) define cyclic operations through ternary order and imply*

$$[\ell_1, \ell_3] + \frac{1}{2}[\ell_2, \ell_2] = 0. \quad (13)$$

This is the standard cyclic homotopy-transfer theorem [3, 4]. Its application is the mathematical argument for the identities. The sparse calculations below are a separate implementation certificate: they test that the declared signs, PBW normalization, carrier maps, and typed pairing actually realize the formal theorem.

The first two transferred Taylor operations are

$$\ell_1 = \pi q_1 \iota, \quad (14)$$

$$\ell_2 = \pi q_2(\iota, \iota). \quad (15)$$

Define the second Taylor component of the inclusion by

$$I_2 = -Sq_2(\iota, \iota). \quad (16)$$

Then the ternary transfer formula is

$$\ell_3 = \pi q_3(\iota, \iota, \iota) + \sum_{\sigma \in \text{Sh}(2,1)} \epsilon(\sigma) \pi q_2(I_2(\iota_{\sigma(1)}, \iota_{\sigma(2)}), \iota_{\sigma(3)}). \quad (17)$$

The first term is the contact contribution. The second line is the exchange contribution $q_2 Sq_2$.

Lemma 3.2 (Exact mixed exchange support for the frozen SDR). *For the mixed gravity–Maxwell ternary operation, the gravity-outer/mixed-inner exchange channel contains 144 outer/inner co-efficient pairs. The three graded unshuffles produce 324 signed contributions and 342 canonical full-complex PBW coefficients. Every one has full output row 38 = Θ^* . The retained projection has no support on that row, so this channel vanishes after projection. The mixed-outer/gravity-inner and mixed-outer/mixed-inner channels have no outer/inner pairs.*

Proof. The exact consumer reconstructs the 96 gravity and 12 mixed coefficients of I_2 , including their full-row supports, before composing them with q_2 . Outer Leibniz expansion can split one signed unshuffle contribution among several derivative distributions before canonical collection; this is why 324 signed contributions yield 342 PBW coefficients. Direct application of π annihilates the 342 row-38 coefficients. As a negative control, adding the single mutant entry

$$\pi_{\text{mut}}(\text{retained row 0, full row 38}) = 1 \quad (18)$$

exposes all 342 coefficients. Thus the zero is caused by the certified projection and is not an empty table hard-coded by the consumer. $\square$

Corollary 3.3 (Contact form for the frozen retained coordinates). *The retained mixed exchange contribution is zero and*

$$\ell_3^{\text{mixed}} = \pi q_3^{\text{mixed}}(\iota, \iota, \iota). \quad (19)$$

This equality is a computed property of the declared contraction, not a general truncation of the homotopy-transfer formula. In particular, a different cyclic homotopy or cyclic L_∞ coordinate change may redistribute a representative between contact and exchange trees.

4 Frozen-SDR nonvanishing and its audit

For a retained basis vector u , write $[\ell_3(x, y, z)]_u^{(0)}$ for the exact coefficient of u with the zero PBW derivative word in every input slot.

Theorem 4.1 (Nonzero retained mixed bracket for the frozen cyclic SDR). *On the frozen positive Berger-clock gravity–Maxwell BV complex, the typed cyclic contraction (5) transfers the action-derived Taylor data through ternary order. Its mixed ternary representative is nonzero, with the two exact evaluations*

$$[\ell_3(\widehat{h}_{13}, A_1, A_3)]_{\widehat{h}_{00}^*}^{(0)} = +1, \quad (20)$$

$$[\ell_3(\widehat{h}_{01}, \widehat{h}_{03}, A_3)]_{A_1^+}^{(0)} = -1. \quad (21)$$

The first is a gravity-equation output with one gravity and two Maxwell inputs; the second is a Maxwell-equation output with two gravity and one Maxwell input. Pairing the outputs with $\widehat{h}_{00}$ and A_1 , respectively, makes both witnesses explicit coefficients in the lowered g, g, A, A quartic sector. All mixed gravity–Maxwell exchange channels contributing to this declared ℓ_3^{mixed} vanish for the frozen SDR by Lemma 3.2. The operations obey the arity-three L_∞ identity (13) and cyclicity under the retained pairing.

Proof. The two displayed coefficients are read from the exact independently replayed retained tensor; their output rows are 13 and 32, and their input rows are (9, 28, 30) and (4, 6, 30). All inputs have degree zero, both outputs have degree one, and the pairing partners are retained rows 3 and 28. Hence the evaluations are typed and nonzero over $\mathbb{Q}(\sqrt{10})$ without reference to a total coefficient count. Lemma 3.2 proves the scoped exchange statement. Proposition 3.1 and the certified cyclic SDR give the abstract L_∞ and cyclic conclusions. $\square$

The gravity labels in Equations (20) and (21) denote dressed retained coordinates. Their exact full-carrier inclusion columns, including every nonminimal gauge-fixing tail, are printed in Table 2. The Maxwell columns are identity columns. This prevents the witness notation from being mistaken for a raw-coordinate inclusion.

Corollary 4.2 (Exact sparse implementation audit). *The independent PBW consumer realizes Theorem 4.1 with the following basis-level audit data.*

1. *The retained coupled binary operation ℓ_2 contains 1,474 canonical coefficients.*
2. *The retained mixed ternary operation ℓ_3 contains 25,950 canonical coefficients on 18 nonzero output rows.*
3. *All mixed gravity–Maxwell exchange channels contributing to the declared mixed ℓ_3 vanish for the support reason in Lemma 3.2.*

4. Direct coefficient comparison of the retained arity-three identity closes on all 36 rows:

$$[\ell_1, \ell_3] + \frac{1}{2}[\ell_2, \ell_2] = 0. \quad (22)$$

5. The mixed bracket has 7,614 gravity-output coefficients with two Maxwell inputs and 18,336 Maxwell-output coefficients with one Maxwell input.

A localized one-coefficient mutation of ℓ_3 creates two exact arity-three defects.

The audit is obtained as follows. Equations (15)–(17) are evaluated in exact sparse PBW normal form over $\mathbb{Q}(\sqrt{10})$. The independent consumer parses all 59,598 coefficients of the full mixed q_3 payload and reproduces all 25,950 retained contact coefficients with no missing, extra, or changed terms. Lemma 3.2 evaluates every possible mixed exchange channel. Direct comparison of (22) gives zero defects on all rows. The input-sector and output-sector ledgers give the two sector counts. Finally, the mutation changes one retained coefficient and produces two normalized defects, preventing acceptance by a consumer that merely assumes the expected answer.

The numbers in Corollary 4.2 are reproducibility metadata, not invariant content: they depend on the retained basis, row ordering, PBW and integration-by-parts normalization, and the chosen SDR. The invariantly stronger questions in Equation (2) remain open. In particular, Theorem 4.1 is weaker than a residual particle vertex because $\mathcal{R}_{36}$ is retained, not minimal or decomposed into its dynamical branches.

5 Independent cyclic implementation audits

Lowering the output of ℓ_3 with the odd pairing gives a four-linear functional. For degree-zero retained inputs, cyclic transposition of the first input requires formal adjunction of every PBW derivative through the invariant volume form. If w is a derivative word, the exact formal adjoint distributes its letters over the other three slots with sign $(-1)^{|w|}$ and the corresponding multinomial multiplicities.

Proposition 5.1 (Independent degree-zero lowered cyclicity). *Every degree-zero coefficient of the lowered retained ℓ_3 satisfies exact cyclic transposition. There are*

$$25,662 = 7,506_{\text{gravity output}} + 18,156_{\text{Maxwell output}} \quad (23)$$

such coefficients, and the defect count is zero on every row. Replacing the Maxwell field-equation weight two by weight one produces 17,108 defects on 14 rows.

Proof. The verifier constructs both sides independently from the retained tensor, the typed pairing, and exact PBW integration by parts. The signed pairing orientations are retained as a separate convention ledger; only their absolute component multiplicities enter the degree-zero field-equation transpose. Equality holds coefficientwise. The weight mutation changes no coefficient of ℓ_3 itself and therefore tests the lowering convention rather than the imported interaction. $\square$

The remaining

$$25,950 - 25,662 = 288 \quad (24)$$

coefficients are the ghost/antifield completion. Their retained output rows are

$$23, 24, 25, 26, 32, 33, 34.$$

If x is the transposed first input and u is the coordinate paired with the output, the suspended-Darboux sign is

$$(-1)^{|x||u|+\epsilon_2(x)+\epsilon_2(u)}, \quad \epsilon_2(v) = \begin{cases} 1, & \deg v = 2, \\ 0, & \text{otherwise.} \end{cases} \quad (25)$$

Here the degree-two rows are precisely the retained ghost-antifield coordinates.

Proposition 5.2 (Independent full-BV quartic cyclicity). *The complete 25,950-coefficient lowered retained ℓ_3 satisfies exact cyclic transposition. Within the 288-coefficient completion, 120 coefficients carry the positive transpose sign and 168 carry the negative sign. The exact defect count is zero. Omitting the degree-two polarization term exposes 132 defects on all seven completion output rows.*

This proposition is a full-BV statement about the pinned classical retained tensor. It does not imply full BV local cohomology, a quantum master equation, or a Lorentzian quantum theory.

6 Causal compatibility through ternary order

The causal statement is not inferred from finite PBW tables. It is supplied by the independently certified unary Green chain contractions and the coupled cyclic Cartan theorem, extending the Berger-clock construction of the companion paper [8]. If

$$\Lambda_{36,\pm} = \Lambda_{26,\pm}^{\text{grav+clock}} \oplus \Lambda_{M,\pm}, \quad \Lambda_{M,\pm} = W_M G_{P_M,\pm}, \quad (26)$$

then the certified complete-carrier lift is

$$\Lambda_{64,\pm} = S + \iota \Lambda_{36,\pm} \pi, \quad q_1 \Lambda_{64,\pm} + \Lambda_{64,\pm} q_1 = \text{id}_{64}. \quad (27)$$

The side conditions imply the explicit retained bridge

$$\pi \Lambda_{64,\pm} \iota = \Lambda_{36,\pm}. \quad (28)$$

The advanced and retarded homotopies have the corresponding one-sided causal supports and formal-adjoint reversal. Equivariance is not inferred from the background label: the exact carrier ledgers verify

$$K_{64} \iota = \iota K_{36}, \quad \pi K_{64} = K_{36} \pi, \quad K_{64} S = S K_{64}. \quad (29)$$

Let $\iota_K^{(1)}$ be the unary Cartan primitive. The higher sources are

$$A_K^{(2)} = [q_2, \iota_K^{(1)}] - L_K^{(2)}, \quad (30)$$

$$A_K^{(3)} = [q_3, \iota_K^{(1)}] + [q_2, \iota_K^{(2)}] - L_K^{(3)}. \quad (31)$$

At the frozen background $L_K^{(2)} = L_K^{(3)} = 0$. The L_∞ identities, graded Jacobi identity, and K -equivariance give

$$[q_1, A_K^{(2)}] = [q_1, A_K^{(3)}] = 0. \quad (32)$$

Raw causal primitives are cyclicized with the exact Reynolds projectors

$$\text{Cyc}_3 = \frac{1 + \tau + \tau^2}{3}, \quad \text{Cyc}_4 = \frac{1 + \tau + \tau^2 + \tau^3}{4}. \quad (33)$$

Theorem 6.1 (Cyclic causal Cartan compatibility). *The coupled 64-row complex and its typed 36-row retained contraction obey the K_{Berger} Cartan identities through arity three. The higher cyclic primitives satisfy*

$$\text{supp}(\iota_K^{(n)}(f_1, \dots, f_n)) \subseteq J\left(\bigcup_i \text{supp } f_i\right), \quad n \leq 3. \quad (34)$$

The Taylor operations are support-local and no spatial inverse or mode projector enters the construction.

The higher cyclic primitives use the two-sided causal hull. They are not claimed to be separately retarded or advanced; the unary homotopies retain that stronger one-sided property. Equations (27)–(29) make the bridge between the complete and retained unary causal data explicit. They do not turn the support-local algebraic homotopy S used in (17) into a causal propagator. Thus Theorem 6.1 is a compatibility result parallel to Theorem 4.1, not a construction of ℓ_3 by causal integration. It does not construct interacting retarded products, a Hadamard state, or a Lorentzian quantum master equation.

7 Residual descent and the projector obstruction

The next calculation determines what the surviving tensor means physically. On any admissible enlarged or alternative carrier, the dynamical sector must contain

$$\mathcal{R}_{\text{dyn}} = \mathcal{R}_{\text{Einstein}} \oplus \mathcal{R}_{\text{extra Weyl}} \oplus \mathcal{R}_{\text{Maxwell}}, \quad (35)$$

together with exact inclusion, projection, pairing, parity, real structure, and K_{Berger} weights. The mixing entries are then

$$M^a_{bcd} = \pi_{\text{dyn}}^a \ell_3(\iota_b^{\text{dyn}}, \iota_c^{\text{dyn}}, \iota_d^{\text{dyn}}). \quad (36)$$

Every claimed zero must carry an exact support or cancellation witness, and graded symmetry and cyclicity must be replayed in branch coordinates.

The canonical attempt on $\mathcal{R}_{36}$ has received a binary negative verdict. With the Einstein image fixed by the certified rough tensor-wave branch, the exact endpoint

$$A_{10} = \square_2^2 + V_2, \quad \text{ord}(V_2) \leq 2, \quad (37)$$

has 92 nondivisible degree-two remainder entries. A normalized first witness is

$$\frac{71p_1^2 + 71p_2^2 + 9p_3^2}{80} \mod (-p_0^2 + p_1^2 + p_2^2 + p_3^2), \quad (38)$$

and the normalized functional $(80/71)[p_1^2]$ evaluates it to one. Hence no canonical finite-order support-local same-bundle complementary Einstein-like/extra-Weyl projector of the declared type exists on $\mathcal{R}_{36}$.

This is a scoped obstruction, not a global no-go. Mixed-bundle mapping cylinders, filtered carriers, higher-rank local carriers, and explicitly REDUCED-MODE nonlocal splittings remain open. Exact symbol counting requires at least four additional BV rows; the smallest natural support-local candidate adds a spatial STF2 prolongation and its cyclic dual, giving rank 46.

That candidate has now been constructed. Let T_{STF} extract

$$h_{12}, \quad h_{13}, \quad h_{23}, \quad h_{11} - h_{22}, \quad h_{11} + h_{22} - 2h_{33}, \quad (39)$$

and set $F = T_{\text{STF}}\square_2$. Introducing $Y = Fh$ and its five cyclic-dual rows gives degree ranks $(4, 19, 19, 4)$. The metric Hessian block is

$$\begin{pmatrix} A_{10} + F^\dagger F & -F^\dagger \\ -F & I_5 \end{pmatrix}, \quad (40)$$

whose Schur complement is exactly A_{10} .

Proposition 7.1 (Exact rank-46 STF2 graph carrier). *The finite-order cotangent shear defining Equation (40) and its inverse produce a cyclic strong deformation retract*

$$(\mathcal{R}_{46}, q_1^{46}, \omega_{46}) \xrightleftharpoons[\iota_{36}^{46}]{\pi_{46}^{36}} (\mathcal{R}_{36}, \ell_1, \omega_{36}), \quad S_{46}, \quad (41)$$

with

$$\begin{aligned} (q_1^{46})^2 &= 0, & \pi_{46}^{36} \iota_{36}^{46} &= I_{36}, & q_1^{46} S_{46} + S_{46} q_1^{46} &= I_{46} - \iota_{36}^{46} \pi_{46}^{36}, \\ S_{46}^2 &= 0, & S_{46} \iota_{36}^{46} &= 0, & \pi_{46}^{36} S_{46} &= 0. \end{aligned}$$

Both chain maps, the induced pairing, unary cyclicity, and homotopy cyclicity hold exactly over $\mathbb{Q}(\sqrt{10})$. The added STF2 complement is contractible, so this construction changes neither the retained cohomology nor, by itself, the physical branch content.

This is the honest enlarged carrier, not yet the branch theorem. It does not supply an Einstein-like/extra-Weyl projector, a q_2, q_3 lift, K_{Berger} equivariance, or causal support on $\mathcal{R}_{46}$. Equation (36) therefore remains a target formula rather than an authorized rank-46 calculation.

The receiving contract also reserves a separate deformation/vertex payload. That deformation complex, its normalized basis, and its Euler–Lagrange map are neither defined nor used in this paper. Their rank and topological statements are therefore deferred to the residual-mixing paper rather than introduced as a second theorem here. In particular, no deformation class is relabelled as a third dynamical particle branch.

The obstruction neither invalidates Theorem 4.1 nor computes a branch mixing table. Together with the graph-carrier construction it sharpens the next gate to a binary rank-46 projector verdict: construct the exact branch projector or return a normalized obstruction, before lifting the nonlinear brackets.

8 Interpretation and nonclaims

At representative level, the retained classical tensor contains both a gravity-equation response to one gravity and two Maxwell inputs and a Maxwell-equation response to two gravity and one Maxwell input. After lowering, both are g, g, A, A quartic coefficients. They remain visible under the frozen exact cyclic transfer and satisfy the ternary homotopy identity. Once the rank-46 graph carrier receives an admissible branch projector and nonlinear lift—or another branch carrier supplies them—Equation (36) will decide whether this representative closes on the Einstein-like branch, couples it to the extra-Weyl branch, or vanishes on residual cohomology.

Several stronger conclusions do not follow.

1. The bracket changes no unary kinetic operator. It therefore introduces no new negative kinetic direction at this order, but this is not a unitarity or positive-Hilbert-space theorem.
2. A nonzero retained bracket is not yet a photon or graviton scattering amplitude. Residual projection, causal state selection, and asymptotic or relational observables are still required.

3. Nonvanishing of this coefficient representative proves neither a nonzero operation on ℓ_1 -cohomology nor a nontrivial cyclic deformation class. No obstruction to a cyclic L_∞ field redefinition has been computed, and no invariance under changing SDR is claimed.
4. Deformation/vertex classes are not one-particle states and are not included in the dynamical branch list used here.
5. The vanishing one-particle cohomology of the separate free conformal cylinder problem is not converted here into an interacting vacuum theorem.
6. No loop coefficient, anomaly cancellation, QME restoration, Hadamard state, renormalized product, or quantum interaction has been constructed.
7. No arity-four or convergent all-orders L_∞ or Cartan theorem is claimed.

The appropriate provisional interpretation is therefore a retained deformation/vertex representative rather than a particle interaction. The mixing table and deformation-cohomology calculation are the two distinct next tests.

9 Reproducibility and certificate map

The algebraic theorem is pinned to the classical hardening commit `e99d0c1d39490de5261fc6ca1dc2aeaa0d149655`. Its principal artifacts are

Artifact	Role
BERGER_COUPLED_K_CARTAN_THROUGH_ARITY_THREE	full/retained cyclic causal K -Cartan theorem
BERGER_RETAINED_MIXED_ELL3_INDEPENDENT_ACCEPTANCE	exact independent $64 \rightarrow 36$ transfer and mutation witnesses
BERGER_RETAINED_MIXED_ELL3_PHYSICAL_CYCLICITY	independent degree-zero lowered cyclicity and pairing-weight mutation
BERGER_RETAINED_MIXED_ELL3_FULL_BV_CYCLICITY	independent ghost/antifield completion and full retained BV cyclicity
BERGER_RESIDUAL_MIXED_ELL3_BRANCH_PROJECTION_READINESS_V2	historical fail-closed input contract for the mixing table
BERGER_RETAINED_36_RESIDUAL_BRANCH_LOCAL_PROJECTOR_OBSTRUCTION_V1	normalized obstruction to the canonical local 36-row projector
BERGER_RETAINED_46_STF2_PROLONGATION_BRANCH_CARRIER_V1	exact local cyclic graph carrier; rank-46 branch projector still open

The scoped reproduction commands are

```
PYTHONPATH=. python3 \
  d_quotient_classical/backreacted_clock/\
  berger_coupled_k_cartan_through_arity_three.py --check --guards
PYTHONPATH=. python3 \
  d_quotient_classical/backreacted_clock/\
  verify_berger_coupled_k_cartan_through_arity_three.py
PYTHONPATH=quantum-weyl python3 -m \
  transfer.berger_retained_mixed_ell3_acceptance_certificate --check
PYTHONPATH=quantum-weyl python3 -m \
  transfer.verify_berger_retained_mixed_ell3_acceptance
```

```

PYTHONPATH=quantum-weyl python3 -m \
    transfer.berger_retained_mixed_ell3_physical_cyclicity_certificate --check
PYTHONPATH=quantum-weyl python3 -m \
    transfer.verify_berger_retained_mixed_ell3_physical_cyclicity
PYTHONPATH=quantum-weyl python3 -m \
    transfer.berger_retained_mixed_ell3_ghost_antifield_cyclicity_certificate --check
PYTHONPATH=quantum-weyl python3 -m \
    transfer.verify_berger_retained_mixed_ell3_ghost_antifield_cyclicity
PYTHONPATH=quantum-weyl python3 -m \
    transfer.berger_residual_ell3_branch_projection_readiness_v2 --check
PYTHONPATH=quantum-weyl python3 -m \
    transfer.verify_berger_residual_ell3_branch_projection_readiness_v2
PYTHONPATH=. python3 \
    d_quotient_classical/backreacted_clock/\
    verify_berger_retained_36_residual_branch_local_projector_obstruction.py
PYTHONPATH=. python3 \
    d_quotient_classical/backreacted_clock/\
    berger_retained_46_stf2_prolongation_branch_carrier.py --check --guards
PYTHONPATH=. python3 \
    d_quotient_classical/backreacted_clock/\
    verify_berger_retained_46_stf2_prolongation_branch_carrier.py

```

The companion machine-readable claim map records the dependency hashes, positive claims, explicit nonclaims, and the fact that this manuscript is a working draft rather than a theorem freeze.

10 Outlook

Paper 11 now has two mathematical continuations. The first is to decide the rank-46 STF2 branch-projector-or-obstruction gate, materialize the cyclic q_2, q_3 lift if that gate is positive, and compute Equation (36). A nonzero Einstein/extra-Weyl entry would turn the additional Weyl physics into an explicit interaction channel; a block-diagonal table would establish nonlinear closure of the Einstein-like direction through this order; and a central topological action would isolate the vertex theory from the dynamical carrier. The second is to test whether the selected mixed representative defines a nonzero cyclic deformation class or operation on ℓ_1 -cohomology.

The algebraic theorem does not need to wait for that verdict. What the mixing table changes is its physical interpretation, not the existence of the nonzero mixed representative for the frozen cyclic SDR.

A Exact inclusion columns for the displayed witnesses

In Table 2, $\alpha = (\alpha_0, \alpha_1, \alpha_2, \alpha_3)$ denotes the ordered PBW derivative word in the four-generator order of the carrier certificate. The table is generated deterministically from the complete 64×36 inclusion record; it contains 106 exact operator coefficients. The five columns shown are the three dressed gravity inputs used by Equations (20)–(21), the gravity-equation output, and its dressed pairing partner.

Table 2: Exact full-carrier inclusion columns for every dressed gravity coordinate used by the two headline witnesses or their pairing partners. Each row denotes the term $c\partial^\alpha e_{\text{full}}$ in $\iota(e_{\text{retained}})$.

Retained ID	Row	Full-carrier ID	α	Exact c
h_hat_13	9	h_hat_13	(0, 0, 0, 0)	1
		bar_c_star_diff_1	(2, 0, 0, 1)	-1
		bar_c_star_diff_1	(0, 2, 0, 1)	1/3
		bar_c_star_diff_1	(0, 1, 1, 0)	-11*sqrt(10)/45
		bar_c_star_diff_1	(0, 0, 2, 1)	1
		bar_c_star_diff_1	(0, 0, 0, 1)	-475/48
		bar_c_star_diff_1	(0, 0, 0, 3)	1
		bar_c_star_diff_2	(0, 0, 0, 0)	-79391*sqrt(10)/43200
		bar_c_star_diff_2	(2, 0, 0, 0)	-31*sqrt(10)/60
		bar_c_star_diff_2	(0, 2, 0, 0)	11*sqrt(10)/30
		bar_c_star_diff_2	(0, 1, 1, 1)	-2/3
		bar_c_star_diff_2	(0, 0, 2, 0)	11*sqrt(10)/90
		bar_c_star_diff_2	(0, 0, 0, 2)	9*sqrt(10)/5
		bar_c_star_diff_3	(2, 1, 0, 0)	-1
		bar_c_star_diff_3	(0, 1, 0, 0)	-329/2160
		bar_c_star_diff_3	(0, 3, 0, 0)	1
		bar_c_star_diff_3	(0, 1, 2, 0)	1
		bar_c_star_diff_3	(0, 1, 0, 2)	1/3
		bar_c_star_diff_3	(0, 0, 1, 1)	2*sqrt(10)/45
		bar_c_star_tau	(1, 1, 0, 1)	-2/3
		bar_c_star_tau	(1, 0, 1, 0)	-71*sqrt(10)/180
		bar_c_star_sigma	(2, 1, 0, 1)	1/3
		bar_c_star_sigma	(2, 0, 1, 0)	71*sqrt(10)/360
		bar_c_star_sigma	(0, 3, 0, 1)	-1/3
		bar_c_star_sigma	(0, 2, 1, 0)	-71*sqrt(10)/360
		bar_c_star_sigma	(0, 1, 2, 1)	-1/3
		bar_c_star_sigma	(0, 1, 0, 1)	34/9
		bar_c_star_sigma	(0, 1, 0, 3)	-1/3
		bar_c_star_sigma	(0, 0, 1, 0)	71*sqrt(10)/81
		bar_c_star_sigma	(0, 0, 3, 0)	-71*sqrt(10)/360
		bar_c_star_sigma	(0, 0, 1, 2)	-13*sqrt(10)/24
h_hat_01	4	h_hat_01	(0, 0, 0, 0)	1
		bar_c_star_diff_1	(1, 0, 0, 0)	2561/720
		bar_c_star_diff_1	(3, 0, 0, 0)	1
		bar_c_star_diff_1	(1, 2, 0, 0)	-1/3
		bar_c_star_diff_1	(1, 0, 2, 0)	-1
		bar_c_star_diff_1	(1, 0, 0, 2)	-1
		bar_c_star_diff_2	(1, 1, 1, 0)	2/3
		bar_c_star_diff_2	(1, 0, 0, 1)	-77*sqrt(10)/60
		bar_c_star_diff_3	(1, 1, 0, 1)	2/3
		bar_c_star_diff_3	(1, 0, 1, 0)	107*sqrt(10)/180
		bar_c_star_tau	(2, 1, 0, 0)	-1/3
		bar_c_star_tau	(0, 1, 0, 0)	-31/9
		bar_c_star_tau	(0, 3, 0, 0)	1
		bar_c_star_tau	(0, 1, 2, 0)	1
		bar_c_star_tau	(0, 1, 0, 2)	1
		bar_c_star_tau	(0, 0, 1, 1)	31*sqrt(10)/30
		bar_c_star_sigma	(3, 1, 0, 0)	-1/3
		bar_c_star_sigma	(1, 1, 0, 0)	-31/27
		bar_c_star_sigma	(1, 3, 0, 0)	1/3
		bar_c_star_sigma	(1, 1, 2, 0)	1/3
h_hat_03	6	h_hat_03	(0, 0, 0, 0)	1
		bar_c_star_diff_1	(1, 1, 0, 1)	2/3
		bar_c_star_diff_1	(1, 0, 1, 0)	-3*sqrt(10)/20
		bar_c_star_diff_2	(1, 1, 0, 0)	3*sqrt(10)/20
		bar_c_star_diff_2	(1, 0, 1, 1)	2/3
		bar_c_star_diff_3	(1, 0, 0, 0)	9/80
		bar_c_star_diff_3	(3, 0, 0, 0)	1

Continued on next page

Table 2 (continued)

Retained ID	Row	Full-carrier ID	α	Exact c
		bar_c_star_diff_3	(1, 2, 0, 0)	-1
		bar_c_star_diff_3	(1, 0, 2, 0)	-1
		bar_c_star_diff_3	(1, 0, 0, 2)	-1/3
		bar_c_star_tau	(2, 0, 0, 1)	-1/3
		bar_c_star_tau	(0, 2, 0, 1)	1
		bar_c_star_tau	(0, 0, 2, 1)	1
		bar_c_star_tau	(0, 0, 0, 3)	1
		bar_c_star_sigma	(3, 0, 0, 1)	-1/3
		bar_c_star_sigma	(1, 2, 0, 1)	1/3
		bar_c_star_sigma	(1, 0, 2, 1)	1/3
		bar_c_star_sigma	(1, 0, 0, 3)	1/3
h_hat_star_00	13	h_hat_star_00	(0, 0, 0, 0)	1
h_hat_00	3	h_hat_00	(0, 0, 0, 0)	1
		bar_c_star_diff_1	(2, 1, 0, 0)	-1/2
		bar_c_star_diff_1	(0, 1, 0, 0)	71/480
		bar_c_star_diff_1	(0, 3, 0, 0)	1/6
		bar_c_star_diff_1	(0, 1, 2, 0)	1/6
		bar_c_star_diff_1	(0, 1, 0, 2)	1/6
		bar_c_star_diff_2	(2, 0, 1, 0)	-1/2
		bar_c_star_diff_2	(0, 2, 1, 0)	1/6
		bar_c_star_diff_2	(0, 1, 0, 1)	$-\sqrt{10}/20$
		bar_c_star_diff_2	(0, 0, 1, 0)	-3/160
		bar_c_star_diff_2	(0, 0, 3, 0)	1/6
		bar_c_star_diff_2	(0, 0, 1, 2)	1/6
		bar_c_star_diff_3	(2, 0, 0, 1)	-1/2
		bar_c_star_diff_3	(0, 2, 0, 1)	1/6
		bar_c_star_diff_3	(0, 0, 2, 1)	1/6
		bar_c_star_diff_3	(0, 0, 0, 1)	3/160
		bar_c_star_diff_3	(0, 0, 0, 3)	1/6
		bar_c_star_tau	(3, 0, 0, 0)	1/2
		bar_c_star_tau	(1, 2, 0, 0)	-5/6
		bar_c_star_tau	(1, 0, 2, 0)	-5/6
		bar_c_star_tau	(1, 0, 0, 2)	-5/6
		bar_c_star_sigma	(2, 2, 0, 0)	1/6
		bar_c_star_sigma	(2, 0, 2, 0)	1/6
		bar_c_star_sigma	(2, 0, 0, 2)	1/6
		bar_c_star_sigma	(0, 2, 0, 0)	-1/3
		bar_c_star_sigma	(0, 4, 0, 0)	-1/6
		bar_c_star_sigma	(0, 2, 2, 0)	-1/3
		bar_c_star_sigma	(0, 2, 0, 2)	-1/3
		bar_c_star_sigma	(0, 1, 1, 1)	$\sqrt{10}/10$
		bar_c_star_sigma	(0, 0, 2, 0)	1/3
		bar_c_star_sigma	(0, 0, 4, 0)	-1/6
		bar_c_star_sigma	(0, 0, 2, 2)	-1/3
		bar_c_star_sigma	(0, 0, 0, 2)	-3/40
		bar_c_star_sigma	(0, 0, 0, 4)	-1/6

References

- [1] I. A. Batalin and G. A. Vilkovisky, *Gauge algebra and quantization*, Phys. Lett. B **102** (1981), 27–31.
- [2] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*, Princeton University Press, 1992.
- [3] H. Kajiura, *Noncommutative homotopy algebras associated with open strings*, Rev. Math. Phys. **19** (2007), 1–99.

- [4] A. S. Arvanitakis, O. Hohm, C. Hull, and V. Lekeu, *Homotopy transfer and effective field theory I: tree level*, arXiv:2007.07942.
- [5] D. Fiorenza and M. Manetti, *L_∞ algebras, Cartan homotopies and period maps*, arXiv:math/0605297.
- [6] A. Kraft and J. Schnitzer, *An introduction to L_∞ -algebras and their homotopy theory*, arXiv:2207.01861.
- [7] M. Benini, G. Musante, and A. Schenkel, *Green hyperbolic complexes on Lorentzian manifolds*, Commun. Math. Phys. (2023), doi:[10.1007/s00220-023-04807-5](https://doi.org/10.1007/s00220-023-04807-5).
- [8] GPT-5.6.sol, *A backreacting phase clock with fixed momentum in pure-Weyl gravity: fixed-coupling rigidity and causal BV Cartan analysis of the helical stabilizer*, companion manuscript, Paper IX (2026).