

Extra Fourth-order Axial Modes under a Direct Lee--Wald Audit

Adjacency bridge note

Bridge note for conformal-gravity, higher-derivative quantization, and Einstein-from-conformal researchers. Status: compact classical theorem, 17 July 2026.

1. Their object and the unresolved question

Three adjacent programmes ask different questions. Maldacena uses a Neumann boundary condition in asymptotically de Sitter or Euclidean anti-de Sitter settings to select the Einstein solution in a tree-level computation ([arXiv:1105.5632](#)). Mannheim argues for a positive PT -based inner product in higher-derivative gravity ([arXiv:2109.12743](#)). Kubo and Kuntz use a BRST/Fock construction and find an indefinite transverse spin-two sector and scattering-unitarity failure ([arXiv:2202.08298](#)). These results concern different boundaries, state spaces, and inner products.

Our narrower classical question is prior to all three quantum conclusions: on one compact common Einstein--Maxwell/Weyl--Maxwell background, are the extra fourth-order roots actual gauge-quotient solution directions, are they radical under the action-derived current, and where does the negative classical current direction lie?

```
H^0(Einstein--Maxwell) --injects--> H^0(Weyl--Maxwell)
|                                     |
| standard image                     +--> Q_extra = two axial directions
|                                     |
+----- direct Lee--Wald matrix -----+
      image signs (1,1), extra signs (2,0)
      full axial signature (3,1)
```

2. Exact dictionary

Item	Adjacent usage	This project	Conversion or caveat
Theory	pure conformal gravity or Einstein branch	tuned Weyl--Maxwell versus Einstein--Maxwell	Matter and compact flux are part of the fixture.
Background	dS/EAdS, flat Fock vacuum, or higher-derivative vacuum	$\mathbb{R} \times S^1 \times S^2$ product with fixed $U(1)$ bundle	No conformal continuation to an asymptotic vacuum is assumed.
Extra mode	fourth-order pole/Jordan/Fock excitation	quotient $Q_{\text{extra}} = H_{WM}^0 / i_* H_{EM}^0$	Quotient solution directions are not yet particles.
Gauge	BRST physical-state quotient or boundary selection	local $\text{Diff} \times \text{Weyl} \times U(1)$, before final residual quotient	Boundary selection is not identified with gauge quotienting.
Form	Dirac, PT , or Fock inner product	classical covariant Lee--Wald current	A classical sign is not a quantum norm.
Selection	boundary condition or quartet mechanism	compact harmonic regularity and fixed bundle	Causal boundary admissibility is open.

3. Reproduced benchmark

The current engine first reproduces the independent Einstein--Maxwell Lee--Wald current, the repository's Bach convention, a pure-Weyl gauge kernel, the flat transverse-traceless zero-restriction control, and exact current conservation. It then compares the reduced self-adjoint Hessian current with a literal variation of the four-dimensional Weyl--Maxwell curvature-momentum current. For arbitrary $\ell \geq 2$ and spherical multiplicity m , their integrated currents agree up to the positive harmonic normalization $N_{\ell m}$.

The replay commands are:

```
python3 -m bridge.einstein_sector.einstein_maxwell_weyl_axial_lee_wald_completion \
--verify bridge/certificates/einstein_maxwell_weyl_axial_lee_wald_completion.json
python3 -m bridge.einstein_sector.weyl_maxwell_axial_general_lee_wald_fixture \
--verify bridge/certificates/weyl_maxwell_axial_general_lee_wald_fixture.json
```

4. Added result

Generic compact axial Lee–Wald theorem. For each physical compact axial harmonic with $\ell \geq 2$, the Weyl–Maxwell solution module strictly contains the Einstein–Maxwell image by two exact polarizations. The direct four-dimensional Lee–Wald form is nondegenerate on the extra quotient with signature $(2, 0)$. The complete generic axial target has signature $(3, 1)$, and its negative direction lies on one Einstein-image master branch rather than on either extra direction.

Algebraically, the invariant factors separate the standard Einstein master polynomial from a doubled extra primary factor. Direct mixed Lee–Wald blocks between the standard and extra primary shells vanish. The extra determinant is positive throughout the physical $\ell \geq 2$ domain. This rules out three simple explanations of the extra root on this fixture: determinant multiplicity only, local gauge, and a presymplectic radical.

The nonlinear compact test adds an important qualification. The real $\ell = 2, k = 0$ extra span has a definite Taub obstruction at fixed bundle charges, so the linearly measurable extra directions are linearization unstable in that declared sector. This is selection by nonlinear constraint, not disappearance from the linear current.

5. Consequence in their language

The result does not choose between PT , Fock–BRST, or boundary-selected quantization. It says that any comparison must account for an explicit classical carrier space on which “extra branch” and “negative direction” are not synonymous. A successful boundary or quantum prescription must explain which of the standard and extra compact directions survive its own quotient and why its inner product is the transported form appropriate to that state space.

For Einstein-from-conformal work, this is a Lorentzian/nonlinear complement: linear Einstein inclusion holds, but the identity inclusion is not symplectic and selected real tangents fail the second-order compact Taub test. That does not refute the EAdS/dS boundary theorem; it asks whether its selection is causally preserved and symplectically appropriate in real time.

6. Scope boundary

The theorem is **LOCAL-ALGEBRAIC/REDUCED-MODE** on a compact product background. It is not a positive-frequency Hilbert norm, a BRST/Fock theorem, a PT metric comparison, a quantum ghost result, or an S-matrix statement. The polar extra branch, final residual quotient, literal second expansion of the four-dimensional action density, and causal/asymptotic boundary phase spaces remain open. The compact negative direction may be removed, retained, or reinterpreted by those later gates.

7. One useful question for adjacent experts

Is there an admissible Lorentzian boundary or BRST selection that removes the negative Einstein-image master direction while treating the two positive extra directions consistently, and can that selection be expressed as a causal symplectic subquotient rather than a condition imposed at both temporal ends?

Reproducibility receipt

```
source papers: arXiv:1105.5632v2; arXiv:2109.12743v1; arXiv:2202.08298v2
project certificate: EINSTEIN_MAXWELL_WEYL_AXIAL_LEE_WALD_COMPLETION
verification: commands in section 3
dependency tags: LOCAL-ALGEBRAIC; REDUCED-MODE
generality level: G2_GENERIC_AXIAL_DIRECT_LEE_WALD_COMPLETION
lifecycle state: CERTIFIED_COMPACT_CLASSICAL_CURRENT
claim flag: EXTRA_AXIAL_NONRADICAL_DIRECT_LEE_WALD
open fields: polar extra; residual descent; causal boundaries; quantum state
```