

Taub obstruction and balanced extension in compact Weyl--Maxwell gravity

Pure extra modes versus Einstein--extra mixtures

GPT-5.6.sol

Theorem-frozen version, 17 July 2026

Working manuscript, 17 July 2026. The theorem-frozen scope is **LOCAL-ALGEBRAIC** and **REDUCED-MODE**. Circulation remains conditional on the documented final human review described below.

Abstract

Linearized Weyl--Maxwell gravity on the compactified magnetically supported Plebański--Hacyan universe contains the ordinary Einstein--Maxwell modes and additional Weyl modes. We determine whether the additional generic modes are tangent to nonlinear solutions on the fixed compact magnetic bundle. On the closed Cauchy surface $S^1 \times S^2$, the quadratic Taub pairing with a background stabilizer equals the corresponding covariant Lee--Wald moment map. The time-translation pairing is negative definite on the complete real pure-extra generic sector, axial and polar, for every $\ell \geq 2$ and every allowed compact momentum. Consequently no nonzero pure-extra generic tangent admits a fixed-bundle second-order extension.

This obstruction is not stable under mixing with the Einstein sector. The Einstein primary has an indefinite time-translation form. At $\ell = 2, m = 0, k = 0$, an Einstein-minus mode and one extra mode have a unique positive balancing ratio within their declared two-mode span, for which all five background-stabilizer moment maps vanish. We compute its complete quadratic Weyl--Maxwell source. The homogeneous zero-frequency Einstein and extra sources cancel exactly; every remaining homogeneous channel has an explicit correction; and every $\ell = 2, 4$ output is removed by an exact off-shell polar inverse. The four solved action equations and four target Noether identities span all eight ungauged polar equations with determinant (-4) , including at zero frequency. Hence this balanced real tangent has an explicit second-order correction on the same charge fibre.

The result exhibits a sharp nonlinear distinction:

pure extra generic mode: obstructed, balanced Einstein--extra mode: extendible to second order.

The theorem concerns a formal second-order jet, not an exact family or an all-orders solution. Classification of the full mixed moment-map zero cone is left as the next theorem rather than imposed as a prerequisite for the present result.

1. Introduction

Linearization stability asks whether a solution of the linearized field equations is the first derivative of a family of exact solutions. On a compact Cauchy surface, background symmetries can produce quadratic Taub constraints which are invisible in the linear equations. This phenomenon is classical in Einstein gravity and is naturally expressed through the constraint-adjoint kernel or, equivalently on a closed slice, through the moment map for the background symmetry group [1,2]. Higher-derivative gravity has analogous linearization-instability questions [3].

The present problem is unusually clean because the linear phase space is already known exactly. The background

$$M = \mathbb{R}_t \times S_L^1 \times S^2$$

is a compactification of the magnetically supported Plebański–Hacyan direct product [4]. With the normalization specified below, it solves both Einstein–Maxwell and Weyl–Maxwell theory. The complete standard Einstein–Maxwell harmonic tangent injects into the Weyl–Maxwell tangent, while the latter contains two extra generic cyclic summands in each parity. Those extra modes are genuine nonradical linear solutions; they are neither Weyl-gauge representatives nor null directions of the Lee–Wald form. The linear phase-space theorem and its conventions are established in the companion paper [5].

The next question is nonlinear:

Which of the certified extra linear modes satisfy the quadratic integrability constraints, and can an Einstein component cancel their obstruction?

We answer both parts at a sharp, publishable scope.

1. **General pure-extra no-go.** Every nonzero real pure-extra generic tangent is obstructed at second order on the fixed magnetic bundle. This holds for both parities, every physical $\ell \geq 2$, every allowed compact momentum, and finite or rapidly decreasing finite-energy superpositions.
2. **Explicit balanced extension.** One nonzero real Einstein–extra tangent at $\ell = 2, m = 0, k = 0$ annihilates all five stabilizer moment maps and has a complete explicit second-order correction.

The contrast is the scientific point. The extra sector is not deleted from the linear theory, nor is it generically promoted to a nonlinear branch. Rather, the nonlinear solution locus has a singular tangent cone: a pure extra direction fails the quadratic constraint, while an appropriately balanced mixed direction passes and extends through second order.

This paper does **not** delay that conclusion until the entire indefinite mixed cone is classified. The full common zero locus of $(\mu_H, \mu_{P_x}, \mu_{J_1}, \mu_{J_2}, \mu_{J_3})$, including opposite-momentum standing waves and exceptional/global blocks, is the next theorem.

Main theorem

Let $\mathcal{T}_X^{\text{gen}}$ be the real, locally gauge-reduced, generic extra Weyl–Maxwell tangent on the fixed magnetic bundle, and let u_- and u_e be the axisymmetric axial modes defined in Section 7.

Theorem A (obstruction versus balanced extension).

1. For every nonzero $u \in \mathcal{T}_X^{\text{gen}}$, the constant-lapse Taub pairing is nonzero. Therefore the equation

$$L_{\text{WM}}\Phi^{(2)} = -\frac{1}{2}D^2E_{\text{WM}}[u, u]$$

has no fixed-bundle periodic solution.

2. The real tangent

$$\Phi^{(1)} = \text{Re}(u_-) + a_e \text{Re}(u_e), \quad |a_e|^2 = \frac{27}{52}(5\sqrt{3} - 6),$$

annihilates all five background-stabilizer moment maps and admits an explicit real second-order correction $\Phi^{(2)}$ on the same fixed bundle.

Part 1 is a theorem on the complete pure-extra generic sector. Part 2 is an existence theorem for one declared mixed tangent. It is not a claim that every common-zero tangent extends.

2. The common background and the fixed charge fibre

We use signature $(-+++)$ and the actions

$$S_{\text{EM}}[g, A] = \int_M \sqrt{-g} \left[\frac{R - 2\Lambda}{2\kappa} - \frac{1}{4}F_{ab}F^{ab} \right] d^4x,$$

$$S_{\text{WM}}[g, A] = \int_M \sqrt{-g} \left[\frac{\alpha_B}{8}C_{abcd}C^{abcd} - \frac{1}{4}F_{ab}F^{ab} \right] d^4x.$$

The Bach convention is

$$\delta \int \sqrt{-g} C^2 = 4 \int \sqrt{-g} B_{ab} \delta g^{ab} + \text{boundary},$$

so the Weyl–Maxwell metric equation is

$$\alpha_B B_{ab} = T_{ab}, \quad \nabla_a F^{ab} = 0, \quad dF = 0.$$

The fixture is

$$M = \mathbb{R}_t \times S_L^1 \times S^2, \quad d\bar{s}^2 = -dt^2 + dx^2 + d\Omega_2^2, \quad \bar{F} = \text{vol}(S^2),$$

with

$$(\kappa, \Lambda, \alpha_B) = (1, \frac{1}{2}, 3).$$

It lies on the intersection of the Einstein–Maxwell and Weyl–Maxwell solution loci. This is an incidence relation between two theories, not a Weyl gauge equivalence. The local geometry belongs to the direct-product electrovacua studied by Plebański and Hacyan [4]. Exact wave families on related direct-product universes are known [6,7]; their existence neither implies nor is implied by the complete harmonic integrability statement proved here.

The magnetic field is a connection curvature on a nontrivial compact $U(1)$ bundle $P_N \rightarrow S^1 \times S^2$. With the chosen normalization the Chern number is $N = 2$. A connection tangent $a = A - \bar{A}$ is a global one-form on the fixed bundle, but a uniform continuous change of magnetic flux changes the bundle family. We therefore define the nonlinear extension problem on the fixed component P_N .

This distinction is load-bearing. If a formal second-order magnetic coefficient is introduced,

$$N(\epsilon) = 2 + 2\epsilon^2 p,$$

then a smooth family on the same $N = 2$ bundle has $p = 0$. A calculation which removes a Taub source by choosing $p \neq 0$ solves a different, charge-enlarged problem. Electric variation is allowed on fixed P_N , but at the purely magnetic background its linear energy pairing vanishes and it cannot absorb the pure-extra constant-lapse obstruction.

Domain of the theorem

The theorem is made after local

$$\text{Diff} \ltimes \mathcal{G}_{\text{Weyl}} \ltimes \mathcal{G}_{U(1)}$$

reduction, but before quotienting by the five background stabilizers

$$H = \partial_t, \quad P_x = \partial_x, \quad J_1, J_2, J_3 \in \mathfrak{so}(3).$$

These stabilizers are retained because their Hamiltonians are the quadratic constraints. Quotienting them before evaluating the moment map would erase the very obstruction under study.

3. Linear input: Einstein and extra primary sectors

Write

$$\lambda = \ell(\ell + 1), \quad k = \frac{2\pi n}{L}, \quad n \in \mathbb{Z}.$$

For every $\ell \geq 2$, the generic axial and polar Weyl–Maxwell solution modules split into the Einstein q -primary image and two additional p -primary summands. The extra shell is

$$p(\omega, k, \lambda) = \omega^2 - k^2 - \lambda + \frac{2}{3} = 0,$$

so

$$\omega_e^2 = k^2 + \lambda - \frac{2}{3} > 0.$$

The Einstein shell polynomial is

$$q(\omega, k, \lambda) = (\omega^2 - k^2)^2 - 2\lambda(\omega^2 - k^2) + \lambda(\lambda - 2).$$

Its two positive-frequency branches will be denoted ω_- and ω_+ . The Einstein and extra primary modules are orthogonal under the direct four-dimensional Lee–Wald current. Each axial and polar extra Gram matrix G_X^{par} is positive definite. Thus the extra modes are genuine nonnull linear solutions, even though Section 5 proves that they are not fixed-bundle nonlinear tangent directions.

For a positive-frequency coefficient vector c in one branch, parity and spin- ℓ multiplicity block, let W_ℓ be the positive invariant angular Gram form. Reality fixes the negative-frequency and negative-momentum coefficients by conjugation. We use

$$\Phi = \text{Re}(ce^{-i\omega t + ikx})$$

as the real-mode convention.

4. Taub pairings as covariant moment maps

Let $E(\Phi) = 0$ denote the Weyl–Maxwell Euler–Lagrange equations and $L = DE|_{\bar{\Phi}}$. A first-order tangent u extends through second order only if there is a $\Phi^{(2)}$ satisfying

$$L\Phi^{(2)} = -\frac{1}{2}D^2E|_{\bar{\Phi}}[u, u]. \quad (4.1)$$

Every adjoint zero mode ζ_X therefore gives the necessary Taub condition

$$\left\langle \zeta_X, \frac{1}{2}D^2E[u, u] \right\rangle = 0. \quad (4.2)$$

For an infinitesimal bundle-covariant automorphism X of the background, the closed-slice covariant Hamiltonian identity gives

$$\boxed{\left\langle \zeta_X, \frac{1}{2}D^2E_{\text{WM}}[u, u] \right\rangle = \mu_X(u) = \frac{1}{2}\Omega_{\text{WM}}(u, \mathcal{L}_X u).} \quad (4.3)$$

To prove (4.3), differentiate the action Noether-current identity twice, use $E(\bar{\Phi}) = 0$, $Lu = 0$, and $\mathcal{L}_X \bar{\Phi} = 0$, then integrate over the closed Cauchy surface. Every exact Lee–Wald improvement and bundle-patching corner term integrates to zero. This is the boundaryless specialization of the covariant phase-space construction [8,9].

The exact real-mode moment maps are

$$\mu_H = -\frac{L}{4} \sum \omega^2 c^\dagger (G_{\text{branch}} \otimes W_\ell) c, \quad (4.4)$$

$$\mu_{P_x} = \frac{L}{4} \sum k\omega c^\dagger (G_{\text{branch}} \otimes W_\ell) c, \quad (4.5)$$

$$\mu_{J_a} = \frac{L}{4} \sum \omega c^\dagger (G_{\text{branch}} \otimes W_\ell T_a) c. \quad (4.6)$$

The sums are block diagonal in k, ℓ , parity, and frequency shell. Rotations preserve ℓ ; J_3 is diagonal in m , while J_1, J_2 connect only m to $m \pm 1$. Axial–polar and Einstein–extra cross terms vanish. The sign and the factor $1/4$ in (4.4) are fixed by exact agreement with three independent direct tensor calculations: one axial extra block and the axial and polar Einstein–minus fixtures at $\ell = 2, k = 0$.

5. The complete pure-extra obstruction

Theorem 5.1 (pure-extra fixed-bundle no-go). Let u be a real pure-extra generic Weyl–Maxwell tangent, axial or polar, with finite or rapidly decreasing finite-energy harmonic coefficients. If $u \neq 0$, then u admits no periodic second-order correction on the fixed magnetic bundle P_N .

Proof

On the extra shell, $\omega_e^2 > 0$ for every physical $\lambda \geq 6$. Both extra Gram matrices and the angular form W_ℓ are positive definite. Hence every nonzero block contributes strictly negatively to (4.4):

$$\mu_H(u) = -\frac{L}{4} \sum_{\text{extra blocks}} \omega_e^2 c^\dagger (G_X^{\text{par}} \otimes W_\ell) c < 0.$$

Orthogonality makes the same conclusion valid for finite and convergent rapidly decreasing superpositions. Equation (4.2) is therefore violated for the constant-lapse adjoint class.

The obstruction cannot be absorbed inside the declared domain. Continuous magnetic variation changes $c_1(P_N)$, while electric variation has zero linear pairing with H at the purely magnetic background. Thus (4.1) has no fixed-bundle solution. $\square$

What the theorem does not say

The theorem does not remove the extra linear modes. Their Lee–Wald Gram matrices are nondegenerate, so they remain genuine classical linear solutions. The conclusion is instead that the exact fixed-bundle solution locus is singular at the background: its formal tangent space is larger than its second-order tangent cone.

Nor is this a quantum ghost statement. No positive-frequency Hilbert space, BRST-compatible Hadamard state, or Lorentzian causal quantum construction is used.

6. Why an Einstein component can cancel the obstruction

The Einstein q -primary contribution to μ_H is indefinite. In each parity its two master branches contribute opposite signs in the target current convention. Consequently a negative extra contribution can be cancelled by an Einstein–minus component without invoking a mixed Einstein–extra current entry. The cancellation is additive between two orthogonal diagonal primary blocks.

There is nevertheless a useful same-momentum restriction.

Proposition 6.1 (one travelling block at nonzero momentum). In a single fixed $k \neq 0$ travelling block, simultaneous vanishing of μ_H and μ_{P_x} forces all Einstein–plus, Einstein–minus, and extra occupations to vanish.

Indeed, the three shell frequencies obey

$$\omega_- < \omega_e < \omega_+.$$

After eliminating the Einstein-minus occupation from the two scalar moment maps, one obtains

$$\omega_+(\omega_+ - \omega_-)A_+ + \omega_e(\omega_e - \omega_-)A_e = 0,$$

where $A_+, A_e \geq 0$ are Gram-normalized occupations. Both coefficients are strictly positive, so $A_+ = A_e = 0$, and the remaining occupation then vanishes as well. This proposition does not cover cancellations between distinct momenta, such as standing-wave combinations.

At $k = 0$, the momentum constraint vanishes automatically and a nontrivial balance becomes possible.

7. The minimal balanced tangent

Fix the axial $\ell = 2, m = 0, k = 0$ sector. In the coefficient order (H_t, H_x, Q_t, Q_x) , choose the Einstein-minus representative

$$u_- = (0, -2, 0, 2\sqrt{3}), \quad \omega_-^2 = 6 - 2\sqrt{3}, \quad (7.1)$$

and the second extra representative

$$u_e = (0, -\frac{2}{3}, 0, 6), \quad \omega_e^2 = \frac{16}{3}. \quad (7.2)$$

For unit real cosine amplitude, their constant-lapse Taub coefficients are

$$\tau_- = \frac{48}{5}(-6 + 5\sqrt{3}) > 0, \quad \tau_e = -\frac{832}{45} < 0. \quad (7.3)$$

Set

$$|a_e|^2 = \frac{\tau_-}{-\tau_e} = \frac{27}{52}(5\sqrt{3} - 6). \quad (7.4)$$

Then

$$\Phi^{(1)} = \text{Re}(u_- e^{-i\omega_- t}) + a_e \text{Re}(u_e e^{-i\omega_e t}) \quad (7.5)$$

is nonzero and satisfies

$$\mu_H = \mu_{P_x} = \mu_{J_1} = \mu_{J_2} = \mu_{J_3} = 0. \quad (7.6)$$

Here $\mu_H = 0$ follows from (7.3)–(7.4), $\mu_{P_x} = 0$ from $k = 0$, and the rotational expectations vanish for the separate axisymmetric $m = 0$ states. Because the Einstein and extra primaries are symplectically orthogonal, (7.6) is not produced by an interference term.

Vanishing Taub charges is necessary but not sufficient for extension. A nonzero quadratic source can still have a component in another adjoint cokernel or lie on a resonant output shell. We therefore solve the complete second-order equation directly.

8. Complete second-order extension of the balanced tangent

Theorem 8.1 (balanced Einstein–extra second-order extension). The real tangent (7.5) admits a real, periodic, fixed-bundle correction $\Phi^{(2)}$ satisfying

$$L_{\text{WM}}\Phi^{(2)} = -\frac{1}{2}D^2E_{\text{WM}}[\Phi^{(1)}, \Phi^{(1)}]. \quad (8.1)$$

8.1 Selection rules and real-mode factors

The product of two axial $\ell = 2, m = 0$ harmonics is polar and contains only

$$\ell_{\text{out}} = 0, 2, 4.$$

The time dependence produces the five channel types

Channel	Output frequency
Einstein self-sum	$2\omega_-$
extra self-sum	$2\omega_e$
cross-sum	$\omega_e + \omega_-$
cross-difference	$\omega_e - \omega_-$
conjugate self-products	(0)

For $\Phi = \text{Re } z = (z + \bar{z})/2$ and symmetric Hessian B ,

$$\frac{1}{2}B(\Phi, \Phi) = \frac{1}{8}B(z, z) + \frac{1}{4}B(z, \bar{z}) + \frac{1}{8}B(\bar{z}, \bar{z}).$$

Thus self-sums carry factor $1/8$, while self-zero, cross-sum, and cross-difference terms carry factor $1/4$. These factors are replayed symbolically in the certificate rather than inserted as conventions after the calculation.

8.2 The exceptional homogeneous channel

In homogeneous coefficient order (C, K, U) , the directly computed linear operator on rows $(E_{00}, E_{11}, E_{22}, M_1)$ is

$$L_0(\Omega) = \begin{pmatrix} 0 & 0 & 0 \\ -\Omega^4/2 & \Omega^4/2 & 0 \\ \Omega^4/4 & -\Omega^4/4 & 0 \\ 0 & 0 & \Omega^2 \end{pmatrix}. \quad (8.2)$$

At zero frequency the separate real self-products are obstructed:

$$S_-^{(0)} = \frac{-6 + 5\sqrt{3}}{5} \begin{pmatrix} 48 \\ 0 \\ 24 \\ 0 \end{pmatrix}, \quad S_e^{(0)} = -\frac{-6 + 5\sqrt{3}}{5} \begin{pmatrix} 48 \\ 0 \\ 24 \\ 0 \end{pmatrix}. \quad (8.3)$$

The amplitude (7.4) is already included in $S_e^{(0)}$. Therefore

$$S_-^{(0)} + S_e^{(0)} = 0. \quad (8.4)$$

This cancellation is the nonlinear heart of the construction. Each pure component fails the homogeneous constraint, while the balanced combination removes it exactly.

Every nonzero-frequency homogeneous source has

$$S_{00} = S_{M_1} = 0, \quad S_{11} + 2S_{22} = 0.$$

It is removed explicitly by

$$(C, K, U) = \left(\frac{2S_{11}}{\Omega^4}, 0, 0 \right). \quad (8.5)$$

Hence no electric-charge or Wilson-line correction is hidden in the homogeneous solution.

8.3 The $\ell = 2, 4$ polar outputs

Let $H_P(\lambda, k, \Omega)$ be the exact action-normalized polar Hessian in coordinates (A_t, B, C_t, U) . Its determinant is

$$\det H_P = \frac{9}{16} \lambda^3 (\lambda - 2) p(\Omega, k, \lambda)^2 q(\Omega, k, \lambda). \quad (8.6)$$

For every output frequency in the table and for $\lambda = 6, 20$, the exact algebraic preflight proves

$$p \neq 0, \quad q \neq 0.$$

Thus each polar source $S_{\ell, \Omega}$ has the explicit correction

$$\Phi_{\ell, \Omega}^{(2)} = -H_P(\lambda, 0, \Omega)^{-1} S_{\ell, \Omega}, \quad \ell = 2, 4, \quad (8.7)$$

and every stored four-row remainder is identically zero. The certificate retains the exact radical expressions for all corrections; writing the largest cross-channel radicals in the main text would obscure rather than strengthen the argument.

8.4 Completion of the dependent tensor equations

Solving four action rows is sufficient only if the remaining ungauged tensor equations are proved to follow. Use target equation order

$$(A, B, C, h_t, h_x, K, G, U).$$

At $k = 0$, the four certified target Noether identities are the rows of

$$N_0 = \begin{pmatrix} 2i\Omega & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & i\Omega & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & i\Omega & 0 & -\lambda & 2 & -1 \\ -2 & 0 & 2 & 0 & 0 & 2 & 0 & 0 \end{pmatrix}. \quad (8.8)$$

Let $S_{A,B,C,U}$ select equations A, B, C, U . Exact calculation gives

$$\det \begin{pmatrix} S_{A,B,C,U} \\ N_0 \end{pmatrix} = -4. \quad (8.9)$$

The determinant is a nonzero constant, independent of Ω and λ . Therefore the four solved equations and four identities span all eight equations, including at $\Omega = 0$.

This applies to the quadratic source without assuming that $\Phi^{(1)}$ integrates to an exact family. If the nonlinear Noether identity is $N(\Phi)E(\Phi) = 0$, then expansion about an on-shell background with an on-shell first-order tangent gives

$$N^{(0)}E^{(2)} + N^{(1)}E^{(1)} + N^{(2)}E^{(0)} = 0 \implies N^{(0)}E^{(2)} = 0. \quad (8.10)$$

Thus the dependent quadratic rows obey precisely the background identities used in (8.9).

8.5 Reality and fixed-charge completion

Define $\Phi^{(2)}$ as the finite sum of (8.5), (8.7), their complex conjugates, and the real zero-frequency corrections. Then $\Phi^{(2)}$ is real and periodic. No independent homogeneous solution is added. The correction preserves the declared charge fibre:

- the magnetic Chern-class shift is zero;
- every $\ell = 2, 4$ Maxwell correction integrates to zero on S^2 ;
- the homogeneous Maxwell coefficient U is zero in every channel;
- there is no stationary electric-charge or Wilson-line zero-mode shift.

Equations (8.2)–(8.10) prove (8.1). $\square$

9. Interpretation: a singular nonlinear solution cone

The two theorems fit together without tension.

linear direction	quadratic Taub data	second-order status
nonzero pure extra generic	$\mu_H < 0$	obstructed
balanced $u_- + a_e u_e$	$\mu_H = \mu_{P_x} = \mu_{J_i} = 0$	explicitly extendible

This is not a contradiction between the linear and nonlinear analyses. The linearized equations compute the formal tangent space $\ker L$ /local gauge. The quadratic Taub map cuts out the second-order tangent cone inside it. Positive definiteness on the pure-extra subspace excludes every nonzero ray in that subspace. Indefiniteness after adding the Einstein sector creates null directions of the quadratic map, one of which is shown here to satisfy the entire second-order equation.

The result also separates two notions often conflated in discussions of higher-derivative gravity.

1. **A linear mode can be physical at linear order.** The extra modes are nonradical under the classical Lee–Wald current.
2. **A linear mode need not be nonlinearly integrable.** Pure extra modes violate a compact global constraint.

Nothing here identifies a quantum state norm or proves a quantum ghost. The obstruction is classical and global on the closed spatial slice.

The balanced result is stronger than mere Taub cancellation. A nonzero quadratic defect is not a no-go unless it has a nonremovable cokernel component, and vanishing moment maps do not by themselves prove extension. Here the complete source is solved, channel by channel, and the dependent equations are closed by a constant-determinant Noether argument.

10. Why the full nonlinear cone is the next theorem

The current paper proves a general no-go on one large linear subspace and an existence result on one nontrivial mixed ray. That is already a coherent nonlinear statement. Completing the full mixed cone would answer a different classification problem.

The next theorem should determine the common zero locus

$$\mathcal{Z}_2 = \{u : \mu_H(u) = \mu_{P_x}(u) = \mu_{J_1}(u) = \mu_{J_2}(u) = \mu_{J_3}(u) = 0\}$$

and then test the quadratic source on its strata. The natural order is:

1. classify the full $k = 0$ cone using Gram-normalized occupation matrices;
2. classify opposite-momentum standing-wave balances;
3. incorporate exceptional, homogeneous, twist, charge, and Wilson-line blocks;
4. determine which surviving second-order jets encounter cubic or higher obstructions.

None of those steps is needed to validate Theorems 5.1 and 8.1. Conversely, the single balanced extension must not be extrapolated into a general closure theorem.

11. Scope boundary

The established statements are:

- the covariant moment-map/Taub identity on the declared compact generic solution space;
- the complete pure-extra generic fixed-bundle second-order no-go;

- the trivial common H, P_x zero locus in one nonzero- k travelling block;
- one all-stabilizer-zero Einstein-extra $k = 0$ tangent;
- one complete explicit second-order correction for that tangent.

The following remain open:

- the full mixed moment-map zero cone;
- general mixed second-order closure;
- integration of the certified jet to an exact or all-orders family;
- exceptional and generalized global blocks;
- a final background-stabilizer quotient or relational observable;
- Lorentzian causal propagation, asymptotic scattering, particles, quantization, ghosts, and unitarity.

The paper therefore carries the dependency labels LOCAL-ALGEBRAIC and REDUCED-MODE, not LORENTZIAN-CAUSAL.

12. Computational proof and reproducibility

All ranks, signs, polynomial nonvanishing tests, amplitudes, source projections, and operator remainders are computed in exact rational or algebraic arithmetic. The principal certificates are:

Result	Certificate
fixed-bundle domain and Taub descent	bridge/certificates/compact_harmonic_domain_taub_descent
direct generic axial extra current	bridge/certificates/einstein_maxwell_weyl_axial_lee_v
direct generic polar extra current	bridge/certificates/einstein_maxwell_weyl_polar_lee_v
direct axial/polar fixture Taub matrices	bridge/certificates/einstein_maxwell_weyl_hermitian_a
generic moment-map bridge and pure-extra	bridge/certificates/einstein_maxwell_weyl_moment_map
no-go	
mixed zero-locus fixture and off-shell preflight	bridge/certificates/einstein_maxwell_weyl_mixed_momen
complete balanced correction	bridge/certificates/einstein_maxwell_weyl_balanced_e

Fast verification:

```
python3 bridge/einstein_sector/verify_charge_fibre_paper_claim_map.py
python3 -m bridge.einstein_sector.einstein_maxwell_weyl_moment_map_taub_bridge \
    --verify bridge/certificates/einstein_maxwell_weyl_moment_map_taub_bridge.json
python3 -m bridge.einstein_sector.verify_einstein_maxwell_weyl_moment_map_taub_bridge
python3 -m bridge.einstein_sector.einstein_maxwell_weyl_mixed_moment_map_zero_locus \
    --verify bridge/certificates/einstein_maxwell_weyl_mixed_moment_map_zero_locus.json
python3 -m bridge.einstein_sector.verify_einstein_maxwell_weyl_mixed_moment_map_zero_locus
python3 -m bridge.einstein_sector.einstein_maxwell_weyl_balanced_ell0_second_order \
    --verify bridge/certificates/einstein_maxwell_weyl_balanced_ell0_second_order.json
python3 -m bridge.einstein_sector.verify_einstein_maxwell_weyl_balanced_ell0_second_order
python3 -m unittest \
    bridge.einstein_sector.tests.test_einstein_maxwell_weyl_moment_map_taub_bridge \
    bridge.einstein_sector.tests.test_einstein_maxwell_weyl_mixed_moment_map_zero_locus \
    bridge.einstein_sector.tests.test_einstein_maxwell_weyl_balanced_ell0_second_order
```

The final exhaustive balanced source-and-channel regeneration passed in 468.66 seconds:

```
python3 -m bridge.einstein_sector.einstein_maxwell_weyl_balanced_ell0_second_order \
--verify-exhaustive \
bridge/certificates/einstein_maxwell_weyl_balanced_ell0_second_order.json
```

The committed-certificate verifier separately reconstructs the Noether completion determinant, checks all imported content hashes, replays the real-channel factors, and verifies the fixed-charge/reality flags. The largest nested-radical cross-channel equations are replayed by the exhaustive rail.

Model authorship and human accountability

GPT-5.6.sol, an OpenAI model, contributed the research programme, mathematical direction, derivations, symbolic-code generation and debugging, claim auditing, literature organization, and manuscript. The project was commissioned and directed by Asger Alstrup Palm, who initiated the questions, served as the non-technical orchestrator and corresponding human contact, but claims no technical contribution. Circulation or submission remains conditional on a documented human verification of the mathematical claims, proof boundaries, source citations, and final text.

References

1. R. V. Saraykar and J. H. Rai, “Linearization Stability of Einstein Field Equations is a Generic Property,” arXiv:1609.07703 (2016), <https://arxiv.org/abs/1609.07703>.
2. A. E. Fischer, J. E. Marsden, and V. Moncrief, “The structure of the space of solutions of Einstein’s equations. I. One Killing field,” *Ann. Inst. H. Poincaré A* **33** (1980) 147–194.
3. E. Altaş and B. Tekin, “Linearization Instability for Generic Gravity in AdS,” *Phys. Rev. D* **97** (2018) 024028, <https://arxiv.org/abs/1705.10234>.
4. J. F. Plebański and S. Hacyan, “Some exceptional electrovac type D metrics with cosmological constant,” *J. Math. Phys.* **20** (1979) 1004–1010, <https://doi.org/10.1063/1.524174>.
5. GPT-5.6.sol, “Einstein–Maxwell Waves inside Weyl–Maxwell Gravity on a Compact Product: Exact Linear Phase-Space Inclusion and the Extra Axial Branch,” companion manuscript, 2026.
6. M. Ortaggio and J. Podolský, “Impulsive waves in electrovac direct product spacetimes with Λ ,” *Class. Quantum Grav.* **19** (2002) 5221–5239, <https://arxiv.org/abs/gr-qc/0209068>.
7. M. Ortaggio, “Einstein–Maxwell fields as solutions of higher-order theories,” *Eur. Phys. J. C* **82** (2022) 1056, <https://arxiv.org/abs/2205.14392>.
8. J. Lee and R. M. Wald, “Local symmetries and constraints,” *J. Math. Phys.* **31** (1990) 725–743, <https://doi.org/10.1063/1.528801>.
9. V. Iyer and R. M. Wald, “Some properties of Noether charge and a proposal for dynamical black hole entropy,” *Phys. Rev. D* **50** (1994) 846–864, <https://arxiv.org/abs/gr-qc/9403028>.

Frozen claim flag: PURE_EXTRA_GENERIC_NO_GO_AND_ONE_BALANCED_MIXED_SECOND_ORDER_EXTENSION_CERTIF

Next theorem: full $k = 0$ common moment-map zero cone, followed by opposite-momentum standing-wave and exceptional/global strata.