

Ghosts, Geometry, and Reality in Fourth-Order Quantum Theories

A guided introduction to positive metrics, Krein completions, residual cohomology, causal transport, relational clocks, and nonlinear selection

GPT-5.6.sol*

Claude Fable 5[†]

Abstract

The phrase “higher-derivative ghost problem” combines four different questions: which real-form completions exist for the local free dynamics, which classes survive local and residual gauge reduction, and whether a chosen completion persists under interaction and quantization. This introduction organizes a core twelve-paper programme around those four levels. Papers I–III classify the Pais–Uhlenbeck and scalar-field completions, distinguish positive metrics from Krein real forms and vacuum covariance, and explain the Jordan limit. Paper IV adds local gauge reduction and Lorentz covariance in Einstein–Weyl gravity. Papers V–VI form the interaction branch: resonant branch conversion obstructs the canonical analytic positive metric in the scalar field and in the open gravitational channel, without excluding every indefinite, nonlocal, or nonperturbative completion.

Papers VII–VIII form a separate pure-Weyl branch on the conformal cylinder. Both use the selected closed-universe problem in which all fifteen residual conformal transformations, including cylinder time translation, are constraints. Paper VII is the state-side theorem. It derives the residual BV–BFV complex, performs the positive-frequency polarization, and computes the absolute residual Chevalley–Eilenberg group

$$H_{\text{res}}^4 = \text{span}\{[W_+^2], [W_-^2]\}, \quad \text{sig } G_{\text{res}} = (2, 0), \quad G_{\text{res}} = I_2.$$

These are ghost-dressed deformation or vertex classes, not one-particle gravitons. Paper VIII is the covariant-side theorem. It constructs a support-local curvature prolongation and causal chain homotopies for the complete free metric BV complex, proves the compact-to-global causal quasi-isomorphism, recovers the residual endpoints, transports the $SO(4, 2)$ action, and identifies the Green, Cauchy, energy-mode, and residual pairings. It imports Paper VII’s cohomology theorem and does not recompute the CE complex; consequently

$$H_{\text{cov}}^4 \cong H_{\text{res}}^4 = \text{span}\{[W_+^2], [W_-^2]\}, \quad \text{sig } G_{\text{cov}} = (2, 0), \quad G_{\text{cov}} = I_2.$$

The combined result is free and formal near the conformal cylinder: its field-side theorem is classical, while its state-side theorem uses a selected completed positive-frequency representation. It is not a positive graviton Hilbert-space theorem, a Hadamard construction,

*OpenAI model and principal author. Developed the research programme, mathematical direction, synthesis, and referee analyses.

[†]Anthropic model and computational coauthor. Contributed the derivations, the machine verification (SymPy, high-precision numerics, Lean 4), and the manuscripts. The project was commissioned and coordinated by Asger Alstrup Palm (asger@area9.dk), who initiated the questions, exercised editorial control, and serves as the corresponding human contact, but is not a technical author.

a nonlinear stability theorem, or a renormalized quantum-master-equation result. The abstract cyclic causal-transfer theorem extracted from this construction now has three scoped consumers: the Berger gravity-clock system, a flat Minkowski doubled-detour portability fixture, and a curved Nariai fixture. For Nariai the repaired ten-block, 310-component parent-detour cone retracts cyclically and support-locally onto the independently certified 26-component metric Bach endpoint, with exact advanced/retarded homotopies and adjoint reversal on all rows. The complete rank-310 retract and its advanced/retarded homotopies now also admit a global support-local cyclic formal first variation along one transverse Einstein tangent. This is a second curved background and a genuine tangent-stability theorem at $\epsilon = 0$, not a smooth nonzero- ϵ family or a uniform theorem on an open family of metrics [39].

Paper IX tests a distinct clock question inside the gauge-reduction level. On an exact positive-energy Berger universe, a rotating standard-sign scalar phase has nonzero momentum while the fixed-coupling compact constraint makes the cylinder-time generator presymplectically null at the linearized level. A complete 54-row causal BV construction and action-derived q_2, q_3 instead give a cyclic causal Cartan contraction through arity three for the background-fixing helical stabilizer $K = D - \omega R$. Raw D is affine in the co-rotating variables and is not covered by that Cartan theorem. This is a scoped classical Taylor theorem, not an all-orders, boundary, Hadamard, or quantum result.

Paper X moves to the compact Plebański-Hacyan product $\mathbb{R} \times S^1 \times S^2$. The complete standard Einstein-Maxwell harmonic tangent injects into Weyl-Maxwell theory with non-degenerate, but non-symplectic, Lee-Wald pullback. The generic axial target contains two additional nonradical directions; the normalized axial current has inertia $(3, 1)$. A post-paper bridge now globalizes the harmonic maps to one natural support-local all-row chain morphism and completes the linear mapping-cofiber triangle, including residual endpoints and three distinct action-derived forms. The triangle is necessarily noncyclic for the standard pairings: the Einstein form is positive on the relevant image while the pulled-back Weyl form has inertia $(1, 1)$, and congruence invariance excludes a cyclic repair. This is a certified linear relative architecture, not nonlinear or physical equivalence. The action-derived Einstein-Maxwell and Weyl-Maxwell Taylor data are now both complete through q_3 . Their strict linear inclusion has a nonzero arity-two defect, and a Taub witness excludes a support-local correcting f_2 on the full smooth periodic fixed-bundle carrier. The defect descends globally to exactly five stabilizer charges, but locality forces the pre-integration receiver to be a horizontal Noether-current complex; the canonical relative Hessian Green-current cone and its local divergence identity are certified, while stabilizer precomposition, Lee-Wald comparison, and cyclic-dual completion remain open. Paper XI returns to the Berger clock and transfers the mixed gravity-Maxwell interaction from 64 rows to a typed cyclic 36-row retained complex. Its nonzero ℓ_3 representative has 25,950 exact coefficients, all arity-three and BV-cyclicity identities close, and the homological exchange channel vanishes because it is supported on a contractible clock-dual row. The theorem is frozen for this retained cyclic-causal representative and for its nonremovability in the declared filtered-cyclic F_2/F_3 class; it is not yet a nonzero operation on cohomology. A successor calculation with the fuller Maxwell ghost structure trivializes the mixed term through first jet, so the older restricted nonremovability result is not promoted to an invariant physical interaction; the second-jet image remains open. A subsequent support-local rank-46 STF2 graph prolongation is an exact cyclic carrier over the retained 36-row complex with contractible complement, but the requested canonical support-local branch projector is excluded by an exact normalized subprincipal obstruction. A physical branch manifest therefore requires a noncontractible filtered or mixed-bundle enlargement, or an explicitly nonlocal reduced-mode split.

A theorem-frozen charge-fibre bridge then proves that every nonzero pure-extra generic Weyl-Maxwell tangent is Taub-obstructed on the fixed compact magnetic bundle, while

one balanced Einstein–extra tangent cancels all five stabilizer moment maps and extends explicitly to second order. Subsequent classified certificates determine the complete generic $k = 0$ and paired opposite-momentum moment-map cones, extend the balanced ray to a two-parameter axial $\ell = 2$ second-order face, and now prove that the complete exceptional $\ell = 1, k = 0$ all- m pure cone is obstructed. More generally, the complete certified finite-support linear inventory—generic Einstein and extra modes, standard and extra exceptional dipoles, the twist Jordan block, and the homogeneous block—has a finite exponential-polynomial second-order correction exactly when its five stabilizer moment maps vanish. Bounded or finite-quasiperiodic continuation has additional certified polynomial-growth and shell-resonance functionals whose common zero locus is not yet solved; infinite-mode completion and causal/retarded correction remain separate gates. On the tuned axisymmetric standard branches the bounded resonance and charge cone is now classified uniformly for every $\ell \geq 2$: it is the origin plus two mixed axial–polar sheets on an exact occupation interval. A separate two-absolute-momentum workload has closed all 108 axisymmetric $L = 4$ and all 56 nonaxisymmetric $L = 1, 3$ basis coefficients, each individually obstructed; the common arbitrary-amplitude zero variety remains open. The standard generalized-zero bounded cone is nevertheless now complete: $a = b = Q_e = 0$ and $B = 0$, with c, d, W_x and the constant twist A free. Moreover $b = B = 0$ is necessary on the entire finite-support bounded cone, leaving the oscillator-coupled polynomial ideal and shell resonances as the next gate. The declared homogeneous/twist- $\ell = 2$ extra common-zero cone is now complete: every nonzero tangent is an $SO(3)$ rotation of one aligned orbit. Passing those global moment maps is not sufficient for bounded continuation. Every nonzero point has an unavoidable quadratic-in-time source, so no bounded finite-quasiperiodic second-order correction exists. On the same complete orbit, a blockwise constructive theorem supplies a real finite exponential-polynomial second-order correction; it is necessarily secular rather than bounded. Causal/retarded continuation remains a separate open gate. The observer programme separately has causal rank-two linear records, six global detector rods, second-order rod-source solvability, a frozen 84-row handoff, a separate-axis rod-gravity unary theorem, and a 72-row memory causal subcomplex. The Berger classical programme also now constructs an actual retarded, spatially global clock-slice Maxwell observable with $1 + z = 2$, a nontrivial reduced probe bracket, and nontrivial relational evolution. Two receiver-adjacent localized conserved emitters have exact causal preparations. Response-specific shell streaming, exact finite kernels, and partition-refined Cauchy covectors now certify a leading rank-two detector response uniformly for the declared clock masses $m_0^2, m_1^2 \in [1, 2]$; the first retarded emitter channel is also executable. This is a finite leading-response theorem, not yet a complete all-shell q_3 feedback or recoil construction. A common localized emission experiment, evaluated recoil, the full apparatus Dirac bracket, and nonlinear backreacted response remain open. The enlarged 108-row apparatus unary is cyclic and nilpotent at zeroth order, but its first clock-radius jet has a certified nonzero square; nonlinear promotion is blocked pending the second clock-map jet and cotangent lift [40].

A separate black-hole programme has classified the static spherical pure-Weyl family in its declared Laurent ansatz, including a horizon-regular non-Einstein fixture. On that static parameter slice a residual-basic field-dependent time generator yields an integrable Hamiltonian, Wald entropy, and the exact first law at every simple horizon. No time-dependent horizon phase space, ringdown, stability, or Hawking theorem follows. The first axial $\ell = 2$ perturbation block now reproduces the Schwarzschild Regge–Wheeler equation exactly. On that Ricci-flat background the Einstein branch $\delta\text{Ric} = 0$ injects into the Bach kernel, while the extra branch is the second-order Lichnerowicz-type field $\psi = \delta\text{Ric}$. The extra branch has a two-parameter ingoing-regular family at the horizon, so horizon regularity alone does not remove it. The exact action-derived flux matrix shows that the pure Einstein/Regge–Wheeler block is symplectically null for conjugate wave pairs,

while the Einstein–extra and extra–extra horizon-flux blocks are nonzero at three exact real-frequency fixtures. For the declared Schwarzschild $m = 1$, $\ell = 2$, $\omega = 3/5$ fixture, the axial and polar metric reconstructions now resolve the repeated characteristics. The sourced extra representatives acquire logarithmic or power-enhanced tails. Their Einstein–extra and extra–extra slice norms diverge, while the Einstein slice norms are finite; the divergence classes cannot be removed by shifting an extra representative by an Einstein solution. Consequently finite symplectic norm at infinity selects exactly the Einstein sector in both parities at this fixture, even though horizon regularity alone retains the extra branches. This is a global phase-space selection theorem at one real frequency, not a symbolic-frequency, all- ℓ , quasinormal-mode, stability, or full exterior-IBVP theorem [41].

Paper XII computes the strict one-loop BV anomaly. Its nonzero even coordinates obstruct the fixed-field-content local Euclidean QME; a formal tau-adic compensator extension contracts the anomaly and restores that QME at one loop. The anomaly-induced Paneitz/Riegert representative and universal $C^2 \log$ coefficient and the raw-to- $\square R = 0$ strict-metric scheme conversion are fixed, while finite C^2 and dressed $R(\hat{g})^2$ counterterms leave the complete renormalized Q_1 underdetermined. In the generic Euclidean ghost sector, all longitudinal insertions now resum into one normalized scalar Schur factor. Its order- -2 correction defines a canonical modified Fredholm $\det_3$ tail, and the critical local residue $\text{Wres}(K^2) = (4\pi)^{-2} \int (R^2 + 2 \text{Ric}^2)/27$ is exact. On the round S^4 benchmark the finite weighted traces, modified determinant, and factorization ratio $5/3$ are exact; on a generic closed background the local weight-raised multiplicative defect is $-\frac{1}{4} \text{Wres}(K^2)$. All eleven generic third-curvature ghost functions—ten pole-three rows and the sole pole-four I_{29} row—are now integrated exactly. On the anisotropic product $S^2(k_1) \times S^2(k_2)$, the normalized longitudinal Schur eigenvalue and its coupled exceptional-mode correction are also exact; the latter contributes the finite factor 3^{-6} that would be lost by priming the vector and scalar determinants separately. Their assembly into the full ghost determinant and the generic physical fourth-order Hessian remain open. Separately, the reduced vacuum-cylinder E/A/L carrier has a compatible complex structure and microlocal Hadamard two-point distributions with signs $(+, -, -)$. This is a Krein carrier, not a full-BV positive graviton Hilbert space, interacting product, or particle theorem [36, 42]. The programme therefore replaces a single ghost verdict by a typed architecture: local completion, residual reduction, covariant causal transport, and interaction stability are related, but none substitutes for the others.

Open-repository authorship and accountability

This manuscript is an open-repository research publication rather than a journal submission. GPT-5.6.sol is the principal author and Claude Fable 5 is the computational coauthor under the contribution statement above. Asger Alstrup Palm commissioned and coordinated the programme, selected its questions, maintained editorial control, and serves as the corresponding human contact; he does not claim technical authorship. Mathematical claims are tied to deterministic certificates, independent verifiers, mutation tests, and explicit lifecycle boundaries in the public repository. This authorship and accountability model is stated on its own terms rather than being adapted to a journal policy that does not govern the publication.

Contents

1 Why higher derivatives create ghosts

6

2	The Pais–Uhlenbeck oscillator as a microscope	10
3	What Bender and Mannheim had already solved	10
4	Paper I: reconstructing the Bender–Mannheim transformation	11
4.1	The exact spectrum of Q	12
4.2	Many metrics, one canonical Gaussian metric	12
5	Paper II: why the canonical metric is the least-distorting one	13
5.1	Cartan projection	13
5.2	Why the theorem is not a routine Kempf–Ness application	13
6	The exceptional point: when two frequencies merge	14
7	From one oscillator to a field	15
8	Metric geometry and vacuum geometry are different	15
9	The universal complex vacuum covariance	15
10	Resonance: the metric diverges but the covariance survives	16
11	What is a Krein space?	16
12	A necessary representation-theoretic distinction	18
12.1	Physical Dyson transport	18
12.2	Identity embedding in an auxiliary standard CCR algebra	18
13	The ultraviolet Fock-sector hierarchy	18
14	The fourth-order vacuum is locally well behaved	19
15	When interactions are switched on	19
15.1	Deforming the positive metric	20
15.2	A finite oscillator fails only at special resonances	20
15.3	The momentum continuum turns exceptional resonances into ordinary scattering	21
15.4	Why the obstruction waits until second order	22
15.5	A different structure survives: two mirror Jordan sectors	22
16	Why gravity is not merely five copies of the oscillator	23
17	Lorentz covariance forbids hybrid solutions	24
18	The gauge-invariant gravitational bridge	24
19	The conformal Jordan boundary	25
20	Paper VI: interaction obstruction in Einstein–Weyl gravity	25

21 Papers VII–VIII: residual states and covariant transport	27
21.1 The selected closed-universe problem	27
21.2 Paper VII: the state-side theorem	27
21.3 Paper VIII: the covariant-side theorem	28
21.4 Relation and combined boundary	29
22 Papers IX–XI and the nonlinear selection bridges	29
22.1 Paper IX: a healthy clock and the correct stabilizer	29
22.2 Paper X: ordinary waves inside the fourth-order theory	30
22.3 Paper XI: the first retained gravity–light bracket	31
22.4 Charge-fibre selection and observable records	31
23 What was already solved, and what is new	34
24 What remains open	35
25 Conclusion	38

1 Why higher derivatives create ghosts

Most familiar equations of motion contain at most two time derivatives. Newton’s equation has the form

$$m\ddot{q} = F(q), \quad (1)$$

and the Klein–Gordon field equation has the form

$$(\square + m^2)\phi = 0, \quad (2)$$

where the empty square is the standard symbol for the wave operator (d’Alembertian), $\square = \partial^2/\partial t^2 - \nabla^2$ — the spacetime analogue of the Laplacian. Specifying a solution requires two pieces of initial data: position and velocity for a particle, or the field and its first time derivative for a field theory.

A fourth-order equation such as

$$\left(\frac{d^2}{dt^2} + \omega_1^2\right)\left(\frac{d^2}{dt^2} + \omega_2^2\right)z(t) = 0 \quad (3)$$

requires four pieces of initial data. Its solution contains two independent frequencies,

$$z(t) = A_1 e^{-i\omega_1 t} + B_1 e^{i\omega_1 t} + A_2 e^{-i\omega_2 t} + B_2 e^{i\omega_2 t}. \quad (4)$$

The extra derivatives have introduced an extra oscillator.

That sounds harmless. The difficulty is that the standard Hamiltonian formulation usually assigns opposite signs to the two oscillators:

$$H \sim H_{\omega_1} - H_{\omega_2}. \quad (5)$$

One branch then appears to carry negative energy. If one changes the quantization convention to keep the energy positive, the same branch may instead acquire negative norm. This is the traditional higher-derivative ghost problem.

The classical theorem of Ostrogradsky [1] explains why the Hamiltonian obtained from a nondegenerate higher-derivative Lagrangian is generically unbounded in its canonical momenta (see [2] for a modern review). But that theorem does not by itself answer every quantum question. The questions divide into three logically distinct levels.

Level 1 — local free dynamics and completion. Can the Hamiltonian be diagonalized? Can a positive inner product be found? Which variables count as real? What happens at the Jordan limit where distinct frequencies or masses coincide?

Level 2 — gauge and residual reduction. After the local gauge constraints have been imposed, which oscillator modes survive as genuine states or observables? On backgrounds with residual symmetries, should those symmetries be retained as global charges or treated as gauge? If they are gauged, what is the resulting BRST cohomology, and does positivity belong to propagating states, to vertex classes, or to neither?

Level 3 — interaction stability. Can a free positive metric or Krein involution be deformed when interactions are added? Can ghost parity remain conserved when particles convert into one another? Do exact energy-conserving processes obstruct the deformation? Does a doubled symmetry survive even if an individual pointed sector fails?

Level 4 — quantum consistency and state selection. Does the local quantum master equation survive regularization? If not, can a declared extension restore it? Does the Lorentzian causal carrier admit a BRST-compatible positive Hadamard state, and can asymptotic particles and unitary probabilities be defined?

These four levels must not be conflated. Solving the free Hamiltonian settles neither the residual gauge problem, the interacting ghost problem, nor the quantum state problem. Residual reduction can remove locally propagating oscillator classes without changing their local equations, while interactions test whether a chosen free completion survives energy exchange. The research programme therefore has four linked branches after the free gravity classification. Papers V–VI study interaction stability. Papers VII–VIII study the state-side and covariant-side residual problems at the pure-Weyl boundary. Paper XII adds the one-loop anomaly/effective-action analysis and the first reduced Lorentzian Hadamard/Krein carrier. Papers IX and XI use a non-conformally-flat Berger clock background to test the D quotient, causal Cartan contraction, and retained gravity–light interactions. Paper X and the charge-fibre bridge study ordinary Einstein–Maxwell waves, additional Weyl–Maxwell modes, and their nonlinear Taub selection on a distinct compact product background.

Series map. The twelve core technical papers have distinct jobs.

1. Paper I reconstructs and classifies the positive free Pais–Uhlenbeck quantization.
2. Paper II characterizes its canonical metric by a minimum-distortion principle.
3. Paper III separates metric, vacuum, representation, and Jordan data for the fourth-order scalar field.
4. Paper IV adds local gauge reduction and Lorentz covariance in free Einstein–Weyl gravity and reaches the pure-Weyl boundary.
5. Paper V formulates the interaction deformation problem and identifies resonant scalar obstructions.

6. Paper VI applies that deformation complex to Einstein–Weyl gravity.
7. Paper VII computes the selected residual state cohomology at the pure-Weyl boundary.
8. Paper VIII constructs the covariant causal bridge to Paper VII’s already computed residual result.
9. Paper IX constructs a healthy S^1 -valued Berger phase-clock, proves fixed-coupling rigidity of its nonzero momentum and linear presymplectic nullity of D , and constructs a causal K -Cartan homotopy through arity three on the rational representative.
10. Paper X embeds the complete standard compact Einstein–Maxwell harmonic phase space into Weyl–Maxwell theory, compares the covariant currents, and classifies the generic axial extra quotient.
11. Paper XI transfers the mixed Berger gravity–Maxwell interaction through ℓ_3 to a typed cyclic retained complex and proves exact algebraic and causal-Cartan compatibility for the frozen retract.
12. Paper XII computes the strict and compensator-extended one-loop BV dispositions, the anomaly-fixed effective-action terms, and the reduced vacuum-cylinder Hadamard/Krein state-space signs.

Thus Papers I–IV form the local free spine, Papers V–VI the interaction branch, Papers VII–VIII and XII the state/covariant/quantum branch, Papers IX and XI the Berger clock and retained-interaction branch, and Paper X the compact incidence of Einstein–Maxwell waves in Weyl–Maxwell theory. Paper VIII does not perform another Chevalley–Eilenberg calculation, Paper IX does not recompute the residual H^4 group, and Paper XI does not yet supply a physical branch mixing table or a final-cohomology operation beyond its declared filtered-cyclic nonremovability theorem; the successor full-ghost carrier trivializes the term through first jet and leaves the second-jet image open. Paper XII does not construct a positive full-BV state, Lorentzian QME, or particle interpretation.

A cross-programme D -quotient validation dossier records the scope of the Berger branch. Paper IX is the resulting classical theorem on Taub reduction, relational clocks, and interaction-order dependence. The first scalar-clock gate is a certified no-go: for one real conformally coupled scalar the invariant homogeneous mode is a branch-local oscillator clock, but its positive improved stress is incompatible with the Bach-flat vacuum cylinder; at the only compatible background, $\bar{T} = 0$, its linearized Diff×Weyl incidence vanishes. A distinct neutral two-field reference sector supplies the minimal homogeneous repair: with internal signature $(+, -)$, the zero-charge condition cancels the improved stress componentwise, the conserved nonzero Wronskian makes the projective angle a global group-valued clock for compact D , and D is a presymplectic null direction on that declared sector. A subsequent local health audit shows that this opposite-sign model cannot be promoted to a globally regular positive clock: after Weyl reduction its ratio mode retains a derivative term, and every neutral winding orbit crosses four zeros at which the kinetic sign changes. The homogeneous reference-clock theorem remains valid. The complete standard positive-sign one-field stealth family, including inhomogeneous configurations, is also obstructed: every reciprocal denominator has a global zero, and no time-dependent member has an everywhere-timelike gradient. The positive-energy backreacted

search now has an exact background theorem. A static Berger cylinder supports two standard-sign rotating conformal scalars with a positive quartic potential, dominant-energy stress, time-like phase, and full raw Diff×Weyl clock incidence on a nonempty open squashing interval. Its phase carries the exact nonzero conserved matter momentum $Q_R = 16\pi^2\alpha_B q\sqrt{1-4q}$. The frozen action/current convention gives $\Omega_{\text{total}}(\delta, \mathcal{L}_D) = \omega\delta Q_R$. At fixed couplings the squashing q is isolated, so the open background interval is not itself a legal charge variation. More strongly, varying the exact lapse before fixing it gives $\delta E_N = -(\alpha_B q^{3/2}/2)\delta Q_R/Q_R$. Compact spatial averaging excludes an inhomogeneous escape, so D is presymplectically null on the declared smooth fixed-coupling linearized Berger phase space. The phase is a genuine rotating clock coordinate whose nonzero momentum is fixed by the compact constraint, rather than a cost-free scalar label or a freely variable residual charge. The minimal temporal/Weyl clock doublets and their BV-dual rows extend to a complete cyclic 54-row gauge-fixed complex with advanced and retarded chain contractions. Action differentiation supplies arbitrary-input four-dimensional q_2 and q_3 , and the legacy local generator recorded in the co-rotating frame is exactly the helical stabilizer $K = D - \omega R$; it admits cyclic causal Cartan primitives through arity three. Raw D has a nonzero zeroth Taylor component and no affine Cartan claim is made. These results close the classical interaction half of the Paper IX gate. They do not prove an all-orders nonlinear quotient, a boundary theorem, Hadamard data, or a quantum master equation.

Paper XI begins only after this base theorem is frozen. It adjoins Maxwell to the complete Berger carrier, imports the repaired action-derived mixed q_2, q_3 , and transfers the ternary operation through the typed cyclic 64-to-36 contraction. It therefore tests interaction survival on a retained carrier rather than reopening Paper IX’s clock-charge theorem. Section 22 states its exact result and the invariant and branch-projection gates that remain open.

Fourth-order theories can admit several inequivalent real-form completions of one complex spectral dynamics. Local free geometry determines which completions are available; residual gauge reduction determines which state and vertex classes survive; physical interactions determine whether the surviving structures remain dynamically compatible.

Strength of the claims. The free classifications are theorems in the classes stated below. In Paper V the 3:1 and 3:2 oscillator obstruction classes are exact, the 5:1 fourth-order result is a computational proposition, and the general resonance hierarchy remains a conjecture. For the field theory what is proved is an analytic metric obstruction on a nonempty open shell subset, not a complex spectrum or the absolute nonexistence of every positive completion. In gravity, Paper VI proves first-order cubic protection and a tree-level second-order obstruction based at the canonical Paper-IV positive real form; it does not prove an all-orders no-go, complex gravitational spectrum, or failure of every nonlocal or differently pointed completion. Its Krein result concerns the natural particle-number-diagonal, cluster-multiplicative lift induced on physical BRST cohomology, not every Poincaré-covariant involution on a multiparticle space. A full-field complex spectrum and a quantum operator implementing the doubled scalar exchange remain open.

Paper VII proves its residual theorem only for the stated formal, closed-universe, positive-frequency polarization. Paper VIII proves a free classical Lorentzian causal transport theorem for the corresponding metric BV complex. Neither paper claims an integrated $SO(4, 2)$ rep-

resentation, a positive graviton Fock space, a BRST-compatible Hadamard state, nonlinear stability, anomaly cancellation, or interacting unitarity. Common-core commutators of unbounded generators do not by themselves provide group integrability [19].

2 The Pais–Uhlenbeck oscillator as a microscope

The Pais–Uhlenbeck (PU) oscillator [3] is the simplest model containing the full fourth-order difficulty. Its Lagrangian can be written

$$L = \frac{\gamma}{2} (\ddot{z}^2 - (\omega_1^2 + \omega_2^2) \dot{z}^2 + \omega_1^2 \omega_2^2 z^2), \quad \omega_1 > \omega_2 > 0. \quad (6)$$

Its equation of motion factorizes:

$$\left(\frac{d^2}{dt^2} + \omega_1^2 \right) \left(\frac{d^2}{dt^2} + \omega_2^2 \right) z = 0. \quad (7)$$

It therefore contains two ordinary frequencies, but their canonical combination has the wrong-sign structure characteristic of a ghost.

The oscillator is useful because it is finite dimensional and exactly solvable. Questions that become difficult analytic problems in a field theory reduce here to matrix algebra:

- Can the Hamiltonian be transformed to a positive normal form?
- Is that transformation unique?
- How many positive metrics exist?
- Is there a preferred one?
- What happens as $\omega_1 \rightarrow \omega_2$?
- Does the limiting Jordan system admit any positive invariant inner product?

A correct answer at the oscillator level does not automatically solve a quantum field theory. But the oscillator provides the local building block for each momentum mode of a free fourth-order field.

3 What Bender and Mannheim had already solved

Bender and Mannheim [5, 6] observed that the PU Hamiltonian should not be quantized using the naive real contour and inner product. After a complex continuation of one canonical coordinate, the Hamiltonian becomes non-Hermitian but PT-symmetric [4].

They constructed an operator

$$Q = \alpha pq + \beta xy \quad (8)$$

and the associated similarity transformation $\rho = e^{-Q/2}$ such that

$$h = \rho H_{\text{PT}} \rho^{-1} \quad (9)$$

is a conventional positive Hamiltonian describing two ordinary oscillators. The original Hamiltonian is then Hermitian not with respect to the naive adjoint, but with respect to the positive metric

$$\eta = \rho^\dagger \rho = e^{-Q}, \quad (10)$$

in the general framework of pseudo-Hermitian quantum mechanics [20, 21]. The physical inner product becomes

$$\langle \psi, \phi \rangle_\eta = \langle \psi, \eta \phi \rangle. \quad (11)$$

With this inner product, the unequal-frequency oscillator has positive norms and a positive spectrum. Thus an important version of the ghost problem was already solved:

The free unequal-frequency Pais–Uhlenbeck oscillator admits a positive-metric quantization.

Three levels must nevertheless be distinguished, and the series addresses each in turn: the *free spectral solution* (the free oscillator maps to a positive two-oscillator Hamiltonian); *perturbative interaction stability* (the metric can be deformed away from on-shell resonances); and *global interacting viability* (energy-conserving conversion processes may obstruct the deformation and, in finite resonant systems, produce complex spectra). In short:

The free ghost problem was solved locally; Paper V determines where that solution survives interaction.

What remained unclear was the mathematical status of the construction. Why does the particular operator Q appear? Is it an ansatz or a canonical object? Is the metric unique? If it is not unique, why should the Bender–Mannheim metric be preferred? What happens when the oscillator becomes a field with infinitely many momentum modes? How does this positive construction relate to an indefinite Krein-space construction [7]? And what survives after gauge reduction in gravity? The four papers [25, 26, 27, 28] address these questions.

4 Paper I: reconstructing the Bender–Mannheim transformation

The first paper [25] reformulates the oscillator as a problem in complex symplectic geometry. Collect the canonical variables into a vector $V = (x, y, p, q)^\top$, with commutation relations encoded by the symplectic matrix J :

$$[V_a, V_b] = iJ_{ab}. \quad (12)$$

A quadratic Hamiltonian has the form $H = \frac{1}{2}V^\top GV$, where G is a symmetric matrix, and the Hamiltonian flow is generated by $A = JG$. The problem is to find a complex symplectic matrix S satisfying

$$S^\top JS = J, \quad S^\top GS = G_0, \quad (13)$$

where G_0 is the positive two-oscillator normal form.

The first result is that, after a specified canonical normalization, this problem has a *unique* Hermitian positive-definite solution S_+ . The complete solution family is the coset

$$S_+H, \quad H \cong SO(2, \mathbb{C}) \times SO(2, \mathbb{C}), \quad (14)$$

where H is the stabilizer of the positive normal form. Thus many complex symplectic transformations diagonalize the Hamiltonian, but exactly one normalized representative is both Hermitian and positive.

The logarithm of this matrix is then converted into a quadratic quantum operator through the complexified metaplectic representation [43]. The resulting operator is precisely $Q = \alpha pq + \beta xy$ with the coefficients found by Bender and Mannheim. The logical direction has therefore been reversed:

$$(G, J, G_0) \longrightarrow S_+ \longrightarrow Q.$$

The operator Q is no longer introduced by guessing a useful quadratic expression. It is reconstructed from the symplectic normal-form problem.

4.1 The exact spectrum of Q

In normalized oscillator coordinates,

$$Q = r (a_1 a_2^\dagger + a_1^\dagger a_2), \quad r = \log \frac{\omega_1 + \omega_2}{\omega_1 - \omega_2}. \quad (15)$$

This is a number-preserving Schwinger $SU(2)$ generator. It commutes with $N_{\text{tot}} = N_1 + N_2$, and on the fixed- N subspace its eigenvalues are $r(-N, -N + 2, \dots, N - 2, N)$. Consequently

$$\text{spec } Q = r \mathbb{Z}, \quad (16)$$

with pure-point spectrum and infinite multiplicities. The metric has spectrum

$$\text{spec}(e^{-Q}) = \{e^{-rn} : n \in \mathbb{Z}\} \cup \{0\}, \quad (17)$$

where zero is an accumulation point rather than an eigenvalue. This provides a natural invariant domain: the algebraic Fock space of finite linear combinations of number states, preserved by Q , $e^{\pm Q/2}$, and the quadratic Hamiltonians appearing in the construction.

4.2 Many metrics, one canonical Gaussian metric

Pseudo-Hermiticity alone does not make the metric unique. If $h = \rho H_{\text{PT}} \rho^{-1}$ and W is a positive operator commuting with h , then

$$\eta_W = \rho^\dagger W \rho \quad (18)$$

also defines an algebraic positive metric form [20, 21]. Thus the complete metric freedom is larger than the finite-dimensional symplectic stabilizer: the stabilizer describes the Gaussian or quadratic part of the ambiguity, while arbitrary positive functions of the normal-mode number operators provide additional non-Gaussian freedom. The first paper therefore distinguishes the *unique normalized positive symplectic diagonalizer* from the *nonunique pseudo-Hermitian metric operator*.

5 Paper II: why the canonical metric is the least-distorting one

Uniqueness inside a restricted class does not by itself explain why S_+ is geometrically natural. The second paper [26] supplies a variational characterization. For an admissible diagonalizer S , define

$$\mathcal{F}(S) = \|\log(S^\dagger S)\|_F^2. \quad (19)$$

The positive matrix $S^\dagger S$ measures how much the transformation stretches and contracts the canonical coordinates; the logarithm converts multiplicative stretching into additive “rapidities,” and the Frobenius norm measures their total size. The theorem is

$$\mathcal{F}(S) \geq \mathcal{F}(S_+) = 4r^2, \quad (20)$$

with equality precisely for S_+ multiplied by a unitary element of the stabilizer — which for the PU oscillator consists of real sign matrices, so all minimizers have the same positive polar factor and determine the same Bender–Mannheim metric. In ordinary language:

S_+ is the least-distorting allowed transformation that makes the Hamiltonian positive.

This is analogous to finding the shortest path from a point to a surface: a shortest path meets a smooth surface orthogonally. Here the “surface” is the orbit generated by transformations that preserve the positive normal form.

5.1 Cartan projection

The geometry is that of the complex symplectic group $\mathrm{Sp}(4, \mathbb{C})$ and its maximal compact subgroup $K = \mathrm{Sp}(4, \mathbb{C}) \cap U(4)$, acting on the space of positive matrices with the affine-invariant metric [44]. The Cartan projection records the logarithmic singular values of a transformation. With the doubled Gram convention used in the papers ($\mu(S) = \log \mathrm{spec}(S^\dagger S)$),

$$\mu(S_+) = (r, r), \quad (21)$$

a point on the wall $x_1 = x_2$ of the C_2 Weyl chamber. The distortion functional is related to the Cartan norm by

$$\mathcal{F}(S) = 2\|\mu(S)\|^2. \quad (22)$$

Thus the minimum-distortion theorem is a Cartan-projection theorem.

5.2 Why the theorem is not a routine Kempf–Ness application

The physical stabilizer $H \cong SO(2, \mathbb{C})^2$ is not compatible with the canonical Cartan involution determined by the physical inner product: generically

$$H \cap K \cong \mathbb{Z}_2 \times \mathbb{Z}_2, \quad (23)$$

rather than the maximal compact $U(1)^2$ of the abstract group. The obstruction belongs to the *embedding* of H , not to its abstract group structure, so the classical nearest-point theory of self-adjoint groups [45, 46] does not apply to H directly.

The resolution is to enlarge H to its smallest compatible (θ -stable) hull. Generically this is

$$SL(2, \mathbb{C}) \times SL(2, \mathbb{C}), \quad (24)$$

the mode-preserving subgroup, whose positive part is a totally geodesic copy of $\mathbb{H}^3 \times \mathbb{H}^3$. The Bender–Mannheim direction is orthogonal to this mode-preserving geometry, so standard nonpositively curved projection theory can be applied to the hull and the result restricted back to the physical stabilizer. The paper proves a general principle:

A positive diagonalizer minimizes Cartan distortion along a compatible subgroup exactly when its Hermitian generator is orthogonal to the subgroup's positive tangent directions.

For a splitting into many oscillator modes, those tangent directions are the mode-preserving deformations, and the normal directions are precisely the inter-mode couplings. The PU oscillator is the simplest nontrivial example: its preferred generator is entirely inter-mode.

6 The exceptional point: when two frequencies merge

The rapidity is $r = \log \frac{\omega_1 + \omega_2}{\omega_1 - \omega_2}$. As $\omega_1 \rightarrow \omega_2$, the denominator tends to zero and $r \rightarrow \infty$. The positive diagonalizer stretches one pair of directions by $e^{r/2}$ and contracts another by $e^{-r/2}$; the corresponding Cartan distance diverges.

This is not merely a poor choice of coordinates. At equal frequency, the dynamical matrix ceases to be diagonalizable and develops a Jordan block — the hallmark of an exceptional point [48, 49]. A 2×2 Jordan block has the form

$$H_J = \begin{pmatrix} \omega & 1 \\ 0 & \omega \end{pmatrix}. \quad (25)$$

Suppose there were a positive-definite Hermitian form η satisfying $H_J^\dagger \eta = \eta H_J$. Then the matrix $\eta^{1/2} H_J \eta^{-1/2}$ would be Hermitian, hence diagonalizable. But similarity transformations preserve Jordan form. This is impossible. Hence a nontrivial Jordan block admits no nondegenerate positive-definite invariant Hermitian form [25, 47]. It may still admit:

- degenerate positive-semidefinite forms;
- nondegenerate indefinite forms;
- or intrinsically Jordan PT-symmetric realizations [6].

What fails is the continuation of the unequal-frequency positive Hilbert metric as a nondegenerate invariant positive form. This distinction matters: the conclusion is not that the equal-frequency theory cannot exist, but that it cannot remain an ordinary diagonalizable positive-metric system of the same kind.

7 From one oscillator to a field

Consider the free fourth-order scalar field

$$(\square + m_1^2)(\square + m_2^2)\phi = 0. \quad (26)$$

After Fourier transforming in space, each spatial momentum k contributes a PU oscillator with frequencies $\omega_i(k) = \sqrt{k^2 + m_i^2}$. The modewise rapidity is

$$r(k) = \log \frac{(\omega_1(k) + \omega_2(k))^2}{m_1^2 - m_2^2} = 2 \log |k| + O(1) \quad (|k| \rightarrow \infty). \quad (27)$$

Thus the modewise similarity transformations become increasingly nonuniform in the ultraviolet.

This is where finite-dimensional and field-theoretic questions diverge. Each individual momentum mode can be treated successfully, while the collection of infinitely many modes may define an unbounded field transformation, a non-equivalent norm on phase space, a different quasifree representation, or a different real form of the field algebra. A modewise solution is therefore necessary but not sufficient for a field theory.

8 Metric geometry and vacuum geometry are different

A central lesson of the field papers [26, 27] is

$$\text{metric distortion} \neq \text{vacuum excitation}.$$

The Cartan rapidity $r(k)$ diverges like $2 \log k$, and the finite-dimensional observable-space metric has eigenvalues $e^{\pm r(k)} \sim k^{\pm 2}$. This describes a strong deformation of the canonical phase-space norm. It does *not* follow that the transformation creates $O(k^2)$ particles.

In normalized coordinates the generator is number preserving:

$$Q_k = r(k)(a_{1,k}a_{2,k}^\dagger + a_{1,k}^\dagger a_{2,k}). \quad (28)$$

It acts inside the complexified beam-splitter algebra, and the canonical vacuum is annihilated by it, even though the associated Cartan norm is arbitrarily large.

The reason is geometric. Positive metrics are points in one quotient space, roughly G/K , while Gaussian vacuum rays belong to a different quotient, G/P_Ω , where P_Ω stabilizes the vacuum line. The vacuum-ray map does not factor through the metric quotient. Two transformations can therefore have very different metric distortion while producing the same vacuum ray. This prevents one from estimating physical particle content directly from the Cartan norm.

9 The universal complex vacuum covariance

For the fourth-order field, the selected two-point function has the spectral form

$$\widetilde{W}_{12}^+(p) = \varepsilon \theta(p^0) \frac{\delta(p^2 - m_1^2) - \delta(p^2 - m_2^2)}{m_1^2 - m_2^2}, \quad (29)$$

where $\varepsilon = \pm 1$ records the overall action convention. This is a divided difference of two ordinary Klein–Gordon two-point functions:

$$W_{12}^+ = \varepsilon \frac{W_{m_1}^+ - W_{m_2}^+}{m_1^2 - m_2^2}. \quad (30)$$

The field paper [27] shows that the admissible positive metrics all select the same complex covariance. They may change the physical adjoint and the representation of observables, but not this underlying spectral two-point distribution. The result should be stated carefully:

The complex covariance is metric-independent; the involution and completion need not be.

A state in the strict algebraic sense requires a fixed involution and a positivity condition. When different completions use different involutions, it is safer to speak first of the common complex quasifree covariance.

10 Resonance: the metric diverges but the covariance survives

Let $m_1^2, m_2^2 \rightarrow m^2$. Then the divided difference has the regular limit

$$\widetilde{W}_{mm}^+(p) = -\varepsilon \theta(p^0) \delta'(p^2 - m^2). \quad (31)$$

In the time domain, per mode,

$$W_{mm}^+(t, k) = -\varepsilon \frac{(1 + i\omega t) e^{-i\omega t}}{4\omega^3}, \quad \omega = \sqrt{k^2 + m^2}. \quad (32)$$

The positive diagonalizing transformation diverges because the dynamics becomes Jordan. Yet the complex vacuum covariance remains finite. This is one of the main conceptual results:

Resonance is a singularity of the metric and dynamics, not necessarily of the vacuum covariance.

At the doubly massless point $m = 0$, the covariance is related, after matching the action sign and infrared convention, to the massless $\square^2$ covariance used in the Krein-space construction of Bateman and Turok [7]. Their sign convention gives the opposite overall covariance to one common PU convention: the relation is an action-sign reversal, not a literal equality before conventions are matched. This bridge identifies the *free complex covariance*; it does not by itself reconstruct Bateman and Turok’s interacting $O(1, 1)$ embedding, ghost symmetry, or indefinite-space probability rule, which are additional structures of their theory.

11 What is a Krein space?

A Hilbert space has a positive inner product: $\langle \psi, \psi \rangle > 0$ for every nonzero state. A Krein space has a nondegenerate but *indefinite* inner product: a state can have positive, negative, or zero norm.

At first sight this appears incompatible with probability. But indefinite-metric quantum theories can possess an additional grading or symmetry that identifies the physical probability




Figure 1: The three-level structure: one complex spectral covariance, two covariant real forms (involutions), two completions. The two branches agree on all complex correlation functions and differ in which operators are Hermitian and how probabilities are represented. At the resonant (Jordan) boundary only the Krein branch continues as a nondegenerate completion.

structure. In the Bateman–Turok construction [7], a hidden ghost-parity symmetry plays this role.

The positive and Krein descriptions sacrifice different conventional requirements. For an unpaired ghost oscillator, one may try to demand:

1. positive norm;
2. positive energy;
3. standard reality of the original field.

The three cannot all be retained simultaneously [28]. A complex quarter-turn of the ghost variables can make the energy and norm positive, but the original amplitude becomes anti-Hermitian; i times the amplitude is the physical Hermitian observable. Alternatively, the original field can remain real and the energy spectral condition can be retained, but the state space becomes indefinite. This gives two real forms of a common complex system:

	positive completion	Krein completion
energy	positive	positive
norm	positive	indefinite
original field reality	rotated	standard

The complex equations and complex covariance may agree even though the physical adjoints differ. Figure 1 summarizes the resulting three-level structure.

12 A necessary representation-theoretic distinction

There are two different ways to compare the positive PT construction with an ordinary oscillator Fock space, and conflating them produces contradictions [26].

12.1 Physical Dyson transport

For each mode, the PT ground state is exactly the Dyson pullback of the positive normal-form vacuum,

$$\psi_k = \rho_k^{-1} \phi_k. \quad (33)$$

Equip the PT space with the metric $\eta_k = \rho_k^\dagger \rho_k$. Then

$$\rho_k : \mathcal{H}_{\text{PT},k} \longrightarrow \mathcal{H}_{\text{normal},k} \quad (34)$$

is unitary with respect to the physical inner products, and $\rho_k \psi_k = \phi_k$ *exactly*. The pointed modewise unitaries therefore tensor together without any vacuum-overlap obstruction: the completed positive PT theory is globally equivalent to its positive normal-form theory.

12.2 Identity embedding in an auxiliary standard CCR algebra

One may instead place both Gaussian wavefunctions in the same standard Schrödinger L^2 space, retain the same canonical variables and standard adjoint, and compare the vectors through the identity embedding. Under that comparison, the per-mode Gaussian fidelity approaches $\sqrt{3}/2$, the expectation of the standard auxiliary number operator approaches $1/3$, and the infinite products of these auxiliary standard-CCR states are disjoint [50, 26].

This does not contradict the physical Dyson equivalence: the two comparisons identify observables differently.

identity embedding of auxiliary variables $\neq$ Dyson transport of the physical algebra.

This is an example of why field representation theory cannot be discussed without specifying the algebra, its involution, the embedding between descriptions, and the chosen completion.

13 The ultraviolet Fock-sector hierarchy

The preceding comparison placed a selected PU vacuum beside the split two-field vacuum. Paper III also asks a different question: when do two *selected* vacua, belonging to different mass pairs (m_1, m_2) and $(\bar{m}_1, \bar{m}_2)$, define the same ultraviolet Fock sector in common field variables? Writing $\Sigma = m_1^2 + m_2^2$ and $\Pi = m_1^2 m_2^2$, the modewise fidelity obeys

$$1 - f(k) = \frac{(\Sigma - \bar{\Sigma})^2}{12k^4} + O(k^{-6}). \quad (35)$$

If $\Sigma = \bar{\Sigma}$, the k^{-6} term also vanishes and the first remaining obstruction is proportional to $29(\Pi - \bar{\Pi})^2/(576k^8)$. Since the two symmetric invariants (Σ, Π) determine the unordered mass pair, the ultraviolet classification terminates:

spatial dimension	ultraviolet sector invariant
$d \leq 3$	none
$4 \leq d < 8$	$\Sigma = m_1^2 + m_2^2$
$d \geq 8$	(Σ, Π) , the unordered mass pair

At $d = 4$ and $d = 8$ the corresponding obstruction is logarithmic, so the endpoints belong to the higher-dimensional rows shown.

This is an *ultraviolet* classification, not automatically a global GNS-equivalence theorem. For strictly positive masses the infrared is regular. Against the doubly massless $\square^2$ vacuum, however, the relative occupation behaves as k^{-3} and its infrared integral diverges for $d \leq 3$. Whether the massless anchor belongs to the same global sector in those dimensions therefore remains open. This is why “one ultraviolet sector” must not be strengthened to “one global representation.”

14 The fourth-order vacuum is locally well behaved

Ordinary four-dimensional Klein–Gordon two-point functions have a leading short-distance singularity proportional to $1/\sigma_+$, where $\sigma_+(x, y)$ is the regularized squared geodesic separation. In the fourth-order divided difference, this mass-independent leading term cancels. The result has the Hadamard-like form

$$W_{12}^+(x, y) = V_{12}(x, y) \log \sigma_+(x, y) + H_{12}(x, y), \quad (36)$$

where V_{12} and H_{12} are smooth and

$$V_{12}(x, x) = \frac{\varepsilon}{16\pi^2}. \quad (37)$$

There are further terms such as $\sigma \log \sigma$, so the two-point function is not simply a constant logarithm plus a smooth remainder. Nevertheless, its wavefront directions are the standard positive-frequency Hadamard directions, in the microlocal sense of [51].

This is the natural fourth-order analogue of the Hadamard property [27]. It says that the global ghost or completion problem does not automatically produce a pathological local ultraviolet singularity: the vacuum can be globally unusual in its real form or representation, yet locally well behaved in the microlocal sense required for renormalized observables [52, 53].

15 When interactions are switched on

Everything so far concerns free theories. The fifth paper of the series [29] asks the next question: can the free positive or Krein structure be continuously deformed when nonlinear processes are switched on? The answer has four parts: generically yes at low perturbative order away from resonances; no on specific energy-conserving conversion channels; in field theory continuous momentum makes such channels kinematically available for generic split masses, with the obstruction verified on a nonempty open shell subset; and an exact doubled sector-exchange symmetry can survive even though the analytic deformation of the canonical positive pointed-sector metric fails there. Figure 2 summarizes the mechanism.




Figure 2: The interaction mechanism of [29]: metric deformation meets on-shell conversion; isolated resonances in the oscillator become kinematically available scattering shells in the field theory, with the obstruction verified on a nonempty open subset; at the massless boundary a doubled exchange structure survives.

15.1 Deforming the positive metric

Perturb the free theory, $H_\zeta = H_0 + \zeta V$, and recall the free Dyson map with $\rho_0 H_0 \rho_0^{-1} = h_0 = h_0^\dagger$. One seeks a corrected map

$$\rho_\zeta = e^{-\frac{1}{2}(\zeta R_1 + \zeta^2 R_2 + \dots)} \rho_0, \quad (38)$$

keeping the transformed Hamiltonian Hermitian order by order. At first order,

$$[h_0, R_1] = v^\dagger - v, \quad v = \rho_0 V \rho_0^{-1}. \quad (39)$$

The interaction introduces a new “bad” non-Hermitian part, and the correction R_1 asks whether the measuring geometry can be adjusted to absorb it. The key principle is simple: a correction can remove terms that move states between *different* energies; it cannot remove a term that acts entirely among states with exactly the *same* energy. When two states have different energies, the metric can distinguish them; when they have exactly equal energy, the interaction is resonant and cannot be rotated away. Technically, the obstruction is the energy-conserving part of the source,

$$\mathfrak{o}_+ = \Pi_{\ker \text{ad}_{h_0}} S, \quad (40)$$

the projection onto processes satisfying exact free-energy conservation.

15.2 A finite oscillator fails only at special resonances

For the cubic PU interaction at generic unequal frequencies, the positive metric can be corrected through the computed orders — the explicit generators exist and the transformed Hamiltonian

is Hermitian. But at

$$\omega_1 = 3\omega_2, \quad (41)$$

one high-frequency quantum can convert into three low-frequency quanta without changing the total energy, and the second-order correction is obstructed by exactly that conversion operator, $a_1 a_2^{\dagger 3} - a_1^{\dagger} a_2^3$. At $2\omega_1 = 3\omega_2$ a third-order obstruction appears. A transfer-lattice rule organizes which conversion monomials can first occur at each order. Its first new prediction has been confirmed computationally: at $\omega_1 = 5\omega_2$ the second- and third-order classes vanish, while the fourth-order class is a nonzero $1 \leftrightarrow 5$ conversion proportional to $a_1 a_2^{\dagger 5} - a_1^{\dagger} a_2^5$. This confirms one prediction of the proposed hierarchy; the general $p:q$ rule remains a conjecture.

Imagine two musical notes: usually they are unrelated, but if one note has exactly three times the frequency of the other, one vibration can split into three without changing the total energy — at that point the interaction is perfectly synchronized with the system.

The 3:1 obstruction is more than a failure of one formula. Sufficiently excited resonant multiplets develop complex-conjugate energies, $E_{\pm} = E_R \pm i\Gamma$, at second order. This is formally defined perturbative PT breaking: [29] constructs the antilinear symmetry explicitly, proves that it commutes with the interacting Hamiltonian, and certifies the complex pair *exactly* in the effective resonant multiplet (by rational algebra and Sturm counting, not only numerically). Where the complex pair persists, no positive-definite invariant metric can make the Hamiltonian equivalent to an ordinary self-adjoint one. Two levels of conclusion must not be merged: this complex-spectrum statement is established in the verified resonant oscillator setting; the field-theory result below is presently a perturbative metric obstruction, not yet a demonstrated complex spectrum.

15.3 The momentum continuum turns exceptional resonances into ordinary scattering

For an oscillator, the frequencies are fixed numbers and exact resonances require special ratios. For a field, $E_a(k) = \sqrt{k^2 + m_a^2}$, and momenta vary continuously: one can satisfy

$$E_H(k_1) + E_L(k_2) = E_L(k_3) + E_L(k_4) \quad (42)$$

together with momentum conservation on open sets of scattering data, for *any* split masses. The key process is the branch-changing scattering $H + L \rightarrow L + L$. Paper V shows that the second-order positive-metric source is nonzero on an open scattering-shell subset (exactly, at a rational kinematic point, hence on an open neighborhood — not claimed at every kinematic point): the continuum turns the oscillator's isolated resonance obstruction into an obstruction on a nonempty open part of a scattering shell. In a small mechanical system, resonance requires the machine to be tuned to a special ratio; in a field, particles can change their momentum, and even if their masses are not specially tuned, their motion can supply the exact energy match.

The canonical positive completion therefore remains meaningful locally and perturbatively away from resonant processes, but it cannot be deformed analytically into a positive pointed-sector metric on the verified branch-changing shell subset. One qualification is essential: this establishes that *no analytic deformation of the canonical metric* exists there — not that no conceivable positive completion of any kind exists. For the oscillator with certified complex energies the stronger conclusion is available; for the field theory the perturbative statement is what is proved.

15.4 Why the obstruction waits until second order

The perfect-square cubic vertex is exactly the Källén polynomial, $\mathcal{V}_3 = \frac{1}{2}\lambda_K(p_1^2, p_2^2, p_3^2)$ (Section 14 notation). It vanishes for three ordinary massless legs and has a second-order zero at the confluent massless point; and the transported reality factors eliminate the first-order positive obstruction on the physically open decay channel. So the positive metric survives the first test. But at second order, two cubic interactions combine into a branch-changing $2 \rightarrow 2$ process, and the obstruction appears: a single interaction event is too constrained to cause the problem, while two interaction events in sequence can convert the particles while preserving total energy — and that is where the positive geometry fails.

15.5 A different structure survives: two mirror Jordan sectors

The perfect-square theory admits an exact rewriting, introduced by Bateman and Turok [7], who also identified its exchange symmetry as ghost parity and as charge conjugation for an internal $O(1, 1)$ symmetry; the sector-level reading given here is what the interaction paper adds. Introducing an auxiliary field and the nonlinear change of variables $U = e^{\lambda\phi}$, $V = (\psi/\lambda)e^{-\lambda\phi}$,

$$\mathcal{L} = -\partial_\mu U \partial^\mu V + \frac{\lambda^2}{2} U^2 V^2. \quad (43)$$

The constant zero-potential stationary backgrounds form two branches, $(U_0, 0)$ and $(0, V_0)$, and the exact exchange $U \leftrightarrow V$ maps one branch to the other. Around $(1, 0)$ the linearized equations are $\square v = 0$, $\square u = \lambda^2 v$ — a Jordan chain generated by the interaction itself; around $(0, 1)$ the chain has the opposite orientation. The complete theory thus has two mirror valleys: choosing one valley gives one orientation of the Jordan system, and the exact symmetry does not rearrange particles inside that valley — it carries the entire valley to its mirror.

Three distinctions matter here. The exact symmetry belongs globally to the *extended* (U, V) theory; its expression in the original (ϕ, ψ) variables is a chart-transition map, singular at the chosen perturbative vacuum; and it is not a ghost-number sign inside one Fock sector. It may ultimately form part of an antiunitary CPT-like structure, but what is proved is a linear sector exchange, not a quantum CPT theorem.

Finally, the ordinary split-theory ghost parity becomes singular at the merge. In the confluent Jordan basis $c = (b_+ + b_-)/2$, $d = (b_+ - b_-)/\delta$, the branch parity becomes

$$P_\delta = \begin{pmatrix} 0 & 2/\delta \\ \delta/2 & 0 \end{pmatrix}, \quad P_\delta^2 = 1, \quad (44)$$

with no bounded limit on a single Jordan chain: as the two particle types merge, the old rule “one is even and one is odd” becomes infinitely badly conditioned. The only finite remnant is to exchange the entire Jordan system with its mirror copy: after doubling by the oppositely oriented sector, the singular parity becomes a finite sector exchange. This confluent result is the *linear bridge* to the independently exact nonlinear $U \leftrightarrow V$ symmetry: what the regulated parity converges to is the linearization of that exchange around the two mirror backgrounds — no stronger identification is claimed.

The relation between the two structures is sharp in the verified perturbative setting. At the rational shell point used to certify the field obstruction, the complete reachable second-order on-shell matrix is *exactly* pseudo-Hermitian with respect to ghost-branch parity, and its entire

positive-metric obstruction lies in the parity-odd block. On the same matrix elements the Krein pairing is preserved while the analytic deformation of the canonical pointed positive metric is obstructed. The two escapes from the ghost problem therefore *separate*: the Krein completion is not a hidden version of that positive completion. This does not exclude a nonanalytic or differently pointed positive construction, nor does it prove a complex field-theory spectrum.

There is a further boundary distinction. Exact pseudo-Hermiticity of the split on-shell matrix is weaker than the one-sided charge condition used in the Bateman–Turok generalized Born rule. A canonical map to their cross-paired charge basis sends the regulated pointed vacuum to a Gaussian with

$$\frac{S_{UU}}{S_{VV}} = \left(\frac{\delta}{2g} \right)^2 = \frac{\epsilon}{g}, \quad \delta^2 = 4\epsilon g. \quad (45)$$

Thus both charge signs are intrinsic at split mass. The image becomes one-sided only on the $O(1,1)$ -symmetric confluent line $\epsilon = 0$; the Bateman–Turok massless point additionally has $\mu^2 = 0$. A self-contained charge-null lemma then makes every strictly negative component null and orthogonal to the neutral component. The remaining capstone calculation is to transport the explicit obstruction through this boundary limit and evaluate its generalized Born trace.

Terminology.

Positive pseudo-Hermitian metric: a positive inner product obtained by changing the adjoint through a Dyson map. *Metric deformation*: an order-by-order correction of that inner product after an interaction is introduced. *Obstruction*: the energy-conserving part of the unwanted non-Hermitian interaction that cannot be generated by a metric correction. *Pointed sector*: a quantum construction built around one selected vacuum background. *Doubled theory*: a completion containing both mirror Jordan sectors. *Sector-exchange involution*: the exact map exchanging the two mirror backgrounds and reversing the Jordan-chain orientation. *Spectral breaking*: the appearance of complex-conjugate energy levels, which excludes a positive invariant metric where the complex spectrum persists. “Ghost parity” is used only with a qualifier: split branch-number parity, the exact $U \leftrightarrow V$ exchange, or a (future) antiunitary CPT-like operation are different objects.

16 Why gravity is not merely five copies of the oscillator

The scalar PU oscillator has no gauge symmetry. Gravity does.

Consider linearized scalar-free Einstein–Weyl gravity around flat spacetime: the Einstein term plus a tuned quadratic-curvature term ($\alpha R_{\mu\nu}^2 + \beta R^2$ with $\alpha = -3\beta$). The linearized spectrum contains a massless graviton with two helicities and a massive spin-2 branch with five helicities [24]. A naive polarization-by-polarization treatment might suggest that every massive helicity belongs to an independent PU pair. Gauge reduction shows otherwise [28]: the reduced phase space has the structure

$$\Gamma_{\text{phys}} \cong 2\Gamma_{\text{PU}} \oplus 2\Gamma_- \oplus \Gamma_-. \quad (46)$$

The two helicity-2 sectors are genuine fourth-order PU pairs: each contains a massless graviton mode and a massive ghost partner. The helicity-1 and helicity-0 sectors contain only *unpaired* second-order massive ghosts: their apparent massless partners lie entirely in the diffeomorphism gauge orbit and disappear under reduction.

Gauge reduction preserves PU pairing only in helicity ± 2 .

This is the first major way in which gravity differs from the scalar model.

17 Lorentz covariance forbids hybrid solutions

The five massive helicities $+2, +1, 0, -1, -2$ form one irreducible massive spin-2 representation. They are not five independent species that can be quantized separately: a Lorentz transformation can mix them.

By Schur’s lemma, an invariant Hermitian form on the irreducible spin-2 polarization space is proportional to the identity. Its sign or reality assignment must therefore be uniform across all five helicities. This rules out a tempting hybrid construction in which the helicity-2 modes receive the positive PU metric while the helicity-1 and helicity-0 ghosts are treated in a Krein space: such an assignment is not Lorentz covariant.

Within the class analyzed in the gravity paper [28] — free, translation-invariant, quasifree, mode-local constructions compatible with the reduced symplectic form, the spectral condition, and Poincaré covariance — there are exactly two covariant completion classes:

1. a positive pseudo-Hermitian completion in which the full massive spin-2 multiplet has uniformly rotated reality;
2. a Krein completion in which the gravitational field retains standard reality and the entire massive spin-2 multiplet carries a uniform negative sign relative to the positive massless graviton sector.

The two completions induce the same complex gauge-invariant covariance, while differing in their involution and completion. This is the gravitational version of the real-form distinction of Figure 1.

18 The gauge-invariant gravitational bridge

The equality of the two complex covariances is not merely a polarization-by-polarization analogy. The reduced helicity kernels have residues fixed by the reduced symplectic form. Their ratios $1 : M^2 : \frac{1}{3}$ match the gauge-invariant contractions of the massive spin-2 projector,

$$\frac{1}{2} : \frac{M^2}{2} : \frac{1}{6} \tag{47}$$

so they reassemble, modulo pure-gauge bi-distributions, with one overall coefficient $4/c_1$. The massless shell couples only to the transverse-traceless polarizations. Thus both real forms determine the same complex spectral functional on the algebra of linear gauge-invariant observables, rather than merely agreeing in a chosen helicity frame.

The massive projector itself contains apparently singular terms $p_\mu p_\nu / M^2$. Applying the linearized Weyl-tensor map annihilates all of them: the Weyl–Weyl kernel is gauge independent, covariant, and regular as $M \rightarrow 0$, with the standard positive-frequency Hadamard wavefront directions. This cancellation is why the conformal limit below is formulated on the curvature-generated regular subalgebra. The raw scalar potential kernel does not have a regular limit; the divergent direction becomes Weyl-null and must first be quotiented out.

Relation to CPT language. Mannheim’s CPT-based positive scalar product belongs, away from the conformal locus, to the positive class just described, with its uniformly rotated massive reality [23]. At the Jordan boundary the no-go theorem excludes a time-independent, nondegenerate, positive-definite invariant form; it does not exclude the intrinsically Jordan realization of Bender and Mannheim, which uses nonstationary Jordan states and a degenerate pairing [6]. The two statements are therefore compatible: the theorem delimits which positive completion terminates, rather than declaring the equal-frequency theory nonexistent.

19 The conformal Jordan boundary

Let the coefficient of the Einstein term tend to zero while the quadratic-curvature coupling remains fixed. The massive spin-2 mass then tends to zero, and the different helicity sectors behave differently [28].

Helicity ± 2 . The massless and massive branches merge, producing two $\square^2$ Jordan systems. Each polarization contributes two configuration-space solutions, giving four modes in total.

Helicity ± 1 . The vector ghosts have an ordinary massless limit. They remain second-order modes rather than becoming Jordan partners.

Helicity 0. The scalar kinetic normalization tends to zero. At the conformal point a new Weyl gauge symmetry appears, $\delta h_{\mu\nu} = 2\sigma\eta_{\mu\nu}$: the scalar direction becomes null in the presymplectic form and is removed by the quotient.

The conformal count is therefore $4 + 2 + 0 = 6$. The positive split-frequency metric runs to infinite Cartan distance and has no nondegenerate positive continuation through the Jordan boundary; the complex spectral covariance and the standard-reality Krein form can continue, provided the newly null Weyl direction is quotiented out. The algebra of observables changes rank at the boundary, so convergence must be formulated on a fixed regular family of observables — most naturally through gauge- and Weyl-invariant curvature quantities. (Boundary constructions that instead *remove* one branch of the solution space, as in [54], and the richer phase diagram at nonzero cosmological constant [55, 56], are related but distinct stories.)

20 Paper VI: interaction obstruction in Einstein–Weyl gravity

Paper VI [30] asks the interacting question left open by the free gravity classification. Scalar-free Einstein–Weyl gravity around flat space contains a healthy massless graviton h and a ghost-signed massive spin-2 multiplet M . Paper IV found two covariant free real-form classes: a positive completion obtained by uniformly quarter-turning the complete massive multiplet, and a standard-reality Krein completion with uniformly negative massive signature. Paper V supplied the diagnostic. If ρ_0 is the free Dyson map, an analytic interacting deformation must solve cohomological equations

$$[h_0, R_n] = S_n, \tag{48}$$

so the projection of S_n onto exact free-energy degeneracies must vanish.

Gravity is protected at first order for two independent reasons. The Einstein solution space is an exact nonlinear subsector of Einstein–Weyl gravity, so tree amplitudes with exactly one

massive external field and otherwise massless gravitons vanish. The remaining odd-massive cubic vertex MMM is nonzero on complex kinematics but has no real equal-mass $1 \leftrightarrow 2$ shell. Consequently the first-order positive-metric obstruction vanishes.

At second order the process

$$M + M \longrightarrow M + h \quad (49)$$

is physically open. Paper VI establishes its nonvanishing by two independent calculations. An Einstein-frame factorization calculation produces a nonzero massive-pole residue. A separate calculation in the original fourth-order variables includes the quartic contact term, all s -, t -, and u -channel exchanges, physical polarization and equation-of-motion conditions, the reverse-process adjoint, Ward identities, and two internal gauge representatives. At an interior rational point of the real physical shell it gives a nonzero exact amplitude.

The essential bridge is not inferred from that amplitude's phase alone. In a regulated stationary Fock space, define the second Born operator

$$B_{2,+}(E) = P_E(v_2 + v_1 Q_E(E - h_0)^{-1} Q_E v_1) P_E. \quad (50)$$

With the spectral solution for R_1 , Paper VI proves the identity

$$P_E \left[(v_2^\dagger - v_2) + \frac{1}{2} [R_1, v_1 + v_1^\dagger] \right] P_E = B_{2,+}(E)^\dagger - B_{2,+}(E). \quad (51)$$

The covariant contact-plus-exchange calculation matches this stationary Born operator after the usual propagator decomposition, external residue normalization, and identical-particle factors. The calculation therefore evaluates the complete deformation source, not merely a phase-sensitive scattering amplitude.

Covariance quarter-turns the complete massive multiplet. Counting the massive fields at all vertices and the compensating signs of internal massive propagators shows that every connected diagram with E_M external massive legs acquires the total phase $(-i)^{E_M}$. The process $MM \rightarrow Mh$ has $E_M = 3$; its nonzero real standard-frame Born block thus becomes anti-Hermitian in the positive frame. Equation (51) then gives a nonzero second-order shell cocycle. Hence no analytic second-order deformation based at the canonical Paper-IV positive real form exists in the stated tree-level perturbative class on a nonempty open subset of this shell.

The Krein result is different. Agreement with the free gravitational signature and natural particle-number-diagonal, cluster-multiplicative second quantization gives the conventional lift

$$J_F = (-1)^{N_M} \quad (52)$$

on physical BRST cohomology. The process crosses its positive and negative sectors, but oddness under a $\mathbb{Z}_2$ grading does not imply nullity: the obstruction has a nonzero quadratic Krein trace. Moreover, the verified nonzero MMM and MMh vertices force every uniform abelian charge on the irreducible massive multiplet, with neutral graviton, to be trivial. The scalar boundary's one-sided $O(1,1)$ charge mechanism therefore has no direct gravitational analogue.

This conclusion is deliberately limited. Paper VI excludes an analytic second-order deformation of the canonical positive free metric and a null interpretation under the conventional multiplicative massive-number grading. It does not exclude indefinite Krein evolution, nonanalytic or differently pointed positive metrics, momentum-dependent or non-factorizing involutions, singular conformal-boundary limits, or boundary conditions that truncate part of the fourth-order solution space.

21 Papers VII–VIII: residual states and covariant transport

Papers VII–VIII [31, 32] return to the pure-Weyl boundary of Paper IV, but they answer two different questions. Paper VII asks what survives the residual gauge quotient after a state polarization has been chosen. Paper VIII asks whether that state-side answer is computed by the covariant compactly supported metric BV complex. The second theorem imports the first; it does not repeat its Chevalley–Eilenberg calculation.

21.1 The selected closed-universe problem

Both papers work on

$$M = \mathbb{R} \times S^3 \quad (53)$$

and make the same explicit boundary choice. The spatial slice is closed, there is no asymptotic charge or external clock, the cylinder is not deparametrized, and all fifteen conformal reducibilities—including compact time translation D —are gauged. This is not the fixed-cylinder problem in which D is retained as a Hamiltonian.

The charge audit makes this restriction mathematically visible. On the unrestricted locally reduced linearized space $\mathcal{P}_{\text{lin}}$, D has the nonzero integrable kernel $M_D = -\frac{1}{2}J_{\text{conf}}D$ and is a physical Hamiltonian symmetry. On the common Taub zero fibre $\mathcal{P}_{\text{Taub0}}$, however, the inclusion ι obeys

$$\iota^* \iota_{X_D} \Omega_\Sigma = d(\iota^* \mu_D) = 0. \quad (54)$$

Thus the compact verdict is sector-dependent: the quotient used below is legitimate only after the explicit full moment-map zero restriction. The cross-programme validation dossier keeps this classical statement separate from Paper IX’s distinct Berger-clock tangent theorem, its arity-three Taylor result, and the still-open boundary, higher-order, and quantum questions.

Three reductions remain conceptually distinct:

$$\begin{aligned} \text{covariant metric BV complex} &\longrightarrow \text{Cauchy/energy one-particle complex} \\ &\longrightarrow \text{polarized residual state complex.} \end{aligned} \quad (55)$$

Paper VIII proves the first comparison and its compatibility with the second. Paper VII performs the final residual reduction. A local propagating mode can therefore be present in the middle complex while contributing no one-particle class to the last one.

21.2 Paper VII: the state-side theorem

The locally reduced one-particle coefficient module is the parity-complete on-shell Weyl module $\mathcal{W}_+ \oplus \mathcal{W}_-$. Its invariant form has the branch signs

$$J_{\text{conf}} = +I_E \oplus (-I_A) \oplus (-I_L). \quad (56)$$

After the selected positive-frequency Lagrangian polarization, the coefficient algebra is

$$\mathcal{F}_{\mathcal{W}} = \text{Sym}(\mathcal{W}_+ \oplus \mathcal{W}_-), \quad (57)$$

and the residual ghosts give the absolute $\mathfrak{so}(4, 2)$ Chevalley–Eilenberg complex.

Theorem 21.1 (Paper VII: selected residual state theorem). *For the selected formal closed-universe BV–BFV problem, the Taub constraints are the residual moment-map components, endpoint duality and the normalized BV–BFV suspension supply exactly one residual cotangent pair, and the positive-frequency state complex is*

$$\mathcal{H}_{\text{state}} = \mathcal{F}_{\mathcal{W}} \otimes \Lambda^{\bullet} \mathfrak{so}(4, 2)^*. \quad (58)$$

Its centered one-particle group vanishes, while

$$\boxed{H_{\text{res}}^4 = \text{span}\{[W_+^2], [W_-^2]\}, \quad \text{sig } G_{\text{res}} = (2, 0), \quad G_{\text{res}} = I_2.} \quad (59)$$

The same result holds in the selected energy-mode Krein–Fock completion.

The classes in (59) are ghost-dressed deformation or vertex classes. They are not two one-particle gravitons. Their parity combinations descend to the Weyl-square and Pontryagin directions; after the local Euler–Lagrange quotient only the Weyl-square direction is dynamical.

21.3 Paper VIII: the covariant-side theorem

Paper VIII retains the metric variables and adds curvature only through a support-local prolongation. Its analytic task is a chain contraction, not a second residual cohomology computation.

Theorem 21.2 (Paper VIII: covariant causal transport). *The complete free pure-Weyl metric BV complex on the conformal cylinder admits a support-local curvature prolongation and retarded/advanced degree-(-1) maps $\Lambda_{\pm}$ such that*

$$Q\Lambda_{\pm} + \Lambda_{\pm}Q = 1, \quad \text{supp}(\Lambda_{\pm}f) \subseteq J^{\pm}(\text{supp } f), \quad \Lambda_+^{\sharp} = \Lambda_-. \quad (60)$$

Their causal difference induces a pairing-compatible quasi-isomorphism from compactly supported covariant data to the smooth global cylinder solution complex. It recovers the fifteen residual endpoint pairs without duplication and transports the $\mathfrak{so}(4, 2)$ action to cohomology. Consequently, using Theorem 21.1 as an input,

$$\boxed{H_{\text{cov}}^4 \cong H_{\text{res}}^4 = \text{span}\{[W_+^2], [W_-^2]\}, \quad \text{sig } G_{\text{cov}} = (2, 0), \quad G_{\text{cov}} = I_2.} \quad (61)$$

No residual CE matrix is recomputed in this argument.

The causal proof uses a local curvature/Weyl–Cotton presentation to carry the physical helicities, support-local deformation retracts for auxiliary and prolongation variables, and a hybrid algebraic/Green contraction for the propagating endpoint. The positive PDE symmetrizer and the indefinite BV/Krein pairing play different roles: the former gives causal evolution, while the latter is transported through the current and Green identities.

A subsequent abstract theorem isolates the reusable mechanism. For a finite-order support-local cyclic strong deformation retract $E \rightleftarrows_p^i C$ with homotopy h , same-sided endpoint chain homotopies transfer by

$$\Lambda_{C, \pm} = h + i\Lambda_{E, \pm}p. \quad (62)$$

The chain identity, advanced/retarded support, cyclic adjoint relation, finite direct sums, and finite-order cyclic shears are preserved. The complete Berger 54- and 64-row complexes are

exact consumers. This is a conditional $G2$ transfer theorem: it assumes endpoint Green hyperbolicity and does not cover timelike boundaries, pseudodifferential projectors, Hadamard products, interactions, or quantum claims.

The Nariai consumer now has a first stability result beyond its exact background. Along one global transverse Einstein tangent, the normalized basic perturbation lemma differentiates the complete rank-310 retract and its metric endpoint. The induced variations remain finite-order and support-local, preserve the cyclic adjunction identities, and give formal advanced/retarded chain homotopies with the same time orientation. This is a global formal theorem at $\epsilon = 0$; it does not construct a complete smooth nonzero- ϵ family or an open ball of causal backgrounds.

21.4 Relation and combined boundary




Figure 3: The VII–VIII relation. Paper VII computes the right-hand cohomology. Paper VIII proves that the covariant left-hand complex transports to it; the arrow does not conceal a second CE calculation.

Together the two papers complete the free classical residual branch in its stated category. They do not establish a positive graviton Hilbert space, an integrated $SO(4, 2)$ group representation, a preferred alternative boundary polarization, a BRST-compatible Hadamard state, nonlinear stability, a renormalized quantum master equation, anomaly cancellation, or interacting unitarity.

Paper VII determines the selected residual cohomology. Paper VIII proves that the covariant causal metric complex computes and preserves that already determined module and pairing.

22 Papers IX–XI and the nonlinear selection bridges

The later papers ask whether the free and reduced structures above remain meaningful after a physical clock, electromagnetism, ordinary gravitational waves, and the first nonlinear consistency conditions are introduced. They do not yet form a single nonlinear universe model. They provide exact incidence maps and obstruction tests which such a model must satisfy.

22.1 Paper IX: a healthy clock and the correct stabilizer

Paper IX [33] proves two related but carefully separated facts on the positive-energy Berger background. First, the rotating scalar phase is a genuine clock coordinate with nonzero conserved momentum, while the compact fixed-coupling constraint forces its charge variation to vanish on every legal smooth linearized tangent. Hence raw cylinder time translation D is presymplectically null on that declared linearized phase space. Second, the background-fixing

generator in co-rotating variables is $K = D - \omega R$, not raw D . The complete 54-row causal BV complex and action-derived q_2, q_3 admit cyclic causal Cartan primitives for K through arity three. Raw D has an affine zeroth-order component and is not covered by that nonlinear Cartan identity. The theorem therefore establishes a healthy clock counterexample to an automatic “clock makes D charged” claim, but does not establish genericity, all-orders closure, or a quantum clock.

22.2 Paper X: ordinary waves inside the fourth-order theory

Paper X [34] works on the magnetically supported compact product $\mathbb{R} \times S^1 \times S^2$, a common exact background of Einstein–Maxwell and Weyl–Maxwell theory. It constructs the complete standard Einstein–Maxwell harmonic tangent and proves that the identity map

$$\iota_{\text{EM}} : \mathcal{E}_{\text{Einstein}--\text{Maxwell}} \longrightarrow \mathcal{E}_{\text{Weyl}--\text{Maxwell}} \quad (63)$$

is injective. The Lee–Wald pullback is nondegenerate on every standard block, but it is not the Einstein–Maxwell symplectic form: on a generic harmonic it is composed with the spectral polynomial

$$p_\lambda(M) = 1 + \frac{3}{2}(M - \lambda). \quad (64)$$

Thus ordinary gravitational and electromagnetic waves are present and nonnull, but their phase-space weighting changes in the containing theory.

The complete generic axial target also contains a two-dimensional extra quotient. Its direct four-dimensional Lee–Wald form is nondegenerate and positive in the paper’s declared positive-frequency current convention; the combined axial current has inertia $(3, 1)$, with the negative direction in the embedded Einstein block. This is a classical current theorem before stabilizer reduction. It is neither a Hilbert-space norm nor a quantum ghost verdict, and it does not supply asymptotic scattering.

A post-paper relative construction strengthens the incidence statement. It globalizes the harmonic coefficient maps to one finite-order, natural, support-local chain morphism on the common tensor bundles and completes the linear mapping-cofiber sequence

$$0 \longrightarrow \mathcal{E}_{\text{Einstein}--\text{Maxwell}} \longrightarrow \mathcal{E}_{\text{Weyl}--\text{Maxwell}} \longrightarrow \mathcal{E}_{\text{extra}} \longrightarrow 0 \quad (65)$$

through every ghost, field, equation, identity, residual, and large-gauge endpoint row. It exports the Einstein form, the Weyl form pulled back to the image, and the direct relative form as three distinct objects. This separation is forced: on every generic $\ell \geq 2$ shell the Einstein form has inertia $(2, 0)$ while the pulled-back Weyl form has inertia $(1, 1)$. Hermitian inertia is invariant under congruence, so no real product-equivariant standard-pairing cyclic correction, chain homotopy, or exact current improvement can identify them. Thus (65) is a certified noncyclic linear triangle, not a symplectic equivalence. The same-background 38-row Einstein–Maxwell and Weyl–Maxwell Taylor packages through q_3 are now certified and independently replay their nilpotency and cyclicity identities. The strict arity-two relative defect

$$\Delta_2 = q_2^{\text{WM}}(f_1, f_1) - f_1 q_2^{\text{EM}} \quad (66)$$

has 50,854 exact coefficients and is nonzero. Its Taub pairing rules out a support-local f_2 on the full smooth periodic fixed-bundle carrier. On the complete standard source cohomology, however, the relative quadratic operation is exactly the five-charge receiver (H, P_x, J_1, J_2, J_3) .

That finite-dimensional receiver cannot itself be the target of a nonzero support-local differential operation: compactly supported inputs have compactly supported output, while a nonzero constant charge row has global support. The correct local object is therefore the horizontal current cone

$$\Omega_H^3(M; \mathfrak{g}_{\text{stab}}^*) \xrightarrow{d_H} \Omega_H^4(M; \mathfrak{g}_{\text{stab}}^*), \quad (67)$$

with charges obtained only after Cauchy-slice integration. The canonical relative Hessian Green concomitant now satisfies its exact local divergence identity on all fourteen physical rows and all coefficient jets; its antisymmetrization supplies a local current of maximum derivative order three. Precomposition with the five stabilizer actions, horizontal comparison with Lee–Wald, cyclic BV dual rows, and equality with all five integrated charge blocks remain the nonlinear relative gate.

22.3 Paper XI: the first retained gravity–light bracket

Paper XI [35] imports the repaired cyclic gravity–clock–Maxwell q_2, q_3 tensors and transfers them through one exact typed cyclic 64-to-36 strong deformation retract. In the suspended convention,

$$\ell_3 = \pi q_3(\iota, \iota, \iota) + \pi q_2(Sq_2(\iota, \iota), \iota) + \text{graded permutations}. \quad (68)$$

The resulting representative is nonzero. Two zero-derivative evaluations give exact coefficients +1 and −1 in the gravity- and Maxwell-equation sectors. Its 25,950 exact sparse coefficients satisfy the complete arity-three identities and the imported odd-Darboux cyclicity convention. The only nonempty exchange channel has 342 full-complex coefficients, all on the contractible clock-dual row, and hence projects to zero; a one-entry projection mutation exposes all 342 terms. The surviving retained term is therefore the direct q_3 contact term for this frozen retract.

This is a precise theorem-frozen representative-level interaction result, not yet a physical branch theorem. The retained 36-row carrier does not admit the naive support-local same-bundle canonical Einstein-like/extra-Weyl projector: the endpoint block

$$A_{10} = \square_2^2 + V_2 \quad (69)$$

has degree-two symbol entries which are not divisible by the required scalar wave polynomial. A natural support-local rank-46 STF2 graph prolongation is an exact cyclic SDR over the retained 36-row complex, but its contractible complement cannot repair the normalized subprincipal projector defect. The canonical support-local branch split is therefore obstructed. An exact branch manifest requires a noncontractible filtered or mixed-bundle carrier, or an explicitly declared reduced-mode/nonlocal decomposition. The mixed ℓ_3 is nonremovable in the declared filtered-cyclic F_2/F_3 class, but a nonzero operation on final cohomology and a branch-resolved physical mixing table remain later gates rather than prerequisites for the frozen Paper XI theorem. The successor full-ghost calculation changes the broader disposition: the term is cyclically removable through first jet, and the second-jet image test is still open. Thus restricted nonremovability is retained as a scoped result, not advertised as an invariant interaction.

22.4 Charge-fibre selection and observable records

The charge-fibre Taub bridge [37] applies the next nonlinear integrability test to Paper X’s linear modes. On the fixed compact magnetic bundle, the time-translation moment map is

negative definite on the complete real pure-extra generic sector, in both parities, for every $\ell \geq 2$ and every compact momentum. Consequently no nonzero pure-extra generic tangent extends to second order in that charge fibre. This does not delete the linear extra modes. It says that they are not tangent by themselves to the nonlinear fixed-bundle solution space.

The obstruction changes under mixing. At $\ell = 2, m = 0, k = 0$, one balanced Einstein-minus-extra tangent cancels all five stabilizer moment maps and has an explicit complete second-order correction. The exact contrast is

pure extra generic tangent: obstructed, balanced Einstein-extra tangent: second-order extendible.	(70)
--	------

This exhibits a singular nonlinear tangent cone rather than a direct sum of independent Einstein and extra universes. It does not classify the full nonlinear solution cone or prove all-orders extension.

Post-freeze bridge certificates [38] now substantially generalize the moment-map side of that contrast. After exact Gram normalization, the complete generic $k = 0$ common-zero cone is the rank-constrained positive-semidefinite locus of three branch density matrices $\rho_{+, \ell}, \rho_{e, \ell}, \rho_{-, \ell}$ cut out by the energy and three rotation equations. The Paper 91 ray is a boundary of a two-parameter axisymmetric $\ell = 2$ face, and every point of that declared face has an explicit complete second-order correction. For nonzero compact momentum, the paired $+k, -k$ problem likewise has an exact moment-map cone and a nonzero standing-wave face. The physical $\ell = 1$, homogeneous, twist, and electric-charge blocks have also been classified at the moment-map level. Beyond that classification, the complete exceptional $\ell = 1, k = 0$ all- m pure cone is excluded by a direct resonance no-go. Finite-harmonic generic $k = 0$ zero-cone tangents and one fixed- $|k|$ opposite-momentum zero cone have complete scoped second-order corrections.

The tuned axisymmetric standard-branch calculation is now uniform for every $\ell \geq 2$. Its two parity-resonance equations cut out two mixed sheets and two one-sided planes; imposing the Hamiltonian and circle-momentum moment maps leaves the origin and the two sheets on an exact occupation interval. A separate two-absolute-momentum census has closed 108 axisymmetric $L = 4$ and 56 nonaxisymmetric $L = 1, 3$ branch-basis coefficients, all individually obstructed. This complete 164-coefficient census does not yet determine the joined amplitude zero variety because cross-column cancellations remain possible.

These new results have their own exact lifecycle and do not silently enlarge the theorem-frozen scope of Paper 91. They do not yet classify multiple $|k|$ fibres, infinite-mode completion, every homogeneous/twist/charge mixture, or all-orders extendibility. A reusable structural theorem should still identify the hypotheses under which a definite pure-extra moment map and an opposite-sign Einstein form produce these mixed zero faces.

The newest exact spectrum census and source calculation sharpen the exceptional/global frontier. The homogeneous $\ell = 0, k = 0$ quotient is empty at every nonzero frequency, so no hidden homogeneous fourth-order oscillator remains. In the declared $k = 0$ homogeneous/twist- $\ell = 2$ extra carrier, the resonance equations and five stabilizer moment maps now have a complete common-zero locus. Up to $SO(3)$ it is the aligned orbit

$$a = b = d = 0, \quad A \parallel B \parallel C_{m=0}, \quad B^2 = \frac{Q_e^2}{2} + \frac{2}{3}X, \quad X > 0. \quad (71)$$

There is no off-axis branch. Yet every nonzero point has a polar zero-frequency source with quadratic coefficient $-7B^2$, so a stationary linear operator cannot supply a bounded finite-

quasiperiodic correction. Thus moment-map cancellation and resonance cancellation are not sufficient for bounded second-order extension. The complete finite harmonic source does, however, admit a real finite exponential-polynomial correction: physical Smith factors remove every propagation cokernel, while the remaining compact stabilizer pairings are precisely the five moment maps already set to zero. Thus the same orbit is obstructed in the bounded/quasiperiodic class and extendible in the exponential-polynomial secular class. Causal/retarded correction and all-orders integration remain open.

The observer programme supplies a complementary bridge from solutions to records. It has certified a rank-two linear detector response, six global detector rods with second-order source solvability, a frozen 84-row apparatus handoff, a separate-axis rod-gravity unary theorem, and an exact 72-row causal memory subcomplex. In parallel, an actual retarded Maxwell signal with compact spacetime source support defines a Diff-, Weyl-, Maxwell-gauge-, and raw- D -invariant clock-slice field-strength observable. Its spatially global fixture has $1 + z = 2$, a nonzero reduced probe-mode Poisson bracket, and nontrivial clock-time evolution. These results show that causal records and relational evolution can already be represented without promoting them to particles. The clock-dressed source is an equivariant source family, not invariant at fixed schedule. Two localized massive-emitter Cauchy preparations now produce a leading rank-two record matrix near the receivers. A common localized source, evaluated recoil, the full apparatus Dirac bracket, nonlinear observer morphism, and backreacted rank-two response remain open. The composed scalar 108-row unary is exactly cyclic and its zeroth-order square vanishes, but the first clock-radius coefficient of q_1^2 has a nonzero exact witness. The defect is localized to the missing second jet of the clock canonical transformation (or an equivalent action-derived completion), so apparatus q_2/q_3 promotion remains blocked without invalidating the linear detector response.

The selected high-mode Maxwell input has since been closed under its exact charge blocks and propagated by exact blockwise temporal functional calculus. A Green-weighted estimate proves that the Maxwell evolution does not amplify the omitted spatial tail. The earlier dense all-shell route is replaced by a response-specific shell stream with fixed detector duals and scalar integrands. Its finite kernels and first retarded emitter channel are exact, and partition-refined leading Cauchy covectors exclude rank loss uniformly on $m_0^2, m_1^2 \in [1, 2]$. The full all-shell massive image, absolute q_3 feedback intervals, evaluated recoil, and restriction to the nonlinear tangent cone remain open.

The late programme has now crossed additional distinct gates: a healthy relational clock, an exact inclusion of ordinary Einstein–Maxwell waves, a nonzero retained gravity–light interaction representative, a nonlinear Taub test that rejects pure extra directions but admits balanced mixed directions, the complete aligned homogeneous/twist common-zero orbit and its bounded continuation obstruction, an actual retarded relational light observable, mass-uniform leading rank-two detector records, a second curved causal consumer, a static black-hole first law, an exact Schwarzschild axial Einstein/extra wave split, nonzero mixed and extra horizon-flux fixtures, two-parity finite-norm Einstein selection at one asymptotic fixture, a free Lorentzian curvature-observable algebra, the first Nariai transverse formal causal variation, and a reduced Hadamard/Krein carrier with signs $(+, -, -)$. These are linked exact results, not yet a complete localized apparatus, an invariant branch-mixing theorem, a cyclic Einstein–Weyl relative equivalence, a dynamical black-hole phase space, a positive full-BV state, or an all-orders interacting quantum universe.

23 What was already solved, and what is new

The series does not claim that earlier positive-metric or Krein quantizations failed. Its contribution is to separate their mathematical roles and connect them through an explicit dependency structure.

Paper I [25] reconstructs the Bender–Mannheim positive Pais–Uhlenbeck transformation and classifies the free metric freedom.

Paper II [26] proves the minimum-distortion selection of the canonical positive metric.

Paper III [27] separates metric geometry, vacuum covariance, representation choice, and the Jordan limit in the scalar field.

Paper IV [28] classifies the locally reduced covariant free Einstein–Weyl real forms and their pure-Weyl boundary.

Paper V [29] formulates interaction deformation complexes and identifies resonant scalar obstructions and the surviving doubled boundary structure.

Paper VI [30] applies the deformation test to Einstein–Weyl gravity and isolates the first open gravitational obstruction channel in the stated perturbative class.

Paper VII [31] constructs the selected polarized residual state complex and computes H_{res}^4 and its Krein pairing.

Paper VIII [32] constructs causal homotopies for the covariant free metric BV complex and transports Paper VII’s theorem, endpoints, symmetry action, and pairing. It does not recompute residual CE cohomology.

Paper IX [33] constructs the positive Berger phase clock, proves fixed-coupling presymplectic nullity of D , and certifies the causal K -Cartan identity through the first interaction orders.

Paper X [34] embeds the complete standard compact Einstein–Maxwell tangent in Weyl–Maxwell theory and classifies the generic axial extra quotient with the direct Lee–Wald current. Its post-paper relative bridge completes the natural support-local noncyclic linear triangle and proves the standard-pairing cyclic obstruction.

Paper XI [35] transfers the repaired mixed gravity–Maxwell interaction through ℓ_3 on one frozen cyclic retract, including exact cyclicity, exchange-channel receipts, and nonremovability in the declared filtered-cyclic class; the fuller ghost carrier trivializes it through first jet and leaves the second-jet image open.

Paper XII [36] computes the coefficient-bearing strict one-loop BV anomaly, restores the formal tau-adic compensator theory’s local Euclidean QME, isolates the anomaly-fixed effective-action terms, and constructs the reduced vacuum-cylinder Hadamard/Krein carrier.

Charge-fibre bridge [37] proves an all-harmonic fixed-bundle second-order obstruction for pure-extra generic tangents and an explicit balanced Einstein–extra second-order extension. Its post-freeze certificate suite [38] classifies the generic $k = 0$ and paired opposite-momentum moment-map cones, the standard exceptional/global blocks, and a two-parameter $\ell = 2$ second-order-extendible face.

The programme has one local free spine and four linked branches. Interactions test whether a free completion survives physical conversion. Residual reduction determines the selected state cohomology, and covariant causal transport proves that the metric BV complex computes that same cohomology and pairing. The relational-clock branch tests whether total time translation is presymplectically null after a physical clock construction is added, while the background-fixing helical stabilizer is Cartan-null-homotopic through the certified interaction order. The Einstein–extra branch locates ordinary waves and then asks which linear directions survive nonlinear compact constraints.

24 What remains open

Paper VIII closes the free classical covariant causal bridge in the selected closed-universe problem. The construction of causal homotopies, recovery of the residual endpoints, equivariant transport, and comparison of the covariant, Cauchy, energy-mode, and residual pairings are therefore no longer open items. Papers IX–XI, the charge-fibre bridge, and the observer certificates have also turned several former aspirations into exact scoped results. The remaining frontier is reported by lifecycle and missing gate:

1. full-BV distributional and Hadamard completion, including ghost and antifield covariances, a BRST-compatible positive state, and renormalized time-ordered products; the reduced vacuum-cylinder E/A/L Krein carrier is certified separately;
2. integration of the infinitesimal $\mathfrak{so}(4, 2)$ action to an $SO(4, 2)$ representation;




Figure 4: Dependency structure of the twelve core technical papers and the charge-fibre bridge. The interaction, residual, relational-clock, and Einstein–extra branches share the free gravity spine but answer different questions. An arrow records mathematical dependence, not equivalence of claim lifecycle or physical interpretation.

3. alternative boundary, clock, relative, and deparametrized polarizations, in which D or other generators may remain charges, together with higher-arity and boundary stability of the positive Berger-clock contraction;
4. five-stabilizer precomposition of the relative Hessian current, Lee–Wald improvement comparison, cyclic-dual rows, and integrated-charge recovery, followed by the correct non-cyclic relative L_∞ extension beyond the now-certified matching Taylor packages, nonzero strict arity-two defect, Taub obstruction to local f_2 , and global five-charge receiver;
5. an alternative noncontractible filtered or mixed-bundle projector separating the Einstein-like, extra-Weyl, and Maxwell branches after the canonical support-local rank-46 route’s exact obstruction, followed by the complete projected ℓ_3 mixing table;
6. the remaining invariant interaction tests: nonzero action on final cohomology or nontriviality beyond the declared filtered-cyclic F_2/F_3 class in which nonremovability is already certified;
7. a common localized emitter preparation and clock-frequency comparison, the evaluated recoil coefficient, full apparatus Dirac bracket, higher polarization/interaction rows, observer morphism, and nonlinear backreacted rank-two detector response, together with completion of the response-specific all-shell stream and absolute q_3 feedback bounds beyond the certified finite mass-uniform leading response; the first missing clock-map second jet must first repair the certified 108-row unary obstruction;
8. a structural Taub theorem generalizing the definite pure-extra versus indefinite Einstein cancellation mechanism, the coefficientwise common zero locus of the certified bounded polynomial-growth and shell-resonance functionals, infinite-mode inputs, charge-mixed cones, the causal/retarded correction class, generalized fourth-order target modes, and higher-order extension tests; the complete certified finite-support inventory is now classified in the finite exponential-polynomial category, the tuned standard cone is uniform in $\ell \geq 2$, and the complete 164-coefficient two-momentum basis census leaves only its joined zero variety open;
9. interaction descent for the two surviving deformation classes and renormalized local composite representatives of W_+^2 and W_-^2 ;
10. the compensator-inclusive classical contraction, coefficient-bearing one-loop Slavnov operator, quantum Cartan defect, and only then residual quantum transfer; the strict local counterterm/anomaly cohomology and the formal tau-adic compensator extension are now classified at dimension four, with distinct one-loop QME dispositions; the generic ghost longitudinal tower is now a single modified-Fredholm Schur problem with exact critical residue, exact round- S^4 finite determinant data, and a generic local multiplicative defect. All eleven generic third-curvature ghost functions are integrated exactly; the product- $S^2 \times S^2$ longitudinal Schur carrier and coupled exceptional factor are exact; their assembly with the generic physical fourth-order Hessian remains open;
11. physical boundary problems beyond the two-parity $\ell = 2, \omega = 3/5$ finite-norm Einstein-selection fixture, especially symbolic-frequency and all- ℓ asymptotics, complex-frequency and quasinormal modes, a full causal exterior initial-boundary problem and stability,

an asymptotically flat Bach phase space, Bondi flux, scattering states, and interacting probability and unitarity.

A Maldacena-type boundary condition may truncate a bulk branch rather than complete the same space [54]; it cannot be modeled by simply deleting one ghost from the selected closed-universe complex. Likewise, classical Green hyperbolicity alone selects neither a Hadamard state nor a quantum product. The local gauge-fixed BV calculation gives $\dim H_{\text{even}}^{1,4} = 2$ and $\dim H_{\text{odd}}^{1,4} = 1$ on the regular Bach locus, with no additional pure-Diff or mixed Diff–Weyl class. The repository elliptic complex, measure, zero-mode, contour, and local heat-kernel receipts fix the regulated Slavnov class to $(199/30)[\omega C^2] - (87/20)[\omega E_4]$, with the odd coordinate zero by the declared parity Ward identity. Both even coordinates are nonzero, so the strict fixed-field-content local Euclidean QME is obstructed at one loop. Standard-sign free conformal matter cannot cancel this vector. In the formal tau-adic compensator extension, however, the exact minimal-BV cotangent lift contracts the Weyl quartet, the dimension-four quotient has $\dim H_{\text{ext}}^{0,4} = (3 \text{ even}, 1 \text{ odd})$ and $H_{\text{ext}}^{1,4} = 0$, and the explicit coefficient-bearing Wess–Zumino counterterm restores the extended local Euclidean QME at one loop. This is not a finite-polynomial, all-loop, or Lorentzian result. The quantum problem must still construct the compensator-inclusive classical contraction and Q_1 before testing residual transfer or the Cartan homotopy. The local Wess–Zumino contribution to Q_1 is fixed, but finite C^2 and $R(\hat{g})^2$ counterterms give an exact rank-two ambiguity in the complete bulk operator. A Paneitz/Riegert anomaly-induced representative under its declared inverse or compatible-source conditions and the universal $-199/60$ curvature-squared logarithmic coefficient are now certified through curvature order two. The exact raw strict-Weyl $\square R$ coefficient and its finite $R(g)^2$ conversion to the repository $\square R = 0$ scheme reproduce the Paneitz/Riegert 29/120 coefficient. The absolute finite C^2 and dressed $R(\hat{g})^2$ normalizations, cubic completion, nonlocal R^2 form factor, and global kernel data remain open. The separately certified curvature-image causal propagator and free CCR algebra identify a Lorentzian observable carrier. On the reduced vacuum-cylinder E/A/L mode carrier, a compatible complex structure and microlocal Hadamard distributions now exist, with E positive and A/L negative-Krein. This does not supply a positive full-BV representation, time-ordered product, Lorentzian QME restoration, or particle space. Proposals for interacting quadratic gravity such as [57] are comparison points, not consequences of Papers VII–VIII.

The immediate classical decisions are whether the extra sector admits causal/retarded and higher-order continuation after its certified split between bounded obstruction and finite exponential-polynomial secular extension, and whether a noncontractible branch carrier gives the retained mixed ℓ_3 physical branch meaning. The immediate quantum decision is whether the restored tau-adic local one-loop theory admits an extended classical contraction and complete Slavnov operator, and whether full BRST reduction removes or retains the reduced carrier’s negative-Krein directions.

25 Conclusion

The programme separates the higher-derivative ghost question into four levels:

1. Local free classification determines the complex dynamics, real forms, metrics, gauge reduction, and Jordan limits.
2. Gauge and residual reduction determine which state, vertex, or observable classes survive, and covariant causal transport compares their realizations.
3. Interaction stability tests whether a selected free or reduced structure survives physical nonlinear conversion.
4. Quantum completion tests anomalies, state positivity, microlocal regularity, interacting products, and whether any surviving branch defines a particle or probability space.

Papers I–IV establish the local free spine. The positive PU metric, the minimum-distortion principle, the universal complex covariance, the Krein boundary, and the covariant gravity classification are compatible parts of that spine rather than competing solutions to one undifferentiated problem. Papers V–VI then show that a valid free completion need not admit the same analytic continuation through an interacting branch-changing channel.

Papers VII–VIII complete the selected free classical pure-Weyl branch. Paper VII derives the polarized residual state complex and computes

$$H_{\text{res}}^4 = \text{span}\{[W_+^2], [W_-^2]\}, \quad \text{sig } G_{\text{res}} = (2, 0), \quad G_{\text{res}} = I_2. \quad (72)$$

Paper VIII constructs the support-local curvature prolongation and causal chain homotopies of the covariant metric BV complex, recovers the residual endpoints and symmetry action, and transports the current and Green pairings. It therefore proves

$$H_{\text{cov}}^4 \cong H_{\text{res}}^4, \quad \text{sig } G_{\text{cov}} = (2, 0), \quad G_{\text{cov}} = I_2, \quad (73)$$

using Paper VII’s CE theorem as an input rather than recomputing it. The two surviving classes are deformation or vertex classes, not positive-norm one-particle gravitons.

Paper IX then tests the compact D quotient in a setting with an actual healthy matter clock. The exact Berger background has nonzero clock momentum, while the fixed-coupling lapse constraint makes its variation vanish on every smooth linearized tangent. The complete causal 54-row BV complex and action-derived q_2, q_3 give the corresponding K -Cartan contraction through arity three, where $K = D - \omega R$ fixes the background. They do not provide the affine zeroth-arity equation for raw D . This is a scoped classical theorem, not an all-orders nonlinear or quantum result.

Paper X establishes the complementary incidence theorem needed to connect the programme to ordinary gravitational physics: the complete standard compact Einstein–Maxwell harmonic tangent is present and nonnull inside Weyl–Maxwell theory. Its Lee–Wald pullback is nondegenerate but nonsymplectic, and the target contains two further nonradical generic axial directions. The subsequent natural all-row construction now completes the linear Einstein–Weyl mapping-cofiber triangle, but its three action forms cannot be collapsed to a standard cyclic triangle because their inertias disagree. Paper XI then proves that one retained mixed gravity–Maxwell ℓ_3 representative survives an exact cyclic reduction, while correctly leaving its cohomological, branch-invariant, and field-redefinition meaning open. The subsequent rank-46 cyclic graph carrier removes one architectural obstruction to asking the branch question, but its physical projector remains the decisive gate.

The charge-fibre bridge shows why the linear inclusion is not the final answer. Every nonzero pure-extra generic tangent fails the fixed-bundle second-order Taub test, yet one balanced Einstein–extra tangent cancels all stabilizer moment maps and extends explicitly through

second order. The subsequent cone certificates show that this ray lies in a larger classified mixed moment-map locus and a two-parameter $\ell = 2$ second-order-extendible face. On the complete certified finite-support inventory, including generic, exceptional, twist, and homogeneous inputs, finite exponential-polynomial second-order continuation is now equivalent to the five moment maps vanishing; bounded continuation has further unresolved polynomial-growth and shell-resonance equations. Its standard generalized-zero subcone is now classified exactly, and $b = B = 0$ is necessary throughout the complete finite-support bounded cone; the exceptional $\ell = 1, k = 0$ all- m pure cone is obstructed. The observer certificates add causal linear records, memory, localized emitter preparations, and a mass-uniform finite leading rank-two response, while the retarded Berger Maxwell construction adds a first spatially global relational light observable, without pretending that a complete localized backreacted apparatus already exists; exact temporal charge blocks and response-specific shell streaming now make the first retarded emitter channel executable while leaving full q_3 feedback and recoil open. The Nariai consumer and its global formal transverse variation show that the causal transfer architecture is not confined to one fixed cylinder. The black-hole axial flux theorem shows that the pure Einstein/Regge–Wheeler block is symplectically null in pure Weyl gravity, while exact fixtures show nonzero Einstein–extra and extra–extra horizon flux. The reconstructed extra metrics develop logarithmic or power-enhanced tails at the $\ell = 2, \omega = 3/5$ fixture, and finite symplectic norm at infinity selects exactly the Einstein sector in both parities there. Horizon regularity and asymptotic phase space therefore impose different branch tests; symbolic-frequency and all- ℓ classification remain open. Static black-hole and free curvature-observable results add horizon thermodynamics and a Lorentzian quantum-algebra carrier without promoting either to dynamics or particles. Together these results replace the naive direct-sum picture by an emerging nonlinear extension problem: the Einstein, extra-Weyl, Maxwell, clock, and observer sectors may mix, source, or obstruct one another.

The resulting position is neither that higher-derivative ghosts are automatically harmless nor that they automatically invalidate a theory. The correct question must name the complex dynamics, real form, observable algebra, local and residual quotient, completion, causal realization, and interaction class being considered.

The twelve-paper core programme classifies the local free completions, tests interaction stability, computes and causally realizes the selected pure-Weyl residual cohomology, introduces a healthy relational clock, embeds ordinary Einstein–Maxwell waves, and transfers the first retained gravity–light bracket, then computes the first coefficient-bearing one-loop obstruction and a reduced Hadamard/Krein carrier. The charge-fibre, causal-transfer, observer, black-hole, and quantum-observable successors then test nonlinear selection, portability, causal records, horizon dynamics, and the free observable algebra. The next gates are the five-stabilizer relative-current comparison, a noncontractible alternative to the obstructed rank-46 branch projector, invariant branch mixing, complete localized observer backreaction, general black-hole finite-norm classification and full exterior stability, physical boundaries, a positive full-BV Hadamard state, the complete renormalized quantum master equation, and unitarity.

Verification. The finite-dimensional, variational, symplectic, causal, and modewise identities quoted here are tied to the verification repositories of Papers I–XII and the charge-fibre bridge [25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42]. Each technical paper states the category and claim boundary of its certificates; a certificate from one level is not evidence for a stronger level.

References

- [1] M. Ostrogradsky, *Mémoire sur les équations différentielles relatives au problème des isopérimètres*, Mém. Acad. St. Pétersbourg **VI** 4 (1850) 385.
- [2] R. P. Woodard, *The theorem of Ostrogradsky*, arXiv:1506.02210.
- [3] A. Pais and G. E. Uhlenbeck, *On field theories with non-localized action*, Phys. Rev. **79** (1950) 145.
- [4] C. M. Bender and S. Boettcher, *Real spectra in non-Hermitian Hamiltonians having PT symmetry*, Phys. Rev. Lett. **80** (1998) 5243.
- [5] C. M. Bender and P. D. Mannheim, *No-ghost theorem for the fourth-order derivative Pais–Uhlenbeck oscillator model*, Phys. Rev. Lett. **100** (2008) 110402.
- [6] C. M. Bender and P. D. Mannheim, *Exactly solvable PT-symmetric Hamiltonian having no Hermitian counterpart*, Phys. Rev. D **78** (2008) 025022, arXiv:0804.4190.
- [7] S. Bateman and N. Turok, *Escape from Ostrogradsky via hidden ghost parity*, arXiv:2607.00096.
- [8] K.-j. Hamada and S. Horata, *Conformal algebra and physical states in non-critical 3-brane on $\mathbb{R} \times S^3$* , Prog. Theor. Phys. **110** (2003) 1169–1210, arXiv:hep-th/0307008.
- [9] K.-j. Hamada, *BRST analysis of physical fields and states for 4D quantum gravity on $\mathbb{R} \times S^3$* , Phys. Rev. D **86** (2012) 124006, arXiv:1202.4538.
- [10] A. R. Gover and L. J. Peterson, *The ambient obstruction tensor and the conformal deformation complex*, Pacific J. Math. **226** (2006) 309–351, arXiv:math/0408229.
- [11] A. Čap, *Overdetermined systems, conformal geometry, and the BGG complex*, in *Symmetries and Overdetermined Systems of Partial Differential Equations*, IMA Vol. Math. Appl. **144** (Springer, 2008) 1–24, arXiv:math/0610225.
- [12] I. Dneprov and M. Grigoriev, *Presymplectic BV–AKSZ formulation of conformal gravity*, arXiv:2208.02933.
- [13] A. S. Cattaneo, P. Mnev and N. Reshetikhin, *Classical BV theories on manifolds with boundary*, Commun. Math. Phys. **332** (2014) 535–603, arXiv:1201.0290.
- [14] C. Bär, *Green-hyperbolic operators on globally hyperbolic spacetimes*, Commun. Math. Phys. **333** (2015) 1585–1615, arXiv:1310.0738.

- [15] M. Benini, G. Musante and A. Schenkel, *Green hyperbolic complexes on Lorentzian manifolds*, Commun. Math. Phys. (2023), arXiv:2207.04069.
- [16] M. Beccaria, X. Bekaert and A. A. Tseytlin, *Partition function of free conformal higher spin theory*, JHEP **08** (2014) 113, arXiv:1406.3542.
- [17] R. R. Metsaev, *Ordinary-derivative formulation of conformal low-spin fields*, JHEP **01** (2008) 064, arXiv:0707.4437.
- [18] T. Nutma and M. Taronna, *On conformal higher spin wave operators*, JHEP **06** (2014) 066, arXiv:1404.7452.
- [19] P. E. T. Jorgensen and F. Tian, *Unbounded operators, Lie algebras, and local representations*, J. Math. Phys. **55** (2014) 111702, arXiv:1406.6966.
- [20] F. G. Scholtz, H. B. Geyer and F. J. W. Hahne, *Quasi-Hermitian operators in quantum mechanics and the variational principle*, Ann. Phys. **213** (1992) 74.
- [21] A. Mostafazadeh, *Pseudo-Hermitian representation of quantum mechanics*, Int. J. Geom. Meth. Mod. Phys. **7** (2010) 1191.
- [22] P. D. Mannheim, *Solution to the ghost problem in higher-derivative gravity*, arXiv:2109.12743.
- [23] P. D. Mannheim, *CPT symmetry without Hermiticity*, arXiv:1611.02100.
- [24] K. S. Stelle, *Classical gravity with higher derivatives*, Gen. Rel. Grav. **9** (1978) 353; Phys. Rev. D **16** (1977) 953.
- [25] *Canonical positive symplectic diagonalization of the Pais–Uhlenbeck oscillator*, companion paper (Paper I) and verification repository, 2026.
- [26] *The Pais–Uhlenbeck metric as a minimum-distortion principle, and the representation problem for the fourth-order field*, companion paper (Paper II), 2026.
- [27] *The universal vacuum of the fourth-order scalar field: metric orbits, Fock sectors, and the Krein boundary*, companion paper (Paper III), 2026.
- [28] *Gauge reduction and the completion problem in fourth-order gravity: PU pairing, covariant real forms, and the conformal Jordan boundary*, companion paper (Paper IV), 2026.
- [29] *Interaction obstructions, resonant PT breaking, and doubled Jordan symmetry in fourth-order theories*, companion paper (Paper V), 2026.
- [30] *Interaction obstructions in Einstein–Weyl gravity: cubic protection, second-order metric failure, and Krein visibility*, companion paper (Paper VI), 2026.
- [31] *Residual cohomology of free pure-Weyl gravity on the conformal cylinder: derived reduction, Weyl-square classes, and Krein completion*, companion paper (Paper VII), 2026.

- [32] *Causal curvature prolongation and covariant BV–BFV transport for free pure-Weyl gravity*, companion paper (Paper VIII), 2026.
- [33] *A backreacting phase clock with fixed momentum in pure-Weyl gravity: fixed-coupling rigidity and causal BV Cartan analysis of the helical stabilizer*, companion paper (Paper IX), 2026.
- [34] *Einstein–Maxwell waves inside Weyl–Maxwell gravity on a compact product: exact linear phase-space inclusion and the extra axial branch*, companion paper (Paper X), 2026.
- [35] *A retained mixed gravity–Maxwell bracket under cyclic homological reduction through ℓ_3 : exact transfer and causal Cartan compatibility*, companion paper (Paper XI), 2026.
- [36] *A one-loop BV obstruction in pure Weyl gravity and its formal tau-adic Wess–Zumino resolution*, companion paper (Paper XII), 2026.
- [37] *Pure-extra Taub obstruction and a balanced second-order extension in compact Weyl–Maxwell gravity: pure extra modes versus Einstein–extra mixtures*, open-repository bridge paper (Paper 91), 2026.
- [38] *Mixed-cone extension verification receipt: generic $k = 0$, opposite-momentum, exceptional/global, and axial $\ell = 2$ second-order certificates*, open-repository certificate suite and independent verifiers, 2026.
- [39] *Abstract cyclic Green transfer with Berger, flat-Minkowski, and curved Nariai consumers*, open-repository certificates, independent verifiers, and Paper 90 bridge note, 2026.
- [40] *Closed-universe observer and apparatus suite: localized emitter preparations, causal transfer, and leading rank-two records*, open-repository certificates and independent verifiers, 2026.
- [41] *Static spherical pure-Weyl backgrounds, normalized charges, entropy, and exact first law*, open-repository black-hole certificates and independent verifiers, 2026.
- [42] *Lorentzian curvature-observable causal propagator and curvature-image presymplectic CCR algebra*, open-repository quantum certificates and independent verifiers, 2026.
- [43] G. B. Folland, *Harmonic Analysis in Phase Space*, Princeton University Press, 1989.
- [44] R. Bhatia, *Positive Definite Matrices*, Princeton University Press, 2007.
- [45] G. D. Mostow, *Self-adjoint groups*, Ann. of Math. **62** (1955) 44.
- [46] G. Kempf and L. Ness, *The length of vectors in representation spaces*, in: Algebraic Geometry, Lecture Notes in Math. **732**, Springer, 1979, 233.
- [47] I. Gohberg, P. Lancaster and L. Rodman, *Indefinite Linear Algebra and Applications*, Birkhäuser, 2005.
- [48] T. Kato, *Perturbation Theory for Linear Operators*, Springer, 1966.
- [49] W. D. Heiss, *The physics of exceptional points*, J. Phys. A **45** (2012) 444016.

- [50] J. von Neumann, *On infinite direct products*, Compos. Math. **6** (1939) 1.
- [51] M. J. Radzikowski, *Micro-local approach to the Hadamard condition in quantum field theory on curved space-time*, Commun. Math. Phys. **179** (1996) 529.
- [52] R. Brunetti and K. Fredenhagen, *Microlocal analysis and interacting quantum field theories*, Commun. Math. Phys. **208** (2000) 623.
- [53] S. Hollands and R. M. Wald, *Local Wick polynomials and time ordered products of quantum fields in curved spacetime*, Commun. Math. Phys. **223** (2001) 289.
- [54] J. Maldacena, *Einstein gravity from conformal gravity*, arXiv:1105.5632.
- [55] H. Lü and C. N. Pope, *Critical gravity in four dimensions*, Phys. Rev. Lett. **106** (2011) 181302, arXiv:1101.1971.
- [56] S. Deser, E. Joung and A. Waldron, *Partial masslessness and conformal gravity*, J. Phys. A **46** (2013) 214019, arXiv:1208.1307.
- [57] J. Kubo and J. Kuntz, *Unitarity through PT symmetry in quantum quadratic gravity*, arXiv:2410.08278.

Paper	Technical role	Claim boundary
I–III	Positive/Krein free scalar geometry	Free oscillator and scalar-field categories
IV	Covariant local free gravity classification	Linearized Einstein–Weyl theory
V–VI	Interaction deformation and obstruction	Stated perturbative orders and channels
VII	Absolute residual H^4 and Krein pairing	Selected algebraic/energy-mode polarization
VIII	Covariant causal BV–BFV transport	Free classical Lorentzian cylinder complex
IX	Backreacting Berger phase clock and causal K -Cartan contraction	Fixed-coupling compact classical sector through arity three
X	Compact Einstein–Maxwell inclusion and extra Weyl branch	Linear local-algebraic/reduced-mode phase space before stabilizer reduction
XI	Retained cyclic gravity–Maxwell ℓ_3	One frozen cyclic retract before physical branch projection

Table 1: Eleven-paper core programme grouped by mathematical role. The charge-fibre Taub bridge and observer constructions connect these core papers without being silently counted as additional stages of the same theorem.