

An Axial $\ell = 2$ Non-Split Regge–Wheeler Self-Extension and a Defective Schwarzschild Resonance in Pure Weyl Gravity

GPT-5.6.sol*

Asger Alstrup Palm[†]

27 July 2026

Abstract

We show that an axial $\ell = 2$ Schwarzschild quasinormal mode becomes a defective resonance in four-dimensional pure Weyl gravity. The reduced six-state Bach system has two spin-two Regge–Wheeler layers and one spin-one Maxwell layer. The repeated spin-two block is not a rational direct sum: it is a non-split self-extension. At a validated simple Regge–Wheeler quasinormal frequency, its connection has Smith type $(0, 0, 2)$, hence one length-two root chain. The geometric root is Einstein, while the generalized root has nonzero gauge-invariant traceless-Ricci carrier.

The algebraic mechanism is an exact projective cocycle. After Liouville reduction,

$$[\mathcal{I}_{\text{Bach}}] = \frac{i\omega}{2} \left[1 - \frac{2}{r} \right].$$

The class in brackets is also the direction selected by the complete coupled axial massive-spin-two system. Transforming that system into its massless Maxwell/Regge–Wheeler basis gives

$$[\mathcal{I}_{\text{phys}}] = \frac{1}{3} \left[1 - \frac{2}{r} \right], \quad [\mathcal{I}_{\text{Bach}}] = \frac{3i\omega}{2} [\mathcal{I}_{\text{phys}}].$$

Thus the isolated shifted-Regge–Wheeler target has the correct direction but not the physical normalization. The rational primitives reproduce the differentiated moving massive phase, while the Coulomb exponent has zero first mass derivative. Matrix Frobenius and sectorial Volterra constructions promote this comparison to the complete endpoint-normalized Jost planes. The certified selector therefore equals the velocity of the complete massive axial QNM branch, up to the displayed normalization, and that velocity is nonzero.

The complete radial pencil also admits a fixed-domain, two-ended $\pi/4$ exterior-complex-scaled realization that is Fredholm of index zero. Its inverse has a nonzero rank-one second-order pole, and its compactly supported matrix elements agree with the exterior outgoing cut-off inverse. This outgoing inverse is the meromorphic continuation of the Laplace transform of the mode-reduced retarded parent response, so the pole belongs to an actual causal transfer operator. Exact axial reconstruction shows that the Einstein range of the principal coefficient has nonzero Bondi shear. A complexified conserved, traceless, radially compact odd source has nonzero adjoint overlap. Thus the isolated local resonance contour contains the expected $te^{i\omega_n t}$ Einstein profile. We do not claim a global retarded quasinormal expansion, a real causal source construction, or standard asymptotic falloff for the time-independent generalized component.

*OpenAI model and principal programme author.

[†]Programme orchestrator and corresponding human contact; Honorary Professor, DTU Compute, Technical University of Denmark. asger@area9.dk.

1 Introduction

Fourth-order linear equations are often described by listing their scalar factors. That description is incomplete when a factor is repeated. The layers may form a direct sum, or they may form a non-semisimple extension. Only the second possibility carries an intrinsic generalized state and can turn a simple resonance of the repeated factor into a defective resonance of the full system. We use the standard Regge–Wheeler and Schwarzschild QNM conventions of Refs. [1, 2], translated to our $e^{+i\omega t}$ Fourier sign.

This distinction is realized by axial $\ell = 2$ Schwarzschild perturbations of pure Weyl gravity. In the factor-adapted radial variables, the Bach equation contains two spin-two Regge–Wheeler layers and one spin-one Maxwell layer, but the two spin-two layers cannot be separated by a rational local field redefinition. The extension is also the critical first jet of the massless and massive spin-two directions after exact reduction of the complete coupled axial system. At a Schwarzschild quasinormal frequency, this confluence produces a length-two root chain rather than two independent Einstein roots.

We use $M = 1$, the Fourier convention $e^{+i\omega t}$, and the negative-real-frequency representative of the reflected Schwarzschild pair. Thus damping means $\text{Im } \omega > 0$, and the familiar fundamental axial frequency is written near $-0.37367 + 0.08896i$. The parameter m below is a signed *squared-mass coefficient*; it is not the positive square root usually called the graviton mass.

The paper has five results.

1. The axial spin-two Bach block is a non-split rational Regge–Wheeler self-extension and an exact first parameter jet.
2. Its projective cocycle has the elementary normal form

$$[\mathcal{I}_{\text{Bach}}] = \frac{i\omega}{2} \left[1 - \frac{2}{r} \right],$$

and the complete coupled massive axial tensor branch represents one third of the same class.

3. A validated QNM calculation selects the defective Smith type $(0, 0, 2)$. The generalized root necessarily crosses the Einstein filtration. The same certificate gives a nonzero velocity of the complete massive axial QNM branch.
4. A fixed-domain global exterior-complex-scaled radial pencil is Fredholm of index zero and has a genuine rank-one pole of order two. Its compactly supported matrix elements equal the exterior outgoing cut-off Green operator and meromorphically continue a mode-reduced retarded causal transfer.
5. Exact metric reconstruction and an explicit conserved, traceless odd source show that the range and source covectors of this pole can both be nonzero.

These statements combine exact algebra with validated numerics. The module reduction, cocycle identities, mass comparison, Green reduction, metric reconstruction, and source construction are exact. The existence and uniqueness of the enclosed QNM, nonvanishing of the projective selector, and spin-one local-unit condition are interval-validated.

Claim boundary

Result	Input	Status
Filtration, partial jet, and nonsplitting	Exact identities and exhaustive rational splitting test	LOCAL-ALGEBRAIC
Complete massive first-jet comparison	Exact coupled-system factorization, cocycle primitives, and moving phases	LOCAL-ALGEBRAIC
Endpoint-compatible massive QNM slope	Matrix Frobenius/Volterra Jost planes, spin-one Schur unit, selector	REDUCED-MODE
Defective QNM and non-Einstein root	Validated Evans winding, selector, spin-one unit	REDUCED-MODE
Exterior cut-off Green double pole	Analytic Jost maps and exact transparent cuts	REDUCED-MODE
Global ECS radial Fredholm pole	Two-ended exponential dichotomy and connection reduction	REDUCED-MODE
Causal Laplace-to-resonance bridge	Normally hyperbolic parent, retarded mass derivative, ECS continuation	LORENTZIAN-CAUSAL
Bondi-shear and compact-source nonannihilation	Exact reconstruction and conserved-source identities	REDUCED-MODE
Global retarded ringdown expansion	Real-axis causal weights, reconstruction, and contour deformation	Not established

The last row is essential. The paper proves a fixed-domain meromorphic radial response on a global complex-scaled contour, its compactly localized exterior matrix elements, and their origin as the meromorphic continuation of a mode-reduced retarded causal transfer. It also proves the polynomial contribution of an isolated local resonance contour. It does not prove a full off-shell metric/BV retarded propagator, a complete causal quasinormal expansion, late-time asymptotics, detector sensitivity, or excitation by a specified astrophysical matter model.

The nearby quadratic-gravity problem has been studied directly as a coupled massive spin-two system [3, 4]. That literature separates the general-relativistic and massive QNM families at nonzero squared mass. Our distinct result concerns their singular pure-Weyl confluence and proves non-semisimplicity of the critical Bach response. We use the complete coupled massive axial equations for the physical first-jet comparison; a single shifted Regge–Wheeler equation is retained only as a convenient graded reference and is not the complete tensor branch.

2 The filtered axial Bach module

2.1 Differential-module conventions

We work over

$$K = \mathbb{C}(r, \omega), \quad D = f \partial_r, \quad f = 1 - \frac{2}{r},$$

with $\omega \neq 0$ unless stated otherwise. The spin-two and spin-one scalar operators for axial $\ell = 2$ are

$$L_2 = D^2 + U_2, \quad U_2 = \omega^2 - f \left(\frac{6}{r^2} - \frac{6}{r^3} \right),$$

$$L_1 = D^2 + U_1, \quad U_1 = \omega^2 - f \frac{6}{r^2}.$$

We use $D(q)$ and $D(U)$, rather than primes, for the radial derivation. A prime on a connection coefficient below denotes ∂_ω .

The Einstein submodule is the kernel of the gauge-invariant traceless-Ricci, equivalently Schouten, carrier. Thus “non-Einstein” means nonzero projection to this carrier quotient, not failure of a coordinate gauge choice.

2.2 Filtration and partial jet

In the certified factor gauge, the six-state system is

$$\begin{aligned} DX &= AX + EY + CZ, \\ DY &= AY + DZ, \\ DZ &= A_1 Z, \end{aligned} \tag{1}$$

where $X, Y, Z \in K^2$, while A and A_1 are the companion matrices of L_2 and L_1 . The rational reconstruction into the axial metric and curvature variables has full row rank on the declared domain.

Theorem 2.1 (Filtered axial Bach module). *The reduced axial $\ell = 2$ Bach module has a filtration with successive quotients*

$$M_2, \quad M_2, \quad M_1.$$

The middle four-state object is an extension

$$0 \longrightarrow M_2 \longrightarrow E_2 \longrightarrow M_2 \longrightarrow 0,$$

and the final quotient is the spin-one Maxwell module M_1 .

This theorem is an exact rational statement. It does not rely on an endpoint normalization or on the QNM calculation.

Introduce the four-state family

$$\mathcal{B}(\tau) = \begin{pmatrix} A + \tau E & D + \tau C \\ 0 & A_1 \end{pmatrix}, \quad \Phi_4(\tau) = \begin{pmatrix} P(\tau) & Q(\tau) \\ 0 & R \end{pmatrix}.$$

Theorem 2.2 (Exact partial jet). *The system (1) is the first spin-two-row jet of $\mathcal{B}(\tau)$ at $\tau = 0$. Its fundamental matrix is*

$$\Phi_6 = \begin{pmatrix} P & \dot{P} & \dot{Q} \\ 0 & P & Q \\ 0 & 0 & R \end{pmatrix}_{\tau=0}, \quad \det \Phi_6 = (\det P)^2 \det R.$$

Every factor-adapted endpoint connection consequently has the form

$$T(\omega) = \begin{pmatrix} a & b & c \\ 0 & a & d \\ 0 & 0 & g \end{pmatrix}, \quad \det T = a^2 g. \tag{2}$$

Although each scalar factor is second order, its chosen horizon or infinity condition selects a one-dimensional endpoint line. The matrix (2) records one unwanted scalar endpoint coefficient for each of the three admissible factor lines; it is therefore 3×3 , rather than the 6×6 fundamental matrix of the first-order radial system.

Proof. Differentiate $D\Phi_4(\tau) = \mathcal{B}(\tau)\Phi_4(\tau)$ at $\tau = 0$. The differentiated first row, the undifferentiated spin-two row, and the spin-one row reproduce (1). The determinant and connection identities follow from block triangularity. $\square$

2.3 Projective extension class

Consider a scalar self-extension

$$Lx = (s_1 D + s_0)y, \quad Ly = 0, \quad L = D^2 + U.$$

Define

$$\mathcal{I} = s_0 - \frac{1}{2}D(s_1), \quad \mathcal{K}_U = D^3 + 4UD + 2D(U).$$

Lemma 2.3 (Triangular gauge). *For $q \in K$, set*

$$Q(q) = qD - \frac{1}{2}D(q).$$

On $\ker L$,

$$[L, Q(q)] = -\frac{1}{2}\mathcal{K}_U(q).$$

Thus the lift change $x \mapsto x + Q(q)y$ changes the scalar density by

$$\mathcal{I} \mapsto \mathcal{I} - \frac{1}{2}\mathcal{K}_U(q).$$

Proof. Expand the commutator by the Leibniz rule and use $D^2y = -Uy$. The coefficient of Dy cancels; the remaining coefficient is $-[D^3q + 4UDq + 2D(U)q]/2$. $\square$

The projective extension class is therefore

$$[\mathcal{I}] \in K/\mathcal{K}_U K.$$

The operator $\mathcal{K}_U$ is the symmetric-square operator of L .

Theorem 2.4 (Mass-direction normal form). *For the axial $\ell = 2$ Bach extension,*

$$\mathcal{I}_{\text{Bach}} = \frac{i(r-2)(2r\omega^2 + 3\omega^2 + 12)}{5r^4\omega}.$$

Let

$$q(r, \omega) = -\frac{i}{120\omega} \left(15r + 13 + \frac{12}{r} + \frac{9}{r^2} \right).$$

Then, for $\omega \neq 0$,

$$\mathcal{I}_{\text{Bach}} = \mathcal{K}_{U_2}q + \frac{i\omega}{2}f, \tag{3}$$

and hence

$$\boxed{[\mathcal{I}_{\text{Bach}}] = \frac{i\omega}{2}[f].}$$

Proof. Substitute $D = f\partial_r$, U_2 , and the displayed q into $\mathcal{K}_{U_2}q$. After clearing $120r^4\omega$, the assertion is a polynomial identity in r and ω . Exact coefficient comparison gives (3). $\square$

Theorem 2.5 (Non-split physical-axis extension). *For every real $\omega > 0$, the extension E_2 is non-split over $\mathbb{C}(r)[D]$.*

Certificate-supported exact proof. The local pole audit permits poles only at $r = 0$ and at the apparent divisor $r\omega = 2i$. Its indicial bounds force the exhaustive ansatz

$$a(r) = -\frac{i}{\omega^2(r\omega - 2i)} + \frac{c_{-2}}{r^2} + \frac{c_{-1}}{r} + c_0.$$

After substitution in $\mathcal{K}_{U_2}a + 2\mathcal{I}_{\text{Bach}} = 0$ and clearing the declared denominators, coefficient comparison gives

$$M(\omega)c = b(\omega), \quad M \in \mathbb{C}^{6 \times 3},$$

with

$$M = \begin{pmatrix} 0 & -4\omega^4 & 0 \\ -8\omega^4 & 8\omega^4 & 24\omega^2 \\ 16\omega^4 & 42\omega^2 & -156\omega^2 \\ 48\omega^2 & -224\omega^2 & 312\omega^2 \\ -208\omega^2 & 364\omega^2 & -192\omega^2 \\ 224\omega^2 & -168\omega^2 & 0 \end{pmatrix}, \quad b = \begin{pmatrix} 0 \\ -2i(5\omega - 6i) \\ i(35\omega - 78i) \\ -6i(5\omega - 26i) \\ 96 \\ 0 \end{pmatrix}.$$

Fixed minors of the coefficient and augmented matrices are

$$\det M_{[0,1,2]} = 3456\omega^{10}, \quad \det[M \mid \text{rhs}]_{[0,1,2,5]} = -645120i\omega^9.$$

They are simultaneously nonzero for every real $\omega > 0$. Hence $\text{rank } M = 3$, whereas $\text{rank}[M \mid b] \geq 4$, so the inhomogeneous coboundary equation is inconsistent. The pole audit and displayed ansatz are what make this finite inconsistency exhaustive rather than a truncated search. $\square$

Theorem 2.5 is a statement about the differential module; Theorem 2.4 identifies a simple representative of its class. Neither theorem asserts that every QNM of the repeated scalar factor must be defective. Defectiveness is a resonant evaluation of the nonzero extension class.

3 Complete massive axial first jet and Jost boundary

3.1 The coupled physical system

The covariant parent family contains the second-order tensor pencil

$$(\mathcal{E} + m\mathcal{A})\phi = 0,$$

where $m = \mu^2$ is the signed squared-mass coefficient. For axial $\ell = 2$ Schwarzschild perturbations, the complete massive tensor equations of Refs. [3, 4] are

$$D^2Q + \left[\omega^2 - f \left(m + \frac{10}{r^2} - \frac{16}{r^3} \right) \right] Q = \frac{8f(r-3)}{r^3} Z, \quad (4)$$

$$D^2Z + \left[\omega^2 - f \left(m + \frac{4}{r^2} + \frac{2}{r^3} \right) \right] Z = \frac{2f}{r^2} Q. \quad (5)$$

This is a coupled tensor/vector system, not a single shifted Regge–Wheeler equation.

At $m = 0$, introduce the tensor and vector master fields (ϕ_2, ϕ_1) , satisfying $L_2\phi_2 = 0$ and $L_1\phi_1 = 0$. The massless Berndtson transformation specializes to

$$Q = \frac{1}{\omega^2} \left(\frac{4f}{3r} \phi_2 + \frac{4}{3} D\phi_2 - \frac{2f}{r} \phi_1 + D\phi_1 \right), \quad (6)$$

$$Z = -\frac{1}{\omega^2} \left(\frac{7r-6}{3r^2} \phi_2 + D\phi_2 + \frac{1}{r} \phi_1 \right). \quad (7)$$

The induced 4×4 first-order state map has determinant ω^{-4} , and direct differentiation verifies its exact intertwining with (4)–(5) at $m = 0$.

Theorem 3.1 (Complete massive tensor first jet). *Transform the full m -derivative of (4)–(5) through (6)–(7). The reverse tensor-to-vector source is rationally removable. The tensor-led diagonal tangent scalarizes to*

$$L_2 \dot{\phi}_2 = (s_1 D + s_0) \phi_2,$$

where

$$s_1 = -\frac{2(r-2)}{r^3 \omega^2}, \quad s_0 = \frac{(r-2)(3r^4 \omega^2 - 14r + 12)}{3r^5 \omega^2}.$$

Its projective density is

$$\mathcal{I}_{\text{phys}} = \frac{(r-2)(3r^4 \omega^2 - 20r + 30)}{3r^5 \omega^2},$$

and obeys the exact normal form

$$\boxed{\mathcal{I}_{\text{phys}} = \mathcal{K}_{U_2} \left(\frac{r}{6\omega^2} \right) + \frac{1}{3} f.} \quad (8)$$

Consequently,

$$\boxed{[\mathcal{I}_{\text{Bach}}] = \frac{3i\omega}{2} [\mathcal{I}_{\text{phys}}].} \quad (9)$$

Proof. In the factor-state order $(\phi_2, D\phi_2, \phi_1, D\phi_1)$, conjugating the full mass tangent through the state map defined by (6)–(7) gives four rational 2×2 blocks. The reverse block scalarizes to

$$L_1 \dot{\phi}_1 = \frac{32f}{9r^3 \omega^2} \phi_2.$$

Since

$$P = -\frac{16}{27\omega^2}, \quad L_1(P\phi_2) = \frac{32f}{9r^3 \omega^2} \phi_2 \quad \text{on } \ker L_2,$$

the change $\hat{\phi}_1 = \phi_1 - mP\phi_2$ removes it. Scalarizing the tensor diagonal block gives the displayed s_1, s_0 . Exact substitution proves (8); combining it with Theorem 2.4 proves (9). $\square$

The factor $1/3$ is intrinsic. If one tests the stronger but false identity $[\mathcal{I}_{\text{phys}}] = [f]$, the local pole audit leaves the exhaustive gauge $c_{-2}/r^2 + c_{-1}/r + c_0$. Its coefficient matrix has rank 3, the augmented matrix has rank 4, and a left-null obstruction is

$$\frac{80(\omega^2 - 3)}{3\omega^2}.$$

Thus the isolated scalar pencil $L_2 - mf$ has the correct projective direction but is three times the complete tensor-branch tangent. At fixed $\omega \neq 0$, the corrected tangent relation is

$$m = \frac{3i\omega}{2} \tau$$

modulo rational triangular gauge. Its spectral dependence means that it is not a global reparametrization of the two operator families.

3.2 Endpoint-analytic massive Jost crosswalk

Because Lemma 2.3 contains a factor $1/2$, the field operator associated with a primitive q is

$$Q(q) = 2qD - D(q), \quad [L_2, Q(q)]|_{\ker L_2} = -\mathcal{K}_{U_2}(q).$$

For the complete physical tangent take

$$q_m = \frac{r}{6\omega^2}.$$

Choose

$$k = (\omega^2 - m)^{1/2}, \quad k(\omega, 0) = \omega,$$

and the common massive infinity phase

$$e^{\sigma i k r} r^{\rho_\sigma(m)}, \quad \rho_\sigma(m) = \sigma i \left(2k + \frac{m}{k} \right).$$

Then

$$\rho'_\sigma(0) = 0, \quad \frac{\partial_m \phi_{2,\sigma}}{\phi_{2,\sigma}} = -\frac{\sigma i}{2\omega} r + O(1).$$

The gauge $Q(q_m)$ adds $\sigma i r / (3\omega) + O(1)$, so the normal-form physical tangent has

$$\frac{\partial_m \widehat{\phi}_{2,\sigma}}{\phi_{2,\sigma}} = -\frac{\sigma i}{6\omega} r + O(1).$$

Multiplication by $3i\omega/2$ gives $\sigma r/4 + O(1)$, exactly the leading Bach tangent produced by the primitive in Theorem 2.4.

Theorem 3.2 (Endpoint-compatible complete mass jet). *On a neighborhood of the certified QNM and $m = 0$, the complete coupled system (4)–(5) has holomorphic two-dimensional future-horizon-ingoing and infinity-outgoing Jost planes. The Bach and complete massive first jets identify these planes modulo holomorphic same-sign changes of basis. In particular, no opposite-Jost component occurs.*

Let $a_{\text{phys}}(\omega, m)$ be the spin-two Schur divisor after eliminating the locally invertible spin-one channel. Then

$$\boxed{b_B(\omega) = \frac{3i\omega}{2} \partial_m a_{\text{phys}}(\omega, 0) + h(\omega) a_{\text{phys}}(\omega, 0),} \quad (10)$$

where h is holomorphic.

Proof. Write $\Psi = (Q, Z)^\top$. The complete equations have the form

$$D^2 \Psi + [(\omega^2 - mf)I_2 + B(r)] \Psi = 0, \quad B(r) = O(r^{-2}),$$

with

$$B = \begin{pmatrix} -f(10/r^2 - 16/r^3) & -8f(r-3)/r^3 \\ -2f/r^2 & -f(4/r^2 + 2/r^3) \end{pmatrix}.$$

At infinity use the exact common phase

$$\Phi_\sigma = e^{\sigma i k x} r^{\sigma i m/k}, \quad k^2 = \omega^2 - m, \quad k(\omega, 0) = \omega.$$

Its logarithmic derivative is

$$D \log \Phi_\sigma = \sigma i \left(k + \frac{mf}{kr} \right),$$

and direct substitution leaves the scalar residual

$$\frac{m}{k^2 r^4} (4k^2 r^2 - \sigma i k r^2 + 6\sigma i k r - 8\sigma i k - m r^2 + 4m r - 4m) = O(r^{-2}).$$

On the certified $\pi/4$ exterior-complex-scaled ray, the selected exponential has a uniform decay margin. A scalar Volterra equation for the displayed residual, followed by a matrix Volterra equation for B , has an integrable $O(r^{-2})$ majorant. After moving the outer cut sufficiently far out, both Neumann series converge uniformly on a smaller (ω, m) -polydisc. They define a holomorphic outgoing Jost matrix. Its Volterra normalization excludes the opposite exponential.

At the horizon set $z = f$. Then

$$D = \frac{z(1-z)^2}{2} \partial_z,$$

while mf and B are $O(z)$. The indicial exponents are $\lambda = \pm 2i\omega$, each twice, and the order- n recursion denominators are

$$\frac{n(n+4i\omega)}{4}, \quad \frac{n(n-4i\omega)}{4}.$$

The certified disk has nonzero real part and avoids every such resonance. The regular-singular majorant theorem therefore gives holomorphic matrix Frobenius solutions. The selected exponent fixes the ingoing plane up to a holomorphic right $GL(2)$ factor.

The differentiated Coulomb exponent vanishes, and the rational gauges match the remaining linear term exactly as computed above. At the horizon they are regular and preserve the selected exponent. The difference between the two first jets is consequently homogeneous in the same endpoint class. Volterra and Frobenius uniqueness allow only same-sign holomorphic basis changes.

It remains to remove same-sign spin-one mixing. Locally write the two-channel connection as

$$\begin{pmatrix} a + ma_1 & mb_1 \\ mc_1 & g + mg_1 \end{pmatrix}, \quad g(\omega_n) \neq 0.$$

The spin-two Schur complement is

$$a + ma_1 - \frac{m^2 b_1 c_1}{g + mg_1},$$

whose m -derivative at zero is a_1 . Thus the off-diagonal mixing does not enter the first divisor derivative. Holomorphic endpoint normalizations contribute only $h(\omega)a(\omega, 0)$, proving (10). $\square$

Corollary 3.3 (Nonzero complete massive QNM velocity). *Let $\omega_n(m)$ denote the simple complete massive axial QNM branch through the certified root. For the signed squared-mass coefficient $m = \mu^2$,*

$$\boxed{\omega'_n(0) = \frac{2i}{3\omega_n} \kappa_n \neq 0.} \tag{11}$$

A conservative propagated enclosure is

$$\operatorname{Re} \omega'_n(0) \in [0.087, 0.251], \quad \operatorname{Im} \omega'_n(0) \in [-0.054, 0.135].$$

Proof. Differentiate $a_{\text{phys}}(\omega_n(m), m) = 0$ and use (10). The certified selector and QNM disk give the displayed interval division; its real part excludes zero. $\square$

The construction is the asymptotically flat analogue of the mass-derivative mechanism behind logarithmic partners in critical gravity [5, 6]. Here the endpoint tangent is polynomial rather than an independent radial $\log r$ mode.

4 The defective Schwarzschild resonance

4.1 Smith dichotomy

At a simple spin-two zero, analytic elimination of the spin-one unit reduces (2) to

$$T_2(\omega) = \begin{pmatrix} a(\omega) & b(\omega) \\ 0 & a(\omega) \end{pmatrix}, \quad a(\omega_n) = 0, \quad a'(\omega_n) \neq 0.$$

All Smith forms in this section are taken over the discrete valuation ring

$$\mathcal{O}_{\omega_n} = \{\text{holomorphic germs at } \omega_n\},$$

whose uniformizer is $z = \omega - \omega_n$. The listed integers are the valuations of the invariant factors. Their sum is the algebraic multiplicity; the number of positive invariant factors is the geometric root-space dimension only after the unit factors have been removed. This is the local analytic-matrix setting underlying Keldysh root chains and nonlinear eigenvalue pencils [9, 10].

Proposition 4.1 (Local Smith dichotomy). *If $g(\omega_n) \neq 0$, then*

$$\text{Smith}(T) = \begin{cases} (0, 0, 2), & b(\omega_n) \neq 0, \\ (0, 1, 1), & b(\omega_n) = 0. \end{cases}$$

The first case has one length-two root chain; the second has two independent length-one roots.

Proof. With $z = \omega - \omega_n$, $a = a_1 z + O(z^2)$, $a_1 \neq 0$. The determinant valuation of the spin-two block is two. If $b(\omega_n) \neq 0$, its first Fitting ideal is the unit ideal and the exponents are $(0, 2)$. If $b(\omega_n) = 0$, all entries vanish to first order and the exponents are $(1, 1)$. The spin-one unit contributes a zero exponent. $\square$

4.2 Validated specialization

Transport the horizon-ingoing and infinity-outgoing spin-two lines to the same ordinary matching section. In the Riccati chart $v = Y_2/Y_1$, define

$$\delta(\omega, \tau) = v_\infty(\omega, \tau) - v_H(\omega, \tau), \quad \delta_\tau = \partial_\tau \delta|_{\tau=0}.$$

The scalar Evans coefficient is $a = u\delta$ for a holomorphic unit u . At a simple zero,

$$\frac{b(\omega_n)}{a'(\omega_n)} = \frac{\delta_\tau(\omega_n)}{\delta_\omega(\omega_n)}.$$

This definition removes arbitrary homogeneous amplitudes; a projective chart change multiplies numerator and denominator by the same nonzero value at the root.

The certified complex-scaled Volterra construction makes both selected lines holomorphic throughout the closed disk bounded by Γ . The disk lies inside the declared continuation strip and avoids every listed Frobenius and normalization pole. Endpoint series are initialized with outward remainder bounds; a failed remainder, chart transition, or panel overlap is a rejection. Consequently δ is holomorphic, not meromorphic, on the argument-principle domain.

The counterclockwise contour

$$\Gamma(t) = \omega_c + \frac{1}{40} e^{2\pi i t}, \quad 0 \leq t \leq 1,$$

is split into 927 validated panels. Every panel excludes zero, the overlapping argument sectors have width below π , and their unique continuous lift changes by 2π . A local Rouché calculation then encloses the unique root in

$$\Omega_n = \{\omega : |\omega - (-0.3736716844180418 + 0.0889623156889357i)| \leq 10^{-7}\}.$$

Projective interval panels exclude zeros on the contour and give winding number +1. The disk therefore contains exactly one simple spin-two zero. On the localized disk,

$$\delta_\tau(\Omega_n) \subset [0.0010932 \pm 7.60 \times 10^{-8}] + i[-0.0002195 \pm 8.89 \times 10^{-8}],$$

which excludes zero. An independent local calculation bounds the spin-one mismatch modulus below by $1/2500$.

The local transport uses outward-rounded `python-flint acb/arb` balls at 60 decimal digits, order-48 Taylor transport, 966 accepted horizon steps, and 260 accepted outgoing steps. The frequency derivative obeys

$$\delta_\omega(\Omega_n) \subset [\pm 4.88 \times 10^{-3}] + i[-0.0100 \pm 1.82 \times 10^{-3}],$$

and therefore excludes zero. Direct interval division gives the dimensionless intrinsic selector

$$\boxed{\operatorname{Re} \kappa_n \in [-0.047, 0.022], \quad \operatorname{Im} \kappa_n \in [0.064, 0.138].} \quad (12)$$

The enclosure is deliberately reported as an inequality rather than a long decimal approximation. This is an interval-validated Evans/argument-principle computation, not a floating-point winding inferred from sampled phases; compare the general Evans-function methodology in Ref. [11].

Theorem 4.2 (Certified defective resonance). *At the unique QNM in Ω_n , the complete axial connection has Smith valuations*

$$\boxed{(0, 0, 2)}.$$

Its spin-two divisor has algebraic multiplicity two, geometric multiplicity one, and one length-two root chain.

Proof. The winding certificate supplies the simple spin-two zero. The selector enclosure gives $b(\omega_n) \neq 0$, and the spin-one bound gives $g(\omega_n) \neq 0$. Proposition 4.1 applies. $\square$

4.3 One resonant invariant

Encode the endpoint conditions in an analytic finite-interval pencil

$$L(\omega) : X \longrightarrow Y.$$

Such holomorphic boundary pencils and their contour/root-chain reductions fit the analytic Fredholm framework of Refs. [12, 13]. Choose right and left states at the simple scalar resonance,

$$L_n u_n = 0, \quad L_n^* \tilde{u}_n = 0,$$

and define, with all augmented endpoint contributions included,

$$\alpha_n = \langle \tilde{u}_n, L_n' u_n \rangle \neq 0.$$

For the repeated block

$$\mathbb{L}(\omega) = \begin{pmatrix} L(\omega) & K(\omega) \\ 0 & L(\omega) \end{pmatrix},$$

set

$$\beta_n = \langle \tilde{u}_n, K_n u_n \rangle.$$

Theorem 4.3 (Resonant evaluation). *In endpoint frames compatible with the partial jet,*

$$\boxed{\kappa_n = \frac{b(\omega_n)}{a'(\omega_n)} = \frac{\beta_n}{\alpha_n} = - \left. \frac{d\omega_n}{d\tau} \right|_0 = - \frac{3i\omega_n}{2} \omega'_n(0).} \quad (13)$$

The ratio is invariant under a common analytic nonzero Evans unit and under triangular changes of splitting.

Proof. Differentiate

$$[L(\omega) + \tau K(\omega)]u(\tau) = 0$$

along the resonance branch. Pairing with $\tilde{u}_n$ gives

$$\dot{\omega}_{n,\tau} \alpha_n + \beta_n = 0, \quad \dot{\omega}_{n,\tau} := \left. \frac{d\omega_n}{d\tau} \right|_0.$$

Differentiating the connection divisor gives

$$a'(\omega_n) \dot{\omega}_{n,\tau} + b(\omega_n) = 0,$$

because the partial jet identifies $b = \partial_\tau a$. The first three equalities do not use the mass crosswalk; the last follows from Theorem 3.2. A triangular change $K \mapsto K + LQ - QL$ has zero resonant pairing, and an Evans unit multiplies numerator and denominator by the same nonzero value. $\square$

The certified enclosure (12) is intrinsic. Its conversion to the complete massive slope uses the endpoint theorem and the fixed-frequency factor $3i\omega/2$, rather than treating that spectral factor as a global parameter redefinition.

Proposition 4.4 (Non-Einstein generalized root). *The geometric root belongs to the Einstein submodule. Every length-two root polynomial has generalized coefficient with nonzero projection to the Ricci-carrier quotient. In the normalized triangular frame, that projection is*

$$\boxed{- \frac{a'(\omega_n)}{b(\omega_n)} = - \frac{\alpha_n}{\beta_n} = - \frac{1}{\kappa_n} \neq 0.}$$

Proof. Write $a = a_1 z + O(z^2)$, $b = b_0 + O(z)$, with $a_1 b_0 \neq 0$. For a root polynomial

$$c(z) = c_0 + z c_1 + O(z^2), \quad T_2(z) c(z) = O(z^2),$$

the order-zero equation gives $c_0 = e_X$, the Einstein vector. Writing $c_1 = x e_X + y e_Y$, the order- z equation is

$$a_1 + b_0 y = 0.$$

Hence $y = -a_1/b_0 \neq 0$. Adding an Einstein vector changes x , not the carrier coefficient y . $\square$

Thus the defective resonance is not the same Einstein QNM counted twice. Its generalized member is the canonical mass-tangent class modulo the ordinary QNM.

5 Exterior Green pole and critical response

5.1 Finite cuts and the exterior inverse

Put $x = r_*$. On the certified QNM neighborhood, let

$$u_H(x, \omega), \quad u_\infty(x, \omega)$$

be analytic spin-two Jost solutions satisfying the future-horizon and outgoing-infinity conditions. Their Wronskian is the scalar Evans coefficient, up to an analytic unit. Our cut-off exterior terminology is consistent with rigorous asymptotically flat resonance definitions through meromorphically continued cut-off resolvents [14, 15]; the present theorem is only the local radial realization described below.

Choose nontrivial smooth source and observation cutoffs χ_s, χ_o and finite cuts

$$x_- < \text{supp } \chi_s \cup \text{supp } \chi_o < x_+.$$

Impose the exact transparent boundary conditions

$$W(u, u_H)|_{x_-} = 0, \quad W(u, u_\infty)|_{x_+} = 0.$$

They are analytic in ω , and uniqueness of the exterior Jost problems identifies the finite-interval inverse between the cutoffs with $\chi_o R_{\text{ext}}(\omega) \chi_s$.

Theorem 5.1 (Exterior cut-off Green pole). *For any such nontrivial compact source and observation cutoffs,*

$$\chi_o R_{\text{ext}}(\omega) \chi_s$$

has a nonzero rank-one pole of order two at ω_n . In the augmented normalization,

$$\boxed{\text{Coeff}_{(\omega - \omega_n)^{-2}}(\chi_o R_{\text{ext}} \chi_s) = -\frac{\beta_n}{\alpha_n^2} (\chi_o u_n) \otimes (\tilde{u}_n \chi_s).} \quad (14)$$

Proof. The repeated block inverse is

$$\mathbb{L}^{-1} = \begin{pmatrix} L^{-1} & -L^{-1} K L^{-1} \\ 0 & L^{-1} \end{pmatrix}.$$

At a simple scalar resonance,

$$L(\omega)^{-1} = \frac{u_n \otimes \tilde{u}_n}{\alpha_n(\omega - \omega_n)} + O(1).$$

The upper-right block therefore has principal coefficient

$$-\frac{\beta_n}{\alpha_n^2} u_n \otimes \tilde{u}_n.$$

The exact transparent conditions identify this finite-interval coefficient with the cut-off exterior coefficient. It is nonzero because $\beta_n \neq 0$, and a nonzero solution of a second-order ODE cannot vanish on a nonempty open interval. $\square$

The full doubled-state principal coefficient maps the carrier channel to the Einstein channel and vanishes on the Einstein channel; it is square-zero on that graded space. Its projection to a single metric or radial block is rank one but is not intrinsically a nilpotent operator after the grading has been forgotten.

5.2 A fixed-domain global complex-scaled pencil

The finite transparent cuts can be replaced by a single global Sobolev domain on a deformed radial contour. Put $\theta = \pi/4$, choose finite matching points $x_H < x_I$, and parameterize a two-ended contour by

$$x_\theta(s) = \begin{cases} x_H + e^{i\theta}(s+1), & s \leq -1, \\ x_c(s), & -1 \leq s \leq 1, \\ x_I + e^{i\theta}(s-1), & s \geq 1, \end{cases}$$

where x_c is a fixed compact connector. On the right tail use the certified inverse-tortoise branch. On the left tail use $z = f$ and

$$\frac{dz}{dx} = \frac{z(1-z)^2}{2};$$

then $z \rightarrow 0$ exponentially as $s \rightarrow -\infty$.

Let $\mathbb{A}_x$ be the complete six-state matrix after passage to tortoise coordinate and define

$$\mathcal{L}_\theta(\omega)Y = \frac{dY}{ds} - x'_\theta(s)\mathbb{A}_x(x_\theta(s), \omega)Y : H^1(\mathbb{R}; \mathbb{C}^6) \longrightarrow L^2(\mathbb{R}; \mathbb{C}^6).$$

The contour and both Banach spaces are independent of ω .

Theorem 5.2 (Global ECS Fredholm pole). *After shrinking to a neighborhood of Ω_n , $\mathcal{L}_\theta(\omega)$ is a holomorphic Fredholm pencil of index zero. Its inverse is meromorphic and has a nonzero rank-one pole of order two at ω_n :*

$$\mathcal{L}_\theta(\omega)^{-1} = \frac{\Pi_{-2}^\theta}{(\omega - \omega_n)^2} + \frac{\Pi_{-1}^\theta}{\omega - \omega_n} + O(1), \quad \text{rank } \Pi_{-2}^\theta = 1.$$

The ordinary resonant state and its differentiated Jost tangent belong to the same fixed H^1 domain. For sources and observations supported on the undeformed compact core, the corresponding matrix elements agree with the exterior cut-off inverse of Theorem 5.1.

Proof. At either end the limiting first-order matrix has characteristic rates

$$\pm ie^{i\theta}\omega$$

with multiplicity three; nontrivial Jordan blocks within a repeated sign do not affect hyperbolicity. On the certified frequency disk,

$$\text{Re}(-ie^{i\theta}\omega) < -\rho < 0, \quad \text{Re}(+ie^{i\theta}\omega) > \rho > 0.$$

Thus the infinity-outgoing space has forward stable dimension three, while the horizon-ingoing space has forward unstable dimension three and decays as $s \rightarrow -\infty$. The asymptotic first-order operator has index

$$\dim E_-^u + \dim E_+^s - 6 = 3 + 3 - 6 = 0.$$

The coefficient remainder decays exponentially in the horizon chart and tends to zero algebraically at infinity. Its multiplication map $H^1 \rightarrow L^2$ is compact: Rellich compactness handles a fixed finite interval, while the tail operator norm tends to zero. Hence the full pencil is Fredholm of the same index and is holomorphic on the fixed domain.

The differentiated outgoing germ is bounded by a constant times $(1 + |s|)e^{-\rho|s|}$, together with the same type of bound for its derivative. In particular,

$$\int_0^\infty (1+t)^2 e^{-2\rho t} dt = \frac{2\rho^2 + 2\rho + 1}{4\rho^3} < \infty.$$

It therefore lies in the same domain as the ordinary selected germ.

Finally, analytic exponential-dichotomy Green operators eliminate the two tails while retaining their selected finite-cut trace spaces. The remaining compact-core pencil has exactly the Jost transparent planes used above, so it is analytically equivalent to invertible factors direct sum the certified 3×3 connection matrix. Smith valuations $(0, 0, 2)$ give the stated Laurent order and principal rank. Uniqueness of the analytic tail solutions gives equality with the exterior outgoing inverse on compactly supported source and observation channels. $\square$

Theorem 5.2 closes the fixed-domain Fredholm problem for the reduced complex-scaled radial pencil. Complex scaling selects and regularizes resonances; it does not by itself supply a causal real-axis inverse or authorize deformation of a physical inverse-Laplace contour.

5.3 Parent normalization and isolated contour

Our normalization of α_W is fixed by the quadratic parent action

$$S_m^{(2)} = 4\alpha_W \int \left[\phi \cdot \mathcal{E}(h) - \frac{1}{2} \phi \cdot \mathcal{A}(\phi) + \frac{m}{2} h \cdot \mathcal{E}(h) \right].$$

The equations are

$$\mathcal{E}(h) = \mathcal{A}(\phi), \quad \mathcal{E}(\phi) + m\mathcal{E}(h) = 0.$$

Eliminating the auxiliary field and inverting the $m = 0$ block gives

$$G_{hh}^W = \frac{1}{4\alpha_W} \mathcal{E}^{-1} \mathcal{A} \mathcal{E}^{-1}.$$

At points where the second-order tensor inverses exist,

$$\partial_m (\mathcal{E} + m\mathcal{A})^{-1} = -(\mathcal{E} + m\mathcal{A})^{-1} \mathcal{A} (\mathcal{E} + m\mathcal{A})^{-1}.$$

Consequently

$$\boxed{G_{hh}^W(\omega) = -\frac{1}{4\alpha_W} \partial_m (\mathcal{E} + m\mathcal{A})^{-1} \big|_{m=0}.} \quad (15)$$

No commutativity assumption is used.

Theorem 5.3 (Causal Laplace-to-resonance bridge). *Restrict the parent family $\mathcal{E}_m = \mathcal{E} + m\mathcal{A}$ to the gauge-invariant axial $\ell = 2$ transverse-traceless spin-two sector on the Schwarzschild domain of outer communications. Let G_m^{ret} denote its retarded Green operator. Then*

$$\boxed{G_{hh,W}^{\text{ret}} = -\frac{1}{4\alpha_W} \partial_m G_m^{\text{ret}} \big|_{m=0} = \frac{1}{4\alpha_W} G_0^{\text{ret}} \mathcal{A} G_0^{\text{ret}}.}$$

This operator is retarded. For compact radial source and observation cutoffs, its Laplace transform, initially defined in a lower half-plane, has a unique meromorphic continuation to Ω_n . That continuation has a nonzero rank-one pole of order two with

$$\boxed{G_{-2} = -\frac{\omega'_n(0)}{4\alpha_W} P_n, \quad \omega'_n(0) = \frac{2i}{3\omega_n} \kappa_n \neq 0.}$$

Proof. After axial transverse-traceless reduction, $\mathcal{E}_m$ is normally hyperbolic with a smooth zeroth-order parameter perturbation. It therefore has unique retarded Green operators on the globally hyperbolic domain of outer communications [8]. Duhamel differentiation on every finite causal slab gives

$$\partial_m G_m^{\text{ret}}|_0 = -G_0^{\text{ret}} \mathcal{A} G_0^{\text{ret}}.$$

The composition is retarded because the composition of two future-causal support relations remains future-causal.

With the convention $e^{+i\omega t}$, use the forward transform

$$\widehat{F}(\omega) = \int_0^\infty e^{-i\omega t} F(t) dt.$$

It converges below the energy-growth bound. Integration by parts changes the reduced wave equation $\partial_t^2 \psi + H_m \psi = F$, with zero past data, into

$$(H_m - \omega^2) \widehat{\psi} = \widehat{F}.$$

In that lower half-plane the square-integrable radial solutions are precisely horizon-ingoing and infinity-outgoing, so compactly cut-off matrix elements equal the outgoing radial transfer.

The complete parameter-analytic Jost construction and Theorem 5.2 continue this same transfer meromorphically to Ω_n . Equality in the initial half-plane makes the continuation unique. Differentiating the moving simple-pole expansion of the massive resolvent gives

$$-\partial_m \frac{P_n(m)}{\omega - \omega_n(m)} \Big|_0 = -\frac{\omega'_n(0) P_n}{(\omega - \omega_n)^2} - \frac{P'_n(0)}{\omega - \omega_n}.$$

The complete Jost crosswalk proves $\omega'_n(0) \neq 0$, which gives the stated nonzero rank-one principal coefficient. $\square$

The theorem is **LORENTZIAN-CAUSAL** only at the declared mode-reduced transverse-traceless level. It is not a full off-shell metric/BV propagator and does not identify a specified real matter source. Most importantly, meromorphic continuation is not yet a deformation of the complete inverse-Laplace contour.

If a fixed-domain massive tensor pencil has the local simple-pole form

$$R_m(\omega) = \frac{P_n(m)}{\omega - \omega_n(m)} + H_n(\omega, m),$$

then (15) gives

$$G_{hh}^W(\omega) = -\frac{1}{4\alpha_W} \left[\frac{\omega'_n(0) P_n(0)}{(\omega - \omega_n)^2} + \frac{P'_n(0)}{\omega - \omega_n} + \partial_m H_n(\omega, 0) \right].$$

Thus the double pole measures QNM motion; the accompanying simple pole measures motion of the residue projector. Theorem 3.2 supplies the endpoint-normalized identification between the certified scalar selector and the complete massive axial QNM velocity.

Corollary 5.4 (Isolated resonance contour). *Write the certified radial inverse as*

$$R_{\text{ext}}(\omega) = \frac{\Pi_{-2}}{(\omega - \omega_n)^2} + \frac{\Pi_{-1}}{\omega - \omega_n} + O(1)$$

between compact source and observation cutoffs. For a contour enclosing only this local resonance,

$$\frac{1}{2\pi i} \oint e^{i\omega t} R_{\text{ext}}(\omega) d\omega = e^{i\omega_n t} (\Pi_{-1} + it \Pi_{-2}).$$

The t -weighted spatial profile is the ordinary Einstein QNM, while the constant generalized component crosses the carrier quotient.

This is a local residue theorem, not a global retarded expansion. The latter would require deformation of the physical Laplace contour, bounded asymptotic reconstruction, and control of thresholds, high frequencies, and non-pole contributions. Theorems 5.2 and 5.3 settle the local resonance realization and its causal origin, but not that global contour shift.

6 Source and observation channels

6.1 Metric reconstruction at null infinity

Let E be the unit outgoing Einstein line and R the unit outgoing carrier line in the exact factor frame. After factoring the common retarded phase $e^{i\omega u}$, their Regge–Wheeler-gauge metric heads are

$$(H_0, H_1)_E = (-r, 2r) + O(1),$$

$$(H_0, H_1)_R = \left(\frac{3}{4}r^2 - \frac{3}{2}r, -\frac{3}{2}r^2 \right) + O(1).$$

Hence

$$(H_0, H_1)_R = -\frac{3}{4}r (H_0, H_1)_E + O(E).$$

In odd radiation gauge,

$$h_u \mapsto h_u - i\omega \xi, \quad h_2 \mapsto h_2 - 2\xi, \quad \xi = \frac{h_u}{i\omega},$$

so $h_2 = 2ih_u/\omega$. Writing $h_{AB} = h_2 X_{AB}$, define the Bondi-shear observation map by

$$\mathcal{O}_{\mathcal{J}^+} h = \lim_{r \rightarrow \infty} \frac{h_{AB}}{r}.$$

Theorem 6.1 (Null-infinity reconstruction). *The outgoing Einstein line has nonzero Bondi shear,*

$$\boxed{\mathcal{O}_{\mathcal{J}^+} E = -\frac{2i}{\omega} X_{AB} \neq 0.} \tag{16}$$

Its strain has the standard radiative falloff

$$\frac{h_{AB}^E}{r^2} = -\frac{2i}{\omega r} e^{i\omega u} X_{AB} + O(r^{-2}).$$

The carrier line instead has

$$\boxed{\frac{h_{AB}^R}{r^2} = \frac{3i}{2\omega} e^{i\omega u} X_{AB} + O(r^{-1}).} \tag{17}$$

Therefore the time-independent generalized component has weakened $O(1)$ strain in this standard odd radiation gauge. The Einstein range of the double-pole coefficient is nevertheless observed nontrivially at future null infinity.

Proof. Substitute the exact outgoing factor-frame series into the algebraic metric reconstruction. The odd gauge law then gives (16) and (17). Adding an Einstein representative cannot cancel the carrier r^2 coefficient, which is one power higher. The trace-free odd tensor harmonic cannot be removed by a Weyl rescaling; its removal would require a large boundary diffeomorphism rather than a standard small gauge transformation. $\square$

Proposition 6.2 (Outgoing-trace bridge). *On the local QNM disk, coefficient extraction from the selected outgoing Jost family defines a holomorphic trace*

$$\mathcal{T}_{\mathcal{J}+} : u \mapsto \text{outgoing asymptotic coefficient}.$$

The algebraic reconstruction followed by $\mathcal{T}_{\mathcal{J}+}$ is therefore an analytic observation map. For compactly supported forcing,

$$\text{Coeff}_{(\omega-\omega_n)^{-2}}(\mathcal{O}_{\mathcal{J}+} R_{\text{ext}} \chi_s) = -\frac{\beta_n}{\alpha_n^2} \mathcal{O}_{\mathcal{J}+}(u_n) \otimes (\tilde{u}_n \chi_s).$$

In particular, Theorem 6.1 implies that the first factor is nonzero.

Proof. The outgoing coefficients and reconstruction matrices are holomorphic on the local disk. Applying a holomorphic map to a Laurent expansion acts coefficientwise on its principal part. Substitute (14) and then (16). $\square$

The gauge parameter used above is allowed only as an algebraic reconstruction into the chosen odd radiation gauge. For the Einstein line it preserves the standard Bondi falloff. For the carrier line it is large relative to the usual asymptotically flat phase space, and (17) is consequently not claimed to define finite Bondi flux or an admissible asymptotic state.

Combining Theorem 5.1 with Proposition 6.2, the principal Bondi-shear coefficient is nonzero. Using Corollary 3.3, its parent-normalized form may be written

$$\mathcal{O}_{\mathcal{J}+} G_{-2} = \frac{i \omega'_n(0)}{2\alpha_W \alpha_n \omega_n} X_{AB} \otimes \tilde{u}_n.$$

The isolated contour therefore contains an ordinary u/r radiative profile multiplied by the adjoint source overlap.

6.2 A complexified conserved, traceless source

Let $\ell \geq 2$, $\mu = (\ell - 1)(\ell + 2)$, and $\omega \neq 0$. Use the Martel–Poisson odd source projections and orientation $\epsilon_{tr} = 1$, $\epsilon^{tr} = -1$ [7].

Theorem 6.3 (Complexified conserved-source nonannihilation). *For any*

$$F \in C_c^\infty((2M, \infty)),$$

define

$$P_t = 0, \quad P_r = \frac{\mu F}{2i\omega r f}, \quad P = \frac{\partial_r(rF)}{2i\omega}.$$

Then

$$\nabla_a P^a + \frac{2}{r} r_a P^a - \frac{\mu}{r^2} P = 0,$$

and the Cunningham–Price–Moncrief source is

$$S_{\text{odd}} = -\frac{2r}{\mu}\epsilon^{ab}\nabla_a P_b = \frac{F}{f}.$$

The associated odd stress tensor

$$T^{aB} = \frac{P^a}{16\pi r^2} X^B, \quad T^{AB} = \frac{P}{8\pi r^4} X^{AB},$$

with all other harmonic components zero, is smooth and radially compact, and satisfies

$$\boxed{\nabla_\mu T^{\mu\nu} = 0, \quad T^\mu{}_\mu = 0.}$$

The complexified conserved-traceless odd source map is therefore onto the smooth compact reduced radial forcings.

Moreover, choose a nonnegative nonzero $\eta \in C_c^\infty((2M, \infty))$ supported where the adjoint state $\tilde{u}_n$ is nonzero and set

$$F = \eta \overline{\tilde{u}_n}.$$

Then the augmented pairing has no endpoint term and

$$\boxed{\langle \tilde{u}_n, F \rangle_{\text{aug}} = \int \eta |\tilde{u}_n|^2 dr_* > 0.}$$

Thus the space of complexified frequency-domain conserved and traceless odd sources does not annihilate the certified double pole.

Proof. Since $\sqrt{-\det g_{ab}} = 1$,

$$P^r = f P_r = \frac{\mu F}{2i\omega r}, \quad \nabla_a P^a = \partial_r P^r.$$

Direct substitution proves the reduced conservation identity and $f S_{\text{odd}} = F$. The remaining spacetime conservation equations vanish by the divergence-free odd vector harmonic, and tracelessness follows from $T^{ab} = 0$ and $\Omega_{AB} X^{AB} = 0$. Compact support removes all endpoint terms from the adjoint pairing, and the final integral is strictly positive. $\square$

Remark 6.4 (What the source theorem does not say). The choice $F = \eta \overline{\tilde{u}_n}$ depends on the complex QNM frequency. It is an existence proof in the complexified frequency-domain source space, not a real, causal, temporally compact matter source. A real formal perturbation can be assembled by adjoining the reflected frequency and harmonic sector, but causality and temporal support still require an inverse-Laplace construction. The theorem also does not identify a specified material trajectory or an energy-condition-satisfying model. A standard massive point-particle stress tensor is not trace-free and cannot be coupled directly to the Bach equation without a conformal completion or compensator.

7 Scope, evidence, and open problem

7.1 The remaining analytic problem

The exterior cut-off theorem and Theorems 5.2 and 5.3 give the compact outgoing matrix elements, a global fixed-domain complex-scaled realization, and their origin as the meromorphic continuation of a retarded mode-reduced transfer. The remaining time-domain problem is narrower but substantive. It must provide:

1. bounded reconstruction through the physical gauge quotient and null-infinity trace on the causal solution;
2. a causal Laplace contour and a justified deformation through the QNM disk;
3. control of threshold, branch-cut, high-frequency, and non-pole contributions.

The weakened falloff in (17) explains why this is not routine. A standard outgoing asymptotically flat function space may contain the Einstein root but exclude its intrinsic generalized tangent. On the ECS contour the differentiated germ is exponentially damped and belongs to the same fixed H^1 domain as the ordinary QNM. This solves the reduced Fredholm-domain issue without deciding whether a standard real-axis asymptotic phase space contains the constant generalized metric component.

7.2 Certificate dependency

Certificate family	Imported statement	Type
Axial triangular preflight	Filtration, factor gauge, and reconstruction rank	Exact
Projective cocycle	Cocycle, primitive, and rational nonsplitting obstruction	Exact
Complete massive axial first jet	Coupled Q, Z factorization and factor-three Bach crosswalk	Exact
Complete massive Jost crosswalk	Matrix Frobenius/Volterra endpoint planes and nonzero QNM velocity	Analytic and validated
Analytic continuation	Analytic horizon and infinity Jost germs on the QNM disk	Exact gate
Full Evans contour	Zero-free contour and winding +1	Validated
Local selector	Nonzero projective tangent on the root disk	Validated
Spin-one local unit	Nonzero spin-one mismatch	Validated
Fredholm promotion	Finite-interval Poisson/trace reduction	Exact reduction
Global ECS Fredholm realization	Fixed H^1 domain, index zero, and connection Smith transfer	Exact analytic reduction
Causal Laplace bridge	Retarded parent derivative and meromorphic continuation	Exact causal reduction
Null-infinity reconstruction	Metric heads, radiation gauge, and Bondi shear	Exact
Conserved source	Conservation, tracelessness, and adjoint overlap	Exact

The generated claim map pins the authority files and their hashes. Its verifier independently checks the displayed cocycle identity, the complete-system factor-three tangent normalization, the Smith and Green formulas, the reconstruction falloffs, the source conservation law, and explicit non-results. Missing data, a stale hash, an unsupported domain, or a failed mutation is a rejection rather than a partial pass.

7.3 Reproduction and trust boundary

The exact certificates use Python integer/rational arithmetic and SymPy 1.14.0 for polynomial reduction. Validated complex numerics use `python-flint` 0.9.0 `arb/acb` balls with directed outward rounding. The reviewed environment uses Python 3.12.13 and `jsonschema` 4.26.0. Certificate parsing, these arithmetic libraries, the Python runtime, and the operating system form the trusted computing base. Displayed decimal intervals are outward enclosures; the machine records retain midpoint and radius strings that round-trip through `arb`.

The principal independent verification commands are:

Claim	Command
Projective cocycle and nonsplitting	<code>python3-mblack_hole_programme.phase3.axial_qnm_projective_cocycle_v1.verify</code>
Complete massive first jet	<code>python3-mblack_hole_programme.phase4.axial_complete_massive_jet_crosswalk_v1.verify</code>
Complete massive Jost crosswalk	<code>python3-mblack_hole_programme.phase4.axial_massive_jost_crosswalk_v1.verify</code>
Evans winding	<code>python3-mblack_hole_programme.phase3.axial_qnm_projective_evans_contour_completion.full_contour_winding_v1.verify</code>
Local selector	<code>python3-mblack_hole_programme.phase3.axial_qnm_projective_evans_contour_completion.local_selector_v1.verify</code>
Spin-one unit	<code>python3-mblack_hole_programme.phase3.axial_qnm_spin_one_local_unit_v1.verify</code>
Radial Green pole	<code>python3-mblack_hole_programme.phase4.axial_qnm_fredholm_promotion_v1.verify</code>
Causal Laplace bridge	<code>python3-mblack_hole_programme.phase4.axial_qnm_causal_laplace_bridge_v1.verify</code>
Null-infinity reconstruction	<code>python3-mblack_hole_programme.phase4.axial_qnm_null_infinity_reconstruction_v1.verify</code>
Complexified source overlap	<code>python3-mblack_hole_programme.phase4.axial_qnm_conserved_source_overlap_v1.verify</code>

The accompanying claim-map JSON records exact paths, SHA-256 hashes, producer and verifier commands, and mutation tests. A Git commit is content-addressed but is not an archival preservation service. An immutable DOI-bearing release of the claim map, certificates, source systems, and environment lockfile is therefore an explicit submission gate rather than a completed claim of this draft.

Explicit non-results. This paper does not establish a full off-shell metric/BV retarded propagator, a complete QNM expansion, a late-time asymptotic formula, a retarded inverse-Laplace deformation, a time-domain instability, finite Bondi flux for the generalized constant component, a standard asymptotically flat domain containing that component, excitation by a specified astrophysical source, detector sensitivity, an overtone tower, an all-multipole Bach nonsplitting theorem, or a quantum positivity/no-go theorem. The polynomially weighted term is proved only for the isolated local resonance contour.

8 Conclusion

The repeated Regge–Wheeler factor in axial Schwarzschild pure Weyl gravity is not a direct sum. It is a non-split self-extension with the elementary projective class

$$[\mathcal{I}_{\text{Bach}}] = \frac{i\omega}{2}[1 - 2/r].$$

Transforming the complete coupled massive axial system into its massless Maxwell/Regge–Wheeler basis gives

$$[\mathcal{I}_{\text{phys}}] = \frac{1}{3}[1 - 2/r], \quad [\mathcal{I}_{\text{Bach}}] = \frac{3i\omega}{2}[\mathcal{I}_{\text{phys}}].$$

The corrected rational primitive matches the differentiated massive phase, and the Coulomb exponent has vanishing first derivative. The matrix Frobenius/Volterra construction promotes the comparison to the complete endpoint-normalized Jost planes and proves

$$\omega'_n(0) = \frac{2i}{3\omega_n}\kappa_n \neq 0.$$

At the certified damped Schwarzschild mode, the scalar QNM is simple but the extension selector is nonzero. The full connection is therefore defective. Its geometric root is Einstein; its generalized root has a nonzero Ricci-carrier quotient. The same intrinsic invariant controls the nonzero rank-one second-order pole of the exterior cut-off outgoing Green operator.

The leading pole survives an asymptotic observation test and a complexified source-space test. Its Einstein range has nonzero Bondi shear, and a radially compact conserved, traceless odd source can have strictly nonzero adjoint overlap. The enhanced isolated-contour profile is therefore an ordinary outgoing Einstein QNM multiplied by time, even though the constant generalized component has weakened asymptotic falloff.

This gives a Schwarzschild counterpart, in asymptotically flat spacetime, of the critical-gravity logarithmic-partner mechanism at the level of the filtered Bach resonance. The remaining work is genuinely global: prove bounded causal metric/Bondi reconstruction and justify the retarded contour deformation with threshold and high-frequency control. The fixed-domain complex-scaled Fredholm realization and the causal Laplace-to-resonance bridge of the reduced radial response are complete.

References

- [1] T. Regge and J. A. Wheeler, Stability of a Schwarzschild singularity, *Physical Review* **108** (1957), 1063–1069, [doi:10.1103/PhysRev.108.1063](https://doi.org/10.1103/PhysRev.108.1063).
- [2] E. W. Leaver, An analytic representation for the quasinormal modes of Kerr black holes, *Proceedings of the Royal Society of London A* **402** (1985), 285–298, [doi:10.1098/rspa.1985.0119](https://doi.org/10.1098/rspa.1985.0119).
- [3] R. Brito, V. Cardoso, and P. Pani, Massive spin-2 fields on black hole spacetimes: Instability of the Schwarzschild and Kerr solutions and bounds on the graviton mass, *Physical Review D* **88** (2013), 023514, [doi:10.1103/PhysRevD.88.023514](https://doi.org/10.1103/PhysRevD.88.023514).
- [4] G. Antoniou, L. Gualtieri, and P. Pani, Gravitational quasinormal modes of black holes in quadratic gravity, *Physical Review D* **111** (2025), 064059, [doi:10.1103/PhysRevD.111.064059](https://doi.org/10.1103/PhysRevD.111.064059).
- [5] H. Lü and C. N. Pope, Critical gravity in four dimensions, *Physical Review Letters* **106** (2011), 181302, [doi:10.1103/PhysRevLett.106.181302](https://doi.org/10.1103/PhysRevLett.106.181302).
- [6] E. A. Bergshoeff, O. Hohm, J. Rosseel, and P. K. Townsend, Modes of log gravity, *Physical Review D* **83** (2011), 104038, [doi:10.1103/PhysRevD.83.104038](https://doi.org/10.1103/PhysRevD.83.104038).
- [7] K. Martel and E. Poisson, Gravitational perturbations of the Schwarzschild spacetime: A practical covariant and gauge-invariant formalism, *Physical Review D* **71** (2005), 104003, [doi:10.1103/PhysRevD.71.104003](https://doi.org/10.1103/PhysRevD.71.104003).
- [8] C. Bär, Green-hyperbolic operators on globally hyperbolic spacetimes, *Communications in Mathematical Physics* **333** (2015), 1585–1615, [doi:10.1007/s00220-014-2097-7](https://doi.org/10.1007/s00220-014-2097-7).
- [9] M. V. Keldysh, On the characteristic values and characteristic functions of certain classes of non-self-adjoint equations, *Doklady Akademii Nauk SSSR* **77** (1951), 11–14.
- [10] I. Gohberg, P. Lancaster, and L. Rodman, *Matrix Polynomials*, Classics in Applied Mathematics 58, SIAM, Philadelphia, 2009, [doi:10.1137/1.9780898719024](https://doi.org/10.1137/1.9780898719024).

- [11] B. Barker, J. Humpherys, G. Lyng, and J. Lytle, Evans function computation for the stability of travelling waves, *Philosophical Transactions of the Royal Society A* **376** (2018), 20170184, [doi:10.1098/rsta.2017.0184](https://doi.org/10.1098/rsta.2017.0184).
- [12] W.-J. Beyn, An integral method for solving nonlinear eigenvalue problems, *Linear Algebra and its Applications* **436** (2012), 3839–3863, [doi:10.1016/j.laa.2011.03.030](https://doi.org/10.1016/j.laa.2011.03.030).
- [13] W.-J. Beyn, Y. Latushkin, and J. Rottmann-Matthes, Finding eigenvalues of holomorphic Fredholm operator pencils using boundary value problems and contour integrals, *Integral Equations and Operator Theory* **78** (2014), 155–211, [doi:10.1007/s00020-013-2117-6](https://doi.org/10.1007/s00020-013-2117-6).
- [14] M. Stucker, Quasinormal modes for the Kerr black hole, arXiv:2407.04612 (2024), [doi:10.48550/arXiv.2407.04612](https://doi.org/10.48550/arXiv.2407.04612).
- [15] D. Gajic and C. Warnick, Quasinormal modes on Kerr spacetimes, arXiv:2407.04098 (2024), [doi:10.48550/arXiv.2407.04098](https://doi.org/10.48550/arXiv.2407.04098).