

# A One-Loop BV Obstruction in Pure Weyl Gravity and Its Formal Tau-Adic Wess–Zumino Resolution

Asger Alstrup Palm\*

Working draft, July 2026

## Abstract

We compute the coefficient-bearing one-loop BV anomaly of strict four-dimensional pure Weyl gravity in a complete local gauge-fixed quotient. The repository Euclidean elliptic complex, measure, zero-mode, contour, parity, and local heat-kernel ledgers give

$$\mathcal{A}^{(1)} = \frac{1}{(4\pi)^2} \left( \frac{199}{30} [\omega C^2] - \frac{87}{20} [\omega E_4] \right).$$

Both coordinates are nonzero in the strict quotient. Hence the fixed-field-content local Euclidean quantum master equation is obstructed at one loop. An exact dual-cone witness further excludes cancellation by any nonnegative combination of standard-sign free conformal scalars, Weyl or Dirac fermions, and gauge vectors.

We then adjoin a shifting scalar  $\tau$  and derive its complete minimal-BV cotangent lift from  $S_{\tau, \min} = \int \tau^* (\mathcal{L}_\xi \tau + \omega)$ . In dressed canonical variables

$$\hat{g} = e^{-2\tau} g, \quad \hat{g}^* = e^{2\tau} g^*, \quad \hat{\tau}^* = \tau^* + 2g_{\mu\nu} g^{*\mu\nu},$$

the Weyl sector is an exact contractible quartet. In the formal tau-adic local analytic jet algebra the complete dimension-four extended quotient is

$$\dim H_{\text{ext}}^{0,4} = (3 \text{ even}, 1 \text{ odd}), \quad H_{\text{ext}}^{1,4} = 0.$$

The coefficient-bearing Wess–Zumino counterterm therefore restores the extended local Euclidean QME at one loop.

The two dispositions refer to different theories and are not contradictory: strict field content is obstructed, while the formal tau-adic compensator extension is locally restored at one loop. Neither statement is an all-loop or Lorentzian QME theorem. No Hadamard state, particle Hilbert space, unitarity theorem, or residual quantum transfer is claimed.

## AI-use disclosure and computational accountability

OpenAI GPT-5 was used substantively for mathematical synthesis, verification code, claim auditing, and manuscript drafting. The human author directed the programme and remains responsible for the claims, citations, and final text. All cohomological ranks and cancellations reported here use exact rational arithmetic. Every headline statement is bound to a deterministic certificate and an independent verifier. Submission remains conditional on final human and literature review.

---

\*Corresponding author: [asger@area9.dk](mailto:asger@area9.dk).

# Contents

|          |                                                |          |
|----------|------------------------------------------------|----------|
| <b>1</b> | <b>Question, result, and claim boundary</b>    | <b>2</b> |
| <b>2</b> | <b>Strict local BV obstruction space</b>       | <b>2</b> |
| <b>3</b> | <b>Coefficient-bearing one-loop breaking</b>   | <b>3</b> |
| <b>4</b> | <b>The compensator cotangent lift</b>          | <b>3</b> |
| <b>5</b> | <b>Extended cohomology and QME restoration</b> | <b>4</b> |
| <b>6</b> | <b>What does not follow</b>                    | <b>5</b> |
| <b>7</b> | <b>Machine certification and reproduction</b>  | <b>5</b> |
| <b>8</b> | <b>Outlook</b>                                 | <b>5</b> |

## 1 Question, result, and claim boundary

Pure Weyl gravity has classical  $\text{Diff} \times \text{Weyl}$  gauge symmetry. The quantum question is not whether a determinant has a nonzero logarithm, but whether the regularized Slavnov breaking is trivial in the relative BV cohomology  $H^{1,4}(s \mid d)$ . We therefore separate three calculations:

1. classify the complete local obstruction space;
2. compute the coefficient vector in one specified analytic route;
3. decide whether that vector is a boundary in the chosen field theory.

The result has two theory labels which must not be suppressed:

| Theory                                            | One-loop local Euclidean disposition |
|---------------------------------------------------|--------------------------------------|
| strict pure Weyl gravity, fixed field content     | obstructed                           |
| formal tau-adic Wess–Zumino compensator extension | QME restored                         |

The second row does not reverse the first. It changes the BV complex.

All local results carry **LOCAL-ALGEBRAIC**; the coefficient computation also carries **EUCLIDEAN-SPECTRAL**. No result in this paper carries **LORENTZIAN-CAUSAL**.

## 2 Strict local BV obstruction space

The local jet algebra tracks ghost number, antifield number, form degree, engineering dimension, Grassmann parity, spacetime parity, and Weyl weight. Tensor symmetries, algebraic and differential Bianchi identities, graded commutativity, dummy-index relabelling, integration by parts, and total derivatives are reduced before exact sparse linear algebra is performed.

On the regular Bach-locus chart, the minimal Koszul–Tate sector contracts at positive antifield number. Ten nonminimal pairs contract explicitly, and an arbitrary invertible local BV-canonical gauge-fixing transformation preserves the quotient. The pure-diffeomorphism and mixed  $\text{Diff}$ –Weyl total complexes add no independent four-dimensional anomaly class.

**Theorem 2.1** (Strict dimension-four anomaly quotient). *In the declared complete gauge-fixed local BV complex,*

$$\dim H_{\text{even}}^{1,4}(s \mid d) = 2, \quad \dim H_{\text{odd}}^{1,4}(s \mid d) = 1.$$

*Representatives may be chosen as*

$$[\omega C^2], \quad [\omega E_4], \quad [\omega C\tilde{C}].$$

*The type-D carrier  $\omega \square R$  is exact with its stored  $R^2$  counterterm and horizontal current.*

Closure, exact primitives, complete basis manifests, and normalized dual nonmembership witnesses are stored independently. The universal diffeomorphism descent and intrinsic Weyl descent are recorded separately; in particular, a nonzero Diff-horizontal tower is not called a nontrivial intrinsic Weyl descent.

### 3 Coefficient-bearing one-loop breaking

The coefficient route combines two independent curvature carriers. A local Ricci-flat carrier exposes  $C^2$ , while round  $S^4$  fixes the Euler coordinate. The complete Euclidean symbol sequence has bundle dimensions  $5 \rightarrow 10 \rightarrow 5$ , exact image–kernel equality, a formal adjoint, and the required Diff/Weyl nonminimal doublets. Determinant multiplicities, the functional measure, conformal-Killing zero modes, the indefinite contour, and the parity Ward identity are included in the accepted integration slice.

**Theorem 3.1** (Repository one-loop Slavnov breaking). *In the normalization*

$$S_W = \alpha_C \int \sqrt{g} C^2,$$

*the repository-regulated breaking has coordinates*

$$(C^2, E_4, C\tilde{C}, \square R) = \left( \frac{199}{30}, -\frac{87}{20}, 0, 0 \right).$$

*After reduction against the complete strict quotient,*

$$\mathcal{A}^{(1)} = \frac{1}{(4\pi)^2} \left( \frac{199}{30} [\omega C^2] - \frac{87}{20} [\omega E_4] \right).$$

*Consequently the strict fixed-field-content local Euclidean QME is obstructed at one loop.*

The odd zero is derived from the parity Ward identity for the declared real tensor-Laplacian regulator. The type-D zero is scheme-dependent but exact; it cannot cancel either nontrivial coordinate. A beta function alone would not prove this theorem: the decisive object is the regulated Slavnov breaking reduced in the complete BV quotient.

**Proposition 3.2** (Standard unitary free matter cannot cancel the vector). *Let the added matter content be any nonnegative combination of real conformal scalars, Weyl or Dirac fermions, and standard gauge vectors. No such combination cancels both even coordinates of the strict breaking.*

The proof is an exact separating dual-cone witness. Interacting fixed points, wrong-sign or nonunitary matter, conformal higher spins, and compensator extensions lie outside this no-go.

## 4 The compensator cotangent lift

Adjoin a scalar with BRST transformation

$$Q\tau = \mathcal{L}_\xi \tau + \omega.$$

Its cotangent rows are not guessed. They are the Euler derivatives of the single minimal master term

$$S_{\tau,\min} = \int \tau^* (\mathcal{L}_\xi \tau + \omega).$$

In addition to the field row this forces

$$\delta\omega^* = 2g_{\mu\nu}g^{*\mu\nu} + \tau^*, \quad \delta\xi_\mu^* = N_{\xi,\mu} + \tau^* \nabla_\mu \tau, \quad \gamma\tau^* = \mathcal{L}_\xi \tau^*,$$

together with the opposite-adjoint Lie-prolonged rows.

**Theorem 4.1** (Exact minimal-BV cotangent lift). *On every strict-plus-compensator atom,*

$$\delta^2 = 0, \quad \delta\gamma + \gamma\delta = 0, \quad Q^2 = 0.$$

*The canonical change*

$$\hat{g} = e^{-2\tau} g, \quad \hat{g}^* = e^{2\tau} g^*, \quad \hat{\tau}^* = \tau^* + 2g_{\mu\nu}g^{*\mu\nu}$$

*preserves the canonical one-form and sends the Weyl Koszul–Tate row to  $\delta\omega^* = \hat{\tau}^*$ . Thus  $(\tau, \omega, \omega^*, \hat{\tau}^*)$  is an exact contractible quartet.*

The exponential change is an automorphism only after declaring the formal tau-adic local analytic completion. Its two inverse series are replayed with exact rational coefficients and obey at every order

$$\sum_{k=0}^n \frac{(-2)^k}{k!} \frac{2^{n-k}}{(n-k)!} = \frac{2^n}{n!} (1-1)^n = \delta_{n0}.$$

No finite-polynomial-in- $\tau$  theorem is inferred.

## 5 Extended cohomology and QME restoration

Quartet contraction reduces the extended complex to pure Diff BV cohomology in the dressed metric  $\hat{g}$ . The dimension-four counterterm inventory is therefore larger than in strict Weyl gravity:  $R(\hat{g})^2$  is now closed.

**Theorem 5.1** (Formal tau-adic extended quotient). *In the declared formal tau-adic local analytic jet algebra,*

$$H_{\text{ext,even}}^{0,4} = \text{span}\{C(\hat{g})^2, E_4(\hat{g}), R(\hat{g})^2\},$$

$$H_{\text{ext,odd}}^{0,4} = \text{span}\{C(\hat{g})\tilde{C}(\hat{g})\}, \quad H_{\text{ext}}^{1,4} = 0.$$

$\square R(\hat{g})$  remains horizontally exact. There is no incoming dimension-four positive-antifield boundary: the only dimension-zero natural symmetric tensor is  $\hat{g}$ , whose image is the vanishing trace of the Bach row.

Explicit primitives for the strict four-candidate anomaly carrier are

$$B_C = \int \sqrt{g} \tau C^2, \quad B_P = \int \sqrt{g} \tau C \tilde{C},$$

together with the Euler–Wess–Zumino functional  $B_E$  and the stored  $R^2$  primitive  $B_\square$ . Their boundary matrix has exact rank four.

**Corollary 5.2** (One-loop local Euclidean QME restoration). *The counterterm*

$$-\frac{\hbar}{(4\pi)^2} \left( \frac{199}{30} B_C - \frac{87}{20} B_E \right)$$

*satisfies*

$$\mathcal{A}^{(1)} + s_{\text{ext}} S_{\text{ct}}^{(1)} = 0 \pmod{\mathfrak{d}}.$$

*Hence the compensator-extended local Euclidean QME is restored at one loop in the declared formal tau-adic algebra.*

## 6 What does not follow

The strict obstruction and extended restoration are local one-loop results. They do not establish:

- an all-loop extended QME;
- a Lorentzian renormalized time-ordered product or Slavnov identity;
- a BRST-compatible global Hadamard state;
- positivity, a particle Hilbert space, scattering, or unitarity;
- survival or removal of either residual deformation class;
- a quantum Cartan identity or residual moment-map algebra.

Residual transfer is specifically forbidden at present. The frozen classical contraction belongs to the strict theory and has no compensator rows. The next gate is a compensator-inclusive classical contraction and the coefficient-bearing one-loop Slavnov operator  $Q_1$ . Bridge 4 also remains inactive until a normalized same-background classical mode, causal Green carrier, and pairing coexist. Bridge 5 has a QME disposition but lacks the independent interaction-to-physical-branch map. No classical tangent obstruction is therefore called ghost removal or a quantum constraint.

## 7 Machine certification and reproduction

The two theorem receipts are

- `quantum-weyl/anomalies/certificates/REGULATED_REPOSITORY_BV_SLAVNOV_BREAKING.json`;
- `quantum-weyl/anomalies/certificates/WESS_ZUMINO_EXTENDED_LOCAL_BV_COHOMOLOGY.json`.

The cotangent-lift and matter-cone receipts are separate dependencies. The companion claim map records their content hashes and all explicit nonclaims.

The scoped replay is:

```
PYTHONPATH=quantum-weyl python3 -m \
    anomalies.verify_wess_zumino_minimal_bv_cotangent_lift
PYTHONPATH=quantum-weyl python3 -m \
    anomalies.verify_wess_zumino_extended_local_bv_cohomology
PYTHONPATH=quantum-weyl python3 -m verify_active_frontier
PYTHONPATH=quantum-weyl python3 -m atlas.verify_quantum_atlas_fragment
python3 paper/verify_12_pure_weyl_one_loop_bv_anomaly_claim_map.py
```

## 8 Outlook

The immediate local-to-residual question is now sharp. Given an extended contraction  $(\iota_{\text{ext}}, \pi_{\text{ext}}, S_{\text{ext}})$  and the restored one-loop Slavnov operator  $Q_1$ , compute

$$q_1 = \pi_{\text{ext}} Q_1 \iota_{\text{ext}}$$

and test first-order nilpotency, the quantum Cartan defect, incoming  $H^3 \rightarrow H^4$ , outgoing  $H^4 \rightarrow H^5$ , operator mixing, and the corrected pairing. This is a new theorem, not a corollary of local QME restoration.

The independent Lorentzian rail must construct renormalized causal products and a compatible state before any unitary or particle conclusion is possible. The present result nevertheless fixes the finite local obstruction vector that every such construction must reproduce or trivialize.

## References

- [1] I. A. Batalin and G. A. Vilkovisky, *Gauge algebra and quantization*, Phys. Lett. B **102** (1981), 27–31.
- [2] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*, Princeton University Press, 1992.
- [3] N. Boulanger, *General solutions of the Wess–Zumino consistency condition for the Weyl anomalies*, arXiv:0704.2472.
- [4] R. J. Riegert, *A nonlocal action for the trace anomaly*, Phys. Lett. B **134** (1984), 56–60.