

What Survives the Ghost Test?

A Four-Level Classification in Pure-Weyl Gravity

GPT-5.6.sol*

Claude Fable 5†

Phase-1 synthesis with scoped Phase-2 update, 22 July 2026

Abstract

Fourth-order gravity predicts more gravitational-wave solutions than general relativity. Some appear with the wrong sign and are therefore called ghosts, but a wrong sign in an equation is a warning rather than a complete physical verdict. We ask which extra solutions remain after gauge symmetry, global constraints, nonlinear consistency, boundary conditions, and quantum consistency are imposed. Some survive early reductions; different later tests exclude different candidates.

Pure Weyl gravity is a fourth-order conformal metric theory. We separate four logically different objects: local solutions; gauge-, charge-, and boundary-reduced classical directions; nonlinearly continuable directions; and quantum states or particles.

On compact Weyl–Maxwell laboratories, additional axial and polar directions survive local gauge reduction and have nonzero induced Lee–Wald pairings. The signature of the associated positive-frequency Hermitian current form is not organized by the simple rule “Einstein positive, additional negative.” A Nariai factorization identifies a related mechanism at the operator level: the two spin-two factors have opposite Lee–Wald residues, although no absolute energy sign or identification with the compact-product family is thereby established. A dyonic-flat-family preflight, by contrast, finds no fixed-coupling stationary sign laboratory because the required global Hamiltonian lift and fixed-Chern tangent conditions fail. At second order, finite-harmonic exponential-polynomial continuation is equivalent to five quadratic stabilizer-charge conditions, whereas bounded continuation obeys further polynomial-growth and characteristic-shell resonance constraints. On Schwarzschild, horizon regularity alone admits additional curvature solutions. The Phase-2 generic-angular calculation overturns the former fixture-level Einstein-selection reading: an axial non-Einstein branch has a finite formal radial pairing for every integer $\ell \geq 2$ and real $\omega \neq 0$, while the polar calculation retains a one-dimensional mixed finite line and an exact exceptional wall. These are formal-infinity, fixed-current-representative results, not a completed asymptotically flat phase space. In the quantum analysis, strict fixed-field-content pure Weyl gravity has a nonzero local Euclidean one-loop BV anomaly. A formal compensator makes the anomaly exact, but changes the theory and does not supply a Lorentzian state space.

An exact PT/CPT pilot sharpens, but does not reverse, this classification. Selected compact reduced blocks admit structured positive pseudo-Hermitian metrics, yet the available data do not define an independent parity/time involution and hence do not construct a genuine Mannheim C operator. On the cylinder, the natural involution fails the BRST chain-map test. The counterflow Hamiltonian–Hopf quartet has complex spectrum and therefore admits no positive pseudo-Hermitian metric. For one complete causal Berger BV complex the local Hadamard singular structure and causal commutator are available, but the full covariance remains unconstructed because an explicit 104-dimensional stationary carrier is missing.

*OpenAI model and principal programme author.

†Anthropic model and computational coauthor. The project was commissioned and coordinated by Asger Alstrup Palm (asger@area9.dk), who initiated the questions, exercised editorial control, and serves as corresponding human contact.

We also analyze a changed two-phase counterflow gravity-clock theory. It has a complete free causal BV complex at a specified Berger background and removes a dressed-trace obstruction. Charge reduction reveals an exact tradeoff: making time translation null removes the relative clock, while retaining the clock makes time translation charged. More decisively, the complete $j = \frac{1}{2}$ reduced block contains a Hamiltonian–Hopf quartet throughout the connected trace-healthy same-field stationary family. Phase 1 therefore ends with a classification rather than a selected theory: pole-level signs are too coarse, but the more complete reductions do not generically rescue the candidates tested here. No positive full-BV state, scattering theory, or unitarity theorem is claimed.

Contents

1	The question and the four levels	2
2	Theories, conventions, and scope	3
2.1	The scope tuple	4
3	What “ghost” can mean	6
4	Which additional linear directions survive reduction?	7
5	Which directions survive nonlinear integrability?	9
6	Which boundary conditions retain the extra solutions?	10
7	Can a positive quantum state be constructed?	11
8	What the counterflow candidate taught us	13
9	Integrated Phase-1 verdict	16
10	Relation to prior work	16
11	Limitations and the next scientific boundary	18
12	Computational accountability	18
13	Conclusion	19

1 The question and the four levels

The metric action of strict pure Weyl gravity is

$$S_W[g] = \alpha_W \int_M \sqrt{-g} C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} d^4x, \quad B_{\mu\nu} = 0, \quad (1)$$

where $B_{\mu\nu}$ is the Bach tensor. Around a fixed background, the linearized Bach equation is fourth order. A formal factorization, a partial fraction decomposition, or a negative residue can reveal an indefinite direction. None of these alone says whether the direction survives gauge reduction, global constraints, physical boundary conditions, nonlinear integrability, or the construction of a quantum state.

This paper organizes the first phase of a computer-assisted research programme by the following hierarchy:

Level 1: local solutions of the linear field equations,	(2)
Level 2: reduced classical directions after gauge, charge, and boundary conditions,	
Level 3: directions tangent to nonlinear solutions in a stated correction class,	
Level 4: quantum states, particles, and a positive physical probability interpretation.	

The arrows between these levels are mathematical problems. They are not terminological promotions. A nonzero classical symplectic pairing at Level 2 is not a positive quantum norm at Level 4. A secular second-order primitive at Level 3 is not nonlinear stability. A local Euclidean anomaly calculation is not a Lorentzian scattering theorem.

The phase-one question is therefore not simply “does the propagator contain a ghost?” It is:

Which additional fourth-order directions survive each declared reduction, which fail nonlinear or boundary tests, and what remains undecided before a particle interpretation can even be posed?

The answer is mixed. Some additional directions survive local gauge and presymplectic reduction. Compact nonlinear constraints remove some of them, while balanced combinations can continue formally with secular terms. A generic-angular Schwarzschild calculation finds finite formal additional or mixed radial directions rather than a parity-uniform selection theorem. A structured positive pseudo-Hermitian metric exists on selected finite reduced blocks, but no genuine BRST-compatible C operator or full-BV positive state has been constructed. The strict quantum gauge theory is locally anomalous. Finally, a changed theory that passes a difficult causal construction fails a physical spectral test that no choice of positive inner product can repair. Different candidates fail at different levels.

2 Theories, conventions, and scope

We use Lorentzian signature $(-, +, +, +)$ and $[\nabla_\mu, \nabla_\nu]v^\rho = R^\rho_{\sigma\mu\nu}v^\sigma$. Classical fields are real; complex Fourier modes are bookkeeping devices and physical pairings use the corresponding real or conjugate-frequency form. The connected local gauge group of the strict theory is

$$\text{Diff}_0(M) \ltimes C^\infty(M, \mathbb{R})_{\text{Weyl}}, \quad g_{\mu\nu} \mapsto e^{2\sigma} g_{\mu\nu}. \quad (3)$$

Unless a theorem card says otherwise, no boundary counterterm or topological density is included in the action. Boundary and support conditions are part of the theorem, not implicit conventions.

Two comparison theories appear. On the compact product we use

$$S_{\text{WM}}[g, A] = S_{\text{W}}[g] - \frac{1}{4} \int_M \sqrt{-g} F_{\mu\nu} F^{\mu\nu} d^4x, \quad (4)$$

$$S_{\text{EM}}[g, A] = \frac{1}{2\kappa} \int_M \sqrt{-g} (R - 2\Lambda) d^4x - \frac{1}{4} \int_M \sqrt{-g} F_{\mu\nu} F^{\mu\nu} d^4x. \quad (5)$$

The magnetic bundle is fixed in that comparison. Einstein–Maxwell is a source theory and comparator; it is not identified with a subsector carrying the same symplectic structure.

The quantum repair uses a formal Wess–Zumino field τ with $s\tau = \omega$, where ω is the Weyl ghost, and the dressed metric $\hat{g} = e^{-2\tau}g$. This is an enlarged BV theory. It is not strict pure Weyl gravity in another gauge.

The later counterflow experiment changes the classical action as well. Its defining matter mechanism is two positive phase fields with an auxiliary diagonal $U(1)$ connection,

$$S_{\text{phase}} = -\frac{1}{2} \int_M \sqrt{-\widehat{g}} [f_1^2 (d\theta_1 - A)^2 + f_2^2 (d\theta_2 - A)^2]. \quad (6)$$

Writing

$$\chi = \frac{f_1^2 \theta_1 + f_2^2 \theta_2}{f_1^2 + f_2^2}, \quad \psi = \theta_1 - \theta_2, \quad \mu^2 = \frac{f_1^2 f_2^2}{f_1^2 + f_2^2}, \quad (7)$$

separates a diagonal gauge phase from a gauge-invariant relative phase with positive bare kinetic coefficient μ^2 . Fixing the harmless positive diagonal normalization to $f_1^2 + f_2^2 = 4$, the complete counterflow action used below is

$$S_{\text{cf}} = \int_M \sqrt{-\widehat{g}} \left[\frac{\alpha_B}{8} \widehat{C}_{\mu\nu\rho\sigma} \widehat{C}^{\mu\nu\rho\sigma} + \frac{M_2}{2} \widehat{R} - V_0 - \frac{\mu^2}{2} (\widehat{\nabla}\psi)^2 - 2(A - d\chi)^2 \right] d^4x. \quad (8)$$

There is no Maxwell kinetic term for A . Besides diffeomorphisms, the local gauge transformations are

$$g \mapsto e^{2\sigma} g, \quad \tau \mapsto \tau + \sigma, \quad \chi \mapsto \chi + \lambda, \quad A \mapsto A + d\lambda, \quad (9)$$

so that $\widehat{g}$ and ψ are gauge invariant. Constant shifts of ψ generate the physical relative symmetry R_{rel} rather than a local gauge symmetry. Equation (8), its BV cotangent lift, and ordinary nonminimal gauge-fixing doublets define the changed theory; a component Hessian is evidence for this action, not its definition. Results for the changed theory are never imported back into the strict theory.

2.1 The scope tuple

The mathematical scope of every result is typed by

$$\boxed{\mathfrak{S} = (\mathcal{T}, M, \bar{\Phi}, \mathcal{G}, \mathcal{X}, \mathcal{B})}, \quad (10)$$

where the entries are the theory, spacetime, background fields, gauge group, function or phase space, and boundary/support class. Evidence is recorded separately as

$$\mathfrak{E} = (\text{claim identifier}, \text{dependency tags}, \text{evidence state}, \text{limitation}). \quad (11)$$

Thus an epistemic label is never treated as part of the mathematical object. Two statements with different entries in $\mathfrak{S}$ are not combined without a constructed map. This matters because the programme uses several controlled laboratories:

Laboratory	Question isolated there
$\mathbb{R} \times S^3$ conformal cylinder	Complete free causal complex, a selected residual zero-charge reduction, and a PT/CPT chain-map test
Biaxial Berger spacetime	Clocks, relational observables, interactions, and the counterflow successor
$\mathbb{R} \times S^1 \times S^2$ magnetic product	Einstein/additional branches, Lee–Wald pairing, Taub constraints, resonances, and a finite-block pseudo-Hermitian pilot
Nariai	Causal transfer and a scoped factor-residue sign mechanism
Schwarzschild and static spherical metrics	Horizons, formal radial-pairing filtrations, and black-hole perturbations
Euclidean S^4 and regular local charts	One-loop determinants and local BV anomaly classes

These are related tests of a common architecture, not successive approximations to one already unified universe.

3 What “ghost” can mean

The literature uses the word *ghost* for inequivalent failures. The distinctions are especially important in a gauge theory with a fourth-order equation. A modern review of the nondegenerate higher-derivative Ostrogradsky theorem is Ref. [4]; gauge degeneracy and physical reduction must be analyzed separately.

Diagnosis	Mathematical question	Phase-1 disposition
Ostrogradsky direction	Is a nondegenerate higher-derivative Hamiltonian unbounded before constraints?	Crosswalks are computed on selected blocks; this is not used as a complete physical quotient.
Negative pole residue	Does a gauge-fixed propagator decompose with a negative coefficient?	Under the standard adjoint it is an initial warning, not a final state classification. A PT/CPT prescription changes the physical-adjoint question and must pass its own structural gates.
Negative classical energy or flux	What is the sign of a declared canonical-energy form, flux, or positive-frequency Hermitian current derived from the real Lee–Wald form?	Computed for selected compact and black-hole sectors. Some signs contradict a naive Einstein/additional split.
Radical or pure gauge	Does the induced presymplectic form vanish on the direction?	Additional compact-product directions are nonradical on the declared pre-residual quotient.
Nonlinear obstruction	Is a linear direction tangent to a second-order solution in a stated function class?	Classified completely on a finite harmonic exponential-polynomial space; bounded continuation is stricter.
Negative quantum norm	Is there a BRST-compatible positive state on the full physical algebra?	Not established for the standard adjoint. Under a PT/CPT prescription, positive η alone is insufficient: genuine C , BRST descent, and causal covariance remain unconstructed.
Failure of unitarity	Does an asymptotic state space satisfy the optical theorem?	Not tested: no complete scattering or particle theory exists.
Alternative pole prescription	Does a structured positive metric, a genuine PT/CPT involution, and BRST descent exist in the declared representation?	Tested on selected reduced blocks. Positive pseudo-Hermitian metrics exist in a compact pilot, but no genuine Mannheim C operator or full-BV descent is established; the counterflow quartet is spectrally excluded.

PT-symmetric treatments of the Pais–Uhlenbeck system, Dirac–Pauli formulations of quadratic gravity, and fakeon prescriptions all demonstrate that “negative residue implies negative-probability asymptotic particle” is not the only proposed logical route [8, 14, 15]. They also demonstrate why the chosen state space and prescription must be declared. The scoped Phase-2 pilot therefore separates four gates that are often conflated: existence of a positive metric η , existence of an independent parity/time involution and corresponding C , compatibility with the BRST differential, and compatibility with a causal covariance. Passing the first gate on a finite block does not imply the other three. Fakeon and Dirac–Pauli prescriptions remain neighboring programmes rather than consequences of these reductions. The perturbative renormalizability and extra massive-spin-two spectrum that made this question central to higher-derivative gravity go back to Stelle’s foundational analyses[5, 6].

4 Which additional linear directions survive reduction?

The compact magnetic product is the cleanest place to compare the Einstein–Maxwell and Weyl–Maxwell linear solution spaces. The computation uses both parities, generic harmonics, exceptional modes, and global blocks.

The sign-bearing object must be distinguished from the underlying presymplectic form. Let Ω_{WM} be the real, antisymmetric Lee–Wald form obtained from the Weyl–Maxwell action [1, 2]. Rank and radical are defined on the real reduced solution space; Ω_{WM} itself has no inertia. After complexifying, fix a common stationary shell

$$e^{-i\omega t}, \quad \omega > 0, \quad k = \frac{2\pi n}{L}, \quad \lambda = \ell(\ell + 1), \quad \ell \geq 2, \quad (12)$$

and impose the conjugation reality condition between positive- and negative-frequency coefficients. Here L is the circumference of S^1 and $N_{\ell m} = \int_{S^2} |Y_{\ell m}|^2 d\Omega > 0$ is the chosen spherical-harmonic normalization. On the positive-frequency space define

$$h_+(u, v) = \frac{\Omega_{\text{WM}}(\bar{u}, v)}{-i\omega L N_{\ell m}}, \quad L N_{\ell m} > 0. \quad (13)$$

Reality and antisymmetry of Ω_{WM} make h_+ Hermitian. A positive change of the circle or harmonic normalization, and a nonzero complex rescaling of any basis vector, act by Hermitian congruence and leave its inertia invariant. The signatures below refer only to this h_+ . Exceptional dipoles, generalized zero modes, and $\omega = 0$ are classified in separate real blocks and are not assigned these frequency-shell signatures.

Result card A: compact Einstein/additional decomposition

Theory. Einstein–Maxwell source and Weyl–Maxwell target.

Background. The magnetically supported compactified Plebanski–Hacyan product $\mathbb{R} \times S^1 \times S^2$, in a fixed magnetic bundle.

Domain. Parity-complete generic linear modes, together with the declared exceptional and global blocks.

Gauge and quotient. Local gauge quotient and maximal pre-residual H^0 exact sequence; final stabilizer reduction is not included.

Boundary condition. Spatial periodicity; no asymptotic scattering condition.

Statement. The Einstein–Maxwell solutions inject into the Weyl–Maxwell solutions. The cofiber contains additional axial and polar blocks. On each generic additional real block Ω_{WM} has zero radical and rank four. On the associated positive-frequency complex two-plane, h_+ has inertia

(2, 0); its restriction to the target Einstein image has inertia (1, 1) and the complete generic target has inertia (3, 1). The independently normalized Einstein–Maxwell source Hermitian current form has inertia (2, 0), so inclusion does not define a cyclic or symplectic equivalence.

Not claimed. No positive energy, particle norm, final residual quotient, causal boundary admissibility, or nonlinear closure follows.

The important conclusion is not that the additional block is “healthy” because the displayed frequency form is positive. The conclusion is that it is not an algebraic duplicate or a radical direction on this quotient, and that the standard pairing does not respect the naive expectation that the Einstein image should inherit the ordinary Einstein sign while the complement should carry the bad sign. The action matters: the Einstein–Hilbert and Weyl Lee–Wald forms are different even on the same metric perturbation. Neither the rank statement nor the inertia of h_+ is by itself a positive-energy or positive-norm quantum theorem. The inertia statements are also relative to the declared conjugation and frequency shell that define h_+ in Eq. (13); a different admissible adjoint would organize the same real symplectic data differently. The failure of the naive split is therefore adjoint-dependent evidence that motivates, but does not prove, alternative inner-product prescriptions of the PT/CPT type[8].

The Phase-2 finite-block pilot makes this last qualification exact. On the selected compact physical blocks, the intertwining equation

$$H^\dagger \eta = \eta H, \quad \eta = \eta^\dagger > 0, \quad (14)$$

has a nonempty symmetry-compatible structured cone. Its distinguished fundamental symmetry is $C_0 = -I_2 \oplus I_2 \oplus I_4$. This is a positive pseudo-Hermitian-metric result, not yet a Mannheim construction: the reduced data do not supply independent operators P and T from which a physical C can be defined, and quotient descent has not been established. On the selected conformal-cylinder carrier the natural candidate C_0 is not a chain map for the BRST differential, so the declared invariant commutant does not yield a BRST-descended positive metric. These conclusions carry the dependency tags **LOCAL-ALGEBRAIC** and **REDUCED-MODE**; they do not establish a Lorentzian state.

A second Phase-2 calculation isolates an operator mechanism on Nariai. The spin-two operator factors as $L_E L_C$ with $L_C = L_E - 2/3$, and the two factor projectors have Lee–Wald residues $-1/3$ and $+1/3$. This predicts a relative sign exchange between the two factors and is robust when both static-patch horizons are retained. It does not fix an absolute energy sign, inertia, or positivity, because Nariai has no declared global timelike generator for that purpose; nor does it identify the factors with the compact magnetic-product Einstein image and additional cofiber. It is a mechanism-level comparison across different scope tuples, not a family theorem.

The attempted dyonic-flat-family extension fails earlier. For nonzero electric parameter the stationary symmetry has no continuous global Hamiltonian lift on the circle, since $\mathcal{L}_H A = E dx$ changes the holonomy. Electromagnetic duality also fails to define a tangent direction in a fixed-Chern sector. Consequently that preflight constructs neither a fixed-coupling open stationary family nor a sign, current, inertia, or signature-wall theorem. Recording this obstruction prevents a formally attractive but ill-typed background scan from being read as evidence for Result card A.

This exact-sequence formulation also prevents an unjustified direct-sum claim. A lift of an additional curvature representative may be shifted by an Einstein solution. The invariant information is the injection, cofiber, induced ranks, radicals, and extension class. A canonical branch split requires more structure than composition of differential operators.

On the conformal cylinder a different reduction occurs. In a selected derived zero-charge sector, all fifteen residual conformal generators are implemented by a BFV/Koszul receiver, and the vacuum and selected residual one-particle cohomologies are acyclic. The first surviving H^4 classes are

deformation or vertex classes, not one-particle gravitons. This is a theorem about that selected compact sector. It is not a universal removal of radiative gravitons and it supplies no asymptotic particle interpretation.

The two laboratories therefore teach complementary lessons: additional directions can survive a compact local-gauge quotient, while a stronger zero-charge residual reduction can remove selected one-particle cohomology. Neither statement transfers to the other without a map.

5 Which directions survive nonlinear integrability?

Linearization stability on compact Cauchy surfaces has a classical history from Brill–Deser through Fischer–Marsden, Moncrief, and the Arms–Marsden–Moncrief momentum-map normal form [19, 20, 21, 22]. The general idea that a symmetric solution has a quadratic momentum-map cone is therefore not new. The contribution here is an exhaustive model-specific calculation for a fourth-order Weyl–Maxwell mode space and a separation of formal secular solvability from bounded solvability.

Let L be the linearized equation and $D^2E[u, u]$ the quadratic source. Two correction spaces are kept distinct:

$$\mathcal{X}_{\text{ep}} = \left\{ \text{finite sums of } t^p e^{-i\Omega t} \text{ with spatially periodic finite harmonic support} \right\}, \quad (15)$$

$$\mathcal{X}_{\text{qp}} = \left\{ \text{finite bounded quasiperiodic sums } e^{-i\Omega t} \right\}. \quad (16)$$

The first allows secular polynomials; the second does not.

Result card B: finite-harmonic second-order cone

Theory. Compact Weyl–Maxwell theory, with the Einstein–Maxwell comparison used to label branches.

Background. $\mathbb{R} \times S^1 \times S^2$ with the fixed magnetic bundle.

Domain. The complete declared finite harmonic space: generic Einstein and additional primaries, both parities, exceptional dipoles, twists, homogeneous generalized modes, electric flux, and Wilson directions.

Obstruction covectors. The five stabilizer covectors generated by time translation H , circle translation P_x , and rotations J_1, J_2, J_3 .

Correction class. Finite exponential-polynomial corrections $\mathcal{X}_{\text{ep}}$.

Statement.

$$\mathcal{Z}_2^{\text{ep}} = \{u : \mu_H(u) = \mu_{P_x}(u) = \mu_{J_1}(u) = \mu_{J_2}(u) = \mu_{J_3}(u) = 0\}.$$

The five moment maps are necessary and sufficient in this correction class.

Not claimed. This is not a theorem for all smooth or Sobolev fields, retarded solutions, bounded dynamics, all-order integration, or stability.

The sufficiency statement follows from blockwise constant-coefficient surjectivity: nonzero invariant factors are surjective on finite exponential-polynomial modules, including at characteristic roots after the appropriate polynomial degree is allowed. The only zero invariant factors are the five stabilizer covectors. This is the explicit exhaustive content beyond the general momentum-map normal form.

Bounded continuation has a larger obstruction ledger. If the quadratic source is expanded by frequency shell and polynomial degree, then a bounded primitive exists only when three types of functional vanish:

$$\{\mu_X\} \cup \{\mathcal{P}_{j,r}\} \cup \{\mathcal{R}_{j,a}\}. \quad (17)$$

Here μ_X are the global zero-block Taub constraints, $\mathcal{P}_{j,r}$ are coefficients that would require positive powers of t , and $\mathcal{R}_{j,a}$ are adjoint-kernel pairings on nonzero characteristic shells. Their common nonlinear zero locus is not fully classified for arbitrary finite support. It is classified on several important subspaces. In particular, on the complete global plus ($\ell = 2, k = 0$) additional-primary space, the bounded cone reduces to the standard global generalized-zero cone: no nonzero additional-primary amplitude survives. At other momenta, balanced sheets can remain.

The distinction is physical. A resonant source can have a formal primitive $te^{-i\omega t}$ and hence define a second-order jet, while failing every bounded quasiperiodic test. The programme therefore does not report “second-order solvable” without naming the correction class.

The same calculation also blocks a common branchwise simplification. Balanced Einstein–additional configurations can have zero total moment map while their projected Einstein and additional pieces carry equal and opposite nonzero charges. Consequently the ambient exact sequence does not descend to separately neutral branch fibers. Nonlinear reduction is intrinsically mixed.

6 Which boundary conditions retain the extra solutions?

Compact charge constraints and black-hole endpoint conditions test different questions. The Schwarzschild analysis uses the Ricci perturbation $\psi_{\mu\nu} = \delta R_{\mu\nu}[h]$ as a second-order curvature variable in the fourth-order Bach system. On a Ricci-flat background the universal linear identity is

$$\delta B_{\mu\nu} = \frac{1}{2}\square\delta R_{\mu\nu} + C_{\mu\rho\nu\sigma}\delta R^{\rho\sigma} - \frac{1}{6}\nabla_\mu\nabla_\nu\delta R - \frac{1}{12}g_{\mu\nu}\square\delta R. \quad (18)$$

The Einstein image lies in $\ker \delta R$; the additional curvature content is visible in ψ .

Horizon analyticity does not force ψ to vanish. Regular ingoing additional curvature solutions exist in the declared real-frequency axial analysis. This already shows that future-horizon regularity alone does not select the Einstein image. It does not show that every local differential initial or boundary condition fails: imposing ψ and its normal derivative to vanish on a Cauchy surface is a local restriction that selects the Einstein image at linear order.

The outer boundary instead gives a filtration-sensitive Phase-2 result. In ingoing Eddington–Finkelstein coordinates $(v, r, \vartheta, \varphi)$ let $F^v[u_1, u_2](r)$ denote the sphere integral of the v -component of the action-derived Lee–Wald current for two formal mode representatives. For a conjugate-frequency pair define the truncated radial slice form

$$\mathcal{N}_R[u_1, u_2] = i \int_2^R F^v[\bar{u}_1, u_2](r) dr, \quad \mathcal{N}_\infty[u_1, u_2] = \lim_{R \rightarrow \infty} \mathcal{N}_R[u_1, u_2], \quad (19)$$

whenever the improper integral exists in the declared formal asymptotic class. “Finite radial slice form” below means finiteness of this quadratic form. It is neither an L^2 norm on monochromatic scattering states nor a wave-packet Hilbert norm, and the polar zero-sector quadratic form can vanish.

The dangerous-layer rule in this class is that a term $r^p(\log r)^q$ is nonintegrable for $p \geq -1$ unless its invariant coefficient vanishes. Correctly retaining differentiated reconstruction forcing changes the old fixture-level conclusion.

Result card C: generic-angular formal radial disposition on Schwarzschild

Theory. Strict pure C^2 gravity.

Background. Schwarzschild with $M > 0$ in ingoing Eddington–Finkelstein coordinates.

Domain. Integer $\ell \geq 2$, real $\omega \neq 0$, formal polyhomogeneous infinity data, and the fixed action-derived pure-Weyl Lee–Wald representative.

Gauge and quotient. Regge–Wheeler/Zerilli-type parity reduction and metric reconstruction in the certified formal radial modules.

Statement. In the axial zero-rate sector, both Einstein directions have finite r^{-2} current and a reconstructed non-Einstein all-orders formal lift X_0 has finite $X_0|X_0$ pairing and nonzero finite $E_0|X_0$ pairing. This remains finite under $X_0 \mapsto X_0 + \beta E_0$, so Einstein-only formal radial selection is false throughout the declared domain. In the polar oscillatory block, the leading $p = 0$ coefficient matrix of the ordered current on (E, X_0, X_1, X_2) has generic rank three. Its one-dimensional filtered radical survives through the $p = 0, -1$ layers as a mixed Einstein/additional line and has a generically nonzero first finite $p = -2$ induced pairing. Its exact exceptional wall is

$$Q_{21}(\ell(\ell + 1), \omega^2) = 0,$$

where Q_{21} is the certified bidegree-(21, 21) polynomial. Exact Sturm counts give respectively 0, 6, 2, 6, 2 real nonzero zeros of this induced $p = -2$ coefficient for $\ell = 2$, $\ell = 3$, $4 \leq \ell \leq 10$, $11 \leq \ell \leq 40$, and $\ell \geq 41$; these are not physical scattering thresholds.

Not claimed. The axial X_2 branch, terminal-only polar prefixes, the deeper filtration on the Q_{21} wall, all-order polar solutions, and horizon-to-infinity matching remain open. No asymptotically flat phase space, Hilbert norm, scattering theory, quasinormal spectrum, ringdown, stability, particle, positivity, or quantum theorem follows.

The old $\ell = 2$, $M\omega = 3/5$ divergence table omitted differentiated c' forcing in the axial reconstruction and is superseded. That point is not on the polar Q_{21} wall, but this does not restore its earlier Einstein-selection interpretation. Existing AdS/Euclidean selection arguments operate in different boundary classes [23, 24]. A physical asymptotically flat theorem still requires a completed Bach phase space, horizon-to-infinity matching, wave-packet flux topology, and a well-posed exterior problem. Covariant phase-space and canonical-energy methods provide the conceptual comparison, but their global hypotheses are not asserted here [2, 3].

7 Can a positive quantum state be constructed?

There are three separate quantum objects in the programme: local BV anomaly cohomology, Euclidean one-loop coefficients, and reduced Lorentzian two-point functions. They are not interchangeable.

In four dimensions a local Weyl breaking at ghost number one and form degree four has essential type-B and type-A representatives of the form

$$\mathcal{A}^{(1)} = \frac{1}{(4\pi)^2} \int_M \sqrt{g} \omega (c_W C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} - a_W E_4 + b_W \square R) d^4x. \quad (20)$$

In the action normalization of Eq. (1) and the repository's $\square R = 0$ finite-counterterm convention, the computed coordinates are

$$\boxed{c_W = \frac{199}{30}, \quad a_W = \frac{87}{20}, \quad b_W = 0.} \quad (21)$$

The $\square R$ coefficient is scheme dependent and may be shifted by a local counterterm. The nonzero C^2 and Euler representatives are the essential type-B and type-A classes, respectively, in the Deser–Schwimmer classification[35]. A one-loop divergence, beta function, trace anomaly, and BV

master-equation breaking are related only after the gauge-fixed determinant, normalization, descent, and quotient maps have been supplied.

The complete local determinant and heat-kernel ledger is:

sector	operator	statistics	Z exponent	c_i	a_i
metric TT, upper	$\Delta_{2,\perp}(4)$	bosonic	$-1/2$	$37/24$	$31/72$
Diff–Weyl scalar ghost	$\Delta_0(-4)'$	fermionic	$+1/2$	$89/120$	$269/360$
metric TT, lower	$\Delta_{2,\perp}(2)$	bosonic	$-1/2$	$29/8$	$181/72$
transverse Diff ghost	$\Delta_{1,\perp}(-3)'$	fermionic	$+1/2$	$29/40$	$79/120$
exact sum				$199/30$	$87/20$

The listed c_i and a_i are net signed contributions: the statistics sign and the displayed determinant exponent have already been applied. They are therefore not the unsigned heat-kernel coefficients of the individual operators. All four factors are second order. Their bundle ranks are 5, 1, 5, 3; the primes delete five scalar and ten transverse conformal-Killing ghost zero modes. York/Hodge Jacobians and the nonminimal doublets are already included. The table lists a_i with the sign convention $c_i C^2 - a_i E_4$.

The map from this Euclidean spectral ledger to a gauge anomaly is a residue and cohomology map, not a comparison of names. If $I_i(d)$ are the dimensionally continued integrated carriers, then

$$s_d I_i(d) = (d-4)A_i^4 + s_d B_i(d) + d C_i(d) + O((d-4)^2). \quad (22)$$

Multiplication by the one-loop pole, replacement of the Weyl parameter by the ghost ω , addition of the diffeomorphism completion, and reduction modulo s and d give

$$\Gamma_{\text{div}}^{(1)}(d) \xrightarrow{s_d, \text{Res}_{d=4}} \mathcal{W}_\sigma^{(1)} \xrightarrow{\sigma \mapsto \omega + \text{Diff}} Z^{1,4}(s \mid d) \longrightarrow H^{1,4}(s \mid d). \quad (23)$$

The nonzero coordinates in Eq. (21) survive the last quotient. This is what promotes the effective-action Weyl anomaly to an obstruction to treating Weyl symmetry as an exact quantum gauge redundancy in the strict fixed-field-content theory. These coordinates are stable under similarity transformations within the declared local operator and regulator class. They are not thereby invariant under a genuinely different complex contour or measure prescription: such a redefinition changes the quantum theory and requires its own regulator and quantum-action-principle analysis before any coefficient comparison.

Result card D: strict and compensated local quantum theories

Strict theory. Fixed-field-content Diff $\ltimes$ Weyl pure C^2 gravity.

Domain. Complete declared gauge-fixed local BV quotient on a regular Bach-locus chart, with Euclidean one-loop coefficient input.

Statement. The regulated breaking has $c_W = 199/30$, $a_W = 87/20$, and vanishing parity-odd and chosen type-D coordinates. Its essential ghost-number-one, form-degree-four Weyl components are nonzero. The strict local Euclidean one-loop QME is obstructed.

Extended theory. In the formal τ -adic Wess–Zumino BV algebra with $s\tau = \omega$, the same cocycle is exact and a local counterterm restores the Euclidean QME at one loop.

Not claimed. The extension changes the BV theory. It gives no unconditional all-loop theorem, regulator/QAP for an arbitrary changed action, Lorentzian QME, Hadamard state, positive physical Hilbert space, particles, scattering, or unitarity.

The compensator statement is an exact BV realization of the familiar dilaton/Wess–Zumino mechanism, whose broader structure is known [38]. Its value is not a claim that a compensator is a

free repair. Indeed, the passive compensator leaves a gauge-invariant dressed trace in the classical causal complex. Repairing that obstruction led to the changed counterflow experiment in Section 8.

Separately, Phase 2 audits the Hadamard gate on one complete causal 54-row Berger BV complex. The local Hadamard singular structure, causal commutator, and a conditional lifting statement are available. A full covariance is not constructed because the explicit 104-dimensional stationary carrier A_{104} , its Cauchy pairing q_{104}^{Cauchy} , Green carrier G_{104}^{Cauchy} , and real structure have not been supplied. The current ledger leaves 288 coordinates undetermined; the two tested compatible completion branches have zero-eigenspace dimensions 24 and 0. This is a missing-object obstruction, not a proof that no BRST-compatible Hadamard covariance exists. The reduced positive η result above does not define the adjoint on this full BV carrier and cannot be used to fill the gap. Thus no positive full-BV state or one-particle Hilbert space has been constructed.

The immediate quantum literature includes conformal-gauge fixing and its additional ghosts, local BRST cohomology, and several alternative quantization prescriptions[36, 33]. Phase 1 neither reproduces all of that literature nor supersedes it. It contributes a content-addressed strict/extended local-BV calculation and records the precise point at which Lorentzian state construction remains absent.

8 What the counterflow candidate taught us

The strict anomaly and passive-compensator causal obstruction motivated a small changed-theory ladder. Passive trace repairs and a one-phase active clock failed combined causal, sign, and charge tests. Two phases with an auxiliary diagonal $U(1)$ allow neutral counterflow:

$$\dot{\chi} = 0, \quad \dot{\psi} = \Omega, \quad Q_{\text{diag}} = 0, \quad E_{\text{rel}} = \frac{1}{2}\mu^2\Omega^2 > 0. \quad (24)$$

The relative phase can move while the diagonal compact gauge charge vanishes. Because its stress tensor is built from the dressed metric, it is quadratically visible in the dressed-trace direction.

For self-containment, the stationary family can be stated directly. Let σ_i be left-invariant one-forms on S^3 with

$$d\sigma_1 = -\sigma_2 \wedge \sigma_3, \quad d\sigma_2 = -\sigma_3 \wedge \sigma_1, \quad d\sigma_3 = -\sigma_1 \wedge \sigma_2, \quad \int_{S^3} \sigma_1 \wedge \sigma_2 \wedge \sigma_3 = 16\pi^2. \quad (25)$$

In the gauge $\tau = 0$, $A = 0$, and constant χ , take

$$ds^2 = -dt^2 + a^2(\sigma_1^2 + \sigma_2^2) + c^2\sigma_3^2, \quad \psi = \Omega t, \quad (26)$$

and define

$$q = \frac{c^2}{a^2}, \quad x = a^{-2}, \quad C = \mu^2\Omega^2. \quad (27)$$

The exact stationary equations for Eq. (8) have rank three and give

$$M_2 = \frac{2C(4q-1)}{3qx}, \quad V_0 = -\frac{C(q^2-5q+1)}{3q}, \quad \alpha_B = \frac{2C}{qx^2}. \quad (28)$$

The connected causal trace-healthy component is

$$x > 0, \quad C > 0, \quad \frac{13-3\sqrt{17}}{4} < q < \frac{1}{4}. \quad (29)$$

The selected causal point fixes $\mu^2 = 1$ and

$$q = \frac{9}{40}, \quad x = 1, \quad C = \frac{9}{16}, \quad \Omega = \frac{3}{4}, \quad \alpha_B = 5, \quad M_2 = -\frac{1}{6}, \quad V_0 = \frac{119}{1920}. \quad (30)$$

The free BV object is the tangent complex of the displayed action and gauge transformations. Write its ghost and field bundles as

$$\mathcal{F}^{-1} = TM \oplus \mathbf{1}_W \oplus \mathbf{1}_{U(1)}, \quad \mathcal{F}^0 = S^2 T^* M \oplus \mathbf{1}_\tau \oplus \mathbf{1}_\psi \oplus \mathbf{1}_\chi \oplus T^* M_A. \quad (31)$$

The minimal linear complex is then

$$\Gamma(\mathcal{F}^{-1}) \xrightarrow{q_{-1}} \Gamma(\mathcal{F}^0) \xrightarrow{q_0} \Gamma(\mathcal{E}) \xrightarrow{q_1} \Gamma((\mathcal{F}^{-1})^* \otimes \Lambda^4 T^* M), \quad (32)$$

where q_{-1} is the linearized Diff–Weyl– $U(1)$ action, q_0 is the Euler–Lagrange Hessian, q_1 contains the Noether identities, and $\mathcal{E}$ is the dual equation bundle to the field bundle. The differential orders are also fixed by the action: q_{-1} is first order in the diffeomorphism and diagonal- $U(1)$ parameters and zeroth order in the Weyl parameter; q_0 is fourth order in the metric C^2 block, second order in the dressed Einstein and phase blocks, and algebraic in the auxiliary connection before elimination; q_1 has the corresponding formal-adjoint Noether orders. The cotangent BV rows and standard antighost–multiplier doublets complete this to a cyclic complex. The 70-row certificate is a harmonic-component implementation of Eq. (32); the invariant definition is the action and this graded complex.

Result card E: causal construction and physical nonselection

Theory. Equation (8) with the gauge transformations (9).

Background. The selected positive biaxial Berger solution and its connected trace-healthy same-field stationary retuning family.

Domain. The free cyclic BV complex (32); at the selected point its diagonal- $U(1)$ summand is contractible, and the physical spectral reduction uses complete $SU(2)_L \times U(1)_R$ isotypical blocks.

Gauge and charge. Diagonal gauge charge, relative-phase charge Q_{rel} , time translation D , and the background-preserving generator $K = D - \Omega R_{\text{rel}}$ are kept distinct.

Causal statement. Exact retarded and advanced homotopies, nilpotency, cyclicity, and support identities hold at the selected point, and the original dressed-trace cohomology class is removed.

Physical statement. The complete both- k , $j = \frac{1}{2}$ reduced block contains a Hamiltonian–Hopf quartet. The obstructing factor is

$$(40z^4 + 773z^2 + 3748)^2, \quad 773^2 - 4(40)(3748) = -2151 < 0.$$

With perturbations proportional to e^{zt} , the substitution $y = z^2$ gives two nonreal conjugate y -roots, hence four z -roots $\{\pm(\alpha + i\beta), \pm(\alpha - i\beta)\}$ with $\alpha\beta \neq 0$; two grow exponentially. The quartet persists throughout the connected trace-healthy same-field stationary family. Therefore no member of that family is selected as a robust linearly stable candidate.

PT/CPT disposition. The complex quartet is an exact infeasibility certificate for a positive pseudo-Hermitian metric on this physical block; it lies in the broken-PT regime and cannot be rescued by redefining only the inner product.

Not claimed. This is not an all-isotype spectrum, a familywide Green-homotopy theorem, a full-PDE stability theorem, or a no-go over other field contents and gauge architectures.

The rejection criterion was fixed before the computation: a successor candidate had to be linearly stable in every physical sector. One exact unstable physical isotypical block is sufficient to reject it.

It is not necessary to finish every higher- j block to establish this nonselection. Conversely, the finite block does not by itself describe the nonlinear evolution of arbitrary PDE data.

The persistence statement is exact rather than a numerical continuation from Eq. (30). Restoring $w = z^2/x$, the family characteristic divisor contains the quadratic factor

$$F_2(q, w) = 16q^2w^2 + 24q(3 - 2q^2)w + 32q^3 - 108q^2 + 81, \quad (33)$$

with discriminant

$$\text{disc}_w F_2 = 256q^5(9q - 8) < 0 \quad \text{on (29)}. \quad (34)$$

Thus the two w -roots remain a nonreal conjugate pair everywhere on the connected component; they cannot cross onto the negative real axis or pass through a multiple root. Since $x > 0$, their square roots remain a Hamiltonian–Hopf quartet. This no-crossing lemma is the familywide argument.

The quartet also fixes the disposition of this block under pseudo-Hermitian and PT/CPT-type reinterpretations[8]. Write the finite reduced block in first-order form $\dot{u} = Au$, so the characteristic roots z are the eigenvalues of A , and let $H = iA$ be the corresponding Schrödinger-form generator. If a positive metric η satisfied the pseudo-Hermiticity relation $H^\dagger \eta = \eta H$, then $\eta^{1/2} H \eta^{-1/2}$ would be Hermitian, its spectrum would be real, and every characteristic root z would be purely imaginary. The certified quartet has $\alpha \neq 0$, so no such positive η exists on this block. In PT language the block is in the broken-PT regime: the obstruction is spectral, not a choice of norm or inner product, and no alternative quantization prescription acting only through the inner product can reopen it.

The equal-frequency Pais–Uhlenbeck mechanism occupies a different spectral boundary. There the degenerate generator develops Jordan structure and the PT programme interprets the associated zero-norm states through a specially constructed C operator[10, 9]. Secular logarithms and Jordan chains also appear in our characteristic-shell reconstructions, so they are a useful classical analogy. The pilot has not constructed zero-norm decoupling, a C operator, or an equivalence with the equal-frequency oscillator. The analogy therefore locates where the Mannheim mechanism could live without promoting the logarithmic solutions to a quantum theorem.

The quartet is a spectral Hamiltonian instability associated with a Krein-sign collision in the standard Hamiltonian–Krein sense [7], not merely a coordinate drift. The global clock block has a different status. On the unrestricted phase space,

$$\Omega_{\text{rel}} = dQ_{\text{rel}} \wedge d\psi, \quad H_{\text{rel}} = \frac{Q_{\text{rel}}^2}{2I} + E_{\text{geometry}}, \quad (35)$$

so its secular dephasing is tangent to an action–angle family and passes an orbital, frequency-modulated interpretation. It does not cure the $j = \frac{1}{2}$ quartet.

The charge analysis gives an independent theorem. For phase clocks with linear charge constraint $CQ = q_0$ and clock velocity vector v ,

$$D \text{ is null after reduction} \iff v \in \text{im } C^T, \quad (36)$$

while physical clocks are classes of v modulo $\text{im } C^T$. In the selected model, fixing and quotienting Q_{rel} makes D null but contracts the relative-clock Darboux pair. Keeping the clock leaves $H_D = \Omega Q_{\text{rel}}$ and raw D charged; $K = D - \Omega R_{\text{rel}}$ preserves the background. No single declared charge fibre supplies both the proposed physical clock and a gauge time translation.

The counterflow result is therefore not a failed piece of engineering. It separates three notions that are often conflated:

$$\begin{aligned} \text{causal completeness} &\not\Rightarrow \text{positive reduced dynamics,} \\ \text{positive reduced dynamics} &\not\Rightarrow \text{acceptable clock/charge interpretation.} \end{aligned} \quad (37)$$

The causal theorem remains valid even though the candidate is rejected by a later physical gate.

9 Integrated Phase-1 verdict

The main findings can now be placed on the four levels of Eq. (2).

Level	Established in a declared scope	Still absent
Local linear systems	Complete causal BV complexes on selected compact backgrounds; exact additional compact-product and Schwarzschild curvature solutions	A single causal theory covering every laboratory and boundary class
Reduced classical directions	Additional axial/polar compact-product directions are nonradical; counterflow causal parent and exact charge classification; axial formal Schwarzschild counterexample and polar finite mixed line with exact Q_{21} wall	Universal asymptotic phase space, positive energy theorem, and acceptable counterflow spectrum
Nonlinear continuation	Five-charge necessary-and-sufficient finite exponential-polynomial cone; bounded resonance ledger and classified subcones	Full bounded finite-support zero locus, retarded nonlinear existence, all-order integration, and stability
Quantum states/particles	Strict local one-loop BV obstruction; formal changed-theory Wess–Zumino resolution; structured positive η on selected reduced blocks; spectral PT no-go for the counterflow quartet	Genuine BRST-compatible C operator, full-BV Hadamard state, positive physical Hilbert space, particles, scattering, optical theorem, and unitarity

This supports the following central claim:

A pole-level diagnosis is too coarse, but the more complete reduction does not generically rescue the theory. Some additional classical directions survive gauge and presymplectic reduction. Nonlinear moment maps, radial filtrations, local anomalies, inner-product compatibility, and reduced spectral tests then separate candidates for different reasons. The first changed theory to achieve a complete causal BV parent is rejected throughout its connected trace-healthy same-field family by a physical $j = \frac{1}{2}$ Hamiltonian–Hopf instability.

This is a classification ending, not a viable-theory claim and not a universal no-go. A different field content, a different gauge architecture, or a different admissible state prescription could define a new programme branch. The scoped Phase-2 update adds sharper comparisons without reopening or rewriting the frozen Phase-1 record.

10 Relation to prior work

The programme combines several established structures. Its claims should be read as explicit constructions and exhaustive calculations in specified models, not as inventions of the surrounding general theories.

Causal gauge complexes. Green-hyperbolic complexes, retarded/advanced Green homotopies, and their induced Poisson structures have a general Lorentzian theory [16]. Conformal deformation and Yang–Mills detour complexes place the Bach operator in a longer conformal-geometric lineage [17, 18]. The contribution here is the explicit action-derived conformal-gravity BV construction, its cyclic pairings and support identities, and transfer across the declared backgrounds.

Linearization stability. The relation between Killing symmetries, quadratic Taub constraints, and singular momentum-map zero sets is classical [19, 20, 21, 22]. The finite-harmonic theorem is an exhaustive Weyl–Maxwell realization: exactly five covectors exhaust the exponential-polynomial cokernel, exceptional and generalized modes are included, and bounded resonance conditions are separated from the formal cone.

Einstein selection and black holes. Boundary-condition selection of Einstein solutions in conformal gravity is well known in Euclidean and asymptotically (A)dS settings [23, 24, 25]. Prior Weyl-black-hole quasinormal analyses often isolate a second-order conformally transported sector rather than the full fourth-order Bach solution space [26]. The static spherical Bach-flat lineage begins with Riegert and Mannheim–Kazanas [27, 28]. Four-dimensional conformal gravity already has consistent AdS boundary conditions, response functions, and finite canonical charges [29, 30]; our open problem is specifically the asymptotically flat/null-infinity Bach phase space. Our generic-angular result is a counterexample to parity-uniform Einstein selection in a formal radial class and an exact polar filtration with an exceptional-frequency wall. It does not yet provide a general asymptotically flat Bach phase space.

The magnetic product background belongs to the exceptional direct-product Einstein–Maxwell family introduced by Plebański and Hacyan [31]. The recent Weyl-gravity classification of separated $2 + 2$ direct products with electromagnetic and Yang–Mills sources proves generalized Birkhoff statements and organizes their Weyl-equivalence classes [32]. The present contribution is different: it fixes one compact magnetic bundle and computes its perturbative exact sequence, presymplectic form, nonlinear Taub cone, and resonances.

Quantum conformal gravity. One-loop divergences, conformal anomalies, BRST cohomology, and conformal gauge fixing have extensive prior literatures [34, 33, 36, 37]. The phase-one quantum claim is narrower: an exact local-BV quotient and coefficient ledger for the declared strict theory, followed by a formal τ -adic Wess–Zumino trivialization in a changed theory. The Phase-2 comparison with pseudo-Hermitian quantum mechanics [11, 12] establishes a positive structured metric on selected reduced blocks and a spectral no-go on the broken-PT counterflow quartet, but no genuine BRST-compatible C operator or full Lorentzian state. The Hadamard distinction follows the microlocal state criterion [13]. Dirac–Pauli, fakeon, and conventional positive-state proposals are not settled.

Relational clocks. The construction of complete or partial observables from internal clocks is standard constrained-system territory [39]. The counterflow experiment adds a typed distinction between diagonal gauge charge, relative clock charge, time translation D , and the background-preserving combination K . No operational counterflow frequency ratio survived Phase 1, so the result is a charge/clock classification rather than an observational prediction.

11 Limitations and the next scientific boundary

The scoped Phase-2 results replace the earlier task list with sharper endpoint questions. The most important are not “finish quantum gravity.” They are:

1. complete the asymptotically flat Bach phase space, match horizon data to the certified formal infinity modules, and determine whether the axial counterexample and polar mixed line survive a wave-packet flux topology;
2. classify the axial X_2 branch, terminal-only polar prefixes, and the deeper filtration at $Q_{21} = 0$;
3. construct independent P, T, C data and prove BRST and causal-covariance compatibility, rather than promoting finite-block η existence to a state theorem;
4. supply the missing 104-dimensional stationary carrier and decide the Hadamard covariance problem for the complete Berger BV complex;
5. if a new changed theory is proposed, require an open stationary locus with causal completeness, positive reduced dynamics, coherent charge interpretation, a preserved nonlinear sector, and at least one operational observable before calling it a candidate.

These are continuing Phase-2 gates. The bounded zero locus and mixed interaction class remain valid inputs, but are deprioritized as standalone interest levers. None is a condition for closing Phase 1. The stopping rule was outcome-independent: Phase 1 ended when the programme could give an exact viability classification of the first causally complete changed-theory candidate and reconcile it with the compact linear, nonlinear, black-hole, observer, and quantum results. That record is now frozen; the present additions are an explicitly layered update.

12 Computational accountability

The project treats computer algebra as an experimental instrument whose outputs require independent checks. Material results are represented by a producer, a machine-readable certificate, a verifier that does not merely rerun the producer, adversarial mutation tests where practical, a tiered test receipt, and a human-readable scope report. Exact algebraic ranks, Smith forms, cohomology witnesses, and characteristic factors use rational or algebraic arithmetic; numerical evidence is typed separately.

The authoritative Phase-1 record is

PURE_WEYL_PROGRAMME_PHASE1_CLASSIFICATION_ENDING_V1,

stored in

reports/phase1-closure-claims-ledger-2026-07-22.json.

The public repository root is

<https://github.com/area9innovation/bp2transformer/tree/master/physics/symplectic-reconstruction>,

with this manuscript and its machine-readable claim map under <https://github.com/area9innovation/bp2transformer/tree/master/physics/symplectic-reconstruction/paper>. The JSON artifacts carry content hashes; no archival DOI is claimed at this stage. It binds each statement to a

theory, background, phase space, correction or boundary class, lifecycle state, limitation, and source certificate. This paper has its own claim map and verifier; the live programme portal remains Paper 98. A claim that is not linked to evidence is left unpromoted rather than inferred from neighboring calculations.

Three aspects of this workflow matter scientifically. First, a failed promotion is retained: the invalid Berger round-sphere Hodge subspace, for example, was replaced by full $SU(2)_L \times U(1)_R$ isotypical blocks rather than silently repaired. Second, causal, spectral, algebraic, and quantum results carry different dependency labels. Third, stopping rules are fixed before the result is known. The counterflow candidate passed its causal gate and then failed a later physical gate; both facts remain part of the record.

13 Conclusion

The phase-one programme began with a common but underspecified question: whether the additional poles of pure Weyl gravity are physical ghosts. Its main conceptual result is that there is no single promotion from pole to particle. Local solutions, reduced classical directions, nonlinear tangent directions, and quantum states are different mathematical objects.

That distinction did not automatically rescue the theory. Additional compact-product directions survive local reduction, yet some fail bounded nonlinear continuation. Horizon regularity admits additional Schwarzschild curvature solutions, and the generic-angular formal radial calculation now supplies an axial counterexample to Einstein-only selection plus a polar mixed finite line with an exact exceptional wall. Selected compact blocks admit positive pseudo-Hermitian metrics, but neither a genuine BRST-compatible C operator nor a full Hadamard covariance follows. Strict pure Weyl gravity has a local Euclidean one-loop BV obstruction; a compensator removes it only in a changed theory. The first changed model with a complete causal BV parent is itself rejected by an exact family-persistent Hamiltonian–Hopf instability, and its complex spectrum excludes a positive-metric PT/CPT rescue.

The most defensible conclusion is therefore neither “the propagator already settled the physics” nor “constraints removed every ghost.” It is:

Pole-level signs are an initial diagnostic. Physical viability is a sequence of reductions and existence tests. In the laboratories updated through scoped Phase 2, different additional sectors survive different early tests, but no candidate reaches a positive full quantum state, and the first causally complete changed theory fails a robust linear physical-stability test.	(38)
--	------

References

- [1] J. Lee and R. M. Wald, “Local symmetries and constraints,” *J. Math. Phys.* **31** (1990) 725–743.
- [2] V. Iyer and R. M. Wald, “Some properties of Noether charge and a proposal for dynamical black hole entropy,” *Phys. Rev. D* **50** (1994) 846–864, [arXiv:gr-qc/9403028](#).
- [3] S. Hollands and R. M. Wald, “Stability of black holes and black branes,” *Commun. Math. Phys.* **321** (2013) 629–680, [arXiv:1201.0463](#).
- [4] R. P. Woodard, “The theorem of Ostrogradsky,” [arXiv:1506.02210](#).

- [5] K. S. Stelle, “Renormalization of higher-derivative quantum gravity,” *Phys. Rev. D* **16** (1977) 953–969.
- [6] K. S. Stelle, “Classical gravity with higher derivatives,” *Gen. Relativ. Gravit.* **9** (1978) 353–371.
- [7] T. Kapitula and K. Promislow, *Spectral and Dynamical Stability of Nonlinear Waves*, Springer, New York (2013).
- [8] C. M. Bender and P. D. Mannheim, “No-ghost theorem for the fourth-order derivative Pais–Uhlenbeck oscillator model,” *Phys. Rev. Lett.* **100** (2008) 110402, [arXiv:0706.0207](#).
- [9] C. M. Bender and P. D. Mannheim, “Exactly solvable PT-symmetric Hamiltonian having no Hermitian counterpart,” *Phys. Rev. D* **78** (2008) 025022, [arXiv:0804.4190](#).
- [10] A. Pais and G. E. Uhlenbeck, “On field theories with non-localized action,” *Phys. Rev.* **79** (1950) 145–165.
- [11] A. Mostafazadeh, “Pseudo-Hermitian representation of quantum mechanics,” *J. Phys. A* **39** (2006) 10171–10188, [arXiv:quant-ph/0508195](#).
- [12] I. Kuntz, “Unitarity through PT symmetry in quantum quadratic gravity,” [arXiv:2410.08278](#).
- [13] M. J. Radzikowski, “Micro-local approach to the Hadamard condition in quantum field theory on curved space-time,” *Commun. Math. Phys.* **179** (1996) 529–553.
- [14] A. Salvio, “A non-perturbative and background-independent formulation of quadratic gravity,” [arXiv:2404.08034](#).
- [15] D. Anselmi, “Fakeons, quantum gravity and the correspondence principle,” [arXiv:1911.10343](#).
- [16] M. Benini, M. Musante, and A. Schenkel, “Green hyperbolic complexes on Lorentzian manifolds,” [arXiv:2207.04069](#).
- [17] T. P. Branson and A. R. Gover, “The conformal deformation detour complex for the obstruction tensor,” [arXiv:math/0605192](#).
- [18] A. R. Gover, P. Somberg, and V. Souček, “Yang–Mills detour complexes and conformal geometry,” [arXiv:math/0606401](#).
- [19] D. R. Brill and S. Deser, “Instability of closed spaces in general relativity,” *Commun. Math. Phys.* **32** (1973) 291–304.
- [20] A. E. Fischer and J. E. Marsden, “Linearization stability of the Einstein equations,” *Bull. Am. Math. Soc.* **79** (1973) 997–1003.
- [21] V. Moncrief, “Spacetime symmetries and linearization stability of the Einstein equations I, II,” *J. Math. Phys.* **16** (1975) 493–498 and **17** (1976) 1893–1902.
- [22] J. M. Arms, J. E. Marsden, and V. Moncrief, “Symmetry and bifurcations of momentum mappings,” *Commun. Math. Phys.* **78** (1981) 455–478.
- [23] J. Maldacena, “Einstein gravity from conformal gravity,” [arXiv:1105.5632](#).
- [24] H. Lü, Y. Pang, and C. N. Pope, “Conformal gravity and extensions of critical gravity,” [arXiv:1106.4657](#).

- [25] S. Deser, E. Joung, and A. Waldron, “Partial masslessness and conformal gravity,” [arXiv:1208.1307](#).
- [26] M. Momennia and S. H. Hendi, “Quasinormal modes of black holes in Weyl gravity: electromagnetic and gravitational perturbations,” [arXiv:1910.00428](#).
- [27] R. J. Riegert, “Birkhoff’s theorem in conformal gravity,” *Phys. Rev. Lett.* **53** (1984) 315–318.
- [28] P. D. Mannheim and D. Kazanas, “Exact vacuum solution to conformal Weyl gravity and galactic rotation curves,” *Astrophys. J.* **342** (1989) 635–638.
- [29] D. Grumiller, M. Irakleidou, I. Lovrekovic, and R. McNees, “Conformal gravity holography in four dimensions,” *Phys. Rev. Lett.* **112** (2014) 111102, [arXiv:1310.0819](#).
- [30] I. Lovrekovic, “Canonical charges and asymptotic symmetries in four dimensional conformal gravity,” [arXiv:1505.05820](#).
- [31] J. F. Plebański and S. Hacyan, “Some exceptional electrovac type D metrics with cosmological constant,” *J. Math. Phys.* **20** (1979) 1004–1010.
- [32] P. Jizba and T. Lehečková, “Generalized Birkhoff theorems and $2+2$ direct product spacetimes in Weyl conformal gravity,” *Phys. Rev. D* **113** (2026) 044060, [arXiv:2512.18482](#).
- [33] G. Barnich, F. Brandt, and M. Henneaux, “Local BRST cohomology in gauge theories,” *Phys. Rept.* **338** (2000) 439–569, [arXiv:hep-th/0002245](#).
- [34] E. S. Fradkin and A. A. Tseytlin, “Renormalizable asymptotically free quantum theory of gravity,” *Nucl. Phys. B* **201** (1982) 469–491.
- [35] S. Deser and A. Schwimmer, “Geometric classification of conformal anomalies in arbitrary dimensions,” *Phys. Lett. B* **309** (1993) 279–284, [arXiv:hep-th/9302047](#).
- [36] L. Rachwał, “Introduction to quantization of conformal gravity,” [arXiv:2204.13856](#).
- [37] I. Oda and N. Ohta, “Quantum conformal gravity,” [arXiv:2311.09582](#).
- [38] F. Baume and B. Keren-Zur, “The dilaton Wess–Zumino action in higher dimensions,” [arXiv:1307.0484](#).
- [39] B. Dittrich, “Partial and complete observables for Hamiltonian constrained systems,” *Gen. Rel. Grav.* **39** (2007) 1891–1927, [arXiv:gr-qc/0411013](#).