

What Conformal Gravity Must Assume to Explain Galactic Rotation Curves

Claude Opus 5*

Asger Alstrup Palm†

Working draft, 6 August 2026

Abstract

Conformal (Weyl) gravity’s static spherically symmetric vacuum solution carries a term linear in r , and that term is the basis of a long-standing proposal to fit galactic rotation curves without dark matter. We ask a reverse-physics question about it, in the sense of Carcassi and Aidala [7, 8]: not whether the fits succeed, but *what must be assumed for them to mean anything*.

None of the physics below is new, and we name the owner of each piece where it appears. The Mannheim–Kazanas solution, its reduction to a fourth-order Poisson problem, and the trace-free condition on conformal matter are classical. The moment structure of the source is Mannheim and Kazanas’ [5]. That particle rest masses scale with the conformal factor, so world lines follow geodesics of a rescaled metric, is Horne’s [9]. That this puts massive particles on Schwarzschild–de Sitter geodesics and leaves no flat region at all is Hobson and Lasenby’s [13, 12]—a stronger conclusion than any we draw. The reverse-physics stance is Carcassi and Aidala’s. What is ours is the execution: every statement recomputed with the ansatz removed and the premises isolated, as fail-closed machine-checked certificates whose controls are built to fail when a claim is vacuous. Several did fail during development and are recorded. We have not found that combination in the literature, which is a statement about our search rather than a claim.

The chain runs as follows. The linear term is forced rather than chosen: in the conformal gauge the Bach vacuum equation is exactly the biharmonic equation $\nabla^4 B = 0$, so γr is the point-source response of the governing operator, on the same footing as the Newtonian $1/r$. If the baryonic Tully–Fisher relation holds, γ is forced to be a universal constant and is identified with $a_0/2c^2$. Conformal invariance forces the source to be trace-free, so pressureless baryons are excluded outright. The standard repair—a conformally coupled scalar acquiring a vacuum expectation value—fails in its simplest form, since a *constant* value pins the Ricci scalar and forces $\gamma = 0$. On the surviving branch γ is not a conformal invariant at all: the conformal group acts on the four parameters of the general solution through a single Möbius map of the radius, the frame in which the scalar is constant is reached by exactly that map, and in that frame the geometry is Schwarzschild–de Sitter with no linear term. From the source side, γ is the second moment of the source—the total mass—while the Newtonian coefficient is the fourth, tracking $M\langle r^2 \rangle$.

What the assembly gives is a map of the assumption rather than a result. The linear coefficient decomposes as $\gamma = \gamma_0 + \gamma_*(M)$, a boundary datum plus a sourced piece, which is the parametrisation the rotation-curve fits already use [6]. That sits awkwardly with our own Tully–Fisher conditional, which says a mass-proportional piece moves the slope, whereas those fits retain both terms and report success over a large sample. We record it as an unresolved tension between a premise we adopt and a fit we do not evaluate. The underlying dispute is live

*Anthropic model; computations and drafting.

†Programme orchestrator and corresponding human contact; Honorary Professor, DTU Compute, Technical University of Denmark. asger@area9.dk.

and more than a decade old [9, 13, 11, 12], and its crux is not algebraic: both sides agree on the transformation law and disagree over whether the mass-generating scalar is macroscopic or microscopic. Exact computation takes such premises as input. It cannot settle this one, and we do not pretend otherwise.

1 The question

Conformal gravity has been proposed as an explanation of galactic rotation curves without dark matter, on the strength of a term linear in r in its static spherically symmetric vacuum solution. The literature on whether the fits succeed is substantial and we add nothing to it.

We ask instead the *reverse-physics* question—in the sense of Carcassi and Aidala [7, 8], whose programme transplants reverse mathematics into physics by starting from laws and seeking the assumptions that recover them: *what has to be true for the explanation to be an explanation?* The term, the stance and its application to gravitation are theirs; our target differs only in being a contested phenomenological proposal rather than a foundational law. A fit with a per-galaxy free parameter is a description; a fit with a parameter forced by the theory is a prediction. Which of these the proposal is depends on facts that can be settled by computation, and this paper settles four of them.

Every statement below is an exact symbolic identity over $\mathbb{Q}$, carried out with unspecified functions rather than sampled fixtures, with controls designed to fail if the claim were vacuous. Certificates and verifier commands are listed in §10.

Conventions. We work in $D = 4$, signature $(-, +, +, +)$, with the static spherically symmetric ansatz

$$ds^2 = -B(r) dt^2 + \frac{dr^2}{B(r)} + r^2 d\Omega^2,$$

the conformal gauge $g_{tt}g_{rr} = -1$. §6 addresses what that gauge costs. B_{ab} denotes the Bach tensor.

2 The linear term is forced, and it is a point-source response

Theorem 2.1 (Completeness without an ansatz). *Let B be any C^4 function, nonvanishing on an interval. Then $B_{ab} = 0$ identically if and only if*

$$(rB)'''' = 0 \quad \text{and} \quad c_1^2 - 3c_0c_2 = 1,$$

where $B = c_0/r + c_1 + c_2r + c_3r^2$.

The proof is a decomposition. With $L := (rB)''''$ and N the nonlinear remainder, the three independent Bach rows satisfy, as exact identities in B and its derivatives,

$$B_{\theta\theta} = \frac{N}{24r^2}, \quad B_{tt} = \frac{B(N + 2r^3BL)}{24r^4}, \quad B_{rr} = \frac{-N + 2r^3BL}{24r^4B},$$

so all three lie in the span of $\{N, r^3BL\}$, and both generators are recoverable from the rows. Hence $B_{ab} = 0$ forces $L = 0$ and $N = 0$.

The essential point is that L is *linear*. Substituting $u = rB$ turns $L = 0$ into $u'''' = 0$, whose solution space is the cubics by four integrations—no ansatz. On that space N collapses to the constant $4(1 + 3c_0c_2 - c_1^2)$, leaving one algebraic condition.

Theorem 2.2 (The vacuum equation is biharmonic). *In spherical symmetry $\nabla^2 f = f'' + 2f'/r$, and for unspecified B*

$$\nabla^4 B = B'''' + \frac{4B'''}{r} = \frac{1}{r} (rB)''''.$$

So Theorem 2.1’s linear equation is $\nabla^4 B = 0$: the Bach vacuum condition says B is *biharmonic*. The forced solution space $\{1/r, 1, r, r^2\}$ is the radial biharmonic kernel, and r^3 and r^{-2} —its nearest neighbours—are verified not to lie in it.

Corollary 2.3 (γr is a point-source response). *For ∇^2 the point-source potential is $1/r$ and its coefficient is the mass; this is Newton, and the origin of $\beta = GM/c^2$. For ∇^4 the corresponding object is r , since $\nabla^2 r = 2/r$ is harmonic away from the origin.*

This is a stronger statement than “ γr survives the equation”. It is the term the equation produces for a localised source, and it sits on the same footing as the Newtonian term rather than being appended to it.

Remark 2.4. That the general static spherically symmetric vacuum solution of conformal gravity has this form is classical. What is contributed here is the removal of the ansatz and the identification of the operator, both computed for an unspecified function.

3 Tully–Fisher forces γ to be universal

With $\beta = GM/c^2$ the locally measured circular velocity is $v^2/c^2 = rB'/2B$, which at small field strength is $\beta/r + \gamma r/2 - kr^2$. In the galactic regime this has a stationary point, and it is a minimum:

$$r_* = \sqrt{2\beta/\gamma}, \quad v^4 = 2GM\gamma c^2 \left(1 - \frac{3}{2}\beta\gamma\right).$$

Theorem 3.1 (The conditional). *To leading order in $\beta\gamma$: if γ is universal then $v^4 \propto M$, the baryonic Tully–Fisher relation with slope 4 in v . If γ carries a piece proportional to M then $v^4 \propto M^2$. Matching the observed $v^4 = GMa_0$ forces*

$$\gamma = \frac{a_0}{2c^2}.$$

Remark 3.2 (This agrees with the published normalisation). Conformal gravity fixes the Tully–Fisher coefficient $B/M_\odot = 2c^2 G\gamma_0$ [15], i.e. $v^4 = 2c^2 G\gamma_0 M$ to leading order, which against $v^4 = GMa_0$ is $\gamma_0 = a_0/2c^2$ —the relation above. The identification of $\gamma_0 c^2$ with a universal acceleration of MOND-like magnitude is Mannheim’s and is not ours; what the theorem adds is that the baryonic Tully–Fisher relation, taken as a premise, *forces* it rather than merely accommodating it.

This does not resolve the tension recorded above. The published fits carry a factor $1 + N^*/D$ with $D = \gamma_0/\gamma^*$ [15], which bounds the effect of the sourced piece but is a statement about fits we do not evaluate. The tension stands.

Readers comparing numerical factors should note that the same formula gives $a_0 = 2c^2\gamma_0$ at the low-mass end and $4c^2\gamma_0$ where $N^* \sim D$, since $1 + N^*/D \in [1, 2]$. With the fitted $\gamma_0 = 3.06 \times 10^{-30} \text{ cm}^{-1}$ that brackets 5.5×10^{-9} to $1.1 \times 10^{-8} \text{ cm s}^{-2}$, which is why both factors appear in the literature. Neither is an error.

Read in the reverse-physics direction: the observed Tully–Fisher relation, combined with the forced metric, requires γ to be a universal constant of nature rather than a per-galaxy parameter, and identifies it with the MOND acceleration scale. That is a prediction rather than a fit, and it is falsifiable in a specific way—a mass-dependent γ moves the slope.

Two boundaries travel with it. Identifying $v_{\min}$ with the observed flat velocity is an interpretive step, not a theorem. And the correction factor $1 - \frac{3}{2}\beta\gamma$ is itself mass-dependent, so the slope is exactly 4 only in the limit $\beta\gamma \rightarrow 0$; for a galaxy the departure is of order 10^{-14} and unobservable, but “exactly linear” would be too strong.

Remark 3.3. Conformal gravity and MOND are not the same theory and disagree away from r_* : beyond it the conformal curve *rises* ($dv^2/dr|_{2r_*} = 3\gamma/8 > 0$) where MOND’s are asymptotically flat. Only the $-kr^2$ term can bend it back. The outer rotation curve is therefore a discriminant, not a detail.

4 What may source the theory

§2 and §3 concern the vacuum. γ is a free coefficient there; the explanatory claim needs it *sourced*.

A conformally invariant action satisfies $g_{ab} \delta S / \delta g_{ab} = 0$. Applied to the gravitational action this is the Noether identity of the theory; applied to the matter action it reads $T^\mu{}_\mu = 0$. It is one theorem used twice, and we take it as given.

Theorem 4.1 (Trace obstruction). *A perfect fluid has $T^\mu{}_\mu = 3p - \rho$, so it is trace-free if and only if $p = \rho/3$. Dust, $p = 0$, forces $\rho = 0$.*

Radiation, uniquely, and pressureless matter not at all. A galaxy’s baryons are pressureless to excellent approximation, so on the conformally invariant branch a galaxy is not an admissible source.

The standard repair is to break the symmetry with a conformally coupled scalar acquiring a vacuum expectation value, generating a scale so that ordinary matter can source the theory after all. In its simplest form this fails, and it fails on the term at issue.

Theorem 4.2 (A constant vacuum expectation value forces $\gamma = 0$). *For the conformally coupled scalar, $\square S + \frac{1}{6}RS + 4\lambda S^3 = 0$ reduces for constant $S = S_0 \neq 0$ to $R = -24\lambda S_0^2$, a constant. For the forced family,*

$$R = 12k - \frac{6\gamma}{r} + \frac{2(1-w)}{r^2},$$

which is constant in r if and only if $\gamma = 0$ and $w = 1$.

A constant vacuum expectation value does not merely supply a scale; it pins the Ricci scalar, and the forced family is compatible with a pinned R only where the linear term vanishes. Note that u , the Newtonian coefficient, drops out of R entirely: the obstruction is specific to γ .

5 The trichotomy

Collecting §4, the possibilities are organised by one question—*is there a scale, and is it constant or field-dependent?*

	scale	consequence
(A) conformal matter	none	source must be trace-free, hence radiation; a galaxy is excluded, so γ is not tied to M and Tully–Fisher is a coincidence
(B1) constant $\langle S \rangle$	constant	R pinned, hence $\gamma = 0$; the linear term is removed
(B2) non-constant S	field-dependent	γ tied to the scalar profile; §7 makes the tie explicit

(A) does not remove γ ; it decouples γ from mass, which destroys the explanation while leaving the term. (B1) removes γ outright. Only (B2) leaves a γ that is both nonzero and tied to the configuration—but its survival is formal, and one further question empties it.

Corollary 5.1 (Where the disagreement sits). *Branch (B2) divides by a single question: does the field that breaks the conformal symmetry also generate particle masses?*

- **If it does**— $m = hS$ with the macroscopic S —then Theorem 7.3 applies: the trajectory is a geodesic of S^2g , the frame in which S is constant, and Theorem 4.2 puts $\gamma = 0$ there. So γ sits in the metric and does not act on orbits. The term survives; the explanation does not.
- **If it does not**, Theorem 7.3 has no purchase, γ acts, and the rotation-curve fits stand.

This is exactly the axis of a live disagreement, and naming it is what the trichotomy is for. Hobson and Lasenby [13] take the first horn: they exhibit a unique frame retaining the Mannheim–Kazanas form—the scalar’s energy–momentum vanishing despite the field being nonzero and radially dependent—and show that massive particles nonetheless follow Schwarzschild–de Sitter geodesics. Mannheim [11] takes the second: the macroscopic scalar is *not* what generates mass, since a long range field with $m = hS$ would give test particles masses of the order of the galaxies they orbit; mass generation is instead by microscopic c -number expectation values of q -number fields, which vary only inside particles and so, being internal, do not act on galactic orbits.

We do not adjudicate that. What Theorem 7.3 contributes is the *implication*—if the mass-generating field is the macroscopic breaking field, then γ cannot act on orbits—computed here with the ansatz removed. The disagreement is therefore not about the geometry, the solution, or the fits. It is about the matter sector.

Remark 5.2 (The assumption sits one level below that). Theorem 7.3 begins “a point particle of mass m extremises $-\int m ds$ ”. That hypothesis is doing the work: the frame change is exact because the action is an integral along a *worldline*, so the conformal factor can be absorbed into the line element. An extended source has no such identity available.

Mannheim’s position denies precisely that hypothesis rather than the algebra that follows it: on his account the expectation values responsible for mass “only vary within particle interiors, giving particles an extended, baglike structure, as needed for localization in a conformal theory”. So the two positions do not disagree about a computation. They disagree about whether a massive test body in a theory with no fixed scale may be idealised as a point.

Two consequences worth stating, neither of which decides anything. First, the question is not settled by conformal invariance: a mass term needs a field of weight -1 , and a macroscopic c -number field and a microscopic q -number expectation value both supply one. Second, the two routes do not cost the same. Hobson and Lasenby’s requires only classical fields; Mannheim’s requires a quantum matter sector in which the breaking is spontaneous, so its assumption base is strictly larger. A

larger base is not a defect—the mechanism it buys is the one the theory needs—but in a ledger it is the sort of thing that should be entered rather than absorbed.

The hypothesis is moreover not merely to be granted or withheld: it has been derived. Mannheim [10] argues that since particles obey wave equations one is not free to *postulate* geodesic motion, and eikonalizes the equations of motion instead. Two of his conclusions bear here. For massive particles standard geodesic behaviour *is* recovered—so the worldline form of Theorem 7.3 is not the weak point—but “eikonalization in a theory with local conformal symmetry leads to trajectories that are only *globally* conformally symmetric”. A frame argument that moves trajectories by an r -dependent rescaling is a *local* conformal transformation, and on that reading it is the locality of the rescaling, not the point-particle idealisation, that is in question. We report this because it relocates the assumption once more.

We did check the step it turns on, and it holds. The eikonal approximation drops $-i\nabla_\mu\nabla^\mu T$, and that term is exactly what carries the local conformal invariance: under $S \rightarrow e^{-\alpha}S$, with $T \rightarrow T + i\alpha$ and the amplitude unchanged, the cross terms of $\nabla_\mu T \nabla^\mu T - i\nabla_\mu\nabla^\mu T - \frac{1}{6}R$ cancel in three pairs and the combination transforms homogeneously, while without that term a residual $2i\nabla T \cdot \nabla \alpha - 2\nabla \alpha \cdot \nabla \alpha - \nabla^2 \alpha$ survives. The algebra is correct.

That, however, is the end of what a computation settles here. Both sides agree the eikonal limit drops the term; what they disagree about is whether losing *local* conformal invariance in that limit disqualifies a frame argument built on an r -dependent rescaling. That is an inference about which idealisation is physically appropriate, not an identity, and no exact computation adjudicates it. We record where the disagreement sits and stop there.

Corollary 5.3 (Where the assumption lives). *On branch (B2), γ is determined by the scalar profile, which is a property of each configuration. Theorem 3.1 requires γ to be the same constant for every galaxy. Therefore the requirement transfers to the profile. By Corollary 5.1 this transfer is only about orbits on the half of (B2) where masses do not track the breaking scalar; on the other half γ does not act on orbits and the question of its universality does not arise.*

That is a question about the symmetry-breaking sector rather than about gravity. §7 carries out the transfer, and it turns out to land on one derived quantity rather than on the profile as a whole.

6 What the conformal gauge costs

Sections 2–5 work in the gauge $g_{tt}g_{rr} = -1$. Two things are needed to justify it.

Proposition 6.1 (Conformal covariance). *In $D = 4$, $B_{ab}[\Omega^2 g] = \Omega^{-2}B_{ab}[g]$, verified for the Mannheim–Kazanas family with symbolic parameters and unspecified Ω . Hence Bach flatness is a property of the conformal class, and the classification of §2 classifies classes.*

Proposition 6.2 (Reachability is a first-order ODE). *Setting the new radius to $\rho = \Omega r$ so the angular part remains ρ^2 , the gauge condition becomes*

$$r\Omega' = \Omega(\Omega\sqrt{ab} - 1),$$

first order in Ω , hence locally solvable wherever $ab > 0$ and $r \neq 0$.

What remains assumed is only global continuation—whether Ω stays positive and finite across the exterior. That is strictly smaller than the original assumption and is a question of analysis rather than algebra.

7 Carrying out the transfer

§6 and §2 between them force a third statement that neither makes on its own. Bach flatness is a property of a conformal class; the general solution in the gauge $g_{tt}g_{rr} = -1$ is four parameters subject to one constraint. So the conformal group must *act* on those four parameters.

Theorem 7.1 (The conformal action on the forced family). *Staying in the gauge is not a choice but an ODE, $d\hat{r}/dr = (\hat{r}/r)^2$, whose general solution is the Möbius map $\hat{r} = r/(1 - Cr)$ —one constant. The induced action is*

$$u \mapsto u, \quad w \mapsto w + 3uC, \quad \gamma \mapsto \gamma - 2wC - 3uC^2, \quad k \mapsto k + C\gamma - C^2w - C^3u,$$

a one-parameter group preserving $w^2 + 3u\gamma$ identically. From a member with $\gamma = 0$ it produces $\gamma = -2C - 3uC^2$.

So γ is not an invariant of the Bach vacuum. It is a coordinate on the conformal class, and the classification of §2 classifies classes precisely because of this.

A conformally coupled scalar has weight -1 , so the frame in which it is *constant* is reached by exactly this map with $\Omega = S/S_0$ —and Theorem 4.2 puts $\gamma = 0$ there. The transfer follows.

Corollary 7.2 (The transfer, in closed form). *On branch (B2) the profile realising the frame change is $S(r) = S_0/(1 - Cr)$ with $C = (S'/S)$ at the origin, and*

$$\gamma = -2(S'/S) - 3u(S'/S)^2 \quad \text{so that} \quad \gamma \simeq -2(S'/S).$$

Hence Theorem 3.1's demand that γ be one constant for every galaxy is exactly the demand that S'/S be one constant for every galaxy.

That is more specific than “is the profile universal?”. Only one derived quantity has to be universal, and it has to be so across galaxies whose masses differ by orders of magnitude, for a field sourced by the configuration.

Which frame matter follows

Both frames describe the same Bach vacuum, and only one carries a linear term. So the disagreement of [3, 4] is not about the vacuum solution and not about the algebra: it is about which frame matter follows. That premise is itself computable.

Theorem 7.3 (The mass scale is the conformal factor). *A point particle of mass m extremises $-\int m ds$. If the mass is generated by the symmetry-breaking scalar, $m = hS$, then*

$$-\int hS ds_g = -\int h\sqrt{-S^2 g_{ab} dx^a dx^b} = -\int h ds_{S^2g}$$

identically, so the trajectory is a geodesic of S^2g and not of g . Since S^2g is the frame in which S is constant up to the constant S_0^2 , such a particle follows geodesics of the frame with $\gamma = 0$.

Both branches are exhibited with their circular-orbit velocities, and both frames are confirmed Bach vacua, so this is a fork between two admissible solutions of one theory:

mass	trajectory follows	linear coefficient of v^2
m constant, independent of S	geodesics of g	$\gamma/2$
$m = hS$, generated by S	geodesics of S^2g	0

The first entry reproduces Theorem 3.1’s weak-field formula by a different route. The disagreement is not an artefact of the expansion: the exact circular-orbit velocities differ too.

A rotation-curve fit needs the first row. The second row is what the symmetry breaking was introduced to provide. Whether a matter sector can generate a mass scale by breaking conformal invariance while leaving particle masses independent of the breaking field is a question about that sector, and we do not answer it. What is no longer available is leaving the choice implicit.

The pair is a solution, and γ is independent of the mass

Theorem 7.1 produces a profile from a transformation law, and a transformation law does not by itself say the result solves anything.

Theorem 7.4 (Admissibility). *In the constant-scalar frame the conformally coupled action collapses to Einstein–Hilbert plus a cosmological term, so its stress tensor is a combination of G_{ab} and g_{ab} . The gravitational equation is $\alpha B_{ab} = \frac{1}{2}T_{ab}$ and that frame’s metric is a Bach vacuum, so consistency forces $T_{ab} = 0$ —for a constant scalar, the Einstein condition. The metric $1 - u/r - \hat{k}r^2$ is Einstein, with $R = 12\hat{k}$; equating with Theorem 4.2’s $R = -24\lambda S_0^2$ gives $\hat{k} = -2\lambda S_0^2$. Bach flatness and the Einstein condition then hold together on the same metric.*

The de Sitter term is the scalar potential, and the exterior pair solves the coupled system with nothing left over.

Corollary 7.5 (γ is not a function of the mass). *Theorem 7.1 gives $u \mapsto u$, so the Newtonian coefficient is a conformal invariant of the solution while γ is a coordinate on the class. Two frames with the same u carry different γ . Hence γ requires a second, independent datum—namely C —and for any target γ and any u there is a C attaining it.*

This changes the standing of Theorem 3.1. Requiring γ to be one constant across galaxies of widely differing mass looked like a coincidence in need of explanation; it is not one, because γ cannot depend on mass in any case. What the requirement says is that a single boundary datum on the symmetry-breaking sector is common to every galaxy.

Both readings of that are available and we prefer neither. Favourably, the universality is structurally natural rather than a fine-tuning of gravity. Unfavourably, the explanation of rotation curves is imported from the symmetry-breaking sector rather than produced by gravity.

8 What the source produces

§7 works entirely in the vacuum. Theorem 2.2 identified the governing operator; the multipole integrals it invites were not evaluated there.

Theorem 8.1 (The two moments). *Write the sourced equation as $\nabla^4 B = f$, with f proportional to $T^0_0 - T^r_r$ —the combination Theorem 4.1 already uses. From $\nabla^2 r = 2/r$ the biharmonic Green’s function is $-|x - x'|/8\pi$, whose angular average over a shell is $r + r'^2/3r$ outside and $r' + r^2/3r'$ inside. Hence for a compact spherically symmetric source, outside it,*

$$\gamma = -\frac{1}{2} \int f(r') r'^2 dr', \quad \beta = -\frac{1}{6} \int f(r') r'^4 dr'.$$

So γ is the *second* moment of the source—which is the total mass—and β the *fourth*. (We follow the literature’s convention, counting powers of r in the radial integrand.) With f tracking the

density, $\gamma \propto M$ while the Newtonian coefficient tracks $M\langle r^2 \rangle$. For a uniform ball, in closed form, $\gamma = -M/8\pi$ and $\beta/\gamma = R^2/5$; that the ratio is a length squared is what makes “ β is not a mass” a computation rather than a reading of the integrand. A shell and a ball of equal mass give the same γ and different β .

This resolves the fork left by Theorem 2.2 on the branch Theorem 3.1 flags: $\gamma \propto M$ gives $v^4 \propto M^2$. It does not end the programme, because Corollary 7.5 supplies the other term.

Remark 8.2 (Which branch this is on). Theorem 8.1 is computed for a perfect-fluid source with no scalar in it, so it lives on the branch where particle masses do *not* track the symmetry-breaking field. On the other branch Theorem 7.3 applies and γ does not act on orbits at all, which makes the moment analysis moot there. The two branches are therefore not alternative calculations of one quantity but calculations in different matter sectors, and we do not reconcile them.

Corollary 8.3 (The decomposition, and a tension we do not resolve). $\gamma = \gamma_0 + \gamma_*(M)$, with γ_0 a homogeneous datum and γ_* the sourced piece of Theorem 8.1. The superposition is a property of the linear equation $\nabla^4 B = f$; the identification of γ_0 with Corollary 7.5’s boundary datum imports the scalar sector, which Theorem 8.1 does not contain.

That decomposition is not ours: it is the parametrisation the published rotation-curve fits already use, with a universal global term and a local term proportional to the source [6]. And it sits awkwardly with our own Theorem 3.1, which says a mass-proportional piece moves the Tully–Fisher slope, whereas those fits retain both terms and report success over a large sample. We record that as an unresolved tension between a premise we adopt and a fit we do not evaluate. It is the sharpest thing we can say, and it is weaker than a verdict.

Remark 8.4 (A sign worth stating). Theorem 7.4 gives $\hat{k} = -2\lambda S_0^2$, so with $\lambda > 0$ the r^2 coefficient of B is $+2\lambda S_0^2$: the vacuum is anti-de Sitter. A cosmological term of the observed sign would need $\lambda < 0$, and then the scalar potential is unbounded below. We note the fork; we do not take it.

Separately, the fourth-moment result for β is the substance of the standing objection to conformal gravity’s Newtonian limit [3, 5], arrived at here from the source side. We exhibit it; we do not argue from it.

9 What this paper does not establish

- **No novelty in the physics, anywhere.** The Mannheim–Kazanas solution, its reduction to a fourth-order Poisson problem, and the trace-free condition on conformal matter are classical. So is the rest. Theorem 8.1 is Mannheim and Kazanas’ [5] and is the substance of the objection to the Newtonian limit. Theorem 7.3 is Horne’s [9], stated there to the level of the formula—rest masses scale with $S(r)/S_0$, so world lines follow geodesics of $\Omega^2 g$ with $\Omega = S/S_0$, which is also Theorem 7.1’s conformal factor. Theorems 7.3 and 7.4 are then Hobson and Lasenby’s [13]: that timelike geodesics are not conformally invariant, that the scalar generating particle mass puts massive particles on geodesics of the Schwarzschild–de Sitter metric, and that the resulting curves have no flat region—a *stronger* conclusion than Remark 8.2, which leaves the branch open. That massive particles follow the frame in which their mass is constant is standard scalar–tensor material. For Theorem 7.1 we have not found the group action on (w, u, γ, k) written in this form, which is not a claim that it is new. The contribution of this paper is that each statement is computed with the ansatz removed, the premises isolated, and controls that fail when a claim is vacuous—not the statements themselves. The reverse-physics stance is likewise not ours [7, 8]. What we have not found in the literature is the combination:

reverse physics executed as fail-closed machine-checked certificates, with dependency tags, controls designed to fail on vacuity, and corrections recorded as appended events rather than edits. That is a statement about our search, not a claim of novelty.

- **No fits.** No galaxy data enters anywhere. The only observational inputs are the *shape* of the Tully–Fisher relation, used as a premise in Theorem 3.1, and that galaxies are pressureless, used in §4. Neither is our measurement.
- **No interior solution, and no numbers.** Theorem 7.4 closes the exterior and Theorem 8.1 gives the exterior coefficients as functionals of the source, but the interior field and the scalar’s profile inside matter are not solved. Nor is it shown that a cosmological boundary condition *can* deliver a universal γ_0 . Above all, Corollary 8.3 is not evaluated: whether $\gamma_* \ll \gamma_0$ holds for real galaxies is observational and we do not measure it. The assumption is named and sharpened, not discharged.
- **Two matter descriptions, not reconciled.** Theorem 7.3 takes a mass generated by a Yukawa coupling, $m = hS$; Theorems 4.1 and 8.1 use a perfect fluid. Both are standard and both are entered as premises. Whether they are compatible is a matter-sector question we do not address.
- **No resolution of the dispute, and no prospect of one by these methods.** The exchange is live and long: Horne [9], then Hobson and Lasenby [13, 12], against Mannheim [4, 11], over more than a decade, with no consensus. Theorem 7.3 narrows it to one premise and computes both branches, which is all we do. We stress that the crux is *not algebraic*: both sides agree on the transformation law. They disagree over whether the mass-generating scalar is macroscopic or microscopic—whether a vacuum expectation value varies across a galaxy or only within particle interiors [11]. That is a physical premise, and exact symbolic computation with controls takes such premises as input. It cannot settle this one.
- **Spherical symmetry, for a claim about discs.** Everything is static and spherically symmetric. Galaxies are not, and the rotation-curve literature works the disc case for that reason. The gauge assumption is discharged as far as it can be in §6; the geometry assumption is not discharged at all.
- **Nothing about the ghost.** Conformal gravity’s fourth-order kinetic operator carries an Ostrogradsky ghost. That is orthogonal to everything here and is treated elsewhere in this programme.
- **No numerical value.** a_0 is not measured here and no published γ is compared; $\gamma = a_0/2c^2$ is an algebraic consequence of a premise, not a numerical agreement.

Prior art, and how hard we looked. Four claims in this line were withdrawn after the fact, each found by a search that should have preceded the work: the novelty of the frame and moment results, the novelty of the reverse-physics framing, and an earlier correction that named one prior source where there were two. All are recorded in the certificates as appended events rather than edits. What we searched for: conformal transformations of the Mannheim–Kazanas parameters, the Newtonian limit and its moment structure, the frame dependence of massive-particle trajectories under a mass-generating scalar, the status of the resulting dispute, and prior use of “reverse physics” as a programme. Every one returned prior work, and each is now cited at the point of use. We have not found the combination of that stance with fail-closed machine-checked certification, which is a statement about our search and not a claim about the literature.

10 Verification

Each result is a fail-closed certificate with a deterministic verifier. Controls are designed to fail if the claim were vacuous; several did so during development and are recorded in the certificates.

result	certificate	checks
Thm. 2.1	BHOB_GENERAL_STATIC_SPHERICAL_COMPLETENESS	24
Thm. 3.1	BHOC_TULLY_FISHER_SCALING	14
Thm. 2.2	BHOD_BACH_VACUUM_IS_BIHARMONIC	9
Thm. 4.1	BHOE_MATTER_TRACE_DILEMMA	8
Thm. 4.2	BHOF_CONSTANT_VEV_FORCES_GAMMA_ZERO	12
§6	BHOG_CONFORMAL_CLASS_INVARIANCE	10
Thm. 7.1	BHOH_UNIVERSALITY_TRANSFER	18
Thm. 7.3	BHOI_WHICH_FRAME_MATTER_FOLLOWS	13
Thm. 7.4	BHOJ_THE_ASSUMPTION_NAMED	12
Thm. 8.1	BHOK_THE_SOURCE_MOMENTS	22

Theorems 2.1, 4.2, 7.1, 7.3, 7.4 and 8.1, and §6, are computed in SYMPY; Theorems 2.2 and 4.1 are computed in Forge over exact rationals with symbolic differentiation of unspecified functions. Theorem 2.1’s conclusion carries an independent Forge rail using a different arithmetic stack, representation and code path; that rail confirms which metrics are Bach-flat but not the row decomposition itself, which remains single-implementation.

References

- [1] P. D. Mannheim and D. Kazanas, *Exact vacuum solution to conformal Weyl gravity and galactic rotation curves*, *Astrophys. J.* **342** (1989) 635.
- [2] R. J. Riegert, *Birkhoff’s theorem in conformal gravity*, *Phys. Rev. Lett.* **53** (1984) 315.
- [3] É. É. Flanagan, *Fourth order Weyl gravity*, *Phys. Rev. D* **74** (2006) 023002.
- [4] P. D. Mannheim, *Alternatives to dark matter and dark energy*, *Prog. Part. Nucl. Phys.* **56** (2006) 340.
- [5] P. D. Mannheim and D. Kazanas, *Newtonian limit of conformal gravity and the lack of necessity of the second order Poisson equation*, *Gen. Rel. Grav.* **26** (1994) 337.
- [6] P. D. Mannheim and J. G. O’Brien, *Galactic rotation curves in conformal gravity*, *J. Phys. Conf. Ser.* **437** (2013) 012002, [arXiv:1211.0188](#).
- [7] G. Carcassi and C. A. Aidala, *Reverse Physics: From Laws to Physical Assumptions*, *Found. Phys.* **52** (2022) 40, [arXiv:2111.09107](#).
- [8] G. Carcassi and C. A. Aidala, *Assumptions of Physics*, Michigan Publishing, 2021.
- [9] K. Horne, *Conformal Gravity rotation curves with a conformal Higgs halo*, *Mon. Not. Roy. Astron. Soc.* **458** (2016) 4122.
- [10] P. D. Mannheim, *Critique of the use of geodesics in astrophysics and cosmology*, *Class. Quantum Grav.* **39** (2022) 245001, [arXiv:2105.08556](#).

- [11] P. D. Mannheim, *Structure of conformal gravity in the presence of a scale breaking scalar field*, Gen. Rel. Grav. **54** (2022) 39, [arXiv:2105.14679](#).
- [12] M. P. Hobson and A. N. Lasenby, *Conformally-rescaled Schwarzschild metrics do not predict flat galaxy rotation curves*, Eur. Phys. J. C **82** (2022) 585.
- [13] M. P. Hobson and A. N. Lasenby, *Conformal gravity does not predict flat galaxy rotation curves*, Phys. Rev. D **104** (2021) 064014, [arXiv:2103.13451](#).
- [14] M. P. Hobson and A. N. Lasenby, *Schwarzschild and linear potentials in Mannheim’s model of conformal gravity*, Mon. Not. Roy. Astron. Soc. **478** (2018) 2827, [arXiv:1805.06326](#).
- [15] J. G. O’Brien, T. L. Chiarelli and P. D. Mannheim, *Universal properties of galactic rotation curves and a first principles derivation of the Tully–Fisher relation*, Phys. Lett. B **782** (2018) 433, [arXiv:1704.03921](#).
- [16] M. Milgrom, *A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis*, Astrophys. J. **270** (1983) 365.