
TASK 1



Geological & Surface Features	Identification & Reasoning	Source
1. **Dune-like sand accumulation**	The surface has a broad curved bedform with fine, repeated textures. Its shape and surface patterns depict dunes or ripples formed by the wind due to loose particle transportation.	NASA, The Draa of Mars
2. **Dune-slope grooves**	Narrow grooves run down the sandy right-hand sandy surface. Their direction suggests gravity-driven sediment. Water may not be required, as these gullies can form through sediment	HiRISE, Gullies on the Dunes of Russell Crater

		avalanches or CO_2 frost activity.	
3.	**Curved sediment lobes**	Curved, elongated narrow margins extend toward the lower-right side of the sandy slope. They may represent redistributed sediment or small deposits associated with downslope movement.	HiRISE, Frosty Sand Dunes of Mars NASA, MRO Spots Sand Stripes
4.	**Aeolian ridges (on the right)**	The lower-right surface contains multiple curved ridges with similar orientations. Repeated ridge patterns are typical of wind-shaped sediment. They may be ripples or transverse aeolian ridges (TARs).	HiRISE, Frosty Sand Dunes of Mars NASA, MRO Spots Sand Stripes
5.	**Aeolian ridges (on the left)**	Several long, narrow, roughly parallel ridges cross the smooth surface. Their systematic arrangement suggests wind-related formation rather than randomly distributed rocks. They could be depositional ridges or wind- eroded remnants.	NASA, Yardangs in Aeolis
6.	**Large erosional hollow**	A broad, irregular, darker depression-like area dominates the left-central region. Its steep-looking boundary and uneven surface suggest significant removal or redistribution of material.	NASA, Craters in Fretted Terrain
7.	**Narrow trough-like depression**	A dark, elongated zone extends downward from the upper center. Its position between rougher surrounding surfaces is consistent with a narrow topographic low. It may reflect erosion or structural	HiRISE, Gully Landforms in Aram Chaos

	differences in the terrain, but its exact origin is unclear.	
8. **Exposed rocky outcrop/scarp**	The center contains a sharply bounded, angular, bright-looking rocky feature adjacent to a dark area. This contrast is consistent with an exposed steep rock face and its shadow. Erosion can expose relatively resistant material beneath surface sediment.	NASA, Exposing the Rock in Impact Craters
9. **Possible slope debris/talus**	Rough, irregular material occurs near the base of the central rocky slope. Material produced by rockfalls or slope erosion often accumulates at the foot of steep terrain.	HiRISE, Gully Landforms in Aram Chaos
10. **Smooth sediment mantle**	A relatively uniform, smooth surface occupies much of the center-right. It contrasts with adjacent rougher rock and may indicate that fine sediment or sand covers underlying terrain. Smoothness supports a mantling interpretation.	NASA, Knobby Terrain in Northern Arabia Terra
11. **Small circular depression / possible crater**	A small circular feature is visible near the right-central lower slope. Circularity is consistent with an impact crater. It could also be due to some impact.	NASA, A Fresh Impact Crater with an Odd Shape
12. **Rough rocky slope/eroded bedrock**	The lower-right-central area contains broken-looking, irregular terrain with prominent textural contrasts. These may represent exposed bedrock, weathered surfaces, or an eroded scarp. The roughness distinguishes it	NASA, Wind Erosion in Aeolis

	from adjacent smooth sediment.	
13. Isolated rocky remnant	A small, rough, raised-looking feature stands apart from the smoother left-hand terrain. It may be a resistant rock remnant left after surrounding material was removed. Without topographic measurements, it could also be a small block or other surface feature.	NASA, Yardangs in Aeolis
14. Scattered small pits	Several small dark circular marks occur across the upper-central smooth area. Their clustering suggests a pitted surface. Some could be small impact craters, but pits can also form through collapse, erosion, or volatile-related processes.	NASA, Pitted Material from Tooting Crater
15. Upper-right surface lineations	Narrow irregular bands appear near the upper-right boundary. They could be erosional grooves, exposed layers, or differences in surface texture. Similar patterns on Mars sometimes reflect sedimentary layering subsequently exposed by erosion.	HiRISE, Sedimentary Layers in Columbus Crater
16. Small rounded depressions	Several small rounded marks appear near the bottom-left-central region. Some resemble shallow, degraded craters, but their weak boundaries make specific origins uncertain. Older impact craters can become partially buried or eroded until only subdued depressions remain.	NASA, Craters in Fretted Terrain

17. Short erosional furrows	Closely spaced grooves occur beside the central rocky outcrop and adjacent sandy-looking region. Their repeated downslope arrangement may indicate local erosion or shallow sediment movement.	HiRISE, Frosty Sand Dunes of Mars
18. Irregular upper-left scarp/possible degraded crater margin	The upper-left area encloses a curved boundary surrounding uneven terrain. This may represent an eroded scarp or a remnant of an older circular depression. Its incomplete shape does not justify classifying it as a confirmed impact crater.	NASA, Craters in Fretted Terrain
19. Curved erosional ridges/possible exposed layers	The center-left circled area contains concentric-looking curved lines and a relatively smooth interior. The pattern could represent erosion of materials with different resistance, or curved surface ridges partly covered by sediment.	HiRISE, Sedimentary Layers in Columbus Crater
20. Upper-right rough-textured surface	The larger upper-right red area covers terrain that is more irregular and granular-looking than the nearby smooth dune-like area. This may be exposed substrate, a partially eroded sediment mantle, or rough wind-modified terrain.	NASA, Wind Erosion in Aeolis
21. Lower-center irregular grooves and pits	The lower-center circled area contains branching-looking dark lines, small pits, and uneven surfaces. These could be fractures in exposed material, erosion grooves, or shadowed sediment boundaries.	HiRISE, Gully Landforms in Aram Chaos

22. Rounded dark slope/possible erosional remnant	The bottom circle-like region has a rounded boundary and darker interior. It could be a shadowed slope, partly buried mound, or eroded terrain feature.	NASA, Knobby Terrain in Northern Arabia Terra
23. Isolated rough patch/possible resistant outcrop	The middle-right red area highlights an irregular rocky-looking patch standing apart from smoother surfaces. It may represent exposed resistant material or a local erosional remnant. Its different texture suggests different erosion or surface-cover conditions.	NASA, Yardangs in Arsinoes Chaos, Mars


TASK 2

A:


1. Baeyer Test — Detection of Unsaturation

Purpose: To identify carbon–carbon double bonds (C=C) or triple bonds (C≡C), which may occur in unsaturated organic compounds, including certain fatty acids and oils.

Scientific principle: Dilute alkaline potassium permanganate is purple. When it reacts with many unsaturated organic compounds, the permanganate is reduced, producing brown manganese dioxide (MnO_2). The disappearance of purple accompanied by brown precipitate formation supports a positive result. However, aldehydes and several other oxidizable compounds can interfere, making this a test for chemical reactivity rather than a specific lipid test.



Initial: Purple ($KMnO_4$ solution)



Positive: Brown (MnO_2 precipitate)

Reagents required: Dilute potassium permanganate solution (approximately 0.5–1%), a suitably prepared soluble sample, water or a compatible solvent, and dilute sodium carbonate or another approved alkaline medium if needed.

Procedure (approximately 5–10 minutes):

1. Place approximately 1 mL of a suitable dissolved sample in a clean test tube.
2. Add 1–3 drops of dilute alkaline $KMnO_4$ solution.
3. Mix gently and observe for a few minutes.
4. A rapid loss of purple color accompanied by brown precipitate formation suggests a positive reaction.

The sample and reagent should be tested against positive and negative controls. Oils may require a validated solvent system because they do not mix readily with water.

Chemical reaction: For an alkene, a simplified balanced reaction under cold, dilute alkaline conditions is:

$$3RCH = CHR' + 2KMnO_4 + 4H_2O \rightarrow 3RCH(OH)CH(OH)R' + 2MnO_2 \downarrow + 2KOH$$

The double bond is oxidized to a vicinal diol, while manganese is reduced from its +7 oxidation state to +4.

Safety precautions: Wear gloves, goggles, and a lab coat. Potassium permanganate is an oxidizer and can stain skin and clothing. Keep it away from combustible materials and dispose of manganese-containing waste according to laboratory procedures.

Assessment: Suitable for demonstrating unsaturation, but not recommended as the primary method for identifying lipids.


SOURCE:
https://chem.libretexts.org/Ancillary_Materials/Laboratory_Experiments/Wet_Lab_Experiments/Organic_Chemistry_Labs/Experiments/8%3A_Identification_of_Unknowns_%28Experiment%29

2. Biuret Test — Detection of Proteins


Purpose: To identify proteins and sufficiently long peptides by detecting peptide bonds.

Scientific principle: In alkaline conditions, copper(II) ions form a violet coordination complex with peptide bonds. The solution changes from blue to violet when sufficient

peptide bonds are present. Free amino acids generally do not produce the characteristic reaction.



Negative: Blue (Biuret reagent)



Positive: Violet (Copper–peptide complex)

Reagents required: Sodium hydroxide (NaOH), copper(II) sulfate ($CuSO_4$), and the test sample. Prepared Biuret reagent may also contain sodium potassium tartrate, which helps stabilize dissolved copper ions.

Procedure (approximately 5–10 minutes):

1. Add 2 mL of sample solution to a clean test tube.
2. Add 3 mL of 10% NaOH and mix.
3. Add two drops of 2% $CuSO_4$ solution.
4. Mix gently and observe the resulting color.
5. A violet or lilac color indicates a positive reaction.

Chemical reaction: This is a coordination-complex formation process, rather than a simple precipitation reaction.

$$\mathrm{Cu^{2+}} + \text{peptide ligands} \xrightarrow{\text{alkaline pH}} \text{violet Cu(II)} - \text{peptide complex}$$

The peptide nitrogen atoms coordinate with copper ions, creating the characteristic color.

Safety precautions: NaOH is corrosive; $CuSO_4$ is harmful if swallowed and hazardous to aquatic life. Avoid skin and eye contact. Collect copper-containing waste separately.

Assessment: One of the strongest tests to retain. It is inexpensive, reasonably fast, visually clear, and directly relevant to protein detection. However, a positive Biuret test does not prove that an enzyme is active.

SOURCE:
https://chem.libretexts.org/Courses/Los_Medanos_College/Chemistry_6_and_Chemi

stry_7_Combined_Laboratory_Manual_%28Los_Medanos_College%29/01%3A_Experiments/1.29%3A_Experiment_729_Qualitative_Testing_of_Amino_Acids_and_Proteins_1_2


https://bio.libretexts.org/Bookshelves/Biotechnology/Bio-OER_%28CUNY%29/02%3A_Chemistry/2.09%3A_Proteins

3. Carbohydrate Test — Molisch's Test

Because the task does not specify which carbohydrate test ARC used**, Molisch's test** is selected here as a general carbohydrate-detection method. This must be confirmed before describing it as ARC's actual previous procedure.

Purpose: To detect many carbohydrates, including monosaccharides, disaccharides, and polysaccharides.

Scientific principle: Concentrated sulfuric acid dehydrates sugars, producing furfural or 5-hydroxymethylfurfural. These products react with α-naphthol to form a violet-colored condensation product at the interface of the two liquid layers.



Negative: No Ring



Positive: Violet Ring

Reagents required: Molisch's reagent (α-naphthol dissolved in ethanol), concentrated sulfuric acid (H_2SO_4), and the aqueous test sample.

Procedure (approximately 5–10 minutes):

1. Add approximately 1 mL of sample solution to a test tube.
2. Add two drops of Molisch's reagent.
3. Under trained supervision, carefully introduce approximately 1 mL of concentrated sulfuric acid down the inner wall to form a separate lower layer.
4. Observe the boundary between the two layers without shaking.
5. A violet ring at the interface indicates a positive result.

Chemical Reactions:

Hexose dehydration:

$$C_6H_{12}O_6 \xrightarrow{\text{conc. } H_2SO_4} C_6H_6O_3 + 3H_2O$$

The product, 5-hydroxymethylfurfural, then reacts with α-naphthol to form violet condensation products. The second stage is a complex reaction and is not represented by a single simple stoichiometric equation.

Safety precautions: Concentrated sulfuric acid is highly corrosive and its mixing with water releases substantial heat. This test should only be carried out by trained laboratory personnel using appropriate eye, face, skin, and chemical-resistant protection, ideally in a fume hood. It is unsuitable for unsupervised student stations.

Assessment: Chemically useful for broad carbohydrate detection, but its use of concentrated sulfuric acid makes it a relatively poor choice for an open, time-constrained student competition.

SOURCE:
https://chem.libretexts.org/Courses/Triton_College/Elementary_Organic_Chemistry_%28Lab_Manual%29/09%3A_Lab_9-_Tests_for_Carbohydrates

4. Carbonate Test — Detection of Carbonate Ions

Purpose: To detect carbonate (CO_3^{2-}) and, under suitable conditions, bicarbonate (HCO_3^-) ions in a sample.

Scientific principle: Carbonates react with dilute acids and release carbon dioxide. The released CO_2 can be passed through limewater, where it forms insoluble calcium carbonate and turns the limewater milky.

Reagents required: Dilute hydrochloric acid (HCl), limewater [aqueous $Ca(OH)_2$], and a carbonate-containing solid or solution.

Procedure (approximately 5–10 minutes):

1. Place approximately 1–2 mL of sample solution, or a small solid sample, in a test tube.
2. Add a few drops of dilute hydrochloric acid.
3. Observe whether gas bubbles develop.
4. Transfer the gas into a second test tube containing approximately 1 mL of limewater using appropriate laboratory apparatus.
5. If the limewater becomes cloudy or milky, CO_2 has been detected.

Chemical Reactions:

Carbonate reacting with acid:

$$CaCO_3 + 2HCl \rightarrow CaCl_2 + CO_2 \uparrow + H_2O$$

CO_2 reacting with limewater:

$$CO_2 + Ca(OH)_2 \rightarrow CaCO_3 \downarrow + H_2O$$

The insoluble $CaCO_3$ forms the visible white suspension.

Safety precautions: Use dilute acid, wear eye protection, and never seal the reaction vessel because CO_2 production can increase pressure. Do not directly inhale the evolved gas. Use proper gas-transfer apparatus.

Assessment: Fast, low-cost, and chemically informative. However, carbonate ions are inorganic species, so this test does not directly identify proteins, carbohydrates, lipids, or enzymes.

SOURCE: https://edu.rsc.org/experiments/testing-for-negative-ions/758.article

5. Sulfate Test — Detection of Sulfate Ions

Purpose: To identify sulfate ions (SO_4^{2-}) in an aqueous solution.

Scientific principle: Barium ions react with sulfate ions to form barium sulfate, a highly insoluble white precipitate. Acidification reduces interference from carbonate ions, which could otherwise also form a white barium precipitate.

Reagents required: Dilute hydrochloric acid and barium chloride ($BaCl_2$) solution, typically around 0.1 M for classroom qualitative testing.

Procedure (approximately 5–10 minutes):

1. Place about 2 mL of sample solution in a clean test tube.
2. Add a few drops of dilute HCl.
3. Add a few drops of barium chloride solution.
4. Mix gently.
5. The formation of a persistent white precipitate indicates sulfate ions, provided relevant interferences have been excluded.

Chemical Reactions:

$$BaCl_2(aq) + Na_2SO_4(aq) \rightarrow BaSO_4(s) \downarrow + 2NaCl(aq)$$

Net ionic equation:

$$Ba^{2+}(aq) + SO_4^{2-}(aq) \rightarrow BaSO_4(s)$$

Safety precautions: Soluble barium salts are toxic if swallowed and must not be handled without appropriate precautions. Avoid exposure to skin and eyes. Collect barium-containing waste for proper disposal rather than pouring it down the sink.

Assessment: Fast, chemically selective under properly controlled conditions, and easy to interpret visually. However, it does not directly detect the four biological targets identified in the assignment.

SOURCE: https://edu.rsc.org/experiments/testing-salts-for-anions-and-cations/464.article

Approximate Cost of the Five Initial ARC Chemical Tests

Chemical Test	Reagents Considered	Source of Chemical Prices
Baeyer Test	Potassium permanganate ($KMnO_4$)	500 g, ৳2,100
Biuret Test	Biuret reagent ($CuSO_4$ and NaOH mixture)	500 mL, ৳600
Carbohydrate Test (Molisch's)	Molisch's reagent and concentrated H_2SO_4	**Molisch's reagent** (100 mL, ৳900); Sulfuric acid (2.5 L, ৳2,600)
Carbonate Test	Dilute HCl and limewater [$Ca(OH)_2$]	HCl (500 mL, ৳600); $Ca(OH)_2$ (500 g, ৳900)
Sulfate Test	Barium chloride ($BaCl_2$) and dilute HCl	$BaCl_2$ (500 g, ৳1,100); HCl (500 mL, ৳600)

B:

The comparison considers speed, experimental simplicity, reagent hazards, accuracy of interpretation, and relevance to the intended biological targets.

Test	Advantages	Disadvantages
Baeyer	Rapid color change; inexpensive reagent; demonstrates oxidation and	Not specific to lipids; other reducing compounds may react; aqueous permanganate and oil samples may not

	carbon–carbon unsaturation; useful for teaching organic reactions.	mix properly; produces manganese-containing waste.
Biuret	Clear violet positive result; simple procedure; no heating necessary; direct relevance to proteins; widely established in teaching laboratories.	Less sensitive to low protein concentrations; does not detect free amino acids reliably; cannot distinguish active enzymes from inactive proteins; alkaline reagent is hazardous.
Molisch	Broad carbohydrate coverage; sensitive; quick reaction; detects many mono-, di-, and polysaccharides.	Concentrated sulfuric acid presents a serious handling hazard; poor specificity for carbohydrate classes; requires careful layering; unsuitable for independent beginner handling.
Carbonate	Inexpensive; simple and visible effervescence; clear supporting confirmation with limewater; good for introductory inorganic chemistry.	Requires gas identification for strong evidence; other gas-evolving reactions may interfere; needs additional apparatus; unrelated to the biomolecule objective.
Sulfate	Rapid white-precipitate formation; established qualitative test; low sample requirement; clear ionic reaction.	Soluble barium compounds are toxic; some other ions may interfere; produces hazardous waste; unrelated to direct biomolecule detection.

Note:

Biuret is the most relevant of the five tests for the stated biological objective. It identifies proteins through coordination chemistry, produces an easily observable color change, and requires little equipment. Its limitation is that it measures the presence of peptide-containing material, not whether an enzyme can still catalyze a reaction.

Baeyer's test has a different problem: it detects a chemical property, unsaturation, rather than a biomolecule class. Some lipids contain carbon–carbon double bonds, but saturated lipids do not. Conversely, many substances that are not lipids contain double bonds or other groups that may react with permanganate. Therefore, it cannot be treated as a general lipid test.

Molisch's test covers a useful range of carbohydrates, but the need for concentrated sulfuric acid creates a substantial safety disadvantage. This is particularly important for ARC if participants conduct the procedures independently.

The carbonate and sulfate tests work well as introductory inorganic chemistry experiments. Their main limitation is not experimental performance but their lack of alignment with the stated objective. If ARC is intended to cover both inorganic qualitative analysis and biomolecule identification, these tests still have educational value. If the objective is exclusively biological molecule testing, their inclusion needs reconsideration.

REFERENCE:

https://edu.rsc.org/experiments/testing-for-negative-ions/758.article

https://edu.rsc.org/experiments/testing-salts-for-anions-and-cations/464.article

https://chem.libretexts.org/Courses/Triton_College/Elementary_Organic_Chemistry_%28Lab_Manual%29/09%253A_Lab_9-_Tests_for_Carbohydrates

C:

A better testing set should not be selected simply because the reaction is faster or more colorful. The replacement must detect the intended substance more directly, produce an interpretable result, and be manageable within the equipment and safety constraints.

1. Recommended alternative tests

Alternative test	Reagents and procedure	Reaction / positive result	Advantages and limitations	Better than existing test?
Benedict's test	Mix approximately 1 mL sample with 2 mL Benedict's reagent; heat in a water bath for around 3–5 minutes.	Reducing sugars reduce Cu^{2+} to Cu_2O, producing green, yellow, orange or brick-red results, depending partly on concentration.	Visually engaging and useful for glucose and other reducing sugars. However, sucrose and starch generally require additional treatment to test positive, and other reducing substances may interfere.	Better than Molisch for routine reducing-sugar demonstrations, but not for detecting all carbohydrates.

Iodine test	Add 2–3 drops of iodine–potassium iodide solution to approximately 2 mL of sample.	Iodine/polyiodide species interact with starch helices, producing a blue-black color.	Very quick, simple and suitable for student testing. Does not detect sugars generally.	Better for safe, rapid starch detection, but narrower than Molisch.
Ethanol emulsion test	Mix a small sample with approximately 2 mL ethanol, then add the extract to around 2 mL water.	Lipids that dissolve in ethanol form fine suspended droplets when diluted with water, creating a milky emulsion. This is a physical process, not a chemical reaction.	Simple and useful for many fats and oils. Requires flammable ethanol, and some complex samples can produce misleading cloudiness.	Better than Baeyer for general fat/oil detection.
Sudan III test	Add a few drops of Sudan III stain to a suitable lipid-containing sample and mix.	The fat-soluble dye partitions preferentially into the lipid phase, making it appear red-orange.	Direct visual identification of lipid-rich regions. Dye handling, staining and solvent formulation are limitations.	Better than Baeyer for lipid visualization.
Catalase activity test	Combine an approved catalase-containing preparation with approximately 1–2 mL of 3% hydrogen peroxide.	$2H_2O_2 \xrightarrow{\text{catalase}} 2H_2O + O_2 \uparrow$. Oxygen bubbles indicate catalase activity under suitable controlled conditions.	Very fast, visible and directly tests enzyme activity. However, some nonenzymatic substances can also accelerate peroxide decomposition.	Better than Biuret for demonstrating activity of catalase, but not for detecting proteins generally.
Amylase–starch test	Combine starch solution and amylase, incubate under controlled conditions, and test small aliquots with	Amylase hydrolyzes starch into shorter carbohydrates; decreasing blue-black iodine–starch color suggests starch breakdown.	Demonstrates enzyme action and can investigate temperature or pH effects. Requires control experiments and careful interpretation.	Better for demonstrating amylase activity, but slower and more condition-sensitive than catalase.

iodine at intervals.

SOURCE:

https://chem.libretexts.org/Courses/Triton_College/Elementary_Organic_Chemistry_%28Lab_Manual%29/09%253A_Lab_9-_Tests_for_Carbohydrates

https://www.fda.gov/food/laboratory-methods-food/bam-r12-catalase-test

https://pubs.acs.org/jceda8/article/103/3/1648/5081497/Rethinking-the-Iodine-Reaction-Methodology-in-the

2. Which tests are better for ARC?

Recommended testing set

All planned for under 30 minutes

Biuret test	Protein identification	Retain
Benedict's test	Reducing sugars	Add
Iodine test	Starch	Add
Ethanol emulsion test	Fats and oils	Add
Catalase activity test	Active catalase	Add

This arrangement would cover proteins, reducing sugars, starch, lipids and one specific active enzyme. It is a more direct match for the objective than the original five tests.

All five can be designed around sample volumes below 20 mL and individual reaction times of less than 30 minutes. This assumes reagents are prepared in advance, suitable controls are available, and any heating equipment is ready.

The revised set has two important limitations. First, Benedict's and iodine tests do not together identify every possible carbohydrate. Second, catalase testing establishes activity of a particular enzyme rather than general detection of all enzymes. These limitations should be explained in the final proposal.

3. Scientific and practical recommendations

For an introductory ARC activity, I would retain Biuret, replace Baeyer with a lipid-specific test, replace Molisch with safer carbohydrate tests where suitable, and introduce a direct enzyme activity experiment.

Benedict's test and the iodine test complement rather than replace each other: the former detects reducing sugars, whereas the latter detects starch. The ethanol emulsion test is more relevant to general lipid screening than Baeyer's unsaturation test. The catalase test provides the enzyme activity component that is entirely missing from the existing list.

Carbonate and sulfate tests should not automatically be discarded. If ARC also evaluates inorganic analytical chemistry, they remain useful. But they should be treated as a separate test category rather than presented as methods for identifying the four specified biological targets.

For reliable operation, each station should use a verified positive control, a negative control, a clearly defined positive observation, and a disposal procedure. Reaction time must be distinguished from total station time, including preparation and cleanup. In addition, testing should use approved laboratory samples rather than unknown biological or environmental materials.

Task 3

01. What is the main scientific purpose of a Mars rover?

The main scientific purpose of a Mars rover is to explore the Martian surface and collect information about the planet's geology, climate, and past environmental conditions. One of the most important questions is whether Mars could have supported life in the past.

As explained in the video, NASA's Curiosity rover was designed to investigate whether Mars once had the conditions necessary for microbial life. It uses cameras, a robotic arm, a drill, and scientific instruments to examine rocks, soil, and the atmosphere. By studying their chemical composition, scientists can identify materials that may have helped support life.

However, Curiosity was not designed to detect life directly. Its main goal was to investigate whether the necessary ingredients and environmental conditions for life had ever existed on Mars.

02. What does “habitability” mean, and what conditions would make an ancient Martian environment potentially habitable?

Habitability means the ability of an environment to support life. In the context of Mars, it refers to whether the planet once had suitable conditions for microorganisms, such as tiny single-celled organisms, to survive.

According to the video, three essential requirements are:

1. Liquid water: Life as we know it depends on water for important biological and chemical processes.
2. A source of energy: Microorganisms need energy to carry out activities necessary for survival and growth.
3. A source of carbon: Carbon is a fundamental building block of organic molecules, which are essential components of living organisms.

Other factors, such as suitable temperatures, a usable supply of nutrients, and environmental conditions that are not excessively harmful, would also influence habitability.

If ancient Mars had liquid water, available energy, carbon, and other necessary nutrients, some environments could have supported microbial life. However, a habitable environment does not necessarily mean that life actually existed there.

03. Why is evidence of ancient water important to Mars scientists?

Evidence of ancient water is important because water is one of the main requirements for life as we know it. Finding signs that liquid water once existed on Mars helps scientists determine whether the planet may have supported living organisms in the past.

Water can also change the Martian landscape. For example, flowing water can transport sediment, shape valleys, and deposit layers of material as we can see in the image of task 1. It can also react with rocks and form minerals that provide clues about past environmental conditions.

By examining these features, scientists can understand how Mars changed over time and identify places that may once have been suitable for microbial life.

Therefore, evidence of ancient water helps scientists investigate both the possibility of past life and the geological history of Mars. However, water alone is not proof that life existed.

04. Why are rocks and their layers useful for reconstructing the history of Mars?

Rocks and their layers are useful because they preserve evidence of the environmental conditions and geological processes that shaped Mars over time.

In the video, John Grotzinger compares studying rock layers to turning the pages of a book. Each layer may contain clues about the period when it formed, including the presence of water, sediment deposition, and changes in environmental conditions.

For example, sedimentary layers may have formed when particles accumulated in ancient lakes, rivers, or wind-shaped environments. By examining their composition, structure, and arrangement, scientists can understand how the Martian environment changed over time.

This is why Curiosity was sent to Gale Crater, where Mount Sharp contains a large sequence of rock layers. Studying these layers allows scientists to investigate whether Mars experienced habitable conditions in the past and how those conditions changed.

References

The answers above are based mainly on the provided NASA/JPL link. The following official NASA sources provide additional scientific support:

https://science.nasa.gov/mission/msl-curiosity/science/

https://science.nasa.gov/wp-content/uploads/2024/03/44709_MSLLithoSet2013.pdf

https://science.nasa.gov/mission/msl-curiosity/science-highlights/

https://www.nasa.gov/news-release/nasas-curiosity-rover-finds-clues-to-how-water-helped-shape-martian-landscape/

05. What is a biosignature, and why would scientists be interested in finding one?

A biosignature is a chemical substance, physical structure, or pattern that may indicate the presence of past or present life. Examples include certain organic molecules, unusual mineral patterns, or structures in rocks that might have been formed by microorganisms.

As Katie Stack Morgan explains in the transcript, NASA's Perseverance rover studies Martian rocks to search for possible signs of ancient life. Scientists are interested in biosignatures because they could help answer one of the biggest questions about Mars: Did life ever exist on the planet?

However, finding a possible biosignature does not automatically confirm life. Some similar features can form through non-biological processes. That is why scientists want to bring Martian rock samples back to Earth, where advanced laboratory equipment can help determine whether life was actually involved in forming them.

06. Why might an ancient river delta be a good place to search for evidence of past life?

An ancient river delta is a good place to search for evidence of past life because deltas form where flowing water carries and deposits sediments, often where a river enters a lake or another body of water. These sediments can trap and preserve organic matter and other possible traces of life.

In the transcript, Katie Stack Morgan explains that Perseverance landed at Jezero Crater because it contains a well-preserved ancient river delta. This suggests that water once flowed through the area and deposited layers of sediment.

Fine-grained sediments are particularly important because they can sometimes protect organic material from being destroyed or broken down. Over time, these sediments may harden into sedimentary rocks, preserving clues about the ancient environment.

Therefore, studying the rocks of Jezero Crater's delta could help scientists understand whether Mars once had conditions suitable for life and whether any evidence of that life has been preserved.

07. Why doesn't finding an organic molecule automatically mean that life existed on Mars?

Finding an organic molecule on Mars does not automatically prove that life existed there because organic molecules can form through both biological and non-biological processes. Although organic molecules contain carbon and are important building blocks of life, they can also be produced through chemical reactions that do not involve living organisms. Some may even arrive on Mars through meteorites.

In the transcript, Katie Stack Morgan explains that scientists look for organic matter preserved in rocks, especially in ancient river deltas. However, discovering such material is only a possible clue, not direct proof of life. Scientists must examine its chemical composition, surrounding minerals, and geological environment to understand how it formed.

This is one reason Perseverance collects rock samples for possible return to Earth. More advanced laboratory analysis could help scientists determine whether organic molecules were produced by living organisms or natural chemical processes.

08. Why does the Science Team need to study a rock's geological context before interpreting an interesting observation?

The Science Team needs to study a rock's geological context because the location, surrounding rocks, and conditions under which a rock formed can help explain what an unusual observation actually means.

For example, finding an organic molecule or an unusual pattern in a rock might be interesting, but scientists cannot immediately assume that it is evidence of ancient life. They first need to understand whether the rock formed in a river, lake, volcanic environment, or somewhere else. They also examine its layers, minerals, and signs of later geological changes that may have affected the original material.

As Katie Stack Morgan explains in the transcript, even individual sand grains can provide information about Mars and its geological evolution. By studying the complete geological setting, scientists can distinguish between features that might have been produced by life and those formed through ordinary geological or chemical processes.

Therefore, geological context helps scientists make more reliable interpretations and avoid drawing incorrect conclusions from a single observation. NASA also explains that studying rock layers and their relationships is essential for reconstructing the environmental history of Jezero Crater.

References

https://science.nasa.gov/mission/mars-2020-perseverance/science-objectives/

https://science.nasa.gov/mission/mars-2020-perseverance/science-highlights/

https://science.nasa.gov/blog/stratigraphic-layers/

https://science.nasa.gov/resource/perseverance-explores-the-jezero-crater-delta/

https://science.nasa.gov/blogs/science-news/2026/02/06/nasa-study-non-biologic-processes-dont-fully-explain-mars-organics/