

Early disease and low baseline damage predicts response to belimumab in patients with systemic lupus erythematosus

Gatto M, Saccon F, Zen M, Regola F, Fredi M, Andreoli L, Tincani A, Urban ML, Emmi G, Ceccarelli F, Conti F, Bortoluzzi A, Govoni M, Tani C, Mosca M, Ubiali T, Gerosa M, Bozzolo E, Canti V, Cardinaletti P, Gabrielli A, Tanti G, Gremese E, De Marchi G, De Vita S, Fasano S, Ciccia F, Pazzola G, Salvarani C, Negrini S, Puppo F, Di Matteo A, De Angelis R, Orsolini G, Rossini M, Faggioli P, Laria A, Piga M, Mathieu A, Scarpato S, Rossi FW, de Paulis A, Brunetta E, Ceribelli A, Selmi C, Prete M, Racanelli V, Vacca A, Bartoloni E, Gerli R, Larosa M, Iaccarino L, Doria A

Arthritis Rheumatol. 2020 Apr 10. doi: 10.1002/art.41253. [Epub ahead of print]

PM-ES-BEL-EDTL-200007 (v1) 07/2020

(▼) Este medicamento está sujeto a seguimiento adicional, es prioritaria la notificación de sospechas de reacciones adversas asociadas a este medicamento.





