

Empirical Framework for Two-Player Repeated Games with Random States

Arkadiusz Szydlowski*

University of Leicester

May 20, 2021

Abstract

We provide methods for the empirical analysis of a class of two-player repeated games with i.i.d. shocks. The number of possible equilibria in these games is large and, usually, theory is silent about which equilibrium will be chosen in practice. Thus, our method remains agnostic about selection among these multiple equilibria, which leads to partial identification of the parameters of the game. We propose a profiled likelihood criterion for building confidence sets for the structural parameters of the game and derive an easily computable upper bound on the critical value. We demonstrate good finite-sample performance of our procedure using a simulation study. We apply our method to study the effect of repealing the Wright Amendment on entry and exit into Dallas airline markets and find that the existing static game approach overestimates the negative effect of the law on entry into these markets.

Keywords: Empirical games, Partial identification, Parameter on boundary, Airline market

1 Introduction

The theory of repeated strategic interactions has been thoroughly studied by economists. However, the empirical analysis of repeated games is scarce. Repeated games usually feature many equilibria

**E-mail:* ams102@le.ac.uk, *Address:* Department of Economics, Finance and Accounting, University of Leicester, University Road, Leicester, LE1 7RH, UK

making their analysis difficult. In this paper we endeavour to provide an empirical framework for a two-player binary repeated game with i.i.d. shocks to payoffs. We leverage the theoretical result that, conditional on the continuation payoff, a repeated game reduces to a form of a static game for identification, develop a convenient maximum likelihood characterization of the identified set and use it to develop a profiled likelihood ratio test for estimating confidence sets for the structural parameters. We derive the asymptotic distribution of the test statistic and provide an upper bound on the critical value. Monte Carlo simulations suggest that this bound leads to only moderately conservative inference. However, this critical value requires a preliminary estimator of the identified set which makes it computationally intensive. Thus, we also suggest an alternative easily computable, but more conservative, upper bound on the critical value.

Our method can be applied in the analysis of entry and exit into procurement auctions. These auctions are usually repeated over time, often with the same parties involved ([Bajari et al. \(2010\)](#) analysed entry decisions in these auctions using a static game model). Another application involves entry and exit into airline markets. [Ciliberto & Tamer \(2009\)](#) analyse entry and exit into the US airline markets using a static game allowing for multiple equilibria. They use data from the second quarter of 2001 and view it as a long-run outcome of strategic interaction between the carriers, thus arguing that the static game approach is appropriate. However, this interaction can also be seen as repeated game, where players repeatedly make decisions about the presence in a particular market. In fact, as [Table 1](#) shows, there is a lot of entry and exit in the US airline market across time. For example, Delta dropped out of 42% of the markets they served in 1990, even more strikingly, in 2010 they came back to 9.5% of their 1990 markets after not serving them in 2000. Thus, the network of served markets is far from stable and carriers often react to changing market conditions by exit and entry.

In terms of theoretical predictions, as a static game Nash equilibrium repeated over time constitutes an equilibrium in the repeated version of such game, allowing repeated interactions between agents enlarges the set of possible equilibria and allows the model to justify wider range of behaviours. On the other hand, behaviour observed in the data may exclude some equilibria. Take as an example a classic repeated prisoners dilemma with no stochastic shocks. If we observe in the data that players cooperate in this game, this rules out them playing defect-defect, which is a unique equilibrium in the static game. This example illustrates that predictions from a static

Table 1: Share of markets by presence of major carriers over time (in %)

Presence in Q2 1990 - Q2 2000 - Q2 2010	American	Delta	United	US Airways	Southwest
in - in - in	21.7	24.8	26.1	13.9	59.9
in - out - in	5.6	9.5	3.3	5.3	15.8
in - out - out	60.3	42.2	50.4	56.5	17.3
in - in - out	12.5	23.5	20.3	24.3	7.1

Note: T100 Market data, flights with fewer than 20 passengers in 1990 dropped. Markets are defined as directional routes between origin and destination airports (irrespective of the number of stops on the way). "In" means that a carrier served at least one flight on the route.

and repeated version of a game can substantially differ. In fact, in Section 8 we re-investigate the policy analysis in [Ciliberto & Tamer \(2009\)](#) of the effect of repealing the Wright Amendment on entry and exit into Dallas airline markets by Southwest and American using our empirical repeated game and find that the existing static game approach overestimates the negative effect of the law on entry into these markets.¹

Our article is related to the literature on estimation of dynamic games with incomplete information. However, it differs from the approach in [Bajari et al. \(2007\)](#) in important aspects. We focus on a simple form of a dynamic game without state dependence and with complete information.² In this subset of dynamic games we do not have to restrict ourselves to Markov perfect equilibria, allowing for a rich set of equilibrium strategies potentially dependent on the whole history of play. In fact, Markov perfect equilibria in this game correspond to stage game equilibria, thus Markov restriction makes the dynamic dimension irrelevant. We stress however that we restrict ourselves to stationary-outcome equilibria. Thus, our approach can be seen as an alternative to the established dynamic games approach in settings where agents face a long-term strategic interaction in a stable environment with well known structure of payoffs.

Our inference approach is similar to [Kline & Tamer \(2016\)](#) and [Chen et al. \(2018\)](#) in that we assume that the likelihood is identified with respect to choice probabilities but is not identified with respect to parameters (see also [Giacomini & Kitagawa \(2020\)](#)). We apply the same profiled likelihood ratio criterion as in [Chen et al. \(2018\)](#) but analyse it differently with a view of obtaining a practical inference method tailored to our repeated game model, as the general procedure in the

¹Recent papers analysing the US airline market include [Goolsbee & Syverson \(2008\)](#), [Gerardi & Shapiro \(2009\)](#), [Berry & Jia \(2010\)](#) and [Ciliberto et al. \(2020\)](#).

²In principle, our method can be extended to more general dynamic games with non-trivial state dependence, however implementing such an extension poses a difficult computational challenge so we leave it for further research.

latter paper is computationally prohibitive in our setup.

Since we focus on providing marginal confidence intervals for the identified structural parameters, our article is related to the literature on marginal inference in partially identified models – see e.g. [Bugni et al. \(2017\)](#) and references therein. Our problem features also likelihood ratio statistic with a parameter on the boundary of the parameter space as in e.g. [Shapiro \(1985\)](#), [Andrews \(2001\)](#) (in point identified model) and in e.g. [Chernozhukov et al. \(2007\)](#) (in partially identified model). Recent contribution to this literature involves e.g. [Al Mohamad et al. \(2020\)](#). The technique in the latter paper could, in principle, be used to bound the critical value for our problem as well. However, their method would work well in settings when only a few boundary constraints are binding, which makes it unattractive in our setup.

An alternative approach to inference in our model would characterize the identified set using moment inequalities like in [Beresteanu et al. \(2011\)](#) or [Galichon & Henry \(2011\)](#) and then employ profiled inference methods ([Romano & Shaikh \(2008\)](#), [Bugni et al. \(2017\)](#) or [Kaido et al. \(2019\)](#)).

2 Repeated Game with Random States

A defining feature of our model is that agents, unlike in a standard repeated game, play a different game every period. The payoffs in the stage game are stochastic and distributed i.i.d. over time (conditional on observables). Payoffs are fully observed by the players but the econometrician observes only the history of play and has only limited information about the non-stochastic part of the payoff. In particular, she does not observe the realization of the i.i.d. shocks.³

We focus on binary games. Let $\mathcal{A} = \{0, 1\}^2$ denote the set of actions with a typical element a . There are two players identified by $i = 1, 2$ playing an infinitely repeated game, $t = 0, 1, 2, \dots, \infty$. Every period a random payoff relevant vector $\varepsilon_t = \{(\varepsilon_{1t,a}, \varepsilon_{2t,a})\}_{a \in \mathcal{A}}$ is drawn from the distribution $F_\varepsilon : \mathcal{E} \rightarrow [0, 1]$ with the following properties:

Assumption ID1. ε_t are i.i.d. across time and $E(\varepsilon_{t,a}) = \mathbf{0}$ for all a and $\varepsilon_{t,\mathbf{0}} = \mathbf{0}$ (normalization).

Denote player i 's payoff in period t by $\tilde{u}_i(a_t, \varepsilon_{it,a_t}; \alpha_i)$, where α_i is a finite dimensional parameter. Define $\alpha = (\alpha_1, \alpha_2)$. In practice some observed characteristics $X \in \mathcal{X} \subset R^K$ will usually enter the payoff function. For now we suppress them from notation for the ease of exposition. We discuss

³For the rest of the paper we will simply refer to our repeated game with random states as a “repeated game”.

the additional issues related to the presence of covariates in Section 6. We assume that the payoffs are additive in the random shock:

$$\tilde{u}_i(a_t, \varepsilon_{it, a_t}; \alpha_i) = u_i(a_t; \alpha_i) - \varepsilon_{it, a_t}$$

This assumption is usually imposed in the empirical dynamic games literature (see Assumption DC in [Bajari et al. \(2007\)](#)). Payoffs in future periods are discounted by a common discount factor $\delta \in (0, 1)$.

Let $\mathcal{H}^t = (\mathcal{E} \times \mathcal{A})^t$ denote the set of period t ex ante histories with a typical element $h^t = \{\varepsilon_{s, a_s}, a_s\}_{s=0}^{t-1}$. Let $\tilde{\mathcal{H}}^t = (\mathcal{E} \times \mathcal{A})^t \times \mathcal{E}$ denote the set of period t ex post histories with a typical element $\tilde{h}^t = (h^t, \varepsilon_t)$. A pure strategy profile, σ , is a pair of mappings from $\tilde{\mathcal{H}}^t$ to \mathcal{A} . Player i 's expected lifetime payoff from playing the game is:

$$U_i(\sigma; \alpha_i) = E^\sigma \left[\sum_{t=0}^{\infty} \delta^t \tilde{u}_i(a_t, \varepsilon_{it, a_t}; \alpha_i) \right]$$

where the expectation is taken over the histories induced by σ .

A normalized continuation payoff of player i after a history \tilde{h}^t is given by:

$$V_i^\sigma(\tilde{h}^t; \alpha_i) = (1 - \delta)u_i(\sigma(\tilde{h}^t); \alpha_i) - (1 - \delta)\varepsilon_{it, \sigma(\tilde{h}^t)} + \delta \int V_i^\sigma(\tilde{h}^{t+1}; \alpha_i) dF_\varepsilon$$

3 Identification

Throughout our analysis we will assume that

Assumption ID2. (a) δ and F_ε are known, (b) $u(a_i, a_{-i}; \alpha) = 0$ for some $a_i \in \mathcal{A}$ and all $a_{-i} \in \mathcal{A}$

This assumption is frequently made in dynamic discrete choice and dynamic games literature. In the case of dynamic discrete choice models, [Magnac & Thesmar \(2002\)](#) showed that the payoff functions are not identified without knowledge of the discount factor and distribution of shocks or without normalizing the payoffs for one of the alternatives. In practice one can conduct the analysis for different values of δ and different distributions F_ε in order to investigate the sensitivity of results to this assumption.

We assume that we observe a finite history of play, h^t , in at least one game (market) and the observed play is generated by some subgame-perfect Nash equilibrium. In principle, characterizing the set of equilibria in our game is complicated since there are numerous strategies that can generate a given equilibrium outcome. Fortunately, we do not have to work with **strategies** directly but rather characterize sufficient and necessary condition(s) for observed **choices** to be made in equilibrium. Let $p_t(a)$ denote the probability of a being played in equilibrium at some t . We will restrict this probability to be stationary. Note that this does not necessarily restrict strategies to be stationary and allows, for example, tit-for-tat or trigger strategies.

Assumption ID3. *Players play a stationary-outcome subgame-perfect Nash equilibrium, i.e. $p_t(a) = p(a)$ for all t .*

In principle, we could work with non-stationary outcome equilibria. However, this would require that we observe the same game being played in many different markets such that we can calculate $p_t(a)$ for every t by looking at probability of a being played across markets. Such data is hard to come by since payoffs usually differ between markets due to different market characteristics. In some games Assumption ID3 rules out some important equilibria, e.g. some efficient symmetric equilibria (see Section 6.3 in [Mailath & Samuelson \(2006\)](#)), thus one needs to be careful if it is not too strong in a particular game of interest.

Before we characterize the empirical content of our model, note that since the researcher observes neither present nor historical ε 's, it is not possible to restrict the set of potential equilibria based on the observed history of play, i.e. based on observing history of actions only. Thus, all the potential equilibrium continuation payoffs have to be considered at each time period and history.

Lemma 1. *Let $\mathcal{V}_S(\alpha)$ denote the set of pairs of expected lifetime payoffs that can be reached in a stationary-outcome equilibrium, i.e. for any subgame-perfect Nash equilibrium strategy σ we have $(U_1(\sigma; \alpha_1), U_2(\sigma; \alpha_2)) \in \mathcal{V}_S(\alpha)$, and $v(a) = (v_1(a), v_2(a)) \in \mathcal{V}_S(\alpha)$ denote expected continuation payoffs associated with playing a in the current period. Define $V = (v(a), v(a'_1, a_2), v(a_1, a'_2), v(a'_1, a'_2))$ with $a_1 \neq a'_1, a_2 \neq a'_2$. Then, for some distribution $F_V : \mathcal{V}_S(\alpha)^4 \rightarrow [0, 1]$:*

$$p(a) = \int_{\mathcal{V}_S(\alpha)^4} P(a \text{ is a Nash equilibrium in the normal form game with payoffs } g_i(a) | V = v) dF_V(v) \quad (1)$$

where:

$$g_i(a) = (1 - \delta)(u_i(a; \alpha_i) - \varepsilon_{i,a}) + \delta v_i(a),$$

and $\{(\varepsilon_{1,a}, \varepsilon_{2,a})\}_{a \in \mathcal{A}}$ are drawn from F_ε .

Lemma 1 follows from Proposition 5.7.3 in [Mailath & Samuelson \(2006\)](#). The distribution F_V can be interpreted as an equilibrium selection function since every $v \in \mathcal{V}_S(\alpha)$ corresponds to a different subgame-perfect equilibrium in our stochastic game. If data comes only from a single game and mixing between different equilibria is not allowed, then F_V is degenerate on some particular V . However, if data comes from different games (markets) and we allow different equilibria to be played in different markets, then F_V may put non-trivial mass on several V 's corresponding to different markets.

Proposition 1 is useful for two reasons. First, it allows us to describe the identified set in the model without dealing with strategy functions, which belong to a complicated functional space. Instead, in order to verify if a given parameter value α can be reconciled with observed probabilities, it is enough to check if this probability can be generated in equilibrium by some (combination of) continuation payoffs in $\mathcal{V}_S(\alpha)$ ⁴. Second, calculating the sets $\mathcal{V}_S(\alpha)$ is relatively easy for a game with discretely supported independent shocks and we can approximate these sets for a game with continuous F_ε by increasing the number of support points of ε . We discuss methods for obtaining these sets in the next section.

Note that the normal form game may have multiple equilibria itself and denote by π the equilibrium selection probability in the ambiguous region. Let us now make explicit the dependence of the choice probabilities on the parameter α and the equilibrium selection probabilities π and F_V by writing $p(a; \alpha, \pi, F_V)$. $\mathcal{F}(\alpha)$ denotes the set of all cumulative distribution functions supported on $\mathcal{V}_S(\alpha)$ ⁴. As a result of Proposition 1 and the standard argument for identification of maximum likelihood (see e.g. [Amemiya \(1986\)](#)), we have the following corollary:

Corollary 1. *Define the identified set Θ_{01}^S as the set of α 's for which observed probabilities are*

consistent with some stationary-outcome Nash subgame-perfect equilibrium. Assume that:

$$\sup_{\pi \in [0,1], F_V \in \mathcal{F}(\alpha)} E_0 |\log p(Y; \alpha, \pi, F_V)| < \infty$$

for all $\alpha \in \Theta_{01}^S$ and that the selection probability π does not depend on ε or V . Then:

$$\Theta_{01}^S = \arg \sup_{\alpha} \sup_{\pi \in [0,1], F_V \in \mathcal{F}(\alpha)} E_0 \log p(Y; \alpha, \pi, F_V) \quad (2)$$

where the expectation is taken with respect to the true distribution of a . This set is sharp.

It is worth mentioning that a sharp characterisation of the identified set can also be obtained through moment inequalities using the random set approach of [Beresteanu et al. \(2011\)](#). However, with that approach it is less clear how to obtain easily computable critical values or moderately conservative approximations to these critical values as in Section 5.1. We compare our approach to the moment inequality approach in Appendix C.

To illustrate the identification argument consider a simple game with the following stage game payoffs (for further reference, we will call this game Σ^S):


		P 2	
		1	0
P 1	1	$\alpha_1 - \varepsilon_1, \alpha_2 - \varepsilon_2$	$-\varepsilon_1, 0$
	0	$0, -\varepsilon_2$	$0, 0$

where ε_1 and ε_2 are identically and independently distributed (with some abuse of notation, let F_ε denote their distribution). The corresponding normal form of the repeated game is given by:

		P 2	
		1	0
P 1	1	$(1 - \delta)(\alpha_1 - \varepsilon_1) + \delta v_1(1, 1), (1 - \delta)(\alpha_2 - \varepsilon_2) + \delta v_2(1, 1)$	$-(1 - \delta)\varepsilon_1 + \delta v_1(1, 0), \delta v_2(1, 0)$
	0	$\delta v_1(0, 1), -(1 - \delta)\varepsilon_2 + \delta v_2(0, 1)$	$\delta v_1(0, 0), \delta v_2(0, 0)$

Let $\alpha_1 = \alpha_2 = \alpha > 0$. By analogy to static binary games (see [Tamer \(2003\)](#)), we may have multiple Nash equilibria in this normal form. For example, assuming that players play a static Nash equilibrium in every period, i.e. $v_i(a) = 0$ for all i and a , both (1,1) and (0,0) can arise in equilibrium when ε_1 and ε_2 are between 0 and α .

Figure 1: Multiple equilibria in the normal form



In general, the region of ε for which we will have multiple equilibria depends on V , or more specifically – on the value contrasts D_a^i :

$$D_{10} = v_1(1,0) - v_1(0,0), \quad D_{01} = v_2(0,1) - v_2(0,0)$$

$$D_{11}^1 = v_1(1,1) - v_1(0,1), \quad D_{11}^2 = v_2(1,1) - v_2(1,0)$$

Figure 1 shows regions of values of $(\varepsilon_1, \varepsilon_2)$ for which different equilibria occur for different values of D_a^i 's. There are several regions where the game has two equilibria. Let us focus on Case 1 and the probability that $(1,1)$ is played in a subgame-perfect equilibrium. For simplicity assume that only symmetric equilibria are played in this game so $D_{11}^1 = D_{11}^2 = D_{11}$, $D_{10} = D_{01}$. Let π denote

the probability that (0,0) is played in equilibrium in the region where both (0,0) and (1,1) can be played. Then, we can write the probability of (1,1) being played in this repeated game, conditional on the continuation values V as:

$$p(1,1|V) = \left[F_\varepsilon \left(\alpha + \frac{\delta}{1-\delta} D_{11} \right) \right]^2 - \pi \left[F_\varepsilon \left(\alpha + \frac{\delta}{1-\delta} D_{11} \right) - F_\varepsilon \left(\frac{\delta}{1-\delta} D_{01} \right) \right]^2$$

Thus, conditional on V the problem boils down to a standard identification problem in static games, where we may have multiple equilibria and we do not observe the equilibrium selection probability π . However, in our repeated game lack of identification is magnified by the fact that we observe neither π nor V . Nevertheless, our application shows that this does not preclude us from obtaining meaningful bounds on the model parameters.

4 Equilibrium continuation payoff sets

In this section we illustrate how one can obtain the sets $\mathcal{V}_S(\alpha)$. In fact, we will show how to obtain an outer approximation to this set, $\mathcal{V}(\alpha)$, since we will allow $\mathcal{V}(\alpha)$ to contain continuation payoffs both from stationary- and non-stationary-outcome equilibria. Having calculated $\mathcal{V}(\alpha)$, we can characterize an outer set of the identified set, Θ_{01} , by replacing $\mathcal{V}_S(\alpha)$ with $\mathcal{V}(\alpha)$ in characterisation (2). Even though $\Theta_{01}^S \subset \Theta_{01}$ the outer set obtained in this way is still quite narrow in our application (and Monte Carlo simulations).


With finite discrete support of shocks we can view games with different draws of ε as separate non-stochastic repeated games and use the algorithms introduced in [Abreu et al. \(1990\)](#) and [Abreu & Sannikov \(2014\)](#) to find the sets of equilibrium payoffs in these games. Let $\mathcal{V}_{\varepsilon^m}(\alpha)$, $m = 1, \dots, M$ denote the collection of equilibrium payoff sets for different values of shocks $\varepsilon^m \in \mathcal{E}$. Then, the set of equilibrium payoffs in the full game can be calculated as a Minkowski sum:

$$\mathcal{V}(\alpha) = \{V : V = V_1 \cdot P(\varepsilon = \varepsilon^1) + \dots + V_M \cdot P(\varepsilon = \varepsilon^M), V_1 \in \mathcal{V}_{\varepsilon^1}(\alpha), \dots, V_M \in \mathcal{V}_{\varepsilon^M}(\alpha)\}$$

(see Remark 5.7.1 in [Mailath & Samuelson \(2006\)](#)). We will sometimes refer to this set of equilibrium payoffs as the ‘ \mathcal{V} set’.

We will illustrate the construction of this set using the game Σ^S and a simplified version of

Figure 2: Set of equilibrium payoffs, \mathcal{V}



Note: $\alpha = 1, \delta = 0.75$. Blue line demarcates the set of payoffs in the non-stochastic version of the game with $\varepsilon_1 = \varepsilon_2 = 0$. Red line delineates the set in the stochastic game.

an entry game in which entrants engage in Cournot competition (see [Tamer \(2003\)](#), henceforth, referred to as "Cournot entry game"), both with discrete support of shocks over $\mathcal{E} = \{-2, 0, 2\}^2$.

The stage game payoffs in the latter game are given by:

		P 2	
		1	0
P 1	1	$\alpha - \varepsilon_1, \alpha - \varepsilon_2$	$\frac{9}{4}\alpha - \varepsilon_1, 0$
	0	$0, \frac{9}{4}\alpha - \varepsilon_2$	$0, 0$

Figure 2 compares the equilibrium payoff sets of the non-stochastic and stochastic version of game Σ^S (left panel) and the Cournot entry game (right panel). Note that due to symmetry the \mathcal{V} set in Σ^S without shocks is just a straight line connecting stage game payoffs in the two Nash equilibria. However, once we add the stochastic shocks the game is not necessarily symmetric and the \mathcal{V} set will have a non-empty interior for some values of ε . As a result, the \mathcal{V} set in the stochastic game has a non-empty interior as well. The \mathcal{V} sets in the stochastic game exclude some low payoff pairs from the non-stochastic version since now players can condition their strategies on the realizations of the shocks and, as a result, achieve better outcomes.

We can view a game with finite support of shocks as an approximation of a game with continuous F_ε where the approximation becomes more precise as the number of support points, M , grows. For

the purpose of inference we can make $M \rightarrow \infty$ as the sample size $T \rightarrow \infty$.

Finally, it is worth noting that, although the probabilities $p(a)$ depend only on the continuation value contrasts, e.g. $D_{11}^1 = v_1(1, 1) - v_1(1, 0)$, $D_{11}^2 = v_2(1, 1) - v_2(0, 1)$, working with the value contrasts as a primitive of the game is not that convenient because the restriction $V \in \mathcal{V}(\alpha)^4$ may put some non-trivial restrictions on the set of equilibrium D's. For example, setting $D_{11}^1 = \max_{(v_1, v_2) \in \mathcal{V}} v_1 - \min_{(v_1, v_2) \in \mathcal{V}} v_1$ and $D_{11}^2 = \max_{(v_1, v_2) \in \mathcal{V}} v_2 - \min_{(v_1, v_2) \in \mathcal{V}} v_2$ may not be feasible since the points $(\max_{(v_1, v_2) \in \mathcal{V}} v_1, \max_{(v_1, v_2) \in \mathcal{V}} v_2)$ and $(\min_{(v_1, v_2) \in \mathcal{V}} v_1, \min_{(v_1, v_2) \in \mathcal{V}} v_2)$ may not belong to the \mathcal{V} set. Thus, we work with the equilibrium payoffs V rather than value contrasts because this allows us to handle the constraints more easily.

5 Inference

We suggest using maximum likelihood for inference. In practical applications some components of the payoff vector, X , are observed. They usually enter payoffs linearly and we are interested in estimating their coefficients, $\beta = (\beta_1, \beta_2)$. Thus, all the probabilities below should condition on X . We suppress this conditioning and discuss adding covariates to the model in Section 6.

We assume that the discount factor δ is known so the goal is to estimate a confidence set for $\theta_1 \equiv (\alpha, \beta)$, $\theta_1 \in \Theta_1 \subset \mathcal{R}^{d_1}$. We allow for any equilibrium selection mechanism $\pi \in [0, 1]$ but restrict F_V to be degenerate:

Assumption INF1. *All observations are generated from a unique V .*

If the data is pulled across markets, the latter restriction imposes that the same dynamic equilibrium is played in different markets (similarly to [Bajari et al. \(2007\)](#)).⁴ Let $\theta_2 \equiv (\pi, V) \in \Theta_2(\theta_1)$ where $\Theta_2(\theta_1)$ is the correspondence mapping values of θ_1 to the sets of corresponding continuation values V and the set of feasible selection probabilities π . Further, $\theta = (\theta_1, \theta_2) \in \Theta \subset \mathcal{R}^d$, $d = d_1 + d_2$. As in [Chen et al. \(2018\)](#) we suggest using supremum of the profiled likelihood ratio statistic for inference and building the confidence set as a collection of points for which the LR statistic falls below the critical value from the asymptotic distribution of this sup-profiled statistic.

We observe a sample of action pairs $\{Y_t\}_{t=1}^T$. Let $p(a|\theta)$ denote the model implied probabilities when parameters are set to θ . Denote $\gamma_a(\theta) = p(a|\theta) - p(a|\theta_0)$ for $\theta_0 \in \Theta_0$, where Θ_0 is the identified

⁴[Otsu et al. \(2016\)](#) develop tests for credibility of pooling markets in dynamic Markov games.

set for θ , and let the vector $\gamma(\theta) \equiv [\gamma_{11}(\theta) \ \gamma_{10}(\theta) \ \gamma_{01}(\theta)]' \in \Gamma$ collect the choice probabilities. Then the likelihood of observing a given Y_t can be written as:

$$p(Y_t, \theta) = \gamma_{11}(\theta)^{Y_{1t}Y_{2t}} \gamma_{10}(\theta)^{Y_{1t}(1-Y_{2t})} \gamma_{01}(\theta)^{(1-Y_{1t})Y_{2t}} (1 - \gamma_{11}(\theta) - \gamma_{10}(\theta) - \gamma_{01}(\theta))^{(1-Y_{1t})(1-Y_{2t})}$$

With some abuse of notation, we also write $p(Y_t, \gamma)$. Define the identified sets for θ and θ_1 as:

$$\Theta_0 = \arg \sup_{\theta \in \Theta} E_0[\log p(Y_t, \theta)] \quad \text{and} \quad \Theta_{01} = \arg \sup_{\theta_1 \in \Theta_1} \sup_{\theta_2 \in \Theta_2(\theta_1)} E_0[\log p(Y_t, \theta)]$$

where the expectation E_0 is taken with respect to the true distribution of Y_t , P_0 .

Once we fix V , Y_t depends only on the realisation of ε_t . Thus, under Assumption [ID1](#) we can write the log-likelihood by $L_T(\theta) = \sum_{t=1}^T \log p(Y_t, \theta)$. For a candidate value $\tilde{\theta}_1$ the profiled likelihood ratio statistic is defined as:

$$LR_T(\tilde{\theta}_1) = 2 \left[\sup_{\theta \in \Theta} L_T(\theta) - \sup_{\theta_2 \in \Theta_2(\tilde{\theta}_1)} L_T(\tilde{\theta}_1, \theta_2) \right] + o_p(1) \quad (3)$$

where the $o_p(1)$ term accommodates approximation error coming from using a discrete grid for ε when calculating \mathcal{V} sets (and optimization error). Similarly to [Chen et al. \(2018\)](#) we will calculate the $100\kappa\%$ confidence set for the identified set Θ_{01} as:

$$\hat{\Theta}_{01}^\kappa = \{\theta_1 : LR_T(\theta_1) \leq c_\kappa\}$$

where c_κ is the κ quantile of the asymptotic distribution of $\sup_{\theta_1 \in \Theta_{01}} LR_T(\theta_1)$.

Obtaining asymptotic distribution of the likelihood ratio statistic in our case is difficult because the second supremum in [\(3\)](#) will often be attained at the boundary of the parameter space $\Theta_2(\tilde{\theta}_1)$ and the asymptotic distribution will depend on the shape of the local parameter space at this boundary (which will in turn depend on the value of θ_1). Given these difficulties we focus on obtaining an upper bound on the critical value.

Note that situation here is different than in the missing data example in [Chen et al. \(2018\)](#) as local parameter space at $\theta \in \Theta$ cannot be characterized as a simple translation of the null parameter space and, as a result, their Assumption 4.7 cannot be verified. Thus, we cannot use χ_1^2 quantile

as an upper bound on our critical value.⁵ Additionally, although their Procedure 2 is applicable in our model, it involves repeated computation of the level sets of the likelihood for each MCMC draw which is prohibitively costly in our setup as the likelihood is nonlinear and each evaluation requires recomputing the constraint set.

Let \mathbb{I}_0 denote the Fischer information matrix:

$$\mathbb{I}_0 = -E_0[\partial^2 \log p(y, \gamma_0)/\partial \gamma^2].$$

where $\gamma_0 = \gamma(\theta_0)$, $\theta_0 \in \Theta_0$. We say that a sequence of sets Γ_T covers a set K if there is a sequence of closed balls of radius $k_T \rightarrow \infty$ centred at the origin, B_{k_T} , such that $\Gamma_T \cap B_{k_T} = K \cap B_{k_T}$ with probability approaching one (see p. 1987 in [Chen et al. \(2018\)](#)).

Assumption INF2. *We have:*

- (a) Θ_1 is a compact, nonempty subset of \mathbb{R} and $\Theta_2(\theta_1)$ is a compact, convex, nonempty subset of \mathbb{R}^{d_2} for every $\theta_1 \in \Theta_1$.
- (b) $\sup_{\theta_2 \in \Theta_2(\theta_1)} L_T(\theta_1, \theta_2)$ is quasi-concave in θ_1 .
- (c) $E_0[\partial \log p(y, \gamma)/\partial \gamma] = 0$ has only one (interior) solution $\gamma_0 = 0$.
- (d) There exists a neighbourhood $\mathcal{N} \subset \text{Int}(\Gamma)$ of γ_0 on which $\log p(y, \gamma)$ is twice continuously differentiable for each y , with first derivative in $L^2(P_0)$, and $\sup_{\gamma \in \mathcal{N}} \|\partial^2 \log p(y, \gamma)/\partial \gamma^2\| \leq \bar{l}(y)$ for $\bar{l} \in L^2(P_0)$.
- (e) There exists a neighbourhood \mathcal{N}_θ of Θ_0 on which $\gamma(\theta)$ is twice continuously differentiable with second derivatives bounded uniformly over \mathcal{N}_θ .
- (f) \mathbb{I}_0 is non-singular and $\mathbb{I}_0 = E_0 \left[\frac{\partial \log p(y, \gamma)}{\partial \gamma} \frac{\partial \log p(y, \gamma)}{\partial \gamma}' \right]$.
- (g) The local parameter space $\Gamma_{oT}(\theta_1) = \{\sqrt{T}\mathbb{I}_0^{1/2} \gamma(\theta_1, \theta_2) : (\theta_1, \theta_2) \in \Theta_{oT}\}$ covers a closed convex cone $\mathcal{K}(\theta_1) \subset \mathbb{R}^3$, respectively, for every $\theta_1 \in \Theta_1$, where Θ_{oT} is a sequence of small neighbourhoods of Θ_0 .

⁵Our Monte Carlo simulations (not reported here) also confirm that χ_1^2 critical value fails to provide required coverage.

Assumption (a) is satisfied in our setup because $\Theta_2(\theta_1) = [0, 1] \times \mathcal{V}(\theta_1)$ and $\mathcal{V}(\theta_1)$ is guaranteed to be convex. Assumption (b) implies that $\sup_{\theta_1 \in \Theta_{01}} LR_T(\theta_1)$ is reached at the endpoints of the marginal identified set Θ_{01} . Denote these endpoints by $\underline{\theta}_1$ and $\bar{\theta}_1$. Assumptions (c)-(g) allow us to obtain quadratic approximation to the likelihood with respect to probabilities γ using Proposition 5.1 in [Chen et al. \(2018\)](#). They imply that the likelihood is identified with respect to the choice probabilities (but not with respect to θ).

Let $\Delta_{\theta_1} = \partial\gamma(\theta_0)/\partial\theta_2'$ for $\theta_0 = (\theta_1, \theta_2) \in \Theta_0$. In the simplest case, which is the leading case in our model, each value of the structural parameter θ_1 in the identified set Θ_{01} corresponds to a unique likelihood maximising value θ_2 . This happens if the identified set is a manifold which is not parallel to any of the axes and implies that the θ_2 in the definition of Δ_{θ_1} is unique.

Assumption INF3. *We have:*

(a) $\theta_2^*(\theta_1) = \arg \max_{\theta_2 \in \Theta_2(\theta_1)} L_T(\theta_1, \theta_2)$ is a singleton for any θ_1 in the neighbourhood of Θ_{01} .

(b) Δ_{θ_1} has full row rank in the neighbourhood \mathcal{N}_{θ} .

A sufficient condition for part (a) is strict quasi-concavity of $L_T(\theta_1, \theta_2)$ in θ_2 . Although this cannot always be easily verified analytically, problems with convergence of the numerical optimisation algorithms would usually signify violation of this condition. Part (b) is usually innocuous in our model as it merely requires that choice probabilities are not collinear in parameters. This assumption is similar to conditions in Lemma 1 in [Kline & Tamer \(2016\)](#).⁶ If Assumption INF3 does not hold, one can still use the chi-square critical value described in Section 5.2 to obtain the desired (conservative) confidence set for θ_1 .⁷

We say that the set Θ is approximated at θ_0 by a cone K^{θ_0} if:

$$\begin{aligned} \inf_{s \in K^{\theta_0}} \|(\theta - \theta_0) - s\| &= o(\|\theta - \theta_0\|) & \theta \in \Theta \\ \inf_{\theta \in \Theta} \|(\theta - \theta_0) - s\| &= o(\|s\|) & s \in K^{\theta_0} \end{aligned}$$

⁶Condition INF3(b) is akin to constraint qualification conditions in the stochastic programming literature. In fact, it is a localized version of the linear independence constraint qualification for equality constraints: $\gamma_0 - [p((1, 1), \theta) \ p((1, 0), \theta) \ p((0, 1), \theta)]' = 0$, where the first component of θ is fixed at some θ_1 in the small neighbourhood of the identified set. See [Kaido et al. \(2020\)](#) for a discussion on how constraint qualifications relate to standard assumptions in the moment inequality literature.

⁷In Appendix D we discuss how to adjust our inference procedure in Section 5.1 if Assumption INF3 does not hold.

Theorem 1. Let $\Theta_2(\underline{\theta}_1)$ and $\Theta_2(\bar{\theta}_1)$ be approximated at θ_2 such that $\theta_0 = (\underline{\theta}_1, \theta_2) \in \Theta_0$ or $\theta_0 = (\bar{\theta}_1, \theta_2) \in \Theta_0$ by cones $K_2(\underline{\theta}_1)$ and $K_2(\bar{\theta}_1)$, respectively. If Assumptions [INF1-INF3](#) hold, then:

$$\sup_{\theta_1 \in \Theta_{01}} LR_T(\theta_1) = \max_{\theta_1 \in \{\underline{\theta}_1, \bar{\theta}_1\}} \left\{ \inf_{s \in K_2(\theta_1)} \|V_T - \Delta_{\theta_1} s\|^2 \right\} + o_p(1) \quad (4)$$

where $V_T = \mathbb{I}_0^{-1/2} \frac{1}{\sqrt{T}} \sum_{t=1}^T \frac{\partial \log p(Y_t, \gamma_0)}{\partial \gamma}$ asymptotically follows as a multivariate standard normal distribution.

Theorem 1 shows that the asymptotic distribution of our statistic is equivalent to the distribution of the maximum of two chi-bar square distributions (cf. Proposition 3.4.1 in [Silvapulle & Sen \(2001\)](#)). Note that this bound is useful only if the probabilities $\gamma(\theta_0)$ are not flat in θ for $\theta_0 = (\underline{\theta}_1, \theta_2)$ or $\theta_0 = (\bar{\theta}_1, \theta_2)$, which is guaranteed in our repeated game model (note that derivatives of γ involve density of the shocks, which is non-zero everywhere under standard assumptions).

5.1 Simulated critical value

Using Theorem 1 for inference on the identified set Θ_{01} presents both computational (calculation of constraint sets) and theoretical (parameter on boundary) challenges. For example, simply bootstrapping our criterion function will not lead to a valid critical value. Thus, in order to provide a feasible inference procedure we make some simplifying assumptions and opt for a moderately conservative critical value.

5.1.1 Simplified \mathcal{V} set

Firstly, note that using our likelihood criterion implies that we have to recalculate the \mathcal{V} set, a convex polytope, for each candidate value of θ_1 . This is computationally heavy, and barely feasible in realistic applications given the current state of computational resources. Thus, we simplify our procedure by enlarging the continuation payoff set \mathcal{V} to a cube:

Assumption INF4. $\Theta_2(\theta_1)$ is a cube in \mathbb{R}^{d_2} .

Although this may result in a larger confidence set than implied by the general procedure in the previous section, it significantly simplifies computation of the critical value. Firstly, it allows us to track only the minimal and maximal values of the \mathcal{V} set which considerably cuts the time needed

to compute the profiled likelihood in (3). Secondly, it allows us to derive a simple upper bound on the asymptotic distribution described in Theorem 1 as now the local parameter space is an orthant in the least favourable case.

With this simplification the quantiles of the chi-bar square random variables described in Theorem 1 can be calculated using the formulas in Kudo (1963). However, the maximum in the formulation of the asymptotic approximation in (4) complicates obtaining the critical values as chi-bar square random variables under the maximum are correlated. Fortunately, with our extended \mathcal{V} set we can derive a conservative critical value.

5.1.2 Bounding critical value from above

The first step in applying Theorem 1 lies in estimating Δ_{θ_1} , for which we need to estimate $\partial\gamma(\theta_0)/\partial\theta$ for $\theta_0 = (\underline{\theta}_1, \theta_2)$ and $\theta_0 = (\overline{\theta}_1, \theta_2)$. Thus, application of our results requires a preliminary estimate of the extreme points of the identified set Θ_0 . We follow Chernozhukov et al. (2007) and suggest estimating these extreme points by:

$$\begin{aligned} (\hat{\underline{\theta}}_1, \hat{\underline{\theta}}_2) : \quad \hat{\underline{\theta}}_1 &= \min \{ \theta_1 : LR_T(\theta_1) \leq e_T \}; \hat{\underline{\theta}}_2 = \arg \sup_{\theta_2 \in \Theta_2(\hat{\underline{\theta}}_1)} L_T(\hat{\underline{\theta}}_1, \theta_2) \\ (\hat{\overline{\theta}}_1, \hat{\overline{\theta}}_2) : \quad \hat{\overline{\theta}}_1 &= \max \{ \theta_1 : LR_T(\theta_1) \leq e_T \}; \hat{\overline{\theta}}_2 = \arg \sup_{\theta_2 \in \Theta_2(\hat{\overline{\theta}}_1)} L_T(\hat{\overline{\theta}}_1, \theta_2) \end{aligned}$$

where $e_T = \log(\log(T))$ or $e_T = \log(T)$.⁸


The next step is to estimate the local cones $K_2(\underline{\theta}_1), K_2(\overline{\theta}_1)$. This part is challenging as any sampling error in estimation of $\underline{\theta}_1$ may lead to incorrect estimate of $K_2(\underline{\theta}_1)$. For example, even if $K_2(\underline{\theta}_1)$ was an orthant in \mathbb{R}^{d_2} , $K_2(\hat{\underline{\theta}}_1)$ will likely be larger than a orthant, e.g. a half-space, even when T is relatively large.

Instead, we choose to approximate K_2 by the closest least favourable local parameter space, namely the orthant corresponding to the corner of the cube closest to $\hat{\underline{\theta}}_1$ or $\hat{\overline{\theta}}_1$.⁹ We illustrate this construction in Figure 3 in a simplified setup in which $\Theta_2(\theta_1)$ is a square in \mathbb{R}^2 and Assumption INF3 holds. In this example $\hat{\underline{\theta}}_2$ lies on the edge of the parameter space so the local parameter

⁸If one believes that the degeneracy condition in Chernozhukov et al. (2007) is satisfied, one can use $e_T = 0$ or $e_T = o(1)$.

⁹This resembles the geometric moment selection procedure in static entry games developed by Bontemps & Kumar (2020). However, their setup is very different than ours.

Figure 3: Worst-case local parameter space



Note: The arcs mark local parameter spaces corresponding to different $\theta_2 \in \Theta_2(\hat{\theta}_1)$

space, $K_2(\hat{\theta}_1)$, is a half-space $\mathbb{R} \times \mathbb{R}_+$. Instead of using $K_2(\hat{\theta}_1)$, we take the local parameter space corresponding to the corner closest to $\hat{\theta}_2$, namely $K^{LF}(\hat{\theta}_1) = \mathbb{R}_+^2$.

This construction produces a critical value that, in general, will be larger than the critical value from the asymptotic distribution in Theorem 1. The intuitive reason for why the bound is only moderately conservative is the following. Consider the game Σ^S . In order to match the observed probabilities of observing (1,1) using a very low value of α we have to set the value contrasts D_{11}^1 and D_{11}^2 high. In other words, in order to entice players to cooperate on 1 when stage game payoffs from doing so are low we need to promise them high continuation values. As a result the lower and upper end of the identified set for α likely correspond to D 's being in the corners of the feasible set for these contrasts.

Define \mathcal{C}_{θ_1} as the set of all corners of the cube $\Theta_2(\theta_1)$. Formally, we define our simulated critical value, c_κ , as the estimate of the κ quantile of the distribution of:

$$Q^{LF} \equiv \max_{\theta_1 \in \{\hat{\theta}_1, \hat{\theta}_1\}} \left\{ \inf_{s \in K^{LF}(\mathcal{C}_{\theta_1})} \|V_T - \Delta_{\theta_1} s\|^2 \right\}$$

where $C_{\theta_1} \in \mathcal{C}_{\theta_1}$ is defined as:

$$C_{\theta_1} = \arg \min_{C \in \mathcal{C}_{\theta_1}} \|C - \theta_2^*(\theta_1)\|$$

for $\theta_2^*(\cdot)$ defined in Assumption [INF3](#). $K^{LF}(C_{\theta_1})$ is an orthant in \mathbb{R}^{d_2} approximating the parameter space $\Theta_2(\theta_1)$ at C_{θ_1} . Display [1](#) summarizes our procedure.^{[10](#)}

Display 1. *Simulation procedure:*

1. Estimate $\underline{\theta}_1$ and $\bar{\theta}_1$ by:

$$\hat{\underline{\theta}}_1 = \min\{\theta_1 : LR_T(\theta_1) \leq e_T\} \quad \hat{\bar{\theta}}_1 = \max\{\theta_1 : LR_T(\theta_1) \leq e_T\}$$

where $e_T = \log \log(T)$ or $e_T = \log(T)$.

2. Estimate $\Delta_{\underline{\theta}_1}$ and $\Delta_{\bar{\theta}_1}$ by $\Delta_{\hat{\underline{\theta}}_1} = \partial\gamma(\hat{\underline{\theta}}_1, \hat{\underline{\theta}}_2)/\partial\theta'_2$ and $\Delta_{\hat{\bar{\theta}}_1} = \partial\gamma(\hat{\bar{\theta}}_1, \hat{\bar{\theta}}_2)/\partial\theta'_2$.

3. Find the corners, $C_{\hat{\underline{\theta}}_1}$ and $C_{\hat{\bar{\theta}}_1}$, of the parameter space closest to $\hat{\underline{\theta}}_2$ and $\hat{\bar{\theta}}_2$.

4. Simulate:

$$\max \left\{ \inf_{s \in K^{LF}(C_{\hat{\underline{\theta}}_1})} \|V - \Delta_{\hat{\underline{\theta}}_1} s\|^2, \inf_{s \in K^{LF}(C_{\hat{\bar{\theta}}_1})} \|V - \Delta_{\hat{\bar{\theta}}_1} s\|^2 \right\}$$

by drawing large number of standard normal vectors V , and obtain the critical value c_κ by taking the κ quantile of the empirical distribution of this statistics across the simulation draws.

5. Estimate the κ confidence set for θ_1 by:

$$\hat{\Theta}_{01}^\kappa = \{\theta_1 : LR_T(\theta_1) \leq c_\kappa\}$$

¹⁰ An alternative bound on the critical value would come from setting $K^{LF}(\hat{\underline{\theta}}_1)$ to a cone with a smaller solid angle out of the conical hulls of $\Delta_{\hat{\underline{\theta}}_1}$ and $\Delta_{\hat{\bar{\theta}}_1}$ (least favourable cone) and setting its mirror image with respect to the origin as $K^{LF}(\hat{\bar{\theta}}_1)$ (least favourable alignment of the cones). This procedure produces similar but slightly more conservative critical values.

Note that $\inf_{s \in K^{LF}(C_{\theta_1})} \|V_T - \Delta_{\theta_1} s\|^2$ is a convex program and can be solved repeatedly quite fast using packages like *CVX* in MATLAB. Additionally, the evaluations of $LR_T(\theta_1)$ in the first step of the procedure can be stored and reused to calculate the confidence set in the final step. Note that if we were to use Procedure 2 in [Chen et al. \(2018\)](#) we would have to calculate numerically the (marginal) level sets of $p(\cdot, (\theta_1, \cdot))$ with respect to the Kullback-Leibler distance for each MCMC draw from the posterior distribution of θ . As it is not uncommon in the Bayesian estimation for the number of required MCMC draws to reach several thousand, application of that procedure is unattractive in our setup. Comparing results of Monte Carlo simulations in [Chen et al. \(2018\)](#) to those in Section 7 we conclude, however, that the simplified computation in our inference procedure comes at a cost of increased conservativeness.

We need to make the following technical assumptions in order to justify our procedure:

Assumption INF5. C_{θ_1} , as a function of θ_1 , is constant in θ_1 in the neighbourhoods of $\underline{\theta}_1$ and $\bar{\theta}_1$.

This high-level assumption basically excludes a knife-edge situation in which $\underline{\theta}_2 = \theta_2^*(\underline{\theta}_1)$ and $\bar{\theta}_2 = \theta_2^*(\bar{\theta}_1)$ are (almost) equidistant from the corners of the parameter space and, as argued above, should be generally satisfied in our setup as the structure of our model suggests that $\underline{\theta}_2$ and $\bar{\theta}_2$ will be located at or close to the corners of the parameter space.

Theorem 2. *If assumptions of Theorem 1 and Assumptions INF3-INF5 hold, then for any $c \geq 0$:*

$$P\left(\sup_{\theta_1 \in \Theta_{01}} LR_T(\theta_1) \leq c\right) \geq P\left(\max_{\theta_1 \in \{\hat{\underline{\theta}}_1, \hat{\bar{\theta}}_1\}} \left\{ \inf_{s \in K^{LF}(C_{\theta_1})} \|V_T - \Delta_{\theta_1} s\|^2 \right\} \leq c\right) + o(1). \quad (5)$$

Theorem 2 confirms that our simulated critical value provides valid, but possibly conservative, inference for the identified set Θ_{01} . Our Monte Carlo simulations reported in Section 7 suggest that in fact this procedure is only mildly conservative, thus we recommend using it in applications.

5.2 Simple conservative critical value

The procedure in the previous section, though feasible, still requires preliminary estimation of the identified set and large number of convex optimisations. Therefore, in this section we suggest an alternative procedure that approximates the critical value from above. This bound is noticeably

more conservative than the one implied by Theorem 2 but instead does not require simulation or additional computation in order to obtain the critical value.

Theorem 3. *If Assumption INF2 holds, then:*

$$\lim_{T \rightarrow \infty} P \left(\sup_{\theta_1 \in \Theta_{01}} LR_T(\theta_1) \leq c_\kappa^\chi \right) \geq \kappa$$

where c_κ^χ is the κ quantile of the chi-square distribution with 3 degrees of freedom.

Theorem 3 is convenient because calculation of the critical value c_κ^χ is straightforward, though as shown in Section 7 this critical value gives much more conservative inference than our simulated critical value c_κ . In practice, one can compare confidence sets resulting from using both critical values for a chosen parsimonious specification of the model and apply the chi-square critical value for the remaining specifications only if the confidence sets do not significantly differ. Additionally, note that Theorem 3 does not require Assumption INF3, thus c_κ^χ can be applied in more general circumstances than c_κ .

6 Model with covariates

The previous discussion did not explicitly include covariates. In practice, however, one may be interested in estimating the effect of observed characteristics of the agents and markets on the payoffs. For example, Ciliberto & Tamer (2009) estimate and simulate the effect of the Wright amendment on entry and exit into the Dallas airline markets. All the inference procedures described above can accommodate presence of covariates. However, doing so entails some conceptual and computational issues.

Firstly, this poses some conceptual dilemma as now the econometrician observes part of the stochastic component of the payoff vectors, $X = (X_1, X_2)$, hence she could potentially use history of the covariates to restrict the continuation payoff set \mathcal{V} , i.e. rule out some equilibria in the game. As this would prohibitively complicate the computation of the likelihood and it is not clear if this would allow us, in fact, to obtain much sharper bounds (note that the unobserved ε 's have unbounded support), we assume that the econometrician does not use this knowledge, i.e. we treat X 's just as ε 's in the computation of the \mathcal{V} set. Also we assume that the covariates are either i.i.d.

or time-invariant (if we observe multiple markets/games in each period).

Secondly, the presence of covariates complicates computation – now \mathcal{V} depends both on (X_1, X_2) and (α, β) , where β is the vector of covariate coefficients. In order to deal with this complication we 1) discretize (X_1, X_2) and 2) approximate the boundaries of \mathcal{V} , i.e. v_{min}, v_{max} , by a polynomial spline in (α, β) . If we take as a point of departure the analysis of a partially-identified static game 1) can hardly be seen as a restriction as it is present also in the empirical analysis of that simple game (see [Ciliberto & Tamer \(2009\)](#)). Also our experience shows that the boundaries v_{min}, v_{max} tend to be smooth functions of (α, β) so the polynomial approximation works very well in practice.

Thirdly, now the “static” selection probability π may depend on X , which increases the dimensionality of the optimisation problems in our inference procedure. In order to address that one may approximate $\pi(X_1, X_2)$ by an auxiliary parametric model, for example probit or logit.¹¹

7 Monte Carlo simulations

7.1 Cournot entry game

In order to check the finite sample performance of our methods we performed a small Monte Carlo study using Cournot entry game example described above with independent shocks following logistic or Normal distribution $N(0, 4)$. As mentioned above, to simplify computation, we do not calculate the full continuation payoff set $\mathcal{V}(\alpha)$ but only calculate the extreme values of this set: $v_{min} = \min\{v : v \in \mathcal{V}(\alpha)\}$ and $v_{max} = \max\{v : v \in \mathcal{V}(\alpha)\}$, and use $[v_{min}, v_{max}]^4$ in place of the continuation payoff set.

In simulations we set the true value $\alpha = 1$. The marginal identified set for α in the logistic model is $\Theta_{01} = [0.92, 1.1]$ for a discount factor $\delta = 0.55$ and $\Theta_{01} = [0.76, 1.89]$ for a discount factor $\delta = 0.75$. For the normal model, these sets are $\Theta_{01} = [0.95, 1.09]$ and $\Theta_{01} = [0.8, 2.03]$. Results are given in [Tables 2-3](#). We analyse performance of both the critical value implied by [Theorem 2](#) (Simulated crit. val.) and [Theorem 3](#) (χ^2_3 crit. val.).

The simulations confirm implications of our theoretical results – both critical values deliver coverage probabilities that are close or above nominal values. MC simulations confirm also that χ^2_3

¹¹Here an attractive alternative may be using the moment inequality approach described in [Appendix C](#) as it does not involve the selection probability π .

Table 2: MC simulations: coverage probabilities, $\delta = 0.55$

Logistic shocks						Normal shocks					
Simulated crit. val.			χ^2_3 crit. val.			Simulated crit. val.			χ^2_3 crit. val.		
90%	95%	99%	90%	95%	99%	90%	95%	99%	90%	95%	99%
$T = 100$											
$\Theta_{01} = [0.92, 1.1]$			0.966	0.985	0.995	0.989	0.995	0.999	$\Theta_{01} = [0.95, 1.09]$		
$\alpha = 0.5$			0.037	0.074	0.231	0.108	0.193	0.387	0.952	0.984	0.997
$\alpha = 1.5$			0.724	0.839	0.937	0.881	0.925	0.978	0.027	0.056	0.192
									0.243	0.321	0.552
									0.41	0.51	0.7
$T = 250$											
$\Theta_{01} = [0.92, 1.1]$			0.916	0.978	0.997	0.989	0.996	1	$\Theta_{01} = [0.95, 1.09]$		
$\alpha = 0.5$			0.001	0.001	0.001	0.001	0.001	0.008	0.931	0.965	0.995
$\alpha = 1.5$			0.512	0.653	0.808	0.702	0.776	0.893	0	0	0.001
									0.025	0.041	0.103
									0.06	0.09	0.21
$T = 500$											
$\Theta_{01} = [0.92, 1.1]$			0.923	0.973	0.986	0.98	0.983	0.997	$\Theta_{01} = [0.95, 1.09]$		
$\alpha = 0.5$			0	0	0	0	0	0	0.942	0.969	0.992
$\alpha = 1.5$			0.321	0.425	0.615	0.457	0.601	0.767	0	0	0
									0	0.001	0.003
									0	0	0.01
$T = 1000$											
$\Theta_{01} = [0.92, 1.1]$			0.929	0.975	0.995	0.978	0.994	0.998	$\Theta_{01} = [0.95, 1.09]$		
$\alpha = 0.5$			0	0	0	0	0	0	0.939	0.966	0.993
$\alpha = 1.5$			0.116	0.156	0.313	0.217	0.299	0.473	0	0	0
									0	0	0
$T = 2000$											
$\Theta_{01} = [0.92, 1.1]$			0.933	0.966	0.992	0.981	0.991	0.999	$\Theta_{01} = [0.95, 1.09]$		
$\alpha = 0.5$			0	0	0	0	0	0	0.941	0.967	0.994
$\alpha = 1.5$			0.007	0.014	0.054	0.025	0.045	0.118	0	0	0
									0	0	0

Note: 2000 Monte Carlo replications

Table 3: MC simulations: coverage probabilities, $\delta = 0.75$

		Logistic shocks			χ^2_3 crit. val.			Normal shocks					
		Simulated crit. val.			90%	95%	99%	Simulated crit. val.			χ^2_3 crit. val.		
								90%	95%	99%	90%	95%	99%

critical value is a valid but, as expected, conservative upper bound on the critical value from the asymptotic distribution of our likelihood ratio statistic.

Our test has good power against values outside the identified set for $\delta = 0.55$ but for $\delta = 0.75$ it lacks power above the upper end of the identified set (especially with logistic shocks), which suggests that marginal likelihood is very flat at this end and resulting confidence sets will be quite wide.

7.2 Cournot entry game with covariates

We extend the Cournot entry game example to include observed covariates:

		P 2	
		1	0
P 1	1	$\alpha + X_1\beta - \varepsilon_1, \alpha + X_2\beta - \varepsilon_2$	$\frac{9}{4}\alpha + X_1\beta - \varepsilon_1, 0$
	0	$0, \frac{9}{4}\alpha + X_2\beta - \varepsilon_2$	$0, 0$

We draw both X_1 and X_2 from the uniform distribution over $\{-1, -0.5, 0, 0.5, 1\}$. As now the \mathcal{V} set depends on α, β and (X_1, X_2) , in order to simplify computation we generate a sample of v_{min} and v_{max} for given (X_1, X_2) and different values of α and β and approximate the relationship between (v_{min}, v_{max}) and α, β by a 6-th order bivariate polynomial. Otherwise the MC designs are the same as in the simple model above.

We report only the results for $\delta = 0.75$ (see Table 4). The marginal identified set for α in the logistic model is $\Theta_{01} = [0.33, 3.25]$ and $\Theta_{01} = [0.38, 2.78]$ in the normal model. The results confirm previous conclusions, namely that our simulation method is not excessively conservative, in fact it produces coverage probabilities that are close to the nominal values, which is encouraging. Using the chi-squared with three degrees of freedom is far more conservative, just as in the model without covariates. Overall, the results in Table 4 confirm that introducing covariates into the model does not essentially change the properties of our inference procedures.

8 Empirical application: Wright Amendment

We employ our framework to re-evaluate the Wright Amendment experiment in [Ciliberto & Tamer \(2009\)](#) (CT henceforth). The Wright Amendment, passed into law in 1979, restricted airline service

Table 4: MC simulations: coverage probabilities, $\delta = 0.75$, model with covariates

Logistic shocks				χ^2 crit. val.				Normal shocks			
Simulated crit. val.				χ^2_3 crit. val.				Simulated crit. val.			
90%	95%	99%		90%	95%	99%		90%	95%	99%	χ^2_3 crit. val.
$T = 100$											
$\Theta_{01} = [0.33, 3.25]$	0.958	0.986	0.998	0.991	0.995	1	$\Theta_{01} = [0.38, 2.78]$	0.937	0.971	0.998	0.98
$\alpha = 0.05$	0.281	0.41	0.675	0.463	0.594	0.813	$\alpha = 0.05$	0.301	0.426	0.693	0.51
$\alpha = 3.75$	0.873	0.927	0.976	0.944	0.969	0.992	$\alpha = 3.25$	0.844	0.909	0.969	0.94
											0.96
											0.99
$T = 250$											
$\Theta_{01} = [0.33, 3.25]$	0.915	0.968	0.998	0.98	0.993	1	$\Theta_{01} = [0.38, 2.78]$	0.91	0.96	0.993	0.98
$\alpha = 0.05$	0.033	0.063	0.178	0.09	0.156	0.334	$\alpha = 0.05$	0.041	0.08	0.214	0.12
$\alpha = 3.75$	0.696	0.788	0.91	0.84	0.895	0.965	$\alpha = 3.25$	0.646	0.757	0.892	0.81
											0.88
											0.96
$T = 500$											
$\Theta_{01} = [0.33, 3.25]$	0.909	0.95	0.993	0.978	0.99	0.999	$\Theta_{01} = [0.38, 2.78]$	0.907	0.95	0.99	0.97
$\alpha = 0.05$	0	0.002	0.007	0.003	0.006	0.025	$\alpha = 0.05$	0.001	0.002	0.011	0
$\alpha = 3.75$	0.454	0.566	0.76	0.644	0.74	0.878	$\alpha = 3.25$	0.399	0.52	0.722	0.59
											0.69
											0.84
$T = 1000$											
$\Theta_{01} = [0.33, 3.25]$	0.921	0.958	0.994	0.98	0.992	1	$\Theta_{01} = [0.38, 2.78]$	0.907	0.951	0.984	0.97
$\alpha = 0.05$	0	0	0	0	0	0	$\alpha = 0.05$	0	0	0	0
$\alpha = 3.75$	0.179	0.265	0.462	0.336	0.436	0.651	$\alpha = 3.25$	0.124	0.195	0.404	0.27
											0.37
											0.58
$T = 2000$											
$\Theta_{01} = [0.33, 3.25]$	0.922	0.966	0.994	0.983	0.992	1	$\Theta_{01} = [0.38, 2.78]$	0.92	0.963	0.992	0.98
$\alpha = 0.05$	0	0	0	0	0	0	$\alpha = 0.05$	0	0	0	0
$\alpha = 3.75$	0.018	0.037	0.108	0.054	0.092	0.211	$\alpha = 3.25$	0.01	0.021	0.069	0.03
											0.06
											0.15

Note: 2000 Monte Carlo replications

from Dallas Love airport in order to stimulate growth of Dallas/Forth Worth, permitting flights only to/from Texas, Louisiana, Arkansas, Oklahoma, New Mexico, Alabama, Kansas and Mississippi. The Amendment was partially repealed starting from 2006 and fully withdrawn in 2014. Using a static game model CT compare the predicted changes in market service out of Dallas Love with and without the amendment and find very large positive effects of repealing the amendment on the number of served markets. As the repeal is predicted to especially benefit Southwest Airlines the authors conclude that the prolonged binding of the Amendment was meant to protect American Airlines monopolies in markets out of Dallas/Fort Worth.

They interpret these effects as short run effects because their model does not involve dynamic dimension. However, dynamic strategic responses are important in this context – the threat of American retaliating against entry into the Dallas market in the future may prevent Southwest from entering today. As a result, the static model may overstate the positive effect of the repeal of the Wright Amendment on market entry (in fact CT acknowledge that the size of their estimated effects may suggest that they are overestimated).

In order to address these concerns we redo the experiment using our repeated game model focusing on the interaction between American (AA) and Southwest (WN) in the markets out of Dallas. We use quarterly DB1B and T100 data from 1993 to 2017. We estimate our model using the following covariates: *market size*, the size of the market calculated as the geometric mean of the population of the endpoint cities, *Wright*, indicating that the market was restricted by the Wright amendment, *airport presence*, which gives the average fraction of markets served from the endpoints by American or Southwest and *cost* which is equal to the difference between the “origin-closest hub-destination” distance and the nonstop “origin-destination” distance divided by the latter distance (see CT for detailed description and justification).

Table 5: Summary statistics

	Mean	Min	Max
market size	3,675,950	1,750,056	10,968,557
Wright	0.356	0	1
AA airport presence	0.763	0.211	0.983
WN airport presence	0.367	0	0.713
AA cost	0.009	0	0.083
WN cost	0.238	0	2.811

Table 5 contains the summary statistics. There are 244 markets in our sample.¹² The average market covered a population of around 3-4 million people. As we focus on Dallas, the Wright amendment affects a large fraction of markets – 35.6% of year-quarter-market observations in our sample were affected by this law. The average AA airport presence is significantly higher and AA cost significantly lower than in CT sample as they focus on a larger set of markets and only use data from second quarter of 2001. These numbers show that AA had a strong presence at Dallas airports and that it mainly operated direct flights to all the destinations outside Dallas, which is in line with the fact that Dallas has been traditionally a major hub for AA.

8.1 Estimation results

We estimate the confidence sets using our profiled likelihood criterion with simulated critical value described in Section 5.1.1. We draw 1000 random standard normal vectors in order to obtain this critical value. The stage game in our model has the form:

		WN	
		1	0
AA	1	$X_1\beta - \varepsilon_1, X_2\beta - \varepsilon_2$	$\alpha + X_1\beta - \varepsilon_1, 0$
	0	$0, \alpha + X_2\beta - \varepsilon_2$	$0, 0$

Thus, we restrict both the competitive effect (α) and the effect of covariates (β) to be the same for AA and WN. We also assume that the selection probability π does not depend on (X_1, X_2) , i.e. firms coordinate “static” equilibrium selection across markets. We discretize all continuous variables as binary (below/above mean) and include a constant in both X_1 and X_2 . We consider both strong ($\delta = 0.75$) and weak ($\delta = 0.95$) discounting of future payoffs by the carriers. Table 6 contains the confidence sets estimated using our procedure.

We observe that the data provides useful information about the underlying model parameters. In particular, we can identify the sign of all coefficients besides the competitive effect and the cost variable. Our estimates are also in the same ballpark as the estimates obtained using the static game approach in CT. Our model, however, does not contain a lot of information about the competitive

¹²Note that we assume that the same repeated game is played in each of these markets, thus we focus on, what we believe to be, a homogenous group of Dallas markets.

Table 6: Estimated confidence sets from repeated game with random states, Dallas market

	$\delta = 0.75$		$\delta = 0.95$	
	90% CS	95% CS	90% CS	95% CS
competitive effect (α)	[-37.26 , 33.92]	[-37.26 , 33.92]	[-60.34 , 33.64]	[-60.34 , 33.64]
market size	[0.34 , 0.42]	[0.32 , 0.43]	[0.33 , 0.43]	[0.32 , 0.44]
Wright	[-0.65 , -0.54]	[-0.66 , -0.53]	[-0.65 , -0.54]	[-0.67 , -0.53]
airport presence	[4.57 , 4.67]	[4.56 , 4.68]	[4.54 , 4.7]	[4.53 , 4.71]
cost	[-0.01 , 0.09]	[-0.03 , 0.11]	[-0.02 , 0.1]	[-0.03 , 0.11]
constant	[3.55 , 3.96]	[3.51 , 4]	[-3.42 , 3.37]	[-3.59 , 3.37]

Note: Critical values for confidence sets calculated using simulated critical value described in Display 1 with 1000 random Normal draws. The sets were built using a grid search with step 0.01. The dataset contains 244 markets out of/to Dallas.

effect as the estimated bounds are very wide and include both positive and negative values.¹³ Interestingly, the bounds on other coefficients are very tight, suggesting that these parameters are close to being point-identified. As expected, the bounds obtained for the discount factor $\delta = 0.95$ are, in general, wider than those for $\delta = 0.75$ which corresponds to the fact that the set of possible equilibrium continuation values \mathcal{V} expands as the discount factor grows. Though, the difference in the width of the confidence set is only pronounced for the competitive effect α , with bounds for the near-to-point-identified parameters being very close for both discount factors.

In order to gain some insight about the importance of dynamic strategic interactions in the Dallas market, we compared our estimated confidence sets to the estimates from a simple logit run for each airline with the presence of the other airline as a dummy variable, including the same covariates as in Table 6. The results suggest that using the logit model, which does not properly account for strategic interactions between firms and the resulting endogeneity, leads to biased estimates of the effect of market size and the Wright amendment on payoffs – logit estimates imply a negative effect of market size on WN payoffs (estimated coefficient of -0.07) and positive effect of Wright amendment on AA payoffs (0.76) whereas our bounds identify the opposite signs for the effects of these variables, in line with the intuition.

¹³Note that the 90% and 95% confidence sets for α are the same. This is because we use a 0.01 step in the grid search for building the confidence sets. The log-likelihood is steep around the CS endpoints so the difference in the endpoints of 90% and 95% sets is less than 0.01.

8.2 Policy experiment

Finally, we use our estimated confidence sets to simulate the effect of repealing the Wright amendment from 2006. We look at the change in predicted probabilities in our model: $p(1,1)$, $p(1,0)$, $p(0,1)$, $p(0,0)$, with the actual policy and the scenario in which the Wright amendment is in operation throughout the sample period 1993-2017. In order to obtain the confidence sets for the policy effects: 1) we take all possible combinations of extreme points of the confidence sets, 2) we randomly draw 10000 parameter values from the estimated confidence sets, and for each point selected in these ways we calculate the change in probabilities.¹⁴ Then we take the union of the sets of estimated policy effects from 1) and 2). As in CT, we focus on markets out of Dallas Love airport, which were restricted by the amendment. We use the 95% confidence sets from Table 6.

Table 7: Estimated effect of repealing the Wright amendment, 95% level

		Change in probability (Wright - non Wright)
Both AA and WN	$p(1,1)$	$[-0.177, 0]$
Only AA	$p(1,0)$	$[-0.138, 0.163]$
Only WN	$p(0,1)$	$[-0.048, 0.116]$
Not served	$p(0,0)$	$[0, 0.103]$

Note: The change in probabilities is calculated as the predicted probability with actual trajectory of *Wright* minus the predicted probability assuming that the Wright amendment is in place throughout the sample period 1993-2017. The other variables are kept at their sample values.

Table 7 shows that keeping the Wright amendment in place would most likely decrease the number of markets served by both AA and WN, even by up to 17.7%, and increase the number of non-served markets by up to 10.3%. However, the results do not provide a decisive prediction for what would happen with markets served only by AA or only by WN, with the possibility that keeping Wright amendment in place would increase the number of these markets. The latter result suggests that keeping the Wright amendment in force might have restricted competition in significant number of markets, preventing entry of competitive airlines.

Comparing our estimated policy effects to results in Section 6 of CT we see that our model predicts much more modest, and thus more plausible, effect of the Wright amendment. CT predict even a 63.84% drop in the number of non-served markets after the repeal of the amendment, with

¹⁴For each configuration of the parameters we take $\theta_2 = (\pi, V)$ that minimizes the log-likelihood (see Assumption INF3).

possibly up to 47.44% of these markets served by American and/or Southwest. As the static game model in CT ignores the fact that entry by one airline today may be retaliated in the future by the other airline, it overestimates the negative effect of restricting the traffic in the Dallas market, which leads to exaggerated estimates of the amount of entry once the restrictions are abandoned.

Overall, our results in this section illustrate the usefulness of our approach to analysing repeated strategic interactions between firms and show that combining the repeated games model with the data may lead to interesting findings and improved analysis of strategic interactions in the US airline market.

9 Discussion

Although we focus on repeated games with random states, our identification and inference approach can be extended to general stochastic games with state dependence. Recent advancements in the analysis of these games (Abreu et al. (2016), Abreu et al. (2020)) provide readily available procedures for computing equilibrium payoff sets, which can be naturally embedded into our econometric approach. Additionally, the assumption of binary action sets made in our exposition can be relaxed as the computation algorithm in Abreu & Sannikov (2014) allows non-binary actions and our inference procedure in Display 1 applies in this case, though it is important to note that the computational burden increases with the dimension of the action space as dimension of the continuation vector V grows exponentially with $|\mathcal{A}|$.

As we allow a wide range of possible equilibria, the confidence sets may be quite large in empirical applications. One may restrict the number of possible equilibria in these games by refining the equilibrium concept, e.g. focusing on strategy-proof equilibria, which should result in smaller continuation payoff sets \mathcal{V} and narrower bounds on the parameters of interest. We leave these extensions for further research.

We focus on entry decisions in the dynamic context in our paper, thus we leave out the pricing decisions. Goolsbee & Syverson (2008) show that airlines may adjust prices in advance when faced with the threat of entry, thus adding pricing to the model would be an important extension. See Ciliberto et al. (2020) for a recent effort to model entry and pricing decisions at the same time using a static game.

Appendix

A Mathematical Proofs

A.1 Proof of Theorem 1

First note that under quasi-concavity:

$$\inf_{\theta_1 \in \Theta_{01}} \sup_{\theta_2 \in \Theta_2(\theta_1)} L_T(\theta_1, \theta_2) = \min_{\theta_1 \in \{\underline{\theta}_1, \bar{\theta}_1\}} \sup_{\theta_2 \in \Theta_2(\theta_1)} L_T(\theta_1, \theta_2) \quad (6)$$

Further, Assumptions [INF2\(c\)-\(g\)](#) allow us to apply Proposition 5.1 and Lemma F.1 in [Chen et al. \(2018\)](#) in order to obtain quadratic expansion of the likelihood in γ :

$$\begin{aligned} \sup_{\theta \in \Theta_{oT}} 2L_T(\theta) &= 2 \sum_{t=1}^T \log p(Y_t, \gamma_0) + \|V_T\|^2 - \inf_{\theta \in \Theta_{oT}} \|\sqrt{T}\mathbb{I}_0^{1/2} \gamma(\theta) - V_T\|^2 + o_p(1) = \\ &= 2 \sum_{t=1}^T \log p(Y_t, \gamma_0) + \|V_T\|^2 + o_p(1) \end{aligned}$$

where $V_T = \mathbb{I}_0^{-1/2} \frac{1}{\sqrt{T}} \sum_{t=1}^T \frac{\partial \log p(Y_t, \gamma_0)}{\partial \gamma}$. Now for the restricted part:

$$\begin{aligned} \sup_{\theta \in \Theta_{oT}(\theta_1)} 2L_T(\theta) &= 2 \sum_{t=1}^T \log p(Y_t, \gamma_0) + \|V_T\|^2 - \inf_{\theta=(\theta_1, \theta_2) \in \Theta_{oT}} \|\sqrt{T}\mathbb{I}_0^{1/2} \gamma(\theta) - V_T\|^2 + o_p(1) = \\ &= 2 \sum_{t=1}^T \log p(Y_t, \gamma_0) + \|V_T\|^2 - \inf_{\kappa \in \Gamma_{oT}(\theta_1)} \|\kappa - V_T\|^2 + o_p(1) = \\ &= 2 \sum_{t=1}^T \log p(Y_t, \gamma_0) + \|V_T\|^2 - \inf_{\kappa \in \mathcal{K}(\theta_1)} \|\kappa - V_T\|^2 + o_p(1) \end{aligned}$$

This implies:

$$LR_T(\theta_1) = 2 \left[\sup_{\theta \in \Theta_{oT}} L_T(\theta) - \sup_{\theta \in \Theta_{oT}(\theta_1)} L_T(\theta) \right] = \inf_{\kappa \in \mathcal{K}(\theta_1)} \|\kappa - V_T\|^2 + o_p(1)$$

Now use Assumption [INF3](#) and proceed similarly to [Shapiro \(1985\)](#). Note that the cone $\mathcal{K}(\theta_1)$ can be approximated by:

$$\left\{ \frac{\partial \gamma(\theta_0)}{\partial \theta'_2} s : s \in K_2(\theta_1), \theta_0 = (\theta_1, \theta_2) \in \Theta_0 \right\} = \{\Delta_{\theta_1} s : s \in K_2(\theta_1)\}$$

for $\theta_1 \in \{\underline{\theta}_1, \bar{\theta}_1\}$, which implies that:

$$LR_T(\theta_1) = \inf_{s \in K_2(\theta_1)} \|V_T - \Delta_{\theta_1} s\|^2 + o_p(1)$$

and together with [\(6\)](#) concludes the proof.

A.2 Proof of Theorem 2

First we will demonstrate that $\hat{\theta}_1 \rightarrow^p \underline{\theta}_1$ and $\hat{\bar{\theta}}_1 \rightarrow^p \bar{\theta}_1$ as $T \rightarrow \infty$. For this purpose we apply Theorem 3.1 in [Chernozhukov et al. \(2007\)](#). Their condition C.1 is satisfied as follows: part (a) follows from our Assumption [INF2\(a\)](#), lower-semicontinuity of $E[\log p(Y_t, \theta)]$ in part (b) holds in the neighbourhood \mathcal{N}_θ of Θ_0 by our Assumptions [INF2\(d\)](#) and [INF2\(e\)](#), part (c) follows from our continuity assumptions on $p(Y_t, \theta)$ and discreteness of Y_t , uniform convergence in part (d) can be shown to hold over \mathcal{N}_θ by applying Jennrich's ULLN with the help of our Assumptions [INF2\(a\),\(d\),\(e\)](#) and noting that $0 \leq p(Y_t, \theta) \leq 1$.

Next recall that B_{k_T} denotes a ball centred at zero with radius $k_T \rightarrow \infty$ and define $KB_{k_T}(\theta_1) = \{\kappa : \kappa = \Delta_{\theta_1} s, s \in B_{k_T}\}$. If $\Delta_{\theta_1} = 0$, then trivially $\inf_{s \in K^{LF}(\theta_1)} \|V_T - \Delta_{\theta_1} s\|^2 = \inf_{s \in K^{LF}(\theta_1) \cap B_{k_T}} \|V_T - \Delta_{\theta_1} s\|^2$ so we focus on the case when $\Delta_{\theta_1} \neq 0$. Note that in this case $KB_{k_T}(\theta_1)$ is an ellipsoid. For θ_1 in the neighbourhoods of $\underline{\theta}_1$ and $\bar{\theta}_1$ we have:

$$P \left(\inf_{s \in K^{LF}(\theta_1)} \|V_T - \Delta_{\theta_1} s\|^2 - \inf_{s \in K^{LF}(\theta_1) \cap B_{k_T}} \|V_T - \Delta_{\theta_1} s\|^2 \neq 0 \right) \leq P(V_T \notin KB_{k_T}(\theta_1))$$

but as the elliptic radi grow with k_T we have that $P(V_T \notin KB_{k_T}(\theta_1)) \rightarrow 0$ as $T \rightarrow \infty$. Thus we can write:

$$\inf_{s \in K^{LF}(\theta_1)} \|V_T - \Delta_{\theta_1} s\|^2 = \inf_{s \in K^{LF}(\theta_1) \cap B_{k_T}} \|V_T - \Delta_{\theta_1} s\|^2 + o_p(1) \quad (7)$$

Assumption [INF5](#) implies that $K^{LF}(\theta_1) \cap B_{k_T}$ is a continuous (compact-valued) correspondence around $\underline{\theta}_1$ and $\bar{\theta}_1$. Additionally Δ_{θ_1} is continuous in θ_1 in this neighbourhood by Assumption [INF2\(e\)](#). Thus, we can apply Berge's maximum theorem to conclude that $\inf_{s \in K^{LF}(\theta_1) \cap B_{k_T}} \|V_T - \Delta_{\theta_1} s\|^2$ is a continuous function of θ_1 . Now [\(7\)](#) and continuous mapping theorem imply:

$$\begin{aligned} \inf_{s \in K^{LF}(\underline{\theta}_1) \cap B_{k_T}} \|V_T - \Delta_{\underline{\theta}_1} s\|^2 &= \inf_{s \in K^{LF}(\underline{\theta}_1)} \|V_T - \Delta_{\underline{\theta}_1} s\|^2 + o_p(1) \\ \inf_{s \in K^{LF}(\hat{\bar{\theta}}_1) \cap B_{k_T}} \|V_T - \Delta_{\hat{\bar{\theta}}_1} s\|^2 &= \inf_{s \in K^{LF}(\bar{\theta}_1)} \|V_T - \Delta_{\bar{\theta}_1} s\|^2 + o_p(1) \end{aligned} \quad (8)$$

as $T \rightarrow \infty$.

Next note that $K^{LF}(\theta_1) \subseteq K_2(\theta_1)$. This is trivially satisfied when $K_2(\theta_1) = \mathbb{R}^{d_2}$ or $K_2(\theta_1)$ is an orthant itself. Consider the remaining case when $K_2(\theta_1) = \mathbb{R}_+^{d_+} \times \mathbb{R}_-^{d_-} \times \mathbb{R}^{d_2 - d_+ - d_-}$ where $0 \leq d_+ + d_- \leq d_2 - 1$. Now we must have $K^{LF}(\theta_1) = \mathbb{R}_+^{d_+} \times \mathbb{R}_-^{d_-} \times \mathbb{R}^{\tilde{d}_+} \times \mathbb{R}^{\tilde{d}_-}$ with $\tilde{d}_+ + \tilde{d}_- = d_2 - d_+ - d_-$. To see that, without loss of generality, suppose that $K^{LF}(\theta_1) = \mathbb{R}_- \times \mathbb{R}_+^{d_+ - 1} \times \mathbb{R}_-^{d_-} \times \mathbb{R}^{\tilde{d}_+} \times \mathbb{R}^{\tilde{d}_-}$ and let \tilde{C} be the corner associated with this parameter space and $\theta_2 = \theta_2^*(\theta_1)$ be the profiled-likelihood-minimising value. Now note that the first coordinate of \tilde{C} , \tilde{C}_1 , has to be different than the first coordinate of θ_2 , $\theta_{2,1}$, but these are the same for the closest corner, i.e. $C_{\theta_1,1} = \theta_{2,1}$. We have:

$$\|\tilde{C} - \theta_2\|^2 = |\tilde{C}_1 - \theta_{2,1}|^2 + \|\tilde{C}_{-1} - \theta_{2,-1}\|^2 > \inf_{C \in \mathcal{C}} \|C_{-1} - \theta_{2,-1}\|^2 = \|C_{\theta_1} - \theta_2\|^2$$

which implies that \tilde{C} cannot be the closest corner to θ_2 .

Finally, we have:

$$\max \left\{ \inf_{s \in K^{LF}(\underline{\theta}_1)} \|V_T - \Delta_{\underline{\theta}_1} s\|^2, \inf_{s \in K^{LF}(\bar{\theta}_1)} \|V_T - \Delta_{\bar{\theta}_1} s\|^2 \right\} \geq \max \left\{ \inf_{s \in K_2(\underline{\theta}_1)} \|V_T - \Delta_{\underline{\theta}_1} s\|^2, \inf_{s \in K_2(\bar{\theta}_1)} \|V_T - \Delta_{\bar{\theta}_1} s\|^2 \right\}$$

which together with [\(8\)](#) concludes the proof.

A.3 Proof of Theorem [3](#)

For a closed convex cone K let K° denote its polar cone (see e.g. Section 14 in [Rockafellar \(1970\)](#)). From the proof of Theorem [1](#) and Moreau's decomposition theorem (note that $\mathcal{K}(\theta_1)$ is a closed

convex cone by Assumption [INF2\(g\)](#)):

$$LR_T(\theta_1) = \|V_T\|^2 - \inf_{\kappa \in \mathcal{K}^o(\theta_1)} \|\kappa - V_T\|^2 + o_p(1)$$

which implies $LR_T(\theta_1) \leq \|V_T\|^2 + o_p(1)$ and

$$\sup_{\theta_1 \in \Theta_{01}} LR_T(\theta_1) \leq \max_{\theta_1 \in \{\underline{\theta}_1, \bar{\theta}_1\}} \{\|V_T\|^2, \|V_T\|^2\} + o_p(1) = \|V_T\|^2 + o_p(1)$$

and the result follows from asymptotic normality of V_T .

B Alternative definitions of market presence

In [Table 1](#) an airline is present in the market if it operates at least one flight from the market origin to market destination. Here we consider other definitions of market presence. First, we use DB1B ticketing data, which contains 10% sample of airline tickets from reporting carriers, and redefine market presence as selling at least 5 tickets for the specified route (see [Table 8](#)). Next, we use T100 Segment data and redefine market as a segment of the trip, for example a flight from ORD to MIA through DCA contains two segments ORD-DCA and DCA-MIA (using previous definition this would only be a single market ORD-MIA).

Table 8: Share of markets by presence of major (ticketing) carriers over time (in %)

Presence in Q2 1993 - Q2 2002 - Q2 2012	American	Delta	United	US Airways	Southwest
in - in - in	43.4	61.1	74.8	42.8	86.8
in - out - in	9	5.9	9.2	12.3	4.8
in - out - out	37	23.1	12.8	27	6
in - in - out	10.6	9.9	3.2	17.9	2.4

Note: DB1B Market data, flights with less than 5 tickets in 1993 dropped. Markets are defined as directional routes between origin and destination airports (irrespective of the number of stops on the way). "In" means that a carrier served at least one flight on the route.

The numbers in [Table 8](#) significantly differ from those in [Table 1](#) as ubiquitous codeshare and interlining agreements drive a wedge between the definitions of operating and ticketing carrier. The differences are smaller between [Table 9](#) and [Table 1](#). Despite these differences the main message remains the same – there is substantial amount of entry and exit across time in the US airline

market.

Table 9: Share of markets by presence of major carriers over time (in %)

Presence in Q2 1990 - Q2 2000 - Q2 2010	American	Delta	United	US Airways	Southwest
in - in - in	34	35.8	38.6	16.5	74.6
in - out - in	5	6.8	2.7	3.5	3.1
in - out - out	46.4	33.4	35.8	53.8	14
in - in - out	14.6	24.1	23	26.3	8.3

Note: T100 Segment data, flights with less than 20 passengers in 1990 dropped. Markets are defined as segments of directional routes between origin and destination airports. "In" means that a carrier served at least one flight on the route.

C Moment inequality characterisation

For a non-empty $A \subseteq \mathcal{A}$ let $\mathcal{L}(A; \alpha, V)$ denote the probability of observing some $a \in A$ in equilibrium in the normal form game under the assumption that in the regions of multiple equilibria an equilibrium in A is always selected. For simplicity let Assumption [INF1](#) hold. Following [Galichon & Henry \(2011\)](#) the marginal identified set for α can be characterised using moment inequalities by:

$$\Theta_{01}^S = \{\alpha : E_0(\mathbb{1}\{a \in A_j\}) \leq \mathcal{L}(A_j; \alpha, V), A_j \subseteq \mathcal{A}, V \in \mathcal{V}_S(\alpha)\} \quad (9)$$

where $j = 1, 2, \dots, J$. Now for inference one can implement either the profiled procedure in [Kaido et al. \(2019\)](#) (KMS) or [Bugni et al. \(2017\)](#) (BCS). We discuss how the computational burden of these procedures compares to our approach as computation is the main obstacle for a practical inference in our model.¹⁵ For this discussion we employ similar notation as in Section 5 in the paper, namely $\theta_1 \equiv \alpha$, $\theta_2 \equiv V$, $\theta = (\theta_1, \theta_2)$.

C.1 KMS inference

Let $G_{n,j}^b$ be a standardised estimator of $E_0(\mathbb{1}\{a \in A_j\})$ evaluated on a bootstrap sample scaled by \sqrt{n} and let $\hat{D}_{n,j}(\theta)$ denote the gradient of $\mathcal{L}(A_j; \alpha, V)$ w.r.t. α and V normalised by the sample

¹⁵To be clear, both KMS and BCS apply to a much wider range of problems than our approach. Thus we only argue that for our specialised model the simulated critical value described in the paper may be preferable to KMS and BCS from a practical perspective.

standard deviation of moment j , $\hat{\sigma}_{n,j}$. Further, let $\hat{\xi}_{n,j}(\theta)$ denote $(\iota_n \hat{\sigma}_{n,j})^{-1} \sqrt{n}$ times the sample estimator of moment j , where $\iota_n \rightarrow \infty$. The KMS critical value is obtained by bootstrapping:

$$\Lambda_n^b(\theta, \rho, c) = \{\lambda \in \sqrt{n}(\Theta - \theta) \cap \rho B^d : G_{n,j}^b + \hat{D}_{n,j}(\theta)\lambda + \psi_j(\hat{\xi}_{n,j}(\theta)) \leq c, j = 1, 2, \dots, J\}$$

where ρB^d imposes a technical “box” constraint on the local parameter space and ψ_j is a Generalised Moment Selection function of [Andrews & Soares \(2010\)](#), and can be calculated as:

$$\hat{c}(\theta) = \inf\{c \in \mathbb{R}_+ : P^*(\Lambda_n^b(\theta, \rho, c) \cap \{\lambda_1 = 0\}) \neq \emptyset\} \geq \kappa\}$$

where P^* denotes the law induced by bootstrap sampling and λ_1 is the first element of λ . Finally, the marginal confidence set is built by finding lowest and highest value of θ_1 for which the sample moment inequalities are satisfied with slackness $\hat{c}(\theta_1, \theta_2)$.

Let us now compare our inference method to KMS. Note that the computationally difficult step in our model is the re-evaluation of the continuation value set $\Theta_2(\theta_1)$ for different values of θ_1 , which will be embedded in evaluating $\sqrt{n}(\Theta - \theta)$ within $\Lambda_n^b(\theta, \rho, c)$ in the KMS procedure. Our procedure in Display 1 controls the number of evaluations of $\Theta_2(\theta_1)$ by controlling the size of the grid for candidate values of θ_1 in the pre-estimation of the identified set in Step 1 and re-using evaluations of the likelihood ratio from Step 1 in building the confidence set in the final Step 5.

Similarly, the first step in the KMS procedure in which candidate values of θ are drawn can be adjusted to include only θ 's on the grid of values for θ_1 to limit number of evaluations of $\Theta_2(\theta_1)$. However, as currently implemented, the KMS procedure proceeds with a smooth iterative algorithm to generate further “good” candidate values of θ (“A-M steps”) and, thus, requires recalculation of $\Theta_2(\theta_1)$ for these newly generated values. As the number of iterations required for this algorithm to converge may differ from application to application, it may be difficult to control the number of evaluations of $\Theta_2(\theta_1)$ in practice without significant changes to this algorithm. Therefore, it seems that using the moment inequality characterisation in (9) and KMS is unlikely to dominate our method in terms of computational convenience.

C.2 BCS inference

The main BCS critical value is obtained by taking a minimum over two profiled bootstrap statistics, $T_n^{DR}(\theta_1)$ and $T_n^{PR}(\theta_1)$, in order to improve power. As our inference procedure is conservative it seems fair to compare it to $T_n^{DR}(\theta_1)$ and $T_n^{PR}(\theta_1)$ separately rather than to the more computationally intensive minimum statistic.

Using the notation from the previous section BCS resampling statistics can be written as:

$$T_n^{DR}(\theta_1) = \inf_{\theta_2 \in \hat{\Theta}_2(\theta_1)} \sum_{j=1}^J \left[G_{n,j}^b + \psi_j(\hat{\xi}_{n,j}(\theta)) \right]_-$$

$$T_n^{PR}(\theta_1) = \inf_{\theta_2 \in \Theta_2(\theta_1)} \sum_{j=1}^J \left[G_{n,j}^b + \hat{\xi}_{n,j}(\theta) \right]_-$$

where $\hat{\Theta}_2(\theta_1)$ is the set of minimizers of the KMS test statistic (see their paper for details). Note that similarly to our approach we can control the number of evaluations of $\Theta_2(\theta_1)$ by imposing a grid on θ_1 . However, note that resampling both $T_n^{DR}(\theta_1)$ and $T_n^{PR}(\theta_1)$ requires repeatedly solving a non-linear non-convex constrained optimisation problem. Also, as argued in the main text, solution to this problem will often be reached on the boundary of the set $\hat{\Theta}_2(\theta_1)$ and $\Theta_2(\theta_1)$.¹⁶ This is much more computationally expensive than repeatedly solving a convex optimisation problem in our simulation procedure in Display 1.

D Inference without Assumption INF3

In this section we discuss how we can adjust our simulated critical value if $\theta^*(\theta_1)$ is not unique and $\partial\gamma(\theta_0)/\partial\theta'_2$ contains zero rows for some θ_0 . This will happen, for example, if the true probabilities, γ_0 , are flat on a set with non-empty interior.

Firstly, the main complication here comes from the fact that now the local parameter space for γ at γ_0 cannot be approximated simply by taking $\left\{ \frac{\partial\gamma(\theta_0)}{\partial\theta'_2} s : s \in K_2(\theta_1), \theta_0 = (\theta_1, \theta_2) \in \Theta_0 \right\}$ (cf. proof of Theorem 1) as this set maps to zero when Θ_0 is a compact set with non-empty

¹⁶If we impose a condition equivalent of Assumption INF3, namely we have $\hat{\Theta}_2(\theta_1) = \{\theta_2^*(\theta_1)\}$ the DR statistic seems computationally attractive compared to our approach. Still it requires evaluation of $\theta_2^*(\theta_1)$ for all the candidate values θ_1 whereas we only need to calculate it for the endpoints of the pre-estimated marginal identified set, $\hat{\theta}_1$ and $\hat{\bar{\theta}}_1$.

interior. However, given that γ is smooth around γ_0 (see Assumption [INF2\(e\)](#)) a simple expansion $\Theta_{0\eta} = \{\theta \in \Theta : \inf_{\theta_0 \in \Theta_0} \|\theta - \theta_0\| \leq \eta\}$ for $\eta > 0$ would allow us to bound the asymptotic distribution of our profiled criterion as:

$$\begin{aligned} \sup_{\theta_1 \in \Theta_{01}} LR_T(\theta_1) &\leq \max_{\theta_1 \in \{\underline{\theta}_1, \bar{\theta}_1\}} \left\{ \kappa \in \bigcup_{(\theta_1, \theta_2) \in \Theta_{0\eta}} \left\{ \inf_{\frac{\partial \gamma(\theta_1, \theta_2)}{\partial \theta'_2} s : s \in K_2^{\theta_2}(\theta_1) \right\} \|V_T - \kappa\|^2 \right\} + o_p(1) \\ &= \max_{\theta_1 \in \{\underline{\theta}_1, \bar{\theta}_1\}} \left\{ \inf_{(\theta_1, \theta_2) \in \Theta_{0\eta}} \inf_{s \in K_2^{\theta_2}(\theta_1)} \left\| V_T - \frac{\partial \gamma(\theta_1, \theta_2)}{\partial \theta'_2} s \right\|^2 \right\} + o_p(1) \end{aligned}$$

where $K_2^{\theta_2}(\theta_1)$ is a cone approximating the local parameter space at $\theta_2 \in \theta_2^*(\theta_1)$.

Now in order to approximate the statistic on the right-hand side above, we can replace $\underline{\theta}_1$ and $\bar{\theta}_1$ with $\hat{\underline{\theta}}_1$ and $\hat{\bar{\theta}}_1$ as before. Next note that the identified set estimator in [Chernozhukov et al. \(2007\)](#) approximates the identified set from “outside”, thus in practice we can replace $\Theta_{0\eta}$ with their estimator $\hat{\Theta}_0$ (note that we only have to estimate a “slice” out of $\hat{\Theta}_0$ for $\hat{\underline{\theta}}_1$ and $\hat{\bar{\theta}}_1$). Finally, note that now the (set of) closest corner(s) to $\theta_2 \in \theta_2^*(\theta_1)$ depends on the value of θ_2 , $C_{(\theta_1, \theta_2)}$, and Assumption [INF5](#) is too strong in this setup. Thus, letting $K^{LF}(C)$ denote the orthant corresponding to the corner $C \in C_{(\theta_1, \theta_2)}$ we can replace $\inf_{s \in K_2^{\theta_2}(\theta_1)} \left\| V_T - \frac{\partial \gamma(\theta_1, \theta_2)}{\partial \theta'_2} s \right\|^2$ above with $\max_{C \in C_{(\theta_1, \theta_2)}} \inf_{s \in K^{LF}(C)} \left\| V_T - \frac{\partial \gamma(\theta_1, \theta_2)}{\partial \theta'_2} s \right\|^2$ in order to simulate a conservative critical value.

Acknowledgments: We would like to thank Mike Abito, Wojciech Olszewski, Subir Bose and Andre Stenzel for useful discussions and suggestions and to Arpita Ghosh and Shuai Zhao for providing competent research assistance. We are grateful to participants in the LSE Joint Econometrics and Statistics Workshop, the (Ce)2 Workshop, the UEA Economics seminar and various conferences for useful comments. This research used the ALICE High Performance Computing Facility at the University of Leicester.

References

- Abreu, D., Brooks, B. & Sannikov, Y. (2016), A “pencil-sharpening” algorithm for two player stochastic games with perfect monitoring. Working Paper.
- Abreu, D., Brooks, B. & Sannikov, Y. (2020), ‘Algorithms for stochastic games with perfect monitoring’, *Econometrica* **88**(4), 1661–1695.
- Abreu, D., Pearce, D. & Stacchetti, E. (1990), ‘Toward a theory of discounted repeated games with imperfect monitoring’, *Econometrica* **58**(5), pp. 1041–1063.

- Abreu, D. & Sannikov, Y. (2014), ‘An algorithm for two-player repeated games with perfect monitoring’, *Theoretical Economics* **9**(2), 313–338.
- Al Mohamad, D., Van Zwet, E. W., Cator, E. & Goeman, J. J. (2020), ‘Adaptive critical value for constrained likelihood ratio testing’, *Biometrika* **107**(3).
- Amemiya, T. (1986), *Advanced econometrics*, Blackwell, Oxford.
- Andrews, D. W. K. (2001), ‘Testing when a parameter is on the boundary of the maintained hypothesis’, *Econometrica* **69**(3), 683–734.
- Andrews, D. W. K. & Soares, G. (2010), ‘Inference for parameters defined by moment inequalities using generalized moment selection’, *Econometrica* **78**(1), 119–157.
- Bajari, P., Benkard, C. L. & Levin, J. (2007), ‘Estimating dynamic models of imperfect competition’, *Econometrica* **75**(5), 1331–1370.
- Bajari, P., Hong, H. & Ryan, S. P. (2010), ‘Identification and estimation of a discrete game of complete information’, *Econometrica* **78**(5), 1529–1568.
- Beresteanu, A., Molchanov, I. & Molinari, F. (2011), ‘Sharp identification regions in models with convex moment predictions’, *Econometrica* **79**(6), 1785–1821.
- Berry, S. & Jia, P. (2010), ‘Tracing the woes: An empirical analysis of the airline industry’, *American Economic Journal: Microeconomics* **2**(3), 1–43.
- Bontemps, C. & Kumar, R. (2020), ‘A geometric approach to inference in set-identified entry games’, *Journal of Econometrics* **218**(2).
- Bugni, F. A., Canay, I. A. & Shi, X. (2017), ‘Inference for subvectors and other functions of partially identified parameters in moment inequality models’, *Quantitative Economics* **8**(1), 1–38.
- Chen, X., Christensen, T. M. & Tamer, E. (2018), ‘Monte Carlo confidence sets for identified sets’, *Econometrica* **86**(6), 1965–2018.
- Chernozhukov, V., Hong, H. & Tamer, E. (2007), ‘Estimation and confidence regions for parameter sets in econometric models’, *Econometrica* **75**(5), 1243–1284.
- Ciliberto, F., Murry, C. & Tamer, E. T. (2020), Market structure and competition in airline markets. Working Paper.
- Ciliberto, F. & Tamer, E. (2009), ‘Market structure and multiple equilibria in airline markets’, *Econometrica* **77**(6), 1791–1828.
- Galichon, A. & Henry, M. (2011), ‘Set identification in models with multiple equilibria’, *The Review of Economic Studies* **78**(4), 1264–1298.
- Gerardi, K. & Shapiro, A. (2009), ‘Does competition reduce price dispersion? New evidence from the airline industry’, *Journal of Political Economy* **117**(1), 1–37.
- Giacomini, R. & Kitagawa, T. (2020), Robust Bayesian inference for set-identified models. CEMMAP Working Paper.
- Goolsbee, A. & Syverson, C. (2008), ‘How do incumbents respond to the threat of entry? Evidence from the major airlines’, *The Quarterly Journal of Economics* **123**(4), 1611–1633.

- Kaido, H., Molinari, F. & Stoye, J. (2019), ‘Confidence intervals for projections of partially identified parameters’, *Econometrica* **87**, 1397–1432.
- Kaido, H., Molinari, F. & Stoye, J. (2020), Constraint qualifications in partial identification. Working Paper.
- Kline, B. & Tamer, E. (2016), ‘Bayesian inference in a class of partially identified models’, *Quantitative Economics* **7**(2), 329–366.
- Kudo, A. (1963), ‘A multivariate analogue of the one-sided test’, *Biometrika* **50**(3/4), 403–418.
- Magnac, T. & Thesmar, D. (2002), ‘Identifying dynamic discrete decision processes’, *Econometrica* **70**(2), pp. 801–816.
- Mailath, G. & Samuelson, L. (2006), *Repeated Games and Reputations: Long-Run Relationships*, Oxford University Press.
- Otsu, T., Pesendorfer, M. & Takahashi, Y. (2016), ‘Pooling data across markets in dynamic Markov games’, *Quantitative Economics* **7**(2), 523–559.
- Rockafellar, R. T. (1970), *Convex analysis*, Princeton Mathematical Series, Princeton University Press.
- Romano, J. P. & Shaikh, A. M. (2008), ‘Inference for identifiable parameters in partially identified econometric models’, *Journal of Statistical Planning and Inference* **138**(9), 2786–2807.
- Shapiro, A. (1985), ‘Asymptotic distribution of test statistics in the analysis of moment structures under inequality constraints’, *Biometrika* **72**(1), 133–144.
- Silvapulle, M. J. & Sen, P. K. (2001), *Constrained Statistical Inference*, John Wiley & Sons, Inc.
- Tamer, E. (2003), ‘Incomplete simultaneous discrete response model with multiple equilibria’, *The Review of Economic Studies* **70**(1), 147–165.