

Kriřtoufek, Ladislav

**Working Paper**

## Classical and modified rescaled range analysis: Sampling properties under heavy tails

IES Working Paper, No. 26/2009

**Provided in Cooperation with:**

Charles University, Institute of Economic Studies (IES)

*Suggested Citation:* Kriřtoufek, Ladislav (2009) : Classical and modified rescaled range analysis: Sampling properties under heavy tails, IES Working Paper, No. 26/2009, Charles University in Prague, Institute of Economic Studies (IES), Prague

This Version is available at:

<https://hdl.handle.net/10419/83365>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

Institute of Economic Studies, Faculty of Social Sciences  
Charles University in Prague

# **Classical and modified rescaled range analysis: Sampling properties under heavy tails**

Ladislav Kristoufek

IES Working Paper: 26/2009



Institute of Economic Studies,  
Faculty of Social Sciences,  
Charles University in Prague

[UK FSV – IES]

Opletalova 26  
CZ-110 00, Prague  
E-mail : [ies@fsv.cuni.cz](mailto:ies@fsv.cuni.cz)  
<http://ies.fsv.cuni.cz>

Institut ekonomických studií  
Fakulta sociálních věd  
Univerzita Karlova v Praze

Opletalova 26  
110 00 Praha 1

E-mail : [ies@fsv.cuni.cz](mailto:ies@fsv.cuni.cz)  
<http://ies.fsv.cuni.cz>

**Disclaimer:** The IES Working Papers is an online paper series for works by the faculty and students of the Institute of Economic Studies, Faculty of Social Sciences, Charles University in Prague, Czech Republic. The papers are peer reviewed, but they are *not* edited or formatted by the editors. The views expressed in documents served by this site do not reflect the views of the IES or any other Charles University Department. They are the sole property of the respective authors. Additional info at: [ies@fsv.cuni.cz](mailto:ies@fsv.cuni.cz)

**Copyright Notice:** Although all documents published by the IES are provided without charge, they are licensed for personal, academic or educational use. All rights are reserved by the authors.

**Citations:** All references to documents served by this site must be appropriately cited.

**Bibliographic information:**

Kristoufek, L. (2009). “ Classical and modified rescaled range analysis: Sampling properties under heavy tails ” IES Working Paper 26/2009. IES FSV. Charles University.

This paper can be downloaded at: <http://ies.fsv.cuni.cz>

# Classical and modified rescaled range analysis: Sampling properties under heavy tails

Ladislav Kristoufek\*

\*IES, Charles University Prague and  
Institute of Information Theory and Automation,  
Academy of Sciences of the Czech Republic, Prague  
E-mail: kristoufek@ies-prague.org

November 2009

## **Abstract:**

Mostly used estimators of Hurst exponent for detection of long-range dependence are biased by presence of short-range dependence in the underlying time series. We present confidence intervals estimates for rescaled range and modified rescaled range. We show that the difference in expected values and confidence intervals enables us to use both methods together to clearly distinguish between the two types of processes. Moreover, both methods are robust against the presence of heavy tails in the underlying process.

**Keywords:** rescaled range, modified rescaled range, Hurst exponent, long-range dependence, confidence intervals

**JEL:** G1, G10, G14, G15

## **Acknowledgements:**

Support from the Czech Science Foundation under grant 402/09/0965 and 402/09/H045 and institutional Support from the Department of Education MSMT 0021620841 are gratefully acknowledged.

# 1 Introduction

Long-range dependence in the financial time series has been discussed in number of research papers (e.g. Peters, 1994; Di Matteo, Aste & Dacorogna, 2005; Di Matteo, 2007; Czarnecki, Grech & Pamula, 2008; Grech & Mazur, 2004; Carbone, Castelli & Stanley, 2004; Matos *et al.*, 2008; Vandewalle, Ausloos & Boveroux, 1997; and Alvarez-Ramirez *et al.*, 2008). However, the majority of the papers interpret the results on the basis of simple comparison of estimated self-similarity parameter – Hurst exponent  $H$  – with its asymptotic limit of 0.5. Hurst exponent of 0.5 indicates two possible processes – either independent (Beran, 1994) or short-range dependent process (Lillo & Farmer, 2004). Autocovariances of the independent process are insignificant for all non-zero lags whereas for the short-range dependent process (e.g. ARIMA processes, for details, see Box & Jenkins, 1970), autocovariances are significant at low lags and insignificant at high lags and decay exponentially. If  $H > 0.5$ , the process has significantly positive correlations at all lags and is said to be long-range dependent with positive correlations (Embrechts & Maejima, 2002) or persistent (Mandelbrot & van Ness, 1968). On the other hand, if  $H < 0.5$ , it has similar properties to the previous case as it has significantly negative correlations at all lags and the process is said to be long-range dependent with negative correlations (Embrechts & Maejima, 2002) or anti-persistent (Barkoulas, Baum & Travlos, 2000). Autocovariances of long-range dependent processes are hyperbolically decaying and are either non-summable for persistent or summable for anti-persistent processes (Beran, 1994).

The efficient distinguishing between the short and the long-range dependence is crucial for portfolio selection as well as option pricing as the long-range dependence is connected to infinite or non-existent variance (Peters, 1994). As the majority of Hurst exponent estimation methods is biased by presence of the short-range dependence in the underlying process (Morariu *et al.*, 2007; and Lo, 1991), we test two methods which are strongly related and one of them is robust against the short-range dependence. These two methods are the rescaled range analysis of Hurst (1951), which is also called the “classical”, and the modified rescaled range analysis of Lo (1991). To be able to distinguish between the two types of dependence, we construct the confidence intervals for both methods as the estimates of Hurst exponent can strongly deviate from the asymptotic limit of 0.5 even for an independent process (Weron, 2002; Grech & Mazur, 2005; and Peters, 1994). However, most of the authors base their simulations on an independent process which is normally distributed. As the markets are known to be non-normal and to have heavy tails (Cont, 2001), we present the results for processes based on both the standard normal distribution and the Cauchy distribution, which possesses the heaviest tails of all stable distributions. It can be said that we test the estimates on two extreme cases – normality and very heavy tails.

The paper is structured as follows. In Section 2, we present and describe both techniques in detail. In Section 3, we present the important aspects of Monte Carlo simulations methodology. In Section 4, we show results of Monte Carlo simulations for time

series lengths from 512 to 16 384 observations for both tested methods and both types of distributions of the underlying process. Both methods are robust to heavy tails in the underlying process and even have thinner confidence intervals than for the process based on standard distribution.

## 2 Classical and modified rescaled range analysis

### 2.1.Procedure

The classical rescaled range analysis (R/S) was developed by Edwin Hurst while working as an engineer in Egypt (Hurst, 1951) and was later applied to financial time series by Mandelbrot (1970). For the time series of length  $T$ , Hurst exponent  $H$  can be estimated from the behavior of rescaled range of the series which is described in the following theorem (for details, see Hurst, 1951; and Taqqu, Teverovsky & Willinger, 1995).

**Theorem.** Let us have time series  $\{V_t\}_{t=-\infty}^{\infty}$  and  $T = N * v$  where  $N \in \mathbb{N}$ ,  $v \in \mathbb{N}$ . Denote  $I_n = \{r_{k,n} = V_{(n-1)v+k}; k = 1, 2, \dots, v\}$ ,  $n = 1, 2, \dots, N$  and put arithmetic mean as  $\bar{r}_n = \sum_{k=1}^v r_{k,n} / v$ . Denote a profile of the series as  $X_{k,n} = \sum_{i=1}^k (r_{i,n} - \bar{r}_n)$ . Let's further put a range of the series as  $R_n = \max_{1 \leq k \leq v} X_{k,n} - \min_{1 \leq k \leq v} X_{k,n}$  and a standard deviation as  $S_n = \left( \sum_{k=1}^v (X_{k,n} - \bar{X}_n)^2 / (v-1) \right)^{0.5}$ . Moreover, put

$$\left( \frac{R}{S} \right)_{I_n} = \frac{R_n}{S_n} \quad \text{and} \quad \left( \frac{R}{S} \right)_v = \frac{1}{N} \sum_{n=1}^N \frac{R_n}{S_n}.$$

Finally, for  $c \in \mathbb{R}$ , rescaled range scales by the means of Hurst exponent  $H$  as

$$\left( \frac{R}{S} \right)_v \approx c * v^H$$

where " $\approx$ " means asymptotic equality.

The linear relationship in double-logarithmic scale indicates a power scaling (Weron, 2002). To uncover the scaling law, we use a simple ordinary least squares regression on logarithms of each side of the previous equation. Thus, we get

$$\log(R/S)_v \approx \log c + H \log v,$$

where  $H$  is Hurst exponent.

R/S was shown to be sensitive to a heteroskedasticity and the short-range dependence in the underlying process (DiMatteo, 2007; Lo & MacKinlay, 1999; and Alfi *et al.*, 2008). To deal with the problem, Lo (1991) proposed a modified rescaled range analysis (M-R/S). The procedure differs in the definition of the standard deviation which deals with both the heteroskedasticity and the short-range dependence. The modified standard deviation is

defined with a use of auto-covariance  $\gamma$  of the selected sub-interval  $I_n$  up to the integer lag  $\xi$  as

$$S_{I_n}^M = \sqrt{S_{I_n}^2 + 2 \sum_{j=1}^{\xi} \gamma_j \left(1 - \frac{j}{\xi+1}\right)}.$$

Thus, R/S turns into a special case of M-R/S with  $\xi = 0$  (Dölger & Ozdemir, 2005). The most problematic and also the crucial issue of the new standard deviation measure is the number of lags which are used for its estimation (Wang *et al.*, 2006). On one hand, if the chosen lag is too low, it omits lags which may be significant and therefore the estimates of rescaled ranges and Hurst exponent are still biased. On the other hand, if the used lag is too high, the estimates of rescaled range differ significantly from the true values (Teverovsky, Taqqu & Willinger, 1999).

One can either estimate rescaled ranges and Hurst exponents for arbitrary values of  $\xi$  (Zhuang, Gree & Maggioni, 2000; and Alptekin, 2006) or use an automatic estimator of  $\xi$  for each sub-interval of the length  $\nu$ . There are two estimators of the optimal lag suggested in the literature. Lo (1991) sets the optimal lag  $\xi^*$  based on the first-order autocorrelation coefficient  $\hat{\rho}(1)$  of the sub-interval  $I_n$  as (where  $[]$  is the nearest lower integer operator)

$$\xi^* = \left\lfloor \left( \frac{3\nu}{2} \right)^{\frac{1}{3}} \left( \frac{2\hat{\rho}(1)}{1 - (\hat{\rho}(1))^2} \right)^{\frac{2}{3}} \right\rfloor.$$

Chin (2008) bases the optimal lag choice on the length of the sub-interval solely as

$$\xi^* = \left\lfloor 4 \left( \frac{\nu}{100} \right)^{\frac{2}{9}} \right\rfloor.$$

Teverovsky, Taqqu & Willinger (1999) showed that M-R/S is biased towards rejecting long-range dependence in the process when high number of lags is used. It implies that in the case of the significant short-range dependence in the process, the method of Lo would lead to biased estimates of  $H$ . Moreover, if the short-term memory is not significant or low, the method of Lo does not significantly differ from the method of Chin. For values of the autocorrelation  $\hat{\rho}(1)$  lower than 0.5, both methods do not differ significantly.

## 2.2. Brief literature review

Finite sample properties of the rescaled range analysis and the modified rescaled range analysis were discussed in several papers. Most importantly, the condition for the process to be characterized as independent, which is  $H = 0.5$ , was shown to hold only asymptotically. For finite samples, the estimated Hurst exponent can differ significantly. To deal with the problem, Anis & Lloyd (1976) showed that for the finite samples, expected value of the rescaled range behaves as

$$E(R/S)_v = \frac{\Gamma\left(\frac{v-1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{v}{2}\right)} \sum_{i=1}^{v-1} \sqrt{\frac{v-i}{i}}.$$

Further, Peters (1994) proposed to adjust the expected value of the rescaled range, which is better for low scales  $v < 50$ , to

$$E(R/S)_v = \frac{v - \frac{1}{2}}{v} \sqrt{\frac{2}{v\pi}} \sum_{i=1}^{v-1} \sqrt{\frac{v-i}{i}}.$$

However, Couillard & Davison (2005) and Grech & Mazur (2005) showed that the expected values based on the Monte Carlo simulations are closer to the method of Anis & Lloyd (1974). Importantly, they also showed that the standard deviation of the estimates does not converge according to the central limit theorem, which is as a square root of the time series length. Such finding is essential for the confidence interval construction, since as the standard deviation converges slower, the confidence intervals are much wider.

### 3 Monte Carlo simulations methodology

The choice of parameters is crucial for final results of the simulations and estimates. The thorough description is needed for possible comparison with other results. In the following section, we base the estimates so that we use the sub-length  $v$  equal to the power of a set integer value (Weron, 2002). Thus, we set a basis  $b$ , a minimum power  $pmin$  and a maximum power  $pmax$  so that we get  $v = b^{pmin}, \dots, b^{pmax}$  where  $b^{pmin} \geq b$  is a minimum scale and  $b^{pmax} \leq T$  is a maximum scale. The minimum scale is set to 32 observations and the maximum as the whole time series length. Such choice avoids biases of the low scales where the standard deviations can be estimated inefficiently (Peters, 1994; Grech & Mazur, 2004; and Matos *et al.*, 2008). Moreover, the used intervals are contingent and non-overlapping (see Ellis, 2007 for discussion).

The majority of research papers dealing with the finite sample properties of the rescaled range base their estimates on simulations of the standardized normal distribution  $N(0,1)$ . However, the returns on financial markets are known to be non-normal and fat-tailed (Cont, 2001). To show whether R/S and M-R/S are robust to fat tails in the underlying process, we show the estimates for both the standardized normal distribution and the Cauchy distribution, which is a stable distribution with the fattest tails.

In the simulations, we simulate 1 000 time series with lengths from  $2^9$  to  $2^{14}$  for the standard normal distribution and estimate Hurst exponent based on R/S and M-R/S for each series. We apply the same procedure for the Cauchy distribution. Both processes are independent as each element of the time series is drawn randomly so that the estimates are expected to be 0.5. Note that we use the optimal lag choice based on Chin (2008) as first-order autocorrelation of independent process is equal to zero so that the method of Lo (1991) would turn into classical rescaled range.



## 4 Results

Table 1 shows the descriptive statistics of all Monte Carlo simulations as well as the Jarque-Bera test for normality (Jarque & Bera, 1981) and 2.5% and 97.5% percentiles. Chart 1 and Chart 2 present histograms of the estimates of Hurst exponents for R/S and M-R/S, respectively. Chart 3 shows confidence intervals based on different percentiles for both methods.

We can see that the expected value for R/S for standard normal process is the highest for the shortest time series length of 512 observations and is equal to 0.5316 and quite steadily decreases to 0.5211 for 16 384 observations. Interestingly, the expected values of R/S estimates for the Cauchy process are lower than the ones for the normal distribution up to time series length 4 096. For the two highest lengths, the estimates are similar. The expected values of Hurst exponent for Cauchy-based process range from 0.5281 to 0.5248 and are thus quite stable. Nevertheless, R/S is robust against presence of heavy tails and its estimates are not affected when compared to the estimates for independent process based on normal distribution. Moreover, the standard deviations are lower for the Cauchy-based estimates (Chart 4).

The results are similar for M-R/S. The expected values range from 0.5192 for 512 observations to 0.5124 for the length 16 384 for standard normal process. Again, expected values for the Cauchy-based estimates of Hurst exponent are lower than the ones of normal-based ones up till length 4 096. However, these expected values are increasing with time series length. Still, the expected values remain very close for both types of underlying processes so that M-R/S is robust against the presence of heavy tails in the underlying process as well. The standard deviations are lower for process based on the Cauchy distribution and almost equal to the ones of R/S for the same type of the process. The results for standard deviations for the simulated Cauchy-based series show that the maximum scale equal to the time series length can be used as even very heavy tails of the Cauchy distribution do not bias the estimates of Hurst exponent for both R/S and M-R/S.

To clearly distinguish between the short and the long-range dependence in the process, we need to estimate Hurst exponent by both methods and compare them with the confidence intervals (Chart 4). This way, we can distinguish between three possibilities. If the estimates of Hurst exponent based on both R/S and M-R/S are inside the corresponding confidence intervals, the process can be either independent or slightly short-range dependent. If the estimates of  $H$  are outside the confidence intervals for R/S but inside the ones for M-R/S, the process is short-range dependent. Finally, if the Hurst exponent estimates are outside of confidence intervals for both methods, the process is long-range dependent.

## 5 Conclusion

In the paper, we researched on behavior of the rescaled range analysis and the modified rescaled analysis under both the normal distribution assumption as well as under the heavy tails represented by the Cauchy distribution. As most of the studies focus on the normal distribution only, we tried to fill this gap as returns on financial markets are known to be non-normal and with heavy tails. Moreover, the short and the long-range dependence have different implications for portfolio selection and options pricing so that clear distinguishing between both processes can be crucial.

Monte Carlo simulations showed that both methods are robust to the presence of the heavy tails in the underlying process. Interestingly, the standard deviations of the estimates for the process with the Cauchy distribution are lower than for the normal case for both methods. We also presented the confidence intervals which enable us to clearly distinguish between the short and the long-range dependence in the process.

## References


- Alfi, V., Coccetti, F., Marotta, M., Petri, A. & Pietronero, L.:** *Exact Results for the Roughness of a Finite Size Random Walk*, Physica A, Vol. 370, No. 1, 2008
- Alptekin, N.:** *Long Memory Analysis of USD/TRL Exchange Rate*, International Journal of Social Sciences, Vol. 1, No. 2, 2006
- Alvarez-Ramirez, J., Alvarez, J., Rodriguez, E. & Fernandez-Anaya, G.:** *Time-varying Hurst exponent for US stock markets*, Physica A, Vol. 387, 2008
- Anis, A. & Lloyd, E.:** *The Expected Value of the Adjusted Rescaled Hurst Range of Independent Normal Summands*, Biometrika, Vol. 63, 1976
- Barkoulas, J., Baum, Ch. & Travlos, N.:** *Long memory in the Greek stock market*, Applied Financial Economics, Vol. 10, 2000
- Beran, J.:** *Statistics for Long-Memory Processes*, Monographs on Statistics and Applied Probability 61, Chapman & Hall, New York, ISBN 978-0412049019, 1994
- Box, G. & Jenkins, G.:** *Time Series Analysis: Forecasting and Control*, Holden Day, San Francisco, 1970
- Carbone, A., Castelli, G. & Stanley, H.:** *Time-dependent Hurst exponent in financial time series*, Physica A, Vol 344, 2004
- Chin, W.:** *Spurious Long-range Dependence: Evidence from Malaysian Equity Markets*, MPRA Paper No. 7914, 2008
- Cont, R.:** *Empirical properties of asset returns: stylized facts and statistical issues*, Quantitative Finance, Vol. 1, No. 2, 2001
- Couillard, M. & Davison, M.:** *A comment on measuring the Hurst exponent of financial time series*, Physica A, Vol. 348, pp. 404-418, 2005
- Czarnecki, L., Grech, D. & Pamula, G.:** *Comparison study of global and local approaches describing critical phenomena on the Polish stock exchange market*, Acta Physica Polonica A, 2008
- Di Matteo, T., Aste, T. & Dacorogna, M.:** *Long-term memories of developed and emerging markets: Using the scaling analysis to characterize their stage of development*, Journal of Banking & Finance, No. 29, pp. 827-851, 2005
- Di Matteo, T.:** *Multi-scaling in Finance*, Quantitative Finance, Vol. 7, No. 1, pp. 21-36, 2007
- Dülger, F. & Ozdemir, Z.:** *Current Account Sustainability in Seven Developed Countries*, Journal of Economic and Social Research, Vol. 7, No. 2, 2005
- Ellis, C.:** *The sampling properties of Hurst exponent estimates*, Physica A, Vol. 375, 2007
- Embrechts, P. & Maejima, M.:** *Selfsimilar Processes*, Princeton University Press, ISBN 978-0691096278, 2002
- Grech, D. & Mazur, Z.:** *Can one make any crash prediction in finance using the local Hurst exponent idea?*, Physica A, Vol. 336, 2004
- Grech, D. & Mazur, Z.:** *Statistical Properties of Old and New Techniques in Detrended Analysis of Time Series*, Acta Physica Polonica B, Vol. 36, No. 8, 2005
- Hurst, H. E.:** *Long term storage capacity of reservoirs*, Transactions of the American Society of Engineers, Vol. 116, 1951
- Jarque, C. & Bera, A.:** *Efficient tests for normality, homoscedasticity and serial independence of regression residuals: Monte Carlo evidence*, Economics Letters, Vol. 7, No. 4, 1981
- Lillo, F. & Farmer, J.:** *The Long Memory of the Efficient Market*, Studies in Nonlinear Dynamics & Econometrics, Vol. 8, No. 3, 2004
- Lo, A. & MacKinlay, C.:** *A Non-Random Walk Down Wall Street*, Princeton University Press, New Jersey, ISBN 0-691-09256-7, 1999

- Lo, A.:** *Long-term memory in stock market prices*, *Econometrica*, Vol. 59, No. 5, 1991
- Mandelbrot, B. & van Ness, J.:** *Fractional Brownian motions, fractional noises and applications*, *SIAM Review*, Vol. 10, No. 422, 1968
- Mandelbrot, B.:** *Analysis of long-run dependence in economics: the R/S technique*, *Econometrica*, Vol. 39, 1970
- Matos, J., Gama, S., Ruskin, H., Sharkasi, A. & Crane, M.:** *Time and scale Hurst exponent analysis for financial markets*, *Physica A*, Vol. 387, No. 15, 2008
- Morariu, V., Buimaga-Iarinca, L., Vamoş, C. & Şoltuz, Ş.:** *Detrended Fluctuation Analysis of Autoregressive Processes*, arXiv:0707.1437, 2007
- Peters, E.:** *Fractal Market Analysis – Applying Chaos Theory to Investment and Analysis*, John Wiley & Sons, Inc., New York, ISBN 978-0471585244, 1994
- Taqqu, M., Teverovsky, W. & Willinger, W.:** *Estimators for long-range dependence: an empirical study*, *Fractals*, Vol. 3, No. 4, 1995
- Teverovsky, V., Taqqu, M. & Willinger, W.:** *A critical look at Lo's modified R/S statistic*, *Journal of Statistical Planning and Inference*, Vol. 80, No. 1-2, 1999
- Vandewalle, N., Ausloos, M. & Boveroux, Ph.:** *Detrended Fluctuation Analysis of the Foreign Exchange Market*, *Econophysics Workshop*, Budapest, Hungary, 1997
- Wang, W., Van Gelder, P., Vrijling, J. & Chen, X.:** *Detecting long-memory: Monte Carlo simulations and application to daily streamflow processes*, *Hydrology and Earth System Sciences Discussions*, Vol. 3, 2006
- Weron, R.:** *Estimating long-range dependence: finite sample properties and confidence intervals*, *Physica A*, Vol. 312, 2002
- Zhuang, Y., Green, Ch. & Maggioni, P.:** *The Great Rebound, The Great Crash, and Persistence in British Stock Prices*, *Economic Research Paper No. 00/11*, Loughborough University, 2000

**Table 1 Results of Monte Carlo simulations**


		<b>512</b>	<b>1024</b>	<b>2048</b>	<b>4096</b>	<b>8192</b>	<b>16384</b>
<b>mean</b>	<i>R/S (normal)</i>	0,5316	0,5345	0,5305	0,5292	0,5222	0,5211
	<i>R/S (Cauchy)</i>	0,5281	0,5250	0,5258	0,5236	0,5244	0,5248
	<i>M-R/S (normal)</i>	0,5192	0,5186	0,5157	0,5155	0,5151	0,5124
	<i>M-R/S (Cauchy)</i>	0,5012	0,5091	0,5122	0,5133	0,5151	0,5167
<b>SD</b>	<i>R/S (normal)</i>	0,0853	0,0608	0,0473	0,0379	0,0328	0,0268
	<i>R/S (Cauchy)</i>	0,0688	0,0528	0,0407	0,0344	0,0292	0,0254
	<i>M-R/S (normal)</i>	0,0743	0,0568	0,0496	0,0392	0,0315	0,0279
	<i>M-R/S (Cauchy)</i>	0,0678	0,0551	0,0421	0,0351	0,0285	0,0251
<b>skewness</b>	<i>R/S (normal)</i>	-2,2315	-0,0812	-0,0881	0,0350	0,0661	-0,0444
	<i>R/S (Cauchy)</i>	0,0928	0,1192	0,0792	0,0254	0,0458	0,0814
	<i>M-R/S (normal)</i>	-0,1319	-0,0525	-0,0010	-0,0669	-0,1155	-0,1057
	<i>M-R/S (Cauchy)</i>	-0,0448	-0,2013	0,0445	0,0000	-0,1094	-0,1583
<b>kurtosis</b>	<i>R/S (normal)</i>	27,9929	-0,1830	-0,0685	-0,0372	-0,0968	-0,2790
	<i>R/S (Cauchy)</i>	0,0540	0,1476	-0,0207	-0,1679	0,1114	0,1629
	<i>M-R/S (normal)</i>	-0,0602	-0,3193	-0,2894	0,0055	-0,0266	0,0083
	<i>M-R/S (Cauchy)</i>	0,3288	1,3961	0,2073	-0,2088	0,1467	0,1341
<b>Jarque-Bera</b>	<i>R/S (normal)</i>	33480,1323	2,4950	1,4902	0,2613	1,1189	3,5705
	<i>R/S (Cauchy)</i>	1,5553	3,2763	1,0630	1,2824	0,8667	2,2100
	<i>M-R/S (normal)</i>	3,0493	4,7071	3,4907	0,7474	2,2539	1,8662
	<i>M-R/S (Cauchy)</i>	4,8401	87,9678	2,1199	1,8164	2,8930	4,9264
<b>p-value</b>	<i>R/S (normal)</i>	0,0000	0,2872	0,4747	0,8775	0,5715	0,1678
	<i>R/S (Cauchy)</i>	0,4595	0,1943	0,5877	0,5267	0,6483	0,3312
	<i>M-R/S (normal)</i>	0,2177	0,0950	0,1746	0,6882	0,3240	0,3933
	<i>M-R/S (Cauchy)</i>	0,0889	0,0000	0,3465	0,4032	0,2354	0,0852
<b>P<sub>97.5</sub></b>	<i>R/S (normal)</i>	0,6779	0,6505	0,6237	0,6037	0,5882	0,5733
	<i>R/S (Cauchy)</i>	0,6598	0,6324	0,6093	0,5902	0,5825	0,5772
	<i>M-R/S (normal)</i>	0,6602	0,6275	0,6144	0,5898	0,5726	0,5636
	<i>M-R/S (Cauchy)</i>	0,6361	0,6162	0,5993	0,5819	0,5677	0,5623
<b>P<sub>2.5</sub></b>	<i>R/S (normal)</i>	0,3740	0,4119	0,4353	0,4581	0,4627	0,4669
	<i>R/S (Cauchy)</i>	0,3982	0,4244	0,4455	0,4549	0,4677	0,4773
	<i>M-R/S (normal)</i>	0,3685	0,4077	0,4217	0,4376	0,4522	0,4554
	<i>M-R/S (Cauchy)</i>	0,3695	0,3998	0,4305	0,4466	0,4582	0,4678

**Chart 1 Distribution of estimated Hurst exponents for R/S**




**Chart 1:** The first two rows show the distributions for Monte Carlo simulations based on standard normal underlying process. The second two rows show the distributions based on Cauchy distribution of the underlying process. Length of the time series starts at 512 observations (a and g) and ends at 16 384 observations (f and l).

**Chart 2 Distribution of estimated Hurst exponents for M-R/S**




**Chart 2:** The first two rows show the distributions for Monte Carlo simulations based on standard normal underlying process. The second two rows show the distributions based on Cauchy distribution of the underlying process. Length of the time series starts at 512 observations (a and g) and ends at 16 384 observations (f and l).

**Chart 3 Confidence intervals for R/S and M-R/S**



**Chart 3:** Confidence intervals for **(a)** R/S with standard normal underlying process, **(b)** R/S with Cauchy distributed underlying process, **(c)** M-R/S with standard normal underlying process and **(d)** M-R/S with Cauchy distributed underlying process. Bold black line represents the 5% and 95% confidence intervals. Dashed black line represents 2.5% and 97.5% confidence intervals. Finally, dashed gray line represents 0.5% and 99.5% confidence intervals.

**Chart 4 Confidence intervals for R/S and M-R/S**



**Chart 4:** **(a)** Standard deviations of estimated Hurst exponents for R/S (black line) and M-R/S (grey line) with standard normal distribution (full line) and Cauchy distribution (dashed line) of the process. **(b)** Comparison of 2.5% and 97.5% confidence intervals for R/S (black) and M-R/S (grey). For each method, the higher value of upper confidence interval from standard normal and Cauchy distribution of the underlying processes is shown as is the lower value for the lower confidence interval.



# IES Working Paper Series

2009

1. František Turnovec : *Fairness and Squareness: Fair Decision Making Rules in the EU Council?*
2. Radovan Chalupka : *Improving Risk Adjustment in the Czech Republic*
3. Jan Průša : *The Most Efficient Czech SME Sectors: An Application of Robust Data Envelopment Analysis*
4. Kamila Fialová, Martina Mysíková : *Labor Market Participation: The Impact of Social Benefits in the Czech Republic*
5. Kateřina Pavloková : *Time to death and health expenditure of the Czech health care system*
6. Kamila Fialová, Martina Mysíková : *Minimum Wage: Labour Market Consequences in the Czech Republic*
7. Tomáš Havránek : *Subsidy Competition for FDI: Fierce or Weak?*
8. Ondřej Schneider : *Reforming Pensions in Europe: Economic Fundamentals and Political Factors*
9. Jiří Witzany : *Loss, Default, and Loss Given Default Modeling*
10. Michal Bauer, Julie Chytilová : *Do children make women more patient? Experimental evidence from Indian villages*
11. Roman Horváth : *Interest Margins Determinants of Czech Banks*
12. Lenka Šťastná : *Spatial Interdependence of Local Public Expenditures: Selected Evidence from the Czech Republic*
13. František Turnovec : *Efficiency of Fairness in Voting Systems*
14. Martin Gregor, Dalibor Roháč : *The Optimal State Aid Control: No Control*
15. Ondřej Glazar, Wadim Strielkowski : *Turkey and the European Union: possible incidence of the EU accession on migration flows*
16. Michaela Vlasáková Baruníková : *Option Pricing: The empirical tests of the Black-Scholes pricing formula and the feed-forward networks*
17. Eva Ryšavá, Elisa Galeotti : *Determinants of FDI in Czech Manufacturing Industries between 2000-2006*
18. Martin Gregor, Lenka Šťastná : *Mobile criminals, immobile crime: the efficiency of decentralized crime deterrence*

19. František Turnovec : *How much of Federalism in the European Union*
20. Tomáš Havránek : *Rose Effect and the Euro: The Magic is Gone*
21. Jiří Witzany : *Estimating LGD Correlation*
22. Linnéa Lundberg, Jiri Novak, Maria Vikman : *Ethical vs. Non-Ethical – Is There a Difference? Analyzing Performance of Ethical and Non-Ethical Investment Funds*
23. Jozef Barunik, Lukas Vacha : *Wavelet Analysis of Central European Stock Market Behaviour During the Crisis*
24. Michaela Krčílková, Jan Zápal : *OCA cubed: Mundell in 3D*
25. Jan Průša : *A General Framework to Evaluate Economic Efficiency with an Application to British SME*
26. Ladislav Kristoufek : *Classical and modified rescaled range analysis: Sampling properties under heavy tails*

All papers can be downloaded at: <http://ies.fsv.cuni.cz>

