Problems # 8

Due Thursday, November 30, 2017

1. **Aliasing.** The Surface Water and Ocean Topography (SWOT) satellite is due to launch in 2021. It has two planned orbits. The initial calibration and validation will be carried out with a fast-sampling orbit, with an exact repeat of about a day, and the subsequent science mission will have about a 21-day repeat. Specific time periods are listed in the table below:

Orbit period (days)

Fast-sampling 0.99349 Science 20.86455

What is the alias period for each of these orbits for the lunar semi-diurnal (M2) and solar diurnal (S1) tidal cycles? How long should the satellite operate in each orbit to provide multiple realizations of the tidal amplitude?

Symbol Name period (hours)

S1 Solar diurnal 24.00 M2 Principal lunar 12.42

- 2. Frequency-wavenumber spectra. Download the monthly NOAA Optimum Interpolation monthly mean sea surface temperature data (available from TritonED). Compute frequency/wavenumber spectra for two slabs of data in the Pacific.
 - a Along the equator: $-0.5^{\circ}\mathrm{S},$ from 219.5 to 269.5°E.
 - b In the subtropics: -29.5°S, from 189.5 to 249.5°E.

For both slabs, plot a time/longitude plot (known as a Hovmoller diagram) and compute frequency/wavenumber spectra using at least some amount of averaging/segmenting. Is there a difference in energy propagation between the tropics and subtropics? What evidence do your figures provide?