

Adopted Levels, Gammas

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Balraj Singh and Jun Chen		NDS 178, 41 (2021).	12-Nov-2021

$Q(\beta^-) = -7171.2$ 15; $S(n) = 11861.9$ 15; $S(p) = 7713.1$ 6; $Q(\alpha) = -3955.7$ 7 2021Wa16

$Q(2\epsilon) = 1095.0$ 7, $S(2n) = 20978.6$ 8, $S(2p) = 13835.5$ 6 (2021Wa16).

Mass excess measurements: 2018Ki21, 2008Go23, 2007Ke09, 2005Ch60, 1977De20, 1976De21.

Other reactions:

$^{59}\text{Co}(^6\text{Li}, n)$ $E = 39.7, 90$ MeV: 1984Bo45: Measured $\sigma(\theta)$, reaction mechanism.

$^{60}\text{Ni}(\alpha, p)$: IAR $E = 7.260$ - 7.450 MeV: 1976Fo06: Measured resonance strengths. Other: 1961No04.

$^{60}\text{Ni}(\alpha, \alpha')$: IAR $E = 14.6$ - 20.9 MeV: 1975Lu06: Si telescope, $\sigma(\theta)$, deduced isospin mixing for ^{64}Zn at excitations of 17.6-23.5 MeV.

$^{61}\text{Ni}(\alpha, n)$ $E = 53$ MeV: 1984Bo45, 1979Bo45: $\sigma(\theta)$, reaction mechanism.

Additional information 1.

$^{64}\text{Zn}(t, t)$ $E = 12$ MeV: 1972Hu06: measured $\sigma(\theta)$. Deduced optical model-parameters.

$^{64}\text{Zn}(^6\text{Li}, ^6\text{Li})$, $(^7\text{Li}, ^7\text{Li})$: 1991Bo48.

$^{27}\text{Al}(^{37}\text{Cl}, X)$ and $^{48}\text{Ti}(^{16}\text{O}, X)$: 1984Mi09 (scission mechanism for excited ^{64}Zn nucleus).

$^{63}\text{Cu}(^{16}\text{O}, ^{15}\text{N})$: 1975We20 $E = 38$ - 51 MeV. Reaction mechanism.

$^{60}\text{Ni}(^{18}\text{O}, ^{14}\text{C})$: 1973RoYT, 1972HeYV.

$^{64}\text{Zn}(\pi^+, \pi^-)$: 1993Be02: $E = 293.4$ MeV, measured $\sigma(\theta)$.

$^{64}\text{Zn}(\pi^+, \pi^-)$: 1997Fo03: $E = 140$ - 230 MeV, measured $\sigma(\theta)$.

$^{64}\text{Zn}(K^-, X)$: 1980De11, calculated atomic level shifts.

$^{64}\text{Zn}(^7\text{Li}, t\alpha)$ $E = 42$ MeV: 2001To07 (also 1999Ut03): measured triton and α spectra, $\sigma(\theta)$, deduced astrophysical S factors.

$^{64}\text{Zn}(^{10}\text{Be}, ^{10}\text{Be})$, $((^{11}\text{Be}, ^{11}\text{Be}), E(c.m.) = 24.5$ MeV: 2014DiZV: measured $\sigma(\theta)$ for elastic scattering at REX-ISOLDE facility of CERN, and analyzed using CDCC with optical model calculations.

Giant-dipole resonances: 1981Do12 ($^{64}\text{Zn}(e, p)$); 1977TaYW ($^{64}\text{Zn}(\gamma, \alpha)$); 1973Ya04 ($^{64}\text{Zn}(\gamma, n)$, (γ, np)); 1972CIZK ($^{64}\text{Zn}(\gamma, p)$); 1970Co25 ($^{64}\text{Zn}(\gamma, n)$, $(\gamma, 2n)$, (γ, np)).

Isotope shifts: 1970Le23. Theory: 1982Fo09.

There are several high-spin studies: $^{12}\text{C}(^{54}\text{Fe}, 2p\gamma)$ from 1994Cr05; $^{40}\text{Ca}(^{28}\text{Si}, 4p\gamma)$ $E = 115$ MeV from 1998Ga11; $^{40}\text{Ca}(^{28}\text{Si}, 4p\gamma)$ $E = 120$ MeV from 1997Fu08; $^{40}\text{Ca}(^{28}\text{Si}, 4p\gamma)$ $E = 122$ MeV from 2004Ka18; $^{51}\text{V}(^{16}\text{O}, p2n\gamma)$, $^{59}\text{Co}(^7\text{Li}, 2n\gamma)$ from 1977We10, 1978We15, and 1977Al14; $^{61}\text{Ni}(\alpha, n\gamma)$, $^{56}\text{Fe}(^{11}\text{B}, 2np\gamma)$ from 1980Si02 and 1978Si02; $^{61}\text{Ni}(\alpha, n\gamma)$, $(\text{HI}, xn\gamma)$ from 1978Ne02 and 1976Ch11. While there is general agreement between all these studies below about 5 MeV excitation, above this energy, there are major differences even when the same reaction is used as in 2004Ka18, 1998Ga11 and 1997Fu08. Since the statistics in $\gamma\gamma$ coincidences is the highest in 2004Ka18, where Gammasphere array has been used, evaluators have adopted the high-spin level scheme from 2004Ka18 with the exception of few cases where results of 2004Ka18 are in clear disagreement with other experiments, the results of which are considered by evaluators as more definitive. Such cases are noted in comments. It should also be mentioned that complete details of data are not available from 2004Ka18. Requests by evaluators for obtaining such details from the authors of 2004Ka18 were unsuccessful. Full details of data are also missing in 1998Ga11 and 1997Fu08, although, some were obtained as priv. comm. (1996GaZZ) from authors of 1998Ga11. Several levels proposed by 1998Ga11, 1997Fu08 and 1994Cr05, but not reported by 2004Ka18 have been omitted here. See individual datasets for details.

In the opinion of the evaluators, there are several incomplete or discrepant aspects of the high-spin portion of the level scheme which need to be resolved in further experiments.

 ^{64}Zn LevelsCross Reference (XREF) Flags

A	^{64}Cu β^- decay (12.7006 h)	N	$^{62}\text{Ni}(^3\text{He}, n)$	Others:
B	^{64}Ga ϵ decay (2.627 min)	O	$^{63}\text{Cu}(p, \gamma)$ $E = 1.3$ - 3.2 MeV	AA $^{64}\text{Zn}(\mu^-, X)$
C	$^{12}\text{C}(^{54}\text{Fe}, 2p\gamma)$	P	$^{63}\text{Cu}(p, \gamma)$ $E = 2050$ keV	AB $^{64}\text{Zn}(n, n')$
D	$^{40}\text{Ca}(^{28}\text{Si}, 4p\gamma)$ $E = 115$ MeV	Q	$^{63}\text{Cu}(p, \gamma)$ $E = 2.1$ - 3.1 MeV	AC $^{64}\text{Zn}(n, n'\gamma)$
E	$^{40}\text{Ca}(^{28}\text{Si}, 4p\gamma)$ $E = 120$ MeV	R	$^{63}\text{Cu}(p, \gamma)$ $E = 2098$ keV	AD $^{64}\text{Zn}(p, p')$, (pol p, p')
F	$^{40}\text{Ca}(^{28}\text{Si}, 4p\gamma)$ $E = 122$ MeV	S	$^{63}\text{Cu}(p, \gamma)$ $E = 3217, 3251$ keV	AE $^{64}\text{Zn}(p, p'\gamma)$
G	$^{51}\text{V}(^{16}\text{O}, p2n\gamma)$, $^{59}\text{Co}(^7\text{Li}, 2n\gamma)$	T	$^{63}\text{Cu}(p, \gamma)$ $E = 3.46$ MeV	AF $^{64}\text{Zn}(d, d')$, (pol d, d)
H	$^{60}\text{Ni}(^6\text{Li}, d)$	U	$^{63}\text{Cu}(p, n)$: resonances	AG $^{64}\text{Zn}(^3\text{He}, ^3\text{He}')$
I	$^{60}\text{Ni}(^7\text{Li}, t)$	V	$^{63}\text{Cu}(d, n)$, (pol d, n)	AH $^{64}\text{Zn}(\alpha, \alpha')$

	J	$^{60}\text{Ni}(^{16}\text{O}, ^{12}\text{C})$	W	$^{63}\text{Cu}(^3\text{He}, \text{d})$	AI	$^{64}\text{Zn}(^{16}\text{O}, ^{16}\text{O}'), (^{12}\text{C}, ^{12}\text{C}')$
	K	$^{61}\text{Ni}(\alpha, \text{n}\gamma), ^{56}\text{Fe}(^{11}\text{B}, 2\text{n}\text{p}\gamma)$	X	$^{63}\text{Cu}(\alpha, \text{t})$	AJ	Coulomb excitation
	L	$^{61}\text{Ni}(\alpha, \text{n}\gamma), (\text{HI}, \text{x}\text{n}\gamma)$	Y	$^{64}\text{Zn}(\gamma, \gamma')$	AK	$^{66}\text{Zn}(\text{p}, \text{t})$
	M	$^{62}\text{Ni}(^{12}\text{C}, ^{10}\text{Be}), (^{16}\text{O}, ^{14}\text{C})$	Z	$^{64}\text{Zn}(\text{e}, \text{e}')$		
E(level) [†]	J^π	$T_{1/2}$ [#]	XREF		Comments	
0.0&	0 ⁺	stable@	ABCDEFGH IJKLMN OQRST VWXYZ		<p>XREF: Others: AB, AC, AD, AE, AF, AG, AH, AI, AJ, AK</p> <p>Evaluated rms charge radius $\langle r^2 \rangle^{1/2} = 3.9283 \text{ fm}$ <i>I5</i> (2013An02).</p> <p>Evaluated $\delta \langle r^2 \rangle (^{66}\text{Zn}, ^{64}\text{Zn}) = -0.162 \text{ fm}^2$ <i>2</i> (2013An02).</p> <p>Measured change in isotope shift $\delta \nu (^{68}\text{Zn}, ^{64}\text{Zn}) = -141.2 \text{ MHz}$ <i>I2(stat)</i> <i>66(syst)</i> (2019Xi07, collinear laser spectroscopy at ISOLDE, CERN), with laser wavelength of 480.7254 nm to match the Doppler shifted transition.</p> <p>Measured change in charge radius $\delta \langle r^2 \rangle (^{68}\text{Zn}, ^{64}\text{Zn}) = -0.279 \text{ fm}^2$ <i>4(stat)</i> <i>34(syst)</i> (2019Xi07, collinear laser spectroscopy at ISOLDE, CERN), with laser wavelength of 480.7254 nm to match the Doppler shifted transition.</p> <p>J^π: hyperfine structure measurements: 1929Sc01, 1931Mu02.</p> <p>$T_{1/2}$: see footnote for lower limits for double β decay.</p> <p>Additional information 2.</p>	
991.54& 5	2 ⁺	1.94 ps 5	BCDEFGH IJKLMN OQRST VWXYZ		<p>XREF: Others: AB, AC, AD, AE, AF, AG, AH, AI, AJ, AK</p> <p>$\beta_2 = 0.260$ <i>I8</i> (1987Ja04)</p> <p>$\mu = +0.89$ <i>I4</i> (2005Le12, 2010Mo14, 2020StZV)</p> <p>$Q = -0.143$ <i>2I</i> (1976Ne06, 1977Ne05, 1981Ko06, 2016St14, 2021StZZ)</p> <p>XREF: J(960)N(1024)AB(920).</p> <p>Additional information 3.</p> <p>E(level): level energy held fixed in least-squares adjustment.</p> <p>J^π: E2 γ to 0⁺.</p> <p>β_2: from (p, p'). Others: see (e, e'); (n, n'); (p, p'); (d, d'); ($^3\text{He}, ^3\text{He}'$) and ($\alpha, \alpha'$). $\beta_2(\text{pol p}, \text{p}') = 0.26, 0.25$ (1993Mo15). Negative sign is indicated by 1991Ku30 from an analysis of $\sigma(\theta)(\alpha, \alpha')$ data.</p> <p>μ: from transient fields in Coul. ex. (2005Le12, 2010Mo14). Others: +0.89 <i>9</i> (2002Ke02), +0.92 <i>20</i> (1979Fa06), +0.84 <i>I8</i>, +1.04 <i>24</i> (1978BeZJ, 1979BrZP). 2010Mo14 reanalyzed their previously measured g factor of +0.45 <i>3</i> in 2005Le12 using a different procedure for precession effect, and obtained the same value. Uncertainty of 0.06 in 2010Mo14 increased to 0.14 in 2020StZV evaluation.</p> <p>Q: from electron scattering (1976Ne06, 1977Ne05, value of -0.124 <i>I1</i> reanalyzed by 1981Ko06 to -0.143 <i>2I</i>). Others: -0.32 <i>6</i> or -0.26 <i>6</i> (1988Sa32, reorientation method in Coulomb excitation; -0.135 <i>I4</i> ((e, e'), 1972Li12); -0.01 +9-5 (Coul. ex., 2003KoZQ).</p> <p>$T_{1/2}$: weighted average of 1.97 ps <i>6</i> (DSA in Coul. Ex., 2005Le12); 1.87 ps <i>6</i> (DSA in Coul. Ex., 2002Ke02); 2.06 ps <i>I7</i> ((γ, γ'), 1981Ca10); 2.8 ps <i>7</i> (RDDS in ($^{16}\text{O}, \text{p}2\text{n}\gamma$), 1977Al14); 1.71 ps <i>2I</i> (line shape in Coul. ex., 1973Fi15) and 2.16 ps <i>I5</i> ((γ, γ'), 1971ImZY). Values deduced from B(E2) values in Coul. Ex. and (e, e') are somewhat lower: 1.76 ps <i>4</i> (from B(E2)=0.168 <i>4</i>, 1988Sa32); 1.82 ps <i>I0</i> (from B(E2)=0.162 <i>9</i> and 1.83 ps <i>I3</i> from B(E2)=0.161 <i>I2</i>, 1975Th01); 1.73 ps <i>I5</i> (from B(E2)=0.170 <i>I5</i>, 1962St02). Weighted average of all the values is 1.84 ps <i>4</i>. 2001Ra27 evaluation quotes 1.86 ps <i>17</i> from weighted average of 15 values (B(E2) in Coul. Ex.: 1988Sa32, 1975Th01, 1962St02, 1960An07, 1956Te26; Doppler-shift method in Coul. Ex.: 1973Fi15; Doppler-shift</p>	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF	Comments
1799.41 ^a 4	2 ⁺	2.0 ps 2	BCDEFGH JKLM OPQRST VW Z	in (^{16}O ,p2n γ): 1977A114; (γ , γ'): 1981Ca10,1977Ca14,1972ArZD,1965Ta13; (e,e'): 1977Ne05,1976Ne06,1970Af04. XREF: Others: AC, AD, AE, AF, AH, AJ, AK XREF: J(1750). J ^π : E2 γ to 0 ⁺ . T _{1/2} : from B(E2)(\uparrow)=0.00170 12 ((e,e'), 1977Ne05). Others (from (α ,n γ), (^{11}B ,2n γ)): 1.4 ps 7 (1977A114), 2.1 ps 14 (1977We10), 1.8 ps +6-3 (1976Ch11), >1.0 ps (1978Si02).
1910.26 4	0 ⁺	0.95 ns 5	B H KL OPQR T W	XREF: Others: AC, AD, AE, AF, AH, AK XREF: AF(1960)AK(1940). J ^π : E0 transition to 0 ⁺ . T _{1/2} : γ ce(t) in (p,p' γ) (1985Pa07). Others (from (α ,n γ),(^{11}B ,2n γ)): 2.4 ps +10-6 (1976Ch11), >1.0 ps (1978Si02).
2306.72 ^{&} 5	4 ⁺	0.776 ps 28	CDEFGH JKLM OPQRST VW Z	XREF: Others: AC, AD, AE, AF, AH, AJ, AK μ =+2.0 6 (2005Le12,2010Mo14,2020StZV) XREF: M(2400)V(2230). μ : from transient fields in Coul. ex. (2005Le12, 2010Mo14 reanalyzed previously measured g factor in 2005Le12 of +0.53 16 using a different procedure for precession effect, and obtained g factor=+0.49 15. J ^π : $\gamma(\theta)$ and $\gamma(\text{lin pol})$ in (α ,n γ),(^{11}B ,2n γ). B(E4)=0.00034 10 from (e,e'). T _{1/2} : from DSA method in Coul. ex. (2005Le12). Others: 0.21 ps +11-8 ((n,n' γ),1985Ko27); 0.29 ps 8 (α ,n γ),(^{11}B ,2n γ),1978Si02; 1.0 ps 6 (1977We10) and 0.8 ps 3 (1976Le31) in (^{16}O ,p2n γ),(^7Li ,2n γ); 0.44 ps 9 (1976Ch11) in (α ,n γ).
2609.52 7	0 ⁺	0.20 ps 8	B H KL OPQR T W	XREF: Others: AC, AD, AE, AK J ^π : E0 transition to 0 ⁺ . T _{1/2} : weighted average of 0.15 ps +6-3 ((n,n' γ), 1985Ko27); 0.36 ps 10 (α ,n γ),(^{11}B ,2n γ), 1978Si02; 1.0 ps +6-4 ((α ,n γ), 1976Ch11).
2736.57 ^a 6	4 ⁺	1.5 ps 3	CDEFGH KL OPQRST w	XREF: Others: AC, AD, AE, AH, AJ, AK XREF: AH(2780). J ^π : 937 $\gamma(\theta,\text{lin pol})$ in (α ,n γ),(^{11}B ,2n γ). T _{1/2} : from (α ,n γ),(^{11}B ,2n γ). Weighted average of 1.2 ps 3 (1980Si02); 3.5 ps 21, 1.7 ps 7 (1977We10); 3.5 ps 14 (1977A114); 2.1 ps 7 (1976Ch11).
2793.5 4	2 ⁺	0.049 ps 14	h KL OPQRST w	XREF: Others: AC, AD, AE, AK J ^π : L(p,t)=2 from 0 ⁺ . T _{1/2} : from (α ,n γ),(^{11}B ,2n γ) (1978Si02). Other: <0.009 ps (1976Ch11).
2979.94 15	3 ⁺	0.30 ps +39-11	jKL nOPQRST	XREF: Others: AC, AD, AE XREF: j(2960)n(2930). J ^π : $\gamma(\theta, \text{lin pol})$ in (α ,n γ),(^{11}B ,2n γ). T _{1/2} : from (n,n' γ) (1985Ko27). Others: >2.6 ps (1976Ch11), >1.0 ps (1978Si02) in (α ,n γ),(^{11}B ,2n γ).
2998.54 ^b 17	3 ⁻	0.152 ps 4	CDEFGH jKL nOP rST wX Z	XREF: Others: AB, AC, AD, AE, AF, AG, AH, AJ, AK μ =+1.5 9 (2005Le12,2020StZV) B(E3) \uparrow =0.040 7 (1976Ne06)

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

<u>E(level)[†]</u>	<u>J^π[‡]</u>	<u>T_{1/2}[#]</u>	<u>XREF</u>	<u>Comments</u>
				$\beta_3=0.235$ 16 (1987Ja04) XREF: H(2980)j(2960)n(2930)X(3040)AB(3300)AK(3020). μ : from transient fields in Coul. ex. (2005Le12). B(E3) \uparrow : from (e,e'). Others: see (e,e') and (α,α'). 2002Ki06 quote 0.034 5 from average of 0.040 7 (1976Ne06) and 0.0307 23 (1970Af04). β_3 : from (p,p'). Others: see (e,e'); (n,n'); (p,p'); (d,d'); ($^3\text{He},^3\text{He}'$); (α,α'). $\beta_3(\text{pol p,p'})=0.22, 0.21$ (1993Mo15). J ^π : L(e,e')=3. Strong population in (p,p') and other inelastic scattering experiments identifies this as an octupole state. T _{1/2} : from DSA method in Coul. ex. (2005Le12). Others: 0.097 ps 21 ((n,n' γ), 1985Ko27); 0.080 ps 21 ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$), 1978Si02; >1.0 ps (1976Ch11). XREF: Others: AC, AD, AE J ^π : $\gamma(\theta, \text{lin pol})$ in ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$). T _{1/2} : weighted average of 0.069 ps +21-14 ((n,n' γ), 1985Ko27); 0.080 ps 21 ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$), 1978Si02; 0.056 ps 15, 0.045 ps 12 (($\alpha,\text{n}\gamma$), 1976Ch11). XREF: Others: AE J ^π : prominent γ to 0 ⁺ ; 1 ⁺ proposed in (p, γ) E=1.3-3.2 MeV. XREF: Others: AC, AD, AH, AJ, AK XREF: AK(3110). J ^π : $\gamma(\theta, \text{lin pol})$ in ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$). Also L(p,p')=L(α,α')=4. T _{1/2} : from DSA in Coul. ex. (2005Le12). Others: 0.42 ps +28-10 ((n,n' γ), 1985Ko27); 0.42 ps 11 ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$), 1978Si02; 1.0 ps 3 ($^7\text{Li},2\text{n}\gamma$), 1977We10; 1.4 ps +10-6 (($\alpha,\text{n}\gamma$), 1976Ch11). XREF: Others: AC, AD, AE XREF: V(3120). J ^π : $\gamma(\theta, \text{lin pol})$ in ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$). J=3 is favored over J=2. T _{1/2} : weighted average of 0.083 ps +21-7 ((n,n' γ), 1985Ko27); 0.087 ps 21 ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$), 1978Si02; 0.13 ps 3 (($\alpha,\text{n}\gamma$), 1976Ch11). XREF: Others: AC, AD, AE J ^π : log ft=5.14 from 0 ⁺ . T _{1/2} : from DSA in (n,n' γ) (1985Ko27). Others: from ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$): 0.26 ps 13 (1980Si02), 0.40 ps +21-12 (1976Ch11). Noting a large discrepancy in T _{1/2} results, 1985Ko27 repeated their measurement and obtained a consistent T _{1/2} =0.042 ps. XREF: Others: AD, AE J ^π : $\gamma(\theta)$ in ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$) for 2205 γ . XREF: Others: AC, AD J ^π : $\gamma(\theta, \text{lin pol})$ in ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$) gives 3 ⁺ (poor fit for 2 ⁺), but a 1295 γ to 0 ⁺ (reported in (p, γ) and (n,n' γ), not in ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$)) is inconsistent with J=3. T _{1/2} : weighted average of 0.16 ps +15-6 ((n,n' γ), 1985Ko27); 0.15 ps 5 ($\alpha,\text{n}\gamma$), ($^{11}\text{B},2\text{npy}$), 1980Si02; 0.33 ps +14-8 (($\alpha,\text{n}\gamma$), 1976Ch11). XREF: Others: AK
3005.73 14	2 ⁺	0.057 ps 8	KL OP r T w	
3071.4 7	(1,2 ⁺)		O	
3077.77 13	4 ⁺	0.55 ps 6	CDEFG KLM OPQ ST w	
3094.64 9	(3) ⁺	0.090 ps 11	KL OPQR T Vw	
3186.84 6	1 ⁺	0.042 ps 10	B KL P R T	
3196.9 4	(2,3)		K P ST	
3205.98 9	(3) ⁺	0.18 ps 5	KL P R T	
3240 20	(0 ⁺)			

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁶⁴ Zn Levels (continued)										
E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF							Comments
3261.94 9	1	0.4 ps +7–2	B	H	KL	P	R	T	W	J ^π : L(p,t)=0. XREF: Others: AC , AD , AE J ^π : log <i>ft</i> =6.1 from 0 ⁺ ; γ to 0 ⁺ . T _{1/2} : from (n,n'γ) (1985Ko27). Others (from (α,nγ),(¹¹ B,2npγ)): 0.042 ps 14 (1980Si02); 0.014 ps 8 (1976Ch11). Noting a large discrepancy in T _{1/2} results, 1985Ko27 repeated their measurement and obtained a consistent T _{1/2} =0.4 ps.
3285 3 3297.17 14	(1 ⁻ to 5 ⁻) (2) ⁺	0.27 ps 5			KL	P	R	T	S	J ^π : 1 ⁻ ,2,3,4,5 ⁻ from primary γ from (3 ⁻). XREF: Others: AC , AD , AK J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ) and L(p,p')=2. T _{1/2} : from (α,nγ),(¹¹ B,2npγ). Weighted average of 0.23 ps 7 (1980Si02), 0.31 ps 7 ((α,nγ) (1976Ch11)).
3306.85 15	(4 ⁺)	0.26 ps 8		F	K	P				XREF: Others: AD , AK XREF: AD(3305). J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ). T _{1/2} : from (¹¹ B,2npγ) 1980Si02 . XREF: B(?).
3321.8? 12	(1)		B							J ^π : weak ε branch (log <i>ft</i> =7.1) from 0 ⁺ ; γ to 0 ⁺ . XREF: Others: AC , AD , AE XREF: L(?)AC(?).
3365.99 6	1 ⁺	0.023 ps 8	B		KL	P	RST	vw	Y	J ^π : log <i>ft</i> =5.04 from 0 ⁺ . T _{1/2} : from (α,nγ),(¹¹ B,2npγ) (1980Si02). Others: 0.026 ps +19–15 (1976Ch11), 0.028 ps 5 (in (γ,γ')).
3369.86 13	3 ⁺	0.35 ps +14–10		K	P	T	vw			XREF: Others: AC , AD , AK XREF: AK(3340). J ^π : γ(θ,lin pol) in (¹¹ B,2npγ); but L(p,p')=(1)+2 for a 3367 doublet and L(p,t)=2 for a 3340 group suggest 2 ⁺ . E(level): there may be an additional 2 ⁺ level near this energy as suggested by L(p,t) and L(p,p').
3414 3	(1 ⁻ to 5 ⁻)							S	v	T _{1/2} : from (α,nγ),(¹¹ B,2npγ) (1980Si02). XREF: Others: AK XREF: AK(3410). J ^π : 1 ⁻ ,2,3,4,5 ⁻ from primary γ from (3 ⁻). XREF: Others: AC , AD , AE XREF: L(?)AC(?).
3425.13 10	1 ⁺	0.031 ps 7	B		KL	PQR	T	v	Y	J ^π : log <i>ft</i> =5.63 from 0 ⁺ . T _{1/2} : from (α,nγ),(¹¹ B,2npγ) (1980Si02). Others: <0.010 ps (1976Ch11), 0.044 ps 11 (in (γ,γ')).
3452.0 10	(1,2 ⁺)							T		XREF: Others: AE XREF: T(3454). J ^π : γ to 0 ⁺ . XREF: Others: AC
3458.66 17	(2,3)	0.24 ps 6		K	P	R	T			J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ). T _{1/2} : from (α,nγ),(¹¹ B,2npγ) (1980Si02). Other: 0.17 ps +42–8 (1985Ko27) in (n,n'γ).
3465 5	(5,4,6) ⁻									XREF: Others: AD J ^π : L(p,p')=5.
3500 10 3538.7? 10	(2 ⁺ to 6 ⁺)								W	XREF: Others: AK XREF: Others: AE

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

^{64}Zn Levels (continued)					
E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF		Comments
3545.9? 9	(0 to 3 ⁺)		P	T	J ^π : 2 ⁺ ,3,4,5,6 ⁺ from γ to 4 ⁺ . XREF: Others: AD , AE
3552.3 3	4 ⁺	>1.0 ps	K	R T	J ^π : 0,1,2,3 ⁺ from γ to 1 ⁺ . XREF: Others: AD
3586.9 21					J ^π : $\gamma(\theta, \text{lin pol})$ in $(\alpha, n\gamma)$, ($^{11}\text{B}, 2n\text{p}\gamma$) and γ to 2 ⁺ ; but J=4 inconsistent with a tentative 3551 γ to 0 ⁺ (in (p, γ)). There may be two different levels near 3552. T _{1/2} : from 1980Si02 . XREF: Others: AD , AE
3597.24 20	(2 ⁺ ,3,4 ⁺)		K	PQ T	XREF: AD(3576). XREF: Others: AC , AD , AE , AK
3601.9 10	(1,2 ⁺)			R	J ^π : γ s to 4 ⁺ , 2 ⁺ . Excitation function in (p, γ) suggests 4 ⁺ , whereas, L(p,t)=(2) suggests 2 ⁺ . XREF: Others: AK
3606.5 5	(0 ⁺ to 4 ⁺)		jK		J ^π : γ to 0 ⁺ . J ^π : 0 ⁺ ,1,2,3,4 ⁺ from γ to 2 ⁺ .
3620.7 10	(2 ⁺ to 6 ⁺)		jK		J ^π : 2 ⁺ ,3,4,5,6 ⁺ from γ to 4 ⁺ .
3628.4 5	(4 ⁺)	0.16 ps 5	jK	T	XREF: Others: AD
3630? 3	(0 ⁺ ,6 ⁻)		j	PQ	J ^π : L(p,p')=4; γ to 2 ⁺ . T _{1/2} : from 1980Si02 . E(level): in (p, γ) E=2050 keV, 1980Er05 adopted this energy from (p,p') (1967Br10 , 1974Au04). This level is most likely different from that in (p,p') (E=3633 5 in 1987Ja04) due to different J ^π values for the two levels.
3680 3	(1 ⁻ to 5 ⁻)		n	S w	J ^π : comparison of measured yield in (p, γ), E=2050 keV with Hauser-Feshbach calculations. J ^π : 1 ⁻ ,2,3,4,5 ⁻ from primary γ from (3 ⁻). XREF: Others: AD , AE , AH
3698.9 7			n	w	XREF: ah(3720). J ^π : L(α, α')=3 for a 3720 group suggests 3 ⁻ for 3699 or 3718.
3701.4 4	1 ⁻	0.025 ps 4	H	n P w Y	XREF: Others: AC XREF: H(3680)Y(3704). J ^π : from $\gamma(\theta)$ in (γ, γ') ; L($^6\text{Li}, d$)=1.
3710.0 7	(2 ⁺)		n	T w	T _{1/2} : from (γ, γ') . XREF: Others: AD , AE
3718.4 3	(0 ⁺ to 4 ⁺)	0.031 ps 10	K	n P T w	XREF: T(3707). J ^π : γ s to 0 ⁺ , 4 ⁺ . XREF: Others: AD , AE , AH
3759 3	(1 ⁻ to 5 ⁻)		n	S	XREF: ah(3720). J ^π : 0 ⁺ ,1,2,3,4 ⁺ from γ to 2 ⁺ ; L(α, α')=3 for a 3720 group suggests 3 ⁻ for 3699 or 3718. T _{1/2} : from 1980Si02 . XREF: Others: AD
3780 10	2 ⁺		n		J ^π : 1 ⁻ ,2,3,4,5 ⁻ from primary γ from (3 ⁻). XREF: Others: AK
3795.03 10	1 ⁺		B	PQR T	J ^π : L(p,t)=2. XREF: Others: AD , AE
3815.4 5	(0 ⁺ to 4 ⁺)		h K	w	J ^π : log <i>f</i> =5.58 from 0 ⁺ . XREF: Others: AD
3819.65 21	(0 ⁺ to 4 ⁺)		h	PQ T w	J ^π : 0 ⁺ ,1,2,3,4 ⁺ from γ to 2 ⁺ . XREF: Others: AD
3850.5 4	(≤ 3) ⁽⁺⁾	<0.7 ps	K1	sT Vw	J ^π : 0 ⁺ ,1,2,3,4 ⁺ from γ to 2 ⁺ . XREF: Others: AC , AD

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁶⁴ Zn Levels (continued)									
E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF				Comments		
									J ^π : γ to 2 ⁺ . L(d,n)=1 from 3/2 ⁻ for a 3850 50 group suggests 0 ⁺ to 3 ⁺ . T _{1/2} : from 1980Si02. XREF: Others: AD
3853.27 21	5 ⁺	>2 ps		K1	P	sT	w		J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ). T _{1/2} : from 1980Si02. XREF: Others: AD
3863.7 10	(2 ⁺ to 6 ⁺)			K			w		J ^π : γ to 4 ⁺ . J ^π : 0 ⁺ ,1,2,3,4 ⁺ from primary γ from (2 ⁺). XREF: Others: AC, AD
3880 3	(0 ⁺ to 4 ⁺)			h		R			J ^π : γs to 2 ⁺ , 4 ⁺ . T _{1/2} : from 1980Si02. XREF: Others: AD
3898.5 3	(2 ⁺ ,3,4 ⁺)	0.038 ps 10		h K	PQR	T			J ^π : γs to 2 ⁺ , 4 ⁺ . T _{1/2} : from 1980Si02. XREF: Others: AD
3924.69 ^b 16	5 ⁻	<1.4 ps	CDEFG	KL	PQ				J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ). Also L(p,p')=5. T _{1/2} : from 1977Al14. Others: <1.7 ps (1977We10), >0.35 ps (1980Si02). T _{1/2} (3924.7 level) not lower than ≈0.7 ps from RUL=1 for B(M2)(W.u.). XREF: Others: AD, AK XREF: AK(3920).
3932.0 4	(4,5)			K					J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ). L(p,t)=(2) is inconsistent. There may be an additional 2 ⁺ level near this energy. XREF: Others: AC, AD, AH, AK XREF: AC(?)AK(3920).
3951.9 6	(4 ⁺ ,3 ⁺)			K		T			J ^π : L(p,p')=4; γ to 2 ⁺ . XREF: Others: AD
3993.36 ^{&} 8	6 ⁺	0.12 ps 3	CDEFG	KL					J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ). Also L(p,p')=6. T _{1/2} : from 1980Si02. Others: 0.15 ps 3 (1977We10), <0.14 ps (1976Le31). XREF: Others: AD, AK XREF: AK(4010).
4020.4 4	(2) ⁺			K	P	T			J ^π : L(p,p')=2 and γ to 4 ⁺ . L(p,t)=(0) is inconsistent. There may be an additional 0 ⁺ level near this energy. XREF: Others: AD
4039.7 4	(0 ⁺ to 4 ⁺)			K	P	T			J ^π : 0 ⁺ ,1,2,3,4 ⁺ from γ to 2 ⁺ . XREF: Others: AD
4076.55 20	(5) ⁺	0.49 ps +24-17	CDEF	KL		T			J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npγ); L(p,p')=4. Assignment of 6 ⁺ by 2004Ka18 in (²⁸ Si,4pγ) on the basis of ΔJ=2 deduced from γγ(θ) of 1340γ is inconsistent with γ(θ,pol) data for 1340γ in 1980Si02 from (α,nγ),(¹¹ B,2n2pγ). T _{1/2} : from 1980Si02. XREF: Others: AD, AK XREF: AD(4107)AK(4120).
4110 3	(2) ⁺						T		E(level): from ⁶³ Cu(p,γ) E=3.46 MeV. J ^π : L(p,p')=2. XREF: Others: AD, AE XREF: AD(4132).
4140 3	(2,1) ⁺			n		T			E(level): from ⁶³ Cu(p,γ) E=3.46 MeV. J ^π : L(p,p')=2; possible γ to 0 ⁺ .

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF				Comments
4153.17 22				n P			XREF: Others: AD
4156.53 19	5 ⁻	0.11 ps 3	CDEF	KLmn	ST	w	XREF: Others: AD , AH XREF: AD(4164)AH(4190). J ^π : ΔJ=2, E2 γ from 7 ⁻ ; ΔJ=1 γ to (4 ⁺); analyzing power in (pol p,p') for a group at 4164 10. T _{1/2} : from 1980Si02 .
4159.5 18	1	7.7 fs 25		n P		w y	XREF: Others: AD XREF: AD(4159). J ^π : from γ(θ) in (γ,γ'). T _{1/2} : from (γ,γ').
4181.7 5			jK	n		w	XREF: Others: AC , AD XREF: AC(?).
4205.2 4	(4,3) ⁺			n PQ	T	w	XREF: Others: AC , AD , AK XREF: AC(?)ak(4230). J ^π : L(p,p')=4; γ to 2 ⁺ .
4219 10	(4) ⁺			n			XREF: Others: AD , AK XREF: ak(4230). J ^π : L(p,p')=4.
4236.71 ^a 10	6 ⁺	0.13 ps 4	CDEFG	KL n			XREF: Others: AD J ^π : γ(θ,lin pol) in (α,nγ),(¹¹ B,2npy). Also L(p,p')=6. T _{1/2} : from 1980Si02 . Others: 1.3 ps 2 (1977We10), 42 ps 21 (1977Al14).
4260 3					T		
4288.6 4	(4) ⁺		K			Z	XREF: Others: AD , AK J ^π : L(p,p')=4. Also L(e,e')=2+4.
4304.1 22	(1 ⁻ to 5 ⁻)				S		XREF: Others: AK J ^π : 1 ⁻ ,2,3,4,5 ⁻ from primary γ from (3 ⁻).
4310 3					T		
4319.1 22	(4,3) ⁺				T	w	XREF: Others: AC , AD , AK XREF: AC(?)AK(4340). J ^π : L(p,p')=4; γ to 2 ⁺ .
4362.1 22	(2,1,3) ⁺				ST	w	XREF: Others: AD , AK XREF: AD(4351). E(level): from ⁶³ Cu(p,γ) E=3.46 MeV. J ^π : L(p,p')=2.
4370 3	3 ⁻				T	w	XREF: Others: AD , AH , AK XREF: AD(4385)AH(4370)AK(4410). E(level): from ⁶³ Cu(p,γ) E=3.46 MeV. J ^π : L(α,α')=3; L(p,t)=(3). L(p,p')=(1).
4380 3					T		
4420 3	(4,3) ⁺				T	vw	XREF: Others: AD E(level): from ⁶³ Cu(p,γ) E=3.46 MeV. J ^π : L(p,p')=4. L(d,n)=1 from 3/2 ⁻ for 4420 50 gives 0 ⁺ to 3 ⁺ .
4454.68 15	1 ⁺	3.2 fs 6	B			vw Y	XREF: Others: AC , AD J ^π : log ft=5.44 from 0 ⁺ ; but L(p,p')=1 for a 4453 10 group gives negative parity, unless an unnatural parity state is populated in (p,p'). T _{1/2} : from (γ,γ').
4470 3	(0 ⁺)				T	vw	XREF: Others: AD , AK XREF: AK(4480). J ^π : L(p,t)=(0).
4488 10	(4,3,5) ⁺					w	XREF: Others: AD J ^π : L(p,p')=4.
4504 10							XREF: Others: AD
4522 10							XREF: Others: AD

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF		Comments
4538 10	(4,3,5) ⁺				XREF: Others: AD
4560 3				T	J ^π : L(p,p')=4. XREF: Others: AC , AD XREF: AC(?). E(level): from $^{63}\text{Cu}(p,\gamma)$ E=3.46 MeV. XREF: Others: AD , AK
4573 10	(1 ⁻ ,0 ⁻ ,2 ⁻)			J	J ^π : L(p,p')=(1).
4608.75 20	(1)		B		XREF: Others: AD XREF: AD(4593). J ^π : log ft≈6.3 from 0 ⁺ .
4615 10	(4,3,5) ⁺				XREF: Others: AD
4626 10					J ^π : L(p,p')=4.
4634.87 9	7 ⁻	94 ps 6	CDEFG	KLM	XREF: Others: AD XREF: Others: AD , AH μ=1.6 3 (1983Ba69 , 2020StZV) XREF: M(4650)AD(4648). J ^π : γ(θ,lin pol) in (α,nγ), (^{11}B ,2npy) and L(p,p')=7. T _{1/2} : weighted average of 105 ps 13, 99 ps 10 (1977We10); 90 ps 10, 80 ps 14 (1977Al14). μ: integral PAC method, recoil into gas and vacuum (1983Ba69).
4638.2 5					XREF: Others: AD
4664 3	(1)	41 fs 12		Y	XREF: Others: AD , AK XREF: Y(?). J ^π : from γ(θ) in (γ,γ').
4668.93 19	(6 ⁻)		CDEF	L	T _{1/2} : from (γ,γ') for %Iγ(to g.s.)=100. XREF: Others: AD , AK J ^π : ΔJ=1 γs to 5 ⁻ and (5 ⁺). Negative parity proposed in (^{28}Si ,4pγ) (2004Ka18), (1998Ga11), but positive parity proposed by 1998Ga11 .
4684 3	(1 ⁻ to 5 ⁻)			S	XREF: Others: AD , AK XREF: AD(4702).
4713.15 21	(1)		B		J ^π : 1 ⁻ ,2,3,4,5 ⁻ from primary γ from (3 ⁻). XREF: Others: AD
4729 10					J ^π : weak ε branch (log ft≈6.0) from 0 ⁺ . XREF: Others: AD
4751 10	(4 ⁺ ,3 ⁺ ,5 ⁺)			w	XREF: Others: AD , AK
4761 10				w	J ^π : L(p,p')=4.
4786 10	(4 ⁺ ,3 ⁺ ,5 ⁺)			w	XREF: Others: AD , AH , AK
4797 10				w	XREF: Others: AD , AK
4816 10	(2 ⁺ ,1 ⁺ ,3 ⁺)			w	J ^π : L(p,p')=4.
4823.5 6	(5,6,7)			L	XREF: Others: AD , AK XREF: AD(4831). J ^π : γ to (5 ⁺) and heavy-ion excitation.
4851 10	(4 ⁺ ,3 ⁺ ,5 ⁺)				XREF: Others: AD , AK
4902 10	(4 ⁺ ,3 ⁺ ,5 ⁺)				J ^π : L(p,p')=4. XREF: Others: AD , AK
4935 10	(3 ⁻ ,2 ⁻ ,4 ⁻)				J ^π : L(p,p')=4. XREF: Others: AD , AK
4947 10	(2 ⁺)				J ^π : L(p,p')=3. XREF: Others: AD , AK XREF: ak(4980). J ^π : L(p,t)=2 for a 4980 30 group.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF		Comments
4980.87 ^b 17	7 ⁻	1.3 ps 4	CDEFG	KL	XREF: Others: AD XREF: AD(4970). J ^π : ΔJ=2, E2 γ to 5 ⁻ ; L(p,p')=7. T _{1/2} : from 1977We10 . Other: 3.1 ps 7 (1977Al14).
5005 10	2 ⁺			w	XREF: Others: AD , AK XREF: AK(4980). J ^π : L(p,p')=2; L(p,t)=2 for a 4980 30 group.
5038 10				w	XREF: Others: AD , AK
5050 10	(0 to 3) ⁽⁺⁾			vw	XREF: Others: AD , AH , AK J ^π : L(d,n)=1 from 3/2 ⁻ for a 5050 50 group.
5066.8 20				vw	XREF: Others: AC , AD , AK XREF: AC(?).
5081 10				w	XREF: Others: AD , AK
5110 10			n	w	XREF: Others: AD , AK
5121 10	(2,1,3) ⁺		n	w	XREF: Others: AD J ^π : L(p,p')=2.
5138 10			n	w	XREF: Others: AD
5151.71 12	(7 ⁻)		C F	L n w	XREF: Others: AD , AK J ^π : ΔJ=2 γ to 5 ⁻ .
5160 10			n	w	XREF: Others: AD , AK
5171 10				w	XREF: Others: AD , AK
5191 10	(3,2,4) ⁻		n	w	XREF: Others: AD , AK J ^π : L(p,p')=3.
5197 10			n		XREF: Others: AD
5211 10			n		XREF: Others: AD , AK
5224 10			n		XREF: Others: AD , AK
5234 10			n		XREF: Others: AD , AK
5256 10			n		XREF: Others: AD
5267 10			n		XREF: Others: AD
5292 10			n	w	XREF: Others: AD
5307 10			M	w	XREF: Others: AD XREF: M(5300).
5319 10				w	XREF: Others: AD
5329 10				w	XREF: Others: AD
5337 10					XREF: Others: AD
5351 10					XREF: Others: AD
5361 10					XREF: Others: AD
5375 10	(3 ⁻)				XREF: Others: AD , AH J ^π : L(α,α')=3 for 5370 45 group.
5384 10					XREF: Others: AD
5398 10					XREF: Others: AD
5413 10					XREF: Others: AD
5425 10					XREF: Others: AD
5443 10					XREF: Others: AD
5457 10					XREF: Others: AD
5474 10					XREF: Others: AD
5485 10	(0 to 3) ⁽⁺⁾			V	XREF: Others: AD J ^π : L(d,n)=(1) from 3/2 ⁻ for 5480 50 group.
5495 10	(4 ⁺)			Z	XREF: Others: AD XREF: Z(5500). J ^π : L=4, E4 excitation in (e,e').
5517 10					XREF: Others: AD
5530 10					XREF: Others: AD
5545 10					XREF: Others: AD
5553 10					XREF: Others: AD
5564 10					XREF: Others: AD

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF		Comments
5576 10					XREF: Others: AD
5588 10					XREF: Others: AD
5601 10					XREF: Others: AD
5613 10					XREF: Others: AD
5623.75 21	(8 ⁻)		CDEF	L	XREF: Others: AD J ^π : ΔJ=2 γ to (6 ⁻).
5642 10					XREF: Others: AD
5652 10					XREF: Others: AD
5665 10					XREF: Others: AD
5676 10					XREF: Others: AD
5689 10					XREF: Others: AD
5699.38 18	(8 ⁻)		C F	LM	XREF: Others: AD XREF: M(5700). J ^π : ΔJ=1 γ to (7 ⁻); ΔJ=2 γ to (6 ⁻).
5719 10					XREF: Others: AD
5729 10					XREF: Others: AD
5737 10					XREF: Others: AD
5760 10					XREF: Others: AD
5770 10					XREF: Others: AD
5780 10					XREF: Others: AD
5792 10					XREF: Others: AD
5812 10					XREF: Others: AD , AH J ^π : L(α,α')=5 for a 5800 45 group suggests 5 ⁻ for one of the levels.
5822 10					XREF: Others: AD
5833 10					XREF: Others: AD
5844 10					XREF: Others: AD
5860 10					XREF: Others: AD
5872 10					XREF: Others: AD
5882 10					XREF: Others: AD
5893 10					XREF: Others: AD
5909 10					XREF: Others: AD
5920 10					XREF: Others: AD
5933 10					XREF: Others: AD
5936.0 7	(8 ⁺)		CDEF		J ^π : ΔJ=(2) γ to 6 ⁺ .
5951.7 5	(9 ⁻)		EF	L	XREF: Others: AD J ^π : ΔJ=2 γ to 7 ⁻ ; γ to (8 ⁻).
6031.5 ^a 4	(8 ⁺)		CDEF		J ^π : ΔJ=2 γ to 6 ⁺ ; γ to 7 ⁻ .
6124.0 4	(8 ⁺)		CDEF		E(level): see comment for 6126 level. J ^π : ΔJ=2 γ to 6 ⁺ ; ΔJ=1 γ to 7 ⁻ .
6124.7 ^b 4	(9 ⁻)		CDE	L	E(level),J ^π : only one level proposed by 2004Ka18 in (²⁸ Si,4pγ) E=122 MeV. 1998Ga11 , 1997Fu08 and 1994Cr05 proposed two levels near this energy with J ^π =8 ⁺ and 9 ⁻ , respectively, with the placement of 1144γ from 9 ⁻ . This placement also proposed by 1978Ne02 in (α,nγ) and ⁵⁶ Fe(¹⁴ N,αpnγ) based on 1144γ(θ) result consistent with ΔJ=2, Q γ to 7 ⁻ level. J ^π : γ to 7 ⁻ .
6262.1 6	(7,8,9 ⁻)		C		
6300 50				M	
6377.0 22	(7,8,9 ⁻)		F	J	XREF: J(6390). J ^π : γ to 7 ⁻ ; yrast pattern of population.
6700 50				M	
6830				J	
6963.0 4	(9)		C		J ^π : ΔJ=1 γ to (8 ⁻); γ to (9 ⁻).
6998.1 5	(11 ⁻)	0.97 ps 21	CDEFG	KL	E(level): in (α,nγ),(¹¹ B,2npγ); (²⁸ Si,4pγ) E=115 MeV; and (⁵⁴ Fe,2pγ), this level corresponds to 5681, 9 ⁻ where

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

E(level) [†]	J ^π [‡]	XREF	Comments
			the placement of 1046γ was differently ordered.
			J ^π : ΔJ=2, E2 γ to (9 ⁻).
			T _{1/2} : from 1977We10. Other: 4.0 ps 5 (1977Al14). This half-life was assigned (1977We10,1977Al14) to 5681, 9 ⁻ level; but with the reassignment of the 1314-1046 cascade, evaluators assign the half-life to 6998 level.
7000	(4 ⁺)		Z J ^π : L=4, E4 excitation in (e,e').
7062.4 4	(10 ⁻)	F	J ^π : ΔJ=2 γ to (8 ⁻).
7118.9 4	(10 ⁺)	CDEF	J ^π : ΔJ=2 γ to (8 ⁺); γ to (9 ⁻).
7212.4 7	(11 ⁻)	F	J ^π : ΔJ=2 γ to (9 ⁻).
7334.7 5	(10 ⁺)	C	J ^π : ΔJ=2 γ to (8 ⁺).
7380		M	Y XREF: M(7400).
7556.2 22	(10 ⁻)	F	J ^π : γ to (9 ⁻).
7579.1 ^b 4	(11 ⁻)	C	J ^π : ΔJ=2 γ to (9 ⁻).
7806.0 10	(10 ⁺)	C	J ^π : ΔJ=2 γ to (8 ⁺).
7900 50		M	
7902 3	(9,10,11 ⁻)	C	J ^π : γ to (9 ⁻).
7946.5 21	(10 ⁺)	C	J ^π : ΔJ=2 γ to (8 ⁺).
8157.1 21	(10 ⁺)	C	J ^π : ΔJ=(2) γ to (8 ⁺).
8181.1 11	(10 ⁻)	F	J ^π : γ to (9 ⁻).
8302.8 6	(12 ⁻)	F	J ^π : γ to (11 ⁻).
8322.1 22	(11)	F	J ^π : γ to (10 ⁺).
8426.1 4	(11 ⁻)	DEF	J ^π : γs to (9 ⁻) and (11 ⁻).
8580.4 5	(12 ⁺)	CDEF	J ^π : ΔJ=2 γ to (10 ⁺); γ to (11 ⁻).
8995.4 10	(12 ⁺)	C	J ^π : ΔJ=2 γ to (10 ⁺).
9363.7 8	(11 ⁻)	F	J ^π : γs to (9 ⁻) and (10 ⁻).
9440.3 6	(11 ⁻)	F	J ^π : γ to (10 ⁺).
9666 3	(14)	EF	J ^π : γ to (12).
9772 2	(2 ⁺)	R	Additional information 4.
			E(level): proton capture state, E(p)(lab)=2098 keV.
			J ^π : γs to 0 ⁺ and 4 ⁺ .
9803.5 7	(11 ⁻)	F	J ^π : γ to (10 ⁻); ΔJ=2 γ from (13 ⁻).
9948.4 ^d 6	(12 ⁻)	F	J ^π : ΔJ=1 γ to (11 ⁻); ΔJ=2 γ to (10 ⁻).
10.31×10 ³ 50		Q	Additional information 5.
			E(level): average proton-resonance, E(p)=2.1-3.1 MeV range in the c.m. system.
10460.2 ^c 6	(13 ⁻)	D F	J ^π : ΔJ=2 γ to (11 ⁻); γ to (12 ⁻).
10872	(3 ⁻)	S	Additional information 6.
			E(level),J ^π : proton resonance state, E(p)=3217 resonance, identified as g _{9/2} IAR of 1546 level in ^{64}Cu , with γ decay similar to the decay of 3251 keV resonance (1976Fo06).
10906	(3 ⁻)	S	Additional information 7.
			E(level),J ^π : proton resonance state, E(p)=3251 resonance, identified as g _{9/2} IAR of 1589 level in ^{64}Cu ; spin from I _γ (90°)/I _γ (0°) of primary transitions. Parity from decay modes and lack of 3 ⁺ in the parent nucleus ^{64}Cu .
11023.4 ^d 6	(14 ⁻)	D F	J ^π : ΔJ=1 γ to (13 ⁻); γ to (12 ⁻).
11120	(2 ⁺)	T	Additional information 8.
			E(level): proton-resonance state from E(p)(lab)=3.46 MeV and S(p)(^{64}Zn)=7713.1 keV 6 (2021Wa16).
			J ^π : γ rays to 0 ⁺ and 4 ⁺ .
11464 ^e 4	(15)	F	J ^π : γ to (14).
11626.4 ^c 7	(15 ⁻)	D F	J ^π : ΔJ=1 γ to (14 ⁻); ΔJ=2 γ to (13 ⁻).
12335.7 ^d 7	(16 ⁻)	D F	J ^π : γs to (14 ⁻) and (15 ⁻).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

^{64}Zn Levels (continued)				
E(level) [†]	J^π [‡]	$T_{1/2}$ [#]	XREF	Comments
12468 ^f 4	(16)		F	J^π : γ to (15).
13082.1 ^c 7	(17 ⁻)		D F	J^π : $\Delta J=1$ γ to (16 ⁻); γ to (15 ⁻).
13324 ^e 4	(17)		F	J^π : γ s to (15) and (16).
13948.1 ^d 8	(18 ⁻)		D F	J^π : $\Delta J=1$ γ to (17 ⁻); $\Delta J=2$ γ to (16 ⁻).
14391 ^f 3	(18)		F	J^π : γ s to (16) and (17).
14857 6			F	J^π : γ from (19).
14862.5 ^c 8	(19 ⁻)		D F	J^π : $\Delta J=1$ γ to (18 ⁻); $\Delta J=2$ γ to (17 ⁻).
15.42 $\times 10^3$ 94	1 ⁻	4.6 MeV +16-15		XREF: Others: AH %EWSR=19 for E1 isoscalar giant dipole resonance (ISGDR) strength.
15423.6 ^e 25	(19)		F	J^π : γ s to (17) and (18).
15.7 $\times 10^3$ 5	2 ⁺	6.43 MeV 65		XREF: Others: AH %EWSR=113 for E2 isoscalar giant quadrupole resonance (ISGQR) strength.
15939 7			F	J^π : γ from (20).
15945.0 ^d 9	(20 ⁻)		D F	J^π : $\Delta J=2$ γ to (18 ⁻); γ to (19 ⁻).
16686.8 ^f 25	(20)		F	J^π : γ s to (18) and (19).
17084 5			F	J^π : γ from (21).
17087.2 ^c 10	(21 ⁻)		D F	J^π : $\Delta J=2$ γ to (19 ⁻); γ to (20 ⁻).
17853 ^e 4	(21)		F	J^π : γ s to (19) and (20).
18.34 $\times 10^3$ 70	0 ⁺	9.21 MeV 114		XREF: Others: AH %EWSR=64 for E0 isoscalar giant monopole resonance (ISGMR) strength.
18483.3 ^d 11	(22 ⁻)		D F	J^π : $\Delta J=2$ γ to (20 ⁻); γ to (21 ⁻).
19365 ^f 4	(22)		F	J^π : γ to (20).
19775.6 ^c 13	(23 ⁻)		D F	J^π : $\Delta J=2$ γ to (21 ⁻); γ to (22 ⁻).
20657 ^e 5	(23)		F	J^π : γ to (21).
21297.5 ^d 15	(24 ⁻)		F	J^π : γ s to (22 ⁻) and (23 ⁻).
22892.7 ^c 17	(25 ⁻)		F	J^π : γ to (23 ⁻).
24868.6 ^d 18	(26 ⁻)		F	J^π : γ to (24 ⁻).
25.6 $\times 10^3$ 12	1 ⁻	12.6 MeV 32		XREF: Others: AH %EWSR=68 for E1 isoscalar giant dipole resonance (ISGDR) strength.

[†] From a least-squares fit to E_γ data for levels populated in γ -ray studies, assuming $\Delta E_\gamma=3$ keV for high-energy γ rays from proton capture and resonance states. Normalized $\chi^2=0.92$. Energies of levels populated only in particle-transfer reactions are primarily from (p,p'). Due to high level density and limited resolution, correspondence of levels, above ≈ 3.5 MeV excitation, from different reactions is somewhat ambiguous.

[‡] In cases where L(p,p') is used, parity is $(-1)^L$ and spin is L for levels up to 3.2 MeV, with the possibility of L-1, L, L+1 for higher levels, although J=L is the most likely choice, which is listed first, followed by less likely J=L-1 and J=L-2. For levels above ≈ 5 MeV populated in in-beam high-spin studies, J^π values are based on $\gamma(\theta)$, $\gamma\gamma(\theta)$ (DCO), band associations, and assumption of ascending spins with excitation energy.

[#] Mainly from DSA method in $(\alpha, n\gamma)$, $(^{11}\text{B}, 2n\gamma)$ (also $(\alpha, n\gamma)$ and $(n, n'\gamma)$). Values quoted from [1977Al14](#) are from recoil distance method (RDDS) in $(^{11}\text{B}, 2n\gamma)$. Above 3458 values are available from $(^{11}\text{B}, 2n\gamma)$ only. For some of the levels, values from different studies are in disagreement and are noted under comments.

[@] Double β decay to ^{64}Ni is possible with Q value=1095.77. From measurements of double β decay, lower limits of half-life for decay to ^{64}Ni g.s. have been determined (generally at 90% confidence level): $T_{1/2}(2\nu 2K)$: $\geq 1.1 \times 10^{19}$ y ([2011Be39](#), also [2010Be41](#), [2009Be27](#), [2008Be02](#)); $T_{1/2}(\epsilon\beta^+)$: $\geq 1.1 \times 10^{18}$ y ([2009Da16](#)), $\geq 1.3 \times 10^{20}$ y ([2007Ki13](#)); $T_{1/2}(2\epsilon)$: $\geq 3.3 \times 10^{17}$ y

Adopted Levels, Gammas (continued) ^{64}Zn Levels (continued)

(2009Da16); $T_{1/2}(2\beta^+)$: $>10\times10^{17}$ y (1952Fr23); $T_{1/2}(0\nu2\varepsilon)$: $\geq3.2\times10^{20}$ y (2011Be39), $\geq1.19\times10^{17}$ y (2007B115), $\geq9.52\times10^{16}$ y (2006Zu02,2006Wi12), $\geq1.0\times10^{18}$ y (2005Da47); $T_{1/2}(0\nu\varepsilon\beta^+)$: $>1.2\times10^{22}$ y (2020Az05), $\geq8.5\times10^{20}$ y (2011Be39), $\geq5.07\times10^{18}$ y (2006Zu02,2006Wi12), $\geq3.6\times10^{18}$ y (2005Da47), $>2.8\times10^{16}$ y (2003Ki08), 1.1×10^{19} y 9 (1995Bi24, from observed 511 keV peak, but systematic effects were not estimated); $T_{1/2}(2\nu\varepsilon\varepsilon)$: $>6.0\times10^{16}$ y (2003Ki08, also 2005Zu01,2001Zu03); $T_{1/2}(2\nu\varepsilon\beta^+)$: $\geq9.4\times10^{20}$ y (2011Be39), $\geq8.9\times10^{18}$ y (2005Da47); $T_{1/2}(0\nu+2\nu,\beta^+\varepsilon)>2.3\times10^{18}$ y (1985No03); $T_{1/2}(0\nu+2\nu,2\varepsilon)>8\times10^{15}$ y (1953Be33). Others $T_{1/2}$: 1999TsZZ, 2002Tr04 (evaluation).

& Band(A): g.s. band.

^a Band(B): Band based on 2^+ .

^b Band(C): Band based on 3^- .

^c Band(D): Collective (strongly coupled) band, $\alpha=1$. Configuration= $\pi[(f_{7/2}^{-1})(p_{3/2}f_{5/2}^2(g_{9/2}^1) \otimes \nu[(p_{3/2}f_{5/2})^4(g_{9/2}^2)]]$; also [11,02] in the notation used by 2004Ka18, implying one proton hole in $f_{7/2}$ and one proton in $g_{9/2}$ orbitals, no neutron hole in $f_{7/2}$ orbital and 2 neutrons in $g_{9/2}$ orbital.

^d Band(d): Collective (strongly coupled) band based, $\alpha=0$. See configuration listed above for its signature partner.

^e Band(E): Strongly coupled band, $\alpha=1$.

^f Band(e): Strongly coupled band, $\alpha=0$.

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$										
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	$\alpha^@$	$I_{(\gamma+ce)}$	Comments
991.54	2 ⁺	991.53 5	100	0.0	0 ⁺	E2				B(E2)(W.u.)=20.0 5 E _γ : NRM weighted average of 10 values from different datasets where uncertainties are given. This procedure increases the uncertainty in one of the discrepant values (991.16 4 in (²⁸ Si,4pγ) E=115 MeV) from 0.04 keV to 0.15 keV. Regular weighted average is 991.37 7, but with reduced $\chi^2=6.9$, while unweighted average is 991.41 8. Removal of the 991.16 4 value gives weighted average of 991.56 5.
1799.41	2 ⁺	807.85 6	100.0 16	991.54	2 ⁺	E2+M1	-3.9 7			B(M1)(W.u.)=0.00099 +47-27; B(E2)(W.u.)=39 4 E _γ : NRM weighted average of ten values. δ : from weighted average of -4.6 10 in (n,n'γ) (1985Ko42); -3.3 7 in (α,nγ),(¹¹ B,2npγ) (1978Si02); -5.5 40 in ⁵¹ V(¹⁶ O,p2nγ), ⁵⁹ Co(⁷ Li,2nγ) (1977We10); and -4.5 15 (1964Se02) in (p,p'γ). Others: -1.3 3 (1978We15), -0.08 3 (1977We10), -0.45 5 (1977Al14) in ⁵¹ V(¹⁶ O,p2nγ), ⁵⁹ Co(⁷ Li,2nγ); -0.57 +13-27 (1976Br12) in (α,nγ),(Hl,xnγ). Evaluators prefer high value of $\delta(\text{E2/M1})$ (dominant E2) for transition from the second 2 ⁺ to first 2 ⁺ from a trend in other even-even nuclei.
		1799.34 11	29.6 7	0.0	0 ⁺	E2				B(E2)(W.u.)=0.225 +25-22 E _γ : weighted average of ten values. I _γ : NRM weighted average of ten values. Weighted average is 30.3 16, but with reduced $\chi^2=15$.
1910.26	0 ⁺	110.7 1	3.4 11	1799.41	2 ⁺	[E2]		0.447		B(E2)(W.u.)=76 24 E _γ ,I _γ : from ⁶⁴ Ga ε decay. B(E2)(W.u.)=0.057 3 E _γ : weighted average of five values. I _γ : from ⁶⁴ Ga ε decay. Mult.: from ce data in (p,p'γ). Mult.: transition seen only in ce data from (p,p'γ). I _(γ+ce) : I(ce+pair). q _K ² (E0/E2)=6.0 5, X(E0/E2)=2.25 19, ρ ² (E0)=0.0038 4 (1985Pa07,1986Pa23,2005Ki02).
		918.77 5	100.0 34	991.54	2 ⁺	E2				B(E2)(W.u.)=12.2 +5-4 E _γ : weighted average of seven values. B(E2)(W.u.)<4.5 E _γ ,I _γ : from (α,nγ),(¹¹ B,2npγ). I _γ <2.0 in (p,p'γ) (1985Pa07). B(E2)(W.u.)=17 +11-5 E _γ : NRM weighted average of five values. Mult.: from ce data in (p,p'γ). Mult.: transition seen only in ce data from (p,p'γ).
		1910		0.0	0 ⁺	E0			0.64 13	
2306.72	4 ⁺	1315.15 5	100	991.54	2 ⁺	E2				
2609.52	0 ⁺	809 ^a	<0.5	1799.41	2 ⁺	[E2]				
		1617.93 19	100	991.54	2 ⁺	E2				
		2610		0.0	0 ⁺	E0			0.0030 6	

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)									Comments
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	$\alpha^@$	
2736.57	4 ⁺	429.77 13	9.3 6	2306.72	4 ⁺	M1+E2	-0.25 9		<p>$I_{(\gamma+ce)}$: I(pair+ce). $q_K^2(E0/E2)=0.027\ 6$, $X(E0/E2)=0.031\ 7$, $\rho^2(E0)=0.015\ 7$ (1985Pa07,1986Pa23,2005Ki02). $B(M1)(W.u.)=0.014\ +4-3$; $B(E2)(W.u.)=8\ +7-5$ E_γ: NRM weighted average of eight values. I_γ: weighted average of eight values. δ: Other: $+1.7\ 5$ or $-0.2\ 3$ (1977We10) in $^{51}\text{V}(^{16}\text{O},p2n\gamma)$. $B(E2)(W.u.)=30\ +8-5$ E_γ: weighted average of nine values. $B(E2)(W.u.)=0.064\ +20-15$ E_γ: weighted average of three values with consistent I_γ values. I_γ: NRM weighted average of three values. Others: 23 6 (n,n'γ); 33 (p,p'γ); 42 4 in (p,γ) $E=2050$ keV; 154 13 in (p,γ) $E=2098$ keV are too high and in severe disagreement. $B(M1)(W.u.)=0.052\ +26-23$; $B(E2)(W.u.)=13\ +13-10$ γ reported in (p,p'γ) only. $B(M1)(W.u.)=0.028\ +17-14$; $B(E2)(W.u.)=0.08\ +16-7$ E_γ: weighted average of three values. $B(M1)(W.u.)=0.0033\ +20-17$; $B(E2)(W.u.)=0.10\ +6-5$ E_γ, I_γ: unweighted average of three values. $B(E1)(W.u.)\approx 0.000062$ E_γ, I_γ: from (n,n'γ). $B(E1)(W.u.)=3.29\times 10^{-4}\ 11$ E_γ: NRM weighted average of eight values. $B(E3)(W.u.)=72\ +44-35$ γ from $^{61}\text{Ni}(\alpha, n\gamma), ^{56}\text{Fe}(^{11}\text{B}, 2n\text{p}\gamma)$ only. $B(E3)(W.u.)$ from $B(E3)=0.040\ 7$ ((e,e'), 1976Ne06). Other: 70 50 from $T_{1/2}$ and E3 γ branching. $B(E2)(W.u.)\approx 12.6$ γ reported in (p,p'γ) only. Considered as uncertain by evaluators. $B(M1)(W.u.)=0.050\ +15-21$; $B(E2)(W.u.)=21\ +24-16$ E_γ, I_γ: weighted average of three values. $B(M1)(W.u.)=0.019\ +5-4$; $B(E2)(W.u.)<0.25$ E_γ: weighted average of three values. $B(E2)(W.u.)=0.69\ +13-11$ E_γ: weighted average of three values. I_γ: weighted average of two values.</p>
2793.5	2 ⁺	1802.1 4 2793.0 ^a 15	100	991.54	2 ⁺	M1+E2	+0.7 5		
2979.94	3 ⁺	1180.58 15	100 4	1799.41	2 ⁺	M1+E2	-0.05 3		
		1987.0 7	61 14	991.54	2 ⁺	M1+E2	+0.26 3		
2998.54	3 ⁻	1197 1	≈ 4	1799.41	2 ⁺	[E1]			
		2007.03 18	100 3	991.54	2 ⁺	(E1)			
		2997	0.5 3	0.0	0 ⁺	[E3]			
3005.73	2 ⁺	1092 ^a 1	7.5	1910.26	0 ⁺	[E2]			
		1206.2 2	77 5	1799.41	2 ⁺	M1+E2	+0.6 5		
		2014.3 2	100 7	991.54	2 ⁺	M1(+E2)	-0.06 10		
		3005.5 4	65 7	0.0	0 ⁺	E2			
3071.4	(1,2 ⁺)	1272 1 3071 1	100 39	1799.41	2 ⁺ 0 ⁺				
3077.77	4 ⁺	341.2 3	11 1	2736.57	4 ⁺	M1(+E2)	<0.5	0.0036 9	<p>$B(M1)(W.u.)=0.055\ +24-20$; $B(E2)(W.u.)<230$ E_γ: weighted average of three values.</p>

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	Comments
								I_γ : from ⁵⁴ Fe,2p γ). Others: 120 13 (α ,n γ),(¹¹ B,2np γ); 38 10 (α ,n γ), (HI,xn γ); 24 6 (p, γ) E=2050 keV. δ : from RUL(E2)<300. In (α ,n γ),(¹¹ B,2np γ) reaction, 1980Si02 quote $\delta=-1.2$ 2 but in a footnote also state that δ for this transition is not determined from experiment. B(M1)(W.u.)=0.045 6; B(E2)(W.u.)=4.6 +45-30 E_γ : weighted average of seven values. δ : others: -0.4 1 (1978Ne02) in (α ,n γ), (HI,xn γ), -0.54 12 (1978We15) in ⁵¹ V(¹⁶ O,p2n γ). B(E2)(W.u.)=0.71 +11-9 E_γ : weighted average of seven values. Level-energy difference=2086.2. I_γ : weighted average of four values. Others: 190 8 in (²⁸ Si,4p γ) E=122 MeV; 113 9 in (α ,n γ),(¹¹ B,2np γ); 141 9 (p, γ) E=2025 keV are in disagreement. B(M1)(W.u.)=0.018 +8-7; B(E2)(W.u.)=18 8 E_γ : weighted average of two values. I_γ : unweighted average of three values. B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.022 +4-3; B(E2)(W.u.)=8.4 +14-11 E_γ : weighted average of two values. δ : +9.4 15 or +0.40 5 (for J=3); +0.6 4 (for J=2). B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.0090 +35-24 E_γ, I_γ : from ⁶⁴ Ga ε only. B(M1)(W.u.)=0.052 +16-10 E_γ : weighted average of five values. I_γ : from ⁶⁴ Ga ε decay. Other three available values are either imprecise or in disagreement. B(M1)(W.u.)=0.085 +26-16; B(E2)(W.u.)=75 +24-15 E_γ : weighted average of four values. I_γ : in (α ,n γ),(¹¹ B,2np γ) value is low by a factor of ≈ 9 . B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.017 +5-3; B(E2)(W.u.)=6.1 +19-12 E_γ : weighted average of five values. I_γ : from ⁶⁴ Ga ε decay. Other three available values are either imprecise or in disagreement. B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=1.1 $\times 10^{-4}$ +4-3 E_γ, I_γ : from ⁶⁴ Ga ε decay. Other I_γ : 41 8 (p, γ) 2098 keV; 18 in (p, γ) are in disagreement. γ reported in (p,p' γ) only.
3077.77	4 ⁺	770.95 15	100 10	2306.72	4 ⁺	M1+E2	-0.19 8	
		2086.8 3	78 9	991.54	2 ⁺	E2		
3094.64	(3) ⁺	1295.1 2	19 9	1799.41	2 ⁺	[M1+E2]		
		2103.1 1	100 7	991.54	2 ⁺	M1+E2		
3186.84	1 ⁺	577.3 1	0.76 15	2609.52	0 ⁺	[M1]		
		1276.52 16	47.6 11	1910.26	0 ⁺	(M1)		
		1387.34 10	100.0 26	1799.41	2 ⁺	[M1+E2]		
		2195.34 10	80.4 18	991.54	2 ⁺	[M1+E2]		
		3186.8 2	1.5 3	0.0	0 ⁺	[M1]		
3196.9	(2,3)	1397.4 4	52 5	1799.41	2 ⁺			
		2205.3 8	100 5	991.54	2 ⁺			
		3200 ^a 3	18	0.0	0 ⁺			

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)									
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	Comments	
3205.98	(3) ⁺	898.8 5	13 2	2306.72	4 ⁺	[M1+E2]		B(M1)(W.u.)=0.017 +7-4; B(E2)(W.u.)=36 +15-8 B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. γ not reported in (α , γ),(¹¹ B,2np γ). It is considered (by evaluators) as highly questionable since it involves M3 transition, with unrealistically large B(M3)(W.u.)=1.9×10 ⁷ +7-4.	
		1295.1 ^a 7	10 5	1910.26	0 ⁺			B(M1)(W.u.)=0.033 +13-8; B(E2)(W.u.)=1.8 +16-10 B(M1)(W.u.)=0.00028 +11-6; B(E2)(W.u.)=0.096 +36-21 γ reported in (α , γ),(¹¹ B,2np γ) only.	
		1406.57 8	100.0 10	1799.41	2 ⁺	M1+E2	-0.25 9	B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. γ reported in (p, γ) E=2098 keV only, highly questionable as required mult=[M3].	
		2214.0 5	3.1 10	991.54	2 ⁺	[M1+E2]		I γ : from ⁶⁴ Ga ε decay. E γ ,I γ : from ⁶⁴ Ga ε decay. This γ was reported in (p, γ) E=2050 keV; and in (p,p' γ), with I γ =24 in the latter work. It fits poorly in the decay scheme. E γ : weighted average of four values.	
3261.94	1	3205 ^a 3		0.0	0 ⁺			E γ ,I γ : from ⁶⁴ Ga ε decay. Other I γ =90 10 (p, γ) 2098 keV; \approx 67 (n,n' γ) are in disagreement.	
		1352 ^a 1	<6	1910.26	0 ⁺			B(M1)(W.u.)=0.0041 +17-15; B(E2)(W.u.)=3.1 +14-12 γ reported only in (p, γ) E=2098 keV.	
		1461.3 ^a 1	<4.5	1799.41	2 ⁺			B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.0039 +11-8; B(E2)(W.u.)=1.24 +35-25 B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(E2)(W.u.)=0.087 +37-34 γ reported only in (p, γ) E=2098 keV.	
		2270.40 10	100 5	991.54	2 ⁺			B(E2)(W.u.)<59 γ from (α , γ),(¹¹ B,2np γ) only.	
		3261.7 2	6.2 4	0.0	0 ⁺			B(M1)(W.u.)=0.084 +39-24; B(E2)(W.u.)<14 E γ : from (p, γ) E=2050 keV. In several in-beam studies (1998Ga11,1994Cr05, 1978Ne02), a 1000 γ deexcited a level at 4078, but in (α , γ),(¹¹ B,2np γ), 1980Si02 reported the placement from 4078 level as incorrect and placed it from 3307 level, as also proposed in (p, γ). In (²⁸ Si,4p γ) study of 2004Ka18, a 999.9 γ is placed from both the 3307 and 4078 levels. δ : from 1980Si02. Other: -0.1 2 (1978Ne02).	
3297.17	(2) ⁺	1498 1	29 11	1799.41	2 ⁺	[M1+E2]			
		2305.54 14	100 14	991.54	2 ⁺	[M1+E2]			
		3299 1	42 19	0.0	0 ⁺	[E2]			
3306.85	(4 ⁺)	512 ^a	0.5 5	2793.5	2 ⁺	[E2]			
		1000.15 15	100	2306.72	4 ⁺	M1(+E2)	+0.07 20		
3321.8?	(1)	1411.3 15	100 29	1910.26	0 ⁺				
		3322 2	58 16	0.0	0 ⁺				
3365.99	1 ⁺	756.58 10	9.0 5	2609.52	0 ⁺	[M1]		B(M1)(W.u.)=0.11 +6-3 γ reported in ⁶⁴ Ga ε decay only. E γ and I γ data for four higher energy γ rays from the 3366 level have been taken from ⁶⁴ Ga ε decay. These are reported in other datasets, but with imprecise energies and intensities, as compared to the data in ε decay. B(M1)(W.u.)=0.023 +12-6 E γ ,I γ : from ⁶⁴ Ga ε decay. This γ reported in (p, γ) E=2050 keV; and (p,p' γ) but with imprecise energies and intensities.	
		1455.84 12	13.9 7	1910.26	0 ⁺	[M1]			

Adopted Levels, Gammas (continued)

 $\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	Comments
3365.99	1 ⁺	1566.50 18	15.7 9	1799.41	2 ⁺	[M1+E2]		B(M1)(W.u.)=0.021 +11-6; B(E2)(W.u.)=14 +8-4 E _γ , I _γ : from ⁶⁴ Ga ε decay. This γ reported in (p,γ) E=2050 keV; (p,p'γ); and (α,nγ),(HI,xnγ), but with imprecise energies and intensities.
		2374.30 10	50.5 13	991.54	2 ⁺	[M1+E2]		B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.019 +10-5; B(E2)(W.u.)=5.7 +30-15 E _γ , I _γ : from ⁶⁴ Ga ε decay. This γ reported in (p,γ) E=2050 keV; (p,γ) E=2098 keV; (p,p'γ); (α,nγ),(¹¹ B,2npγ) and (α,nγ),(HI,xnγ), but with imprecise energies and intensities.
		3365.80 10	100 3	0.0	0 ⁺	[M1]		B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.013 +7-4 E _γ , I _γ : from ⁶⁴ Ga ε decay. This γ reported in (p,γ) E=2050 keV; (p,γ) E=2098 keV; (p,p'γ); (α,nγ),(¹¹ B,2npγ) and (α,nγ),(HI,xnγ), but with imprecise energies and intensities.
3369.86	3 ⁺	633.40 15	25 6	2736.57	4 ⁺	[M1+E2]		B(M1)(W.u.)=0.034 +16-12; B(E2)(W.u.)=1.4×10 ² +7-5 γ reported in (p,γ) E=2.05 MeV only.
		1570.3 2	100 5	1799.41	2 ⁺	M1+E2	-0.40 6	B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2.
		2377.8 6	57 5	991.54	2 ⁺	[M1+E2]		B(M1)(W.u.)=0.0077 +31-22; B(E2)(W.u.)=0.84 +42-31 B(M1)(W.u.)=0.0015 +6-4; B(E2)(W.u.)=0.44 +18-13 I _γ : weighted average of values from (α,nγ),(¹¹ B,2npγ); (p,γ) E=2050 keV; and (n,n'γ).
3425.13	1 ⁺	419.5 ^a 4	0.5 5	3005.73	2 ⁺	[M1+E2]		B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)<0.092; B(E2)(W.u.)<890 γ reported in (α,nγ),(¹¹ B,2npγ) only.
		1514.7 2	4.8 8	1910.26	0 ⁺	[M1]		B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.0068 +23-16 E _γ , I _γ : from ⁶⁴ Ga ε decay. γ reported in (p,p'γ).
		1625.87 20	24.9 17	1799.41	2 ⁺	[M1+E2]		B(M1)(W.u.)=0.029 +9-6; B(E2)(W.u.)=18 +6-4 E _γ , I _γ : from ⁶⁴ Ga ε decay. Other: I _γ : 71 5 in (p,γ) E=2.05 MeV. Tentative γ also reported in (n,n'γ).
		2433.6 2	13.2 12	991.54	2 ⁺	[M1+E2]		B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.0045 +14-9; B(E2)(W.u.)=1.3 +4-3 E _γ , I _γ : from ⁶⁴ Ga ε decay. γ reported in (p,γ) E=2098 keV.
		3424.97 15	100 5	0.0	0 ⁺	[M1]		B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2. B(M1)(W.u.)=0.0123 +37-23 E _γ , I _γ : from ⁶⁴ Ga ε decay. γ reported in (p,γ) E=2050 keV; (p,γ) E=2098 keV; (p,p'γ); (α,nγ),(¹¹ B,2npγ) and (α,nγ),(HI,xnγ), but with imprecise energies.
3452.0	(1,2 ⁺)	1542 1	100	1910.26	0 ⁺			
3458.66	(2,3)	1659.2 2	100 7	1799.41	2 ⁺			
		2467.1 3	82 9	991.54	2 ⁺			I _γ : from (α,nγ),(¹¹ B,2npγ). Other: 144 11 in (p,γ) E=2.05 MeV.

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	Comments
3458.66	(2,3)	3455 ^a 1		0.0	0 ⁺			γ reported only in (p, γ) E=2.098 keV as the strongest transition from a 3458 level. The placement is considered tentative by evaluators.
3538.7?	(2 ⁺ to 6 ⁺)	1232 1	100	2306.72	4 ⁺			
3545.9?	(0 to 3 ⁺)	284 1	52	3261.94	1			
		359 2	100	3186.84	1 ⁺			
		1747.65 ^a 15		1799.41	2 ⁺			
3552.3	4 ⁺	1246.7 4	100 6	2306.72	4 ⁺	M1+E2	-0.16 10	γ reported only in (p, γ) E=2.05 MeV. B(M1)(W.u.)<0.0057; B(E2)(W.u.)<0.39 γ reported in (¹¹ B,2np γ). This γ may correspond to a 1245 γ (unplaced) in (p,p' γ) and a 1247.2 2 (unplaced) in (p, γ) E=2050 keV.
		2559.7 4	85 6	991.54	2 ⁺	[E2]		B(E2)(W.u.)<0.15 This γ may correspond to a 2560 γ (unplaced) in (p,p' γ). γ reported in (p, γ) E=2.098 keV. Evaluators treat this γ as highly questionable as ΔJ^π requires mult=E4 and an unrealistic large B(E4)(W.u.)<2.1 $\times 10^6$.
		3551 ^a 1	42 15	0.0	0 ⁺			
3586.9		390 2	100	3196.9	(2,3)			
3597.24	(2 ⁺ ,3,4 ⁺)	860.5 3	57 14	2736.57	4 ⁺			γ reported only in (p, γ) E=2.05 MeV.
		1290.5 3	100 9	2306.72	4 ⁺			
		2606.0 5	82 9	991.54	2 ⁺			
3601.9	(1,2 ⁺)	3602 1	100	0.0	0 ⁺			
3606.5	(0 ⁺ to 4 ⁺)	2614.9 5	100	991.54	2 ⁺			
3620.7	(2 ⁺ to 6 ⁺)	1314.0 10	100	2306.72	4 ⁺			
3628.4	(4 ⁺)	2636.8 5	100	991.54	2 ⁺	[E2]		B(E2)(W.u.)=1.8 +8-5
3698.9		502.0 5	100	3196.9	(2,3)			
3701.4	1 ⁻	3701.3 4	100	0.0	0 ⁺	[E1]		B(E1)(W.u.)=3.3 $\times 10^{-4}$ +6-5
3710.0	(2 ⁺)	1099 1	7	2609.52	0 ⁺			
		1406 1	100	2306.72	4 ⁺			E_γ : fits poorly. E_γ =1403 from level-energy difference.
		\approx 3710		0.0	0 ⁺			
3718.4	(0 ⁺ to 4 ⁺)	2726.8 3	100	991.54	2 ⁺			
3795.03	1 ⁺	1185.4 1	4.4 24	2609.52	0 ⁺			γ reported in ⁶⁴ Ga ε decay only.
		1995.9 2	100 6	1799.41	2 ⁺			E_γ, I_γ : from ⁶⁴ Ga ε decay. γ reported in (p, γ) E=2050 keV.
		2803.3 3	39 4	991.54	2 ⁺			E_γ, I_γ : from ⁶⁴ Ga ε decay. γ reported in (p, γ) E=2098 keV.
		3795.1 3	74 5	0.0	0 ⁺			E_γ, I_γ : from ⁶⁴ Ga ε decay. γ reported in (p, γ) E=2050 keV and (p, γ) E=2098 keV.
3815.4	(0 ⁺ to 4 ⁺)	2016.0 5	100	1799.41	2 ⁺			
3819.65	(0 ⁺ to 4 ⁺)	2020.2 2	100	1799.41	2 ⁺			
3850.5	(≤ 3) ⁽⁺⁾	1116 ^a		2736.57	4 ⁺			γ reported in ($\alpha, n\gamma$), (¹¹ B,2np γ). γ reported in ($\alpha, n\gamma$), (¹¹ B,2np γ).
		2051.0 4	22.0 25	1799.41	2 ⁺			
		2859.2 6	100 4	991.54	2 ⁺			
3853.27	5 ⁺	1116.7 2	100.0 5	2736.57	4 ⁺	M1+E2	-1.00 15	B(M1)(W.u.)<0.0046; B(E2)(W.u.)<6.1 B(M1)(W.u.)<3.0 $\times 10^{-5}$; B(E2)(W.u.)<0.021 γ reported in ($\alpha, n\gamma$), (¹¹ B,2np γ). B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2.
		1547 ^a	0.5 5	2306.72	4 ⁺	[M1+E2]		

Adopted Levels, Gammas (continued)

 $\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	Comments
3863.7	(2 ⁺ to 6 ⁺)	1557.0 10	100	2306.72	4 ⁺			
3898.5	(2 ⁺ ,3,4 ⁺)	1162.5 5	20 7	2736.57	4 ⁺			γ reported in (p, γ) E=2.09 MeV.
		2906.5 4	100 13	991.54	2 ⁺			
3924.69	5 ⁻	617.9 5	8.3 5	3306.85	(4 ⁺)	[E1]		B(E1)(W.u.)>6.7×10 ⁻⁵
		848		3077.77	4 ⁺	[E1]		
		926.2 5	19 4	2998.54	3 ⁻	E2		B(E2)(W.u.)>4.1
		1187.4 10	16 1	2736.57	4 ⁺	[E1]		B(E1)(W.u.)>1.8×10 ⁻⁵
		1618.4 5	100 1	2306.72	4 ⁺	E1+M2	+0.12 4	B(E1)(W.u.)>4.6×10 ⁻⁵ ; B(M2)(W.u.)>0.53 Mult., δ : from 1978We15 in ⁵¹ V(¹⁶ O,p2n γ). RUL(M2)=1 for B(M2)(W.u.) suggests T _{1/2} (3924.7 level) not lower than \approx 0.7 ps.
3932.0	(4,5)	935 ^a		2998.54	3 ⁻			
		1625.3 4	100	2306.72	4 ⁺			
3951.9	(4 ⁺ ,3 ⁺)	2960.4 6	100	991.54	2 ⁺			
3993.36	6 ⁺	1686.60 6	100	2306.72	4 ⁺	E2		B(E2)(W.u.)=23 +8-5
4020.4	(2 ⁺)	1283.4 7	100 20	2736.57	4 ⁺			γ reported in (p, γ) E=2.05 MeV.
		3029.0 5	100 20	991.54	2 ⁺			
4039.7	(0 ⁺ to 4 ⁺)	3048.1 4	100	991.54	2 ⁺			
4076.55	(5 ⁺)	999.7 ^a 6		3077.77	4 ⁺			I γ : <39 from 2004Ka18 in (²⁸ Si,4p γ). Others: 33 5 (1994Cr05), 77 19 (1998Ga11), 127 21 (1978Ne02) in (α ,n γ),(HI,xn γ) where a 999.7 3 γ was assigned from only the 4077 level. But 1980Si02 in (α ,n γ),(¹¹ B,2np γ) do not support the placement of this γ from 4077 level, based on the absence of (1000 γ)(771 γ) coincidences. B(M1)(W.u.)=0.015 +8-5; B(E2)(W.u.)=3.4 +22-16 E γ : unweighted average of all available values. I γ : 26 4 (1994Cr05,2004Ka18), 100 19 (1998Ga11). But this γ was not detected by 1980Si02 in (α ,n γ) and (¹¹ B,2np γ), the authors gave an upper limit of 0.1 for branching ratio. Placement or existence of 1773.2 10 γ in 2004Ka18 is considered uncertain. If M1, B(M1)(W.u.)=0.0014 4. If E2, B(E2)(W.u.)=0.8 3.
		1340.2 4	100.0 16	2736.57	4 ⁺	M1+E2	-0.49 11	
		1771.5 ^a 2		2306.72	4 ⁺			
4140	(2,1) ⁺	\approx 4140		0.0	0 ⁺			
4156.53	5 ⁻	851		3306.85	(4 ⁺)			
		1079.2 4	43 11	3077.77	4 ⁺	(E1)		B(E1)(W.u.)=0.00070 +30-20
		1159.0 ^a 4	46 10	2998.54	3 ⁻	[E2]		I γ : unweighted average of two available intensities. B(E2)(W.u.)=39 +16-11 E γ : reported in (α ,n γ),(¹¹ B,2np) (1980Si02) only, considered as uncertain (evaluators). B(E1)(W.u.)=0.00032 +13-7 E γ : unweighted average of available values.
		1850.4 8	100 10	2306.72	4 ⁺	[E1]		E γ ,I γ : from (γ , γ'). If E1, B(E1)(W.u.)=0.00041 12. If M1, B(M1)(W.u.)=0.021 6.
4159.5	1	3168	85	991.54	2 ⁺			E γ ,I γ : from (γ , γ').
		4159	100	0.0	0 ⁺			γ reported in (α ,n γ),(¹¹ B,2np γ).
4181.7		1875.0 5		2306.72	4 ⁺			

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)							Comments
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	
4181.7		2381 ^a 2		1799.41	2 ⁺		γ reported in (n,n' γ).
4205.2	(4,3) ⁺	3213.6 4	100	991.54	2 ⁺		
4236.71	6 ⁺	1500.1 3	100.0 22	2736.57	4 ⁺	E2	B(E2)(W.u.)=26 +12-6
		1935.3 ^a 7	45 3	2306.72	4 ⁺	(E2)	B(E2)(W.u.)=3.3 +15-8 E γ : γ from ($^{28}\text{Si},4\text{p}\gamma$) study of 2004Ka18 only. It is treated as uncertain due to poor fit in level scheme (level-energy difference gives 1930.0) and non-observation in other γ -ray studies.
4288.6	(4) ⁺	1552.0 4	100	2736.57	4 ⁺		
4319.1	(4,3) ⁺	3327 3		991.54	2 ⁺		E γ : from (n,n' γ).
4454.68	1 ⁺	2544.4 2	20 4	1910.26	0 ⁺	[M1]	B(M1)(W.u.)=0.052 +16-12
		2655.2 2	34 6	1799.41	2 ⁺	[M1+E2]	B(M1)(W.u.)=0.078 +22-17; B(E2)(W.u.)=19 +5-4 B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2.
		3462.4 10	5.8 12	991.54	2 ⁺	[M1+E2]	B(M1)(W.u.)=0.0060 +19-15; B(E2)(W.u.)=0.85 +27-21 B(M1)(W.u.) for pure M1; B(E2)(W.u.) for pure E2.
		4454.3 10	100 6	0.0	0 ⁺	[M1]	B(M1)(W.u.)=0.049 +12-8 B(M1)(W.u.)=0.049 +12-8
4560		3564 ^a 3		991.54	2 ⁺		
4608.75	(1)	3617.1 2	100 18	991.54	2 ⁺		
		4609 2	\approx 32	0.0	0 ⁺		
4634.87	7 ⁻	398.17 6	20.7 9	4236.71	6 ⁺	E1	B(E1)(W.u.)=1.23 \times 10 ⁻⁵ 9
		641.48 5	100.0 6	3993.36	6 ⁺	E1	B(E1)(W.u.)=1.42 \times 10 ⁻⁵ 9
4664	(1)	4664		0.0	0 ⁺		If E1, B(E1)(W.u.)=0.00010 3. If M1, B(M1)(W.u.)=0.0053 13.
4668.93	(6 ⁻)	512.2 2	19 4	4156.53	5 ⁻	D	I γ : from ($^{28}\text{Si},4\text{p}\gamma$) E=115 MeV. Other: 102 15 in ($^{54}\text{Fe},2\text{p}\gamma$).
		592.4 1	67 11	4076.55	(5) ⁺	D(+Q)	I γ : unweighted average of two values.
		744.8 6	100 8	3924.69	5 ⁻	D(+Q)	E γ : unweighted average of available values.
4713.15	(1)	2103.6 2	100 23	2609.52	0 ⁺		
		2913 ^a 2	6	1799.41	2 ⁺		
		4712 2	\approx 11	0.0	0 ⁺		
4823.5	(5,6,7)	746.9 5		4076.55	(5) ⁺		
4980.87	7 ⁻	743.5 10	7.0 9	4236.71	6 ⁺	[E1]	B(E1)(W.u.)=4.3 \times 10 ⁻⁵ +21-11
		824.7 2	21.0 11	4156.53	5 ⁻	E2	B(E2)(W.u.)=12 +6-3 I γ : other: 100 10 in ($^{54}\text{Fe},2\text{p}\gamma$).
		1056.1 1	100.0 17	3924.69	5 ⁻	E2	B(E2)(W.u.)=17 +8-4
5066.8		2760 ^a 2		2306.72	4 ⁺		
5151.71	(7 ⁻)	516.8 1	57 7	4634.87	7 ⁻		I γ : from ($^{54}\text{Fe},2\text{p}\gamma$).
		1227.3 2	100 14	3924.69	5 ⁻	Q	
5623.75	(8 ⁻)	954.8 1	100 3	4668.93	(6 ⁻)	Q	
		990 1		4634.87	7 ⁻		
5699.38	(8 ⁻)	547.8 2	7 2	5151.71	(7 ⁻)	D	I γ : from ($^{54}\text{Fe},2\text{p}\gamma$). Other: 46 2 in ($^{28}\text{Si},4\text{p}\gamma$) E=122 MeV.
		1030.4 2	88 11	4668.93	(6 ⁻)	Q	I γ : unweighted average of two values.
		1064.0 4	100 5	4634.87	7 ⁻		
5936.0	(8 ⁺)	1698.5 10	33 7	4236.71	6 ⁺		
		1942.8 10	100 6	3993.36	6 ⁺	(Q)	

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	Comments
5951.7	(9 ⁻)	328		5623.75 (8 ⁻)			
		971.1 5	≈12	4980.87 7 ⁻		Q	
		1313.50 [#] 16	100	4638.2			E_γ : from 1996GaZZ. Other: 1315 (2004Ka18).
6031.5	(8 ⁺)	1395.6 10	22 3	4634.87 7 ⁻			I_γ : from (²⁸ Si,4p γ) E=115 MeV. Other: 74 3 in (²⁸ Si,4p γ) E=122 MeV.
		1794.5 17	67 6	4236.71 6 ⁺			I_γ : from (⁵⁴ Fe,2p γ). Other: 112 6 in (²⁸ Si,4p γ) E=115 MeV.
		2038.9 5	100 6	3993.36 6 ⁺		Q	
6124.0	(8 ⁺)	502		5623.75 (8 ⁻)			
		1488.5 10	47 3	4634.87 7 ⁻			
		1887.0 4	19 4	4236.71 6 ⁺		Q	I_γ : from (⁵⁴ Fe,2p γ). Other: 51 3 in (²⁸ Si,4p γ) E=115 MeV.
		2130.6 6	100 4	3993.36 6 ⁺		Q	
6124.7	(9 ⁻)	1143.8 4	100	4980.87 7 ⁻		Q	Mult.: $\Delta J=2$ γ from $\gamma(\theta)$ in (α ,n γ), (Hl,xn γ) (1978Ne02) and DCO ratio (1994Cr05) in (⁵⁴ Fe,2p γ). 2004Ka18, in (p, γ) E=122 MeV, propose $\Delta J=1$ from $\gamma\gamma(\theta)$ (DCO) and γ (anisotropy) ratio, but their results seem consistent with $\Delta J=2$ also.
6262.1	(7,8,9 ⁻)	1627.2 6	100	4634.87 7 ⁻			
6377.0	(7,8,9 ⁻)	1741		4634.87 7 ⁻			
6963.0	(9)	838.3 3	100 11	6124.7 (9 ⁻)			
		1263.4 5	36 7	5699.38 (8 ⁻)		D	Mult.: from DCO ratio.
6998.1	(11 ⁻)	1046.45 [#] 10	100	5951.7 (9 ⁻)		E2	B(E2)(W.u.)=31 +9-6
7062.4	(10 ⁻)	1363.0 4	100	5699.38 (8 ⁻)		Q	
7118.9	(10 ⁺)	993.0 ^{&} 10	134 ^{&} 6	6124.0 (8 ⁺)			I_γ : from (²⁸ Si,4p γ) E=115 MeV. $I_\gamma<25$ for 997.6 γ in (⁵⁴ Fe,2p γ).
		993.0 ^{&} 10	101 ^{&} 5	6124.7 (9 ⁻)			I_γ : from (²⁸ Si,4p γ) E=115 MeV. $I_\gamma<25$ for 997.6 γ in (⁵⁴ Fe,2p γ).
		1088.0 5	100 5	6031.5 (8 ⁺)		Q	E_γ : unweighted average of two values.
		1166		5951.7 (9 ⁻)			
		1181.8 13	29 1	5936.0 (8 ⁺)		Q	E_γ : unweighted average of two values.
							I_γ : from (⁵⁴ Fe,2p γ). Other: 55 3 in (²⁸ Si,4p γ) E=115 MeV.
7212.4	(11 ⁻)	1260.3 7	100	5951.7 (9 ⁻)		Q	
7334.7	(10 ⁺)	1210.7 3	100	6124.0 (8 ⁺)		Q	
7556.2	(10 ⁻)	1605		5951.7 (9 ⁻)			
7579.1	(11 ⁻)	1454.3 2	100	6124.7 (9 ⁻)		Q	
7806.0	(10 ⁺)	1869.9 7	100	5936.0 (8 ⁺)		Q	
7902	(9,10,11 ⁻)	1776.9		6124.7 (9 ⁻)			
7946.5	(10 ⁺)	1915 2	100	6031.5 (8 ⁺)		Q	
8157.1	(10 ⁺)	2221 2	100	5936.0 (8 ⁺)		(Q)	
8181.1	(10 ⁻)	2229.8		5951.7 (9 ⁻)			
8302.8	(12 ⁻)	1304.9 5	100	6998.1 (11 ⁻)			
8322.1	(11)	1204		7118.9 (10 ⁺)			
8426.1	(11 ⁻)	1307.15 9	100 3	7118.9 (10 ⁺)			
		1429.1 10	16.7 18	6998.1 (11 ⁻)			E_γ : γ not reported in (²⁸ Si,4p γ) E=122 MeV.
		2048		6377.0 (7,8,9 ⁻)			
8580.4	(12 ⁺)	154.30 5	2.5 3	8426.1 (11 ⁻)			
		259		8322.1 (11)			

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	Comments
8580.4	(12 ⁺)	1462.5 10	100 4	7118.9	(10 ⁺)	Q	E _γ : unweighted average of available values.
		1583.3 [#] 10	87 3	6998.1	(11 ⁻)		
8995.4	(12 ⁺)	1189.4 3	100	7806.0	(10 ⁺)	Q	
9363.7	(11 ⁻)	1808		7556.2	(10 ⁻)		
		3414		5951.7	(9 ⁻)		
9440.3	(11 ⁻)	2321.9 9	100	7118.9	(10 ⁺)		
9666	(14)	1086		8580.4	(12 ⁺)		
9772	(2 ⁺)	5873	7.4 16	3898.5	(2 ⁺ ,3,4 ⁺)		
		5892	10 4	3880	(0 ⁺ to 4 ⁺)		
		5975	11 3	3795.03	1 ⁺		
		6172	11 4	3601.9	(1,2 ⁺)		
		6224	17.4 23	3552.3	4 ⁺		
		6313	11 4	3458.66	(2,3)		
		6344	12.3 23	3425.13	1 ⁺		
		6407	11 3	3365.99	1 ⁺		
		6469	16 4	3297.17	(2) ⁺		
		6514	8.0 28	3261.94	1		
		6569	19.6 22	3205.98	(3) ⁺		
		6585	12.4 13	3186.84	1 ⁺		
		6681	21 6	3094.64	(3) ⁺		
		6768	20.7 14	3005.73	2 ⁺		
		6795	20.9 26	2979.94	3 ⁺		
		6977	17.1 14	2793.5	2 ⁺		
		7040	9.7 26	2736.57	4 ⁺		
		7162	7.9 29	2609.52	0 ⁺		
		7464	16 6	2306.72	4 ⁺		
		7861	10.0 7	1910.26	0 ⁺		
		7972	93 29	1799.41	2 ⁺		
		8782	100 9	991.54	2 ⁺		
		9772	36 7	0.0	0 ⁺		
9803.5	(11 ⁻)	1622.5 8	100 62	8181.1	(10 ⁻)		
		2743		7062.4	(10 ⁻)		
9948.4	(12 ⁻)	508.1 5		9440.3	(11 ⁻)		
		584.8 5	86 5	9363.7	(11 ⁻)	D	
		2886 1	25 2	7062.4	(10 ⁻)	Q	
		2950 1	100 5	6998.1	(11 ⁻)	D	
10.31×10 ³		6110	6.9 12	4205.2	(4,3) ⁺		
		6390	3.4 11	3924.69	5 ⁻		
		6410	8.6 14	3898.5	(2 ⁺ ,3,4 ⁺)		
		6490	12.2 16	3819.65	(0 ⁺ to 4 ⁺)		
		6510	9.5 19	3795.03	1 ⁺		
		6680	2.3 9	3630?	(0 ⁺ ,6 ⁻)		
		6710	11.2 17	3597.24	(2 ⁺ ,3,4 ⁺)		

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)						
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]
10.31×10^3		6890	12.4 19	3425.13	1 ⁺	
		7220	22 4	3094.64	(3) ⁺	
		7230	14 4	3077.77	4 ⁺	
		7330	25 4	2979.94	3 ⁺	
		7520	25 3	2793.5	2 ⁺	
		7570	20 3	2736.57	4 ⁺	
		7700	6.9 12	2609.52	0 ⁺	
		8000	26 3	2306.72	4 ⁺	
		8400	11.0 17	1910.26	0 ⁺	
		8510	60 7	1799.41	2 ⁺	
		9320	100 12	991.54	2 ⁺	
		10310	45 6	0.0	0 ⁺	
		10460.2	512.0 5	9948.4	(12 ⁻)	
		656.7 5	50.0 17	9803.5	(11 ⁻)	Q
10872	(3 ⁻)	1020.0 5	100 3	9440.3	(11 ⁻)	
		2158.3 9	72 3	8302.8	(12 ⁻)	
		3247 1	48.3 17	7212.4	(11 ⁻)	Q
		3461 1	92 3	6998.1	(11 ⁻)	Q
		6568	26	4304.1	(1 ⁻ to 5 ⁻)	
		6719	16	4156.53	5 ⁻	
		7113	11	3759	(1 ⁻ to 5 ⁻)	
		7192	16	3680	(1 ⁻ to 5 ⁻)	
		7458	21	3414	(1 ⁻ to 5 ⁻)	
		7513	11	3365.99	1 ⁺	
		7588 ^a		3285	(1 ⁻ to 5 ⁻)	
		7674	26	3196.9	(2,3)	
		7799	32	3077.77	4 ⁺	
		7871	74	2998.54	3 ⁻	
10906	(3 ⁻)	7891	26	2979.94	3 ⁺	
		8080	47	2793.5	2 ⁺	
		8134	42	2736.57	4 ⁺	
		8568	26	2306.72	4 ⁺	
		9073	47	1799.41	2 ⁺	
		9881	100	991.54	2 ⁺	
		10872 ^a	5.3	0.0	0 ⁺	
		6222	7.7	4684	(1 ⁻ to 5 ⁻)	
		6541	12	4362.1	(2,1,3) ⁺	
		6601	15	4304.1	(1 ⁻ to 5 ⁻)	
		6752	19	4156.53	5 ⁻	
		7051	12	3853.27	5 ⁺	
		7061	12	3850.5	(≤ 3) ⁽⁺⁾	
		7621	12	3285	(1 ⁻ to 5 ⁻)	
		7707	15	3196.9	(2,3)	

E_γ : 7506 from level-energy difference.

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	Comments
10906	(3 ⁻)	7832	19	3077.77	4 ⁺		
		7904	58	2998.54	3 ⁻		
		7924	12	2979.94	3 ⁺		
		8113	19	2793.5	2 ⁺		
		8167	12	2736.57	4 ⁺		
		8601	15	2306.72	4 ⁺		
		9106	38	1799.41	2 ⁺		
		9914	100	991.54	2 ⁺		
		10906 ^a	3.9	0.0	0 ⁺		
11023.4	(14 ⁻)	563.3 3	100.0 5	10460.2	(13 ⁻)	D	
		1074.7 5	24.3 6	9948.4	(12 ⁻)		
		2720 1	1.68 15	8302.8	(12 ⁻)		
11120	(2 ⁺)	6560	≥6.1	4560			
		6650	≥5.3	4470	(0 ⁺)		
		6700	≥3.4	4420	(4,3) ⁺		
		6740	≥2.6	4380			
		6750	≥5.3	4370	3 ⁻		
		6760	≥2.5	4362.1	(2,1,3) ⁺		
		6800	3.4 2	4319.1	(4,3) ⁺		
		6810	≥1.9	4310			
		6860	≥6.7	4260			
		6910	9.9 3	4205.2	(4,3) ⁺		
		6960	6 4	4156.53	5 ⁻		
		6980	≥2.3	4140	(2,1) ⁺		
		7010	≥1.1	4110	(2) ⁺		
		7040	1.5 3	4076.55	(5) ⁺		
		7080	11.5 24	4039.7	(0 ⁺ to 4 ⁺)		
		7100	4 4	4020.4	(2) ⁺		
		7170	1.29 14	3951.9	(4 ⁺ ,3 ⁺)		
		7220	10.6 10	3898.5	(2 ⁺ ,3,4 ⁺)		
		7268	≥2.5	3853.27	5 ⁺		
		7270	9 8	3850.5	(≤3) ⁽⁺⁾		
		7300	11.0 24	3819.65	(0 ⁺ to 4 ⁺)		
		7320	6.2 24	3795.03	1 ⁺		
		7400	≥6.7	3718.4	(0 ⁺ to 4 ⁺)		
		7420	9.5 4	3710.0	(2 ⁺)		
		7490	3.2 13	3628.4	(4) ⁺		E _γ : 7410 from level-energy difference.
		7520	7 5	3597.24	(2 ⁺ ,3,4 ⁺)		
		7570	7 5	3552.3	4 ⁺		
		7575	8 6	3545.9?	(0 to 3 ⁺)		
		7660	13 4	3458.66	(2,3)		
		7670	≥1.8	3452.0	(1,2 ⁺)		
		7690	8.6 24	3425.13	1 ⁺		

Adopted Levels, Gammas (continued)

 $\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	Comments
11120	(2 ⁺)	7750	29 11	3369.86	3 ⁺		
		7755	16 3	3365.99	1 ⁺		
		7820	37 3	3297.17	(2) ⁺		
		7860	15.3 24	3261.94	1		
		7910	32 7	3205.98	(3) ⁺		
		7920	3.5 3	3196.9	(2,3)		
		7930	13.4 19	3186.84	1 ⁺		
		8020	25 11	3094.64	(3) ⁺		
		8040	20 8	3077.77	4 ⁺		
		8110	25 3	3005.73	2 ⁺		
		8120	35 16	2998.54	3 ⁻		
		8140	36.4 24	2979.94	3 ⁺		
		8330	37.0 6	2793.5	2 ⁺		
		8380	19.6 15	2736.57	4 ⁺		
		8510	6.5 13	2609.52	0 ⁺		
		8810	36.8 15	2306.72	4 ⁺		
		9210	27 4	1910.26	0 ⁺		
		9320	57 9	1799.41	2 ⁺		
		10130	100 4	991.54	2 ⁺		
		11120	48 3	0.0	0 ⁺		
11464	(15)	1792 ^a		9666	(14)		
11626.4	(15 ⁻)	603.0 3	100.0 4	11023.4	(14 ⁻)	D	
		1166 1	51.0 4	10460.2	(13 ⁻)	Q	
12335.7	(16 ⁻)	709.5 3	100 6	11626.4	(15 ⁻)		
		1312.5 9	88 6	11023.4	(14 ⁻)		
12468	(16)	1003		11464	(15)		
13082.1	(17 ⁻)	746.4 3	100.0 5	12335.7	(16 ⁻)	D	
		1455.2 5	100 6	11626.4	(15 ⁻)		
13324	(17)	856		12468	(16)		
		1860		11464	(15)		
13948.1	(18 ⁻)	865.8 5	45.0 5	13082.1	(17 ⁻)	D	
		1613.3 6	100.0 19	12335.7	(16 ⁻)	Q	
14391	(18)	1067		13324	(17)		
		1924		12468	(16)		
14862.5	(19 ⁻)	914.5 5	97.1 10	13948.1	(18 ⁻)	D	
		1779.6 6	100.0 10	13082.1	(17 ⁻)	Q	
15423.6	(19)	561		14862.5	(19 ⁻)		
		1032		14391	(18)		
		2099		13324	(17)		
15945.0	(20 ⁻)	1082.1 5	45.8 10	14862.5	(19 ⁻)		
		1997.4 6	100.0 13	13948.1	(18 ⁻)	Q	
16686.8	(20)	742		15945.0	(20 ⁻)		
		1263		15423.6	(19)		

I_γ: other: 122 10 in (²⁸Si,4pγ) E=115 MeV.

Adopted Levels, Gammas (continued)

$\gamma(^{64}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	Comments
16686.8	(20)	2296		14391	(18)		
17087.2	(21 ⁻)	1142.0 5	19 4	15945.0	(20 ⁻)		I_γ : other: 68 16 in (²⁸ Si,4p γ) E=115 MeV.
		2225.1 10	100.0 18	14862.5	(19 ⁻)	Q	
17853	(21)	769		17087.2	(21 ⁻)		
		1167		16686.8	(20)		
		2429		15423.6	(19)		
18483.3	(22 ⁻)	1395.6 9	28.4 14	17087.2	(21 ⁻)		
		2538.6 10	100.0 20	15945.0	(20 ⁻)	Q	
19365	(22)	2678		16686.8	(20)		
19775.6	(23 ⁻)	1292		18483.3	(22 ⁻)		
		2688.5 10	100 5	17087.2	(21 ⁻)	Q	
20657	(23)	2804		17853	(21)		
21297.5	(24 ⁻)	1523		19775.6	(23 ⁻)		
		2814 1	100 4	18483.3	(22 ⁻)		
22892.7	(25 ⁻)	3117 1	100	19775.6	(23 ⁻)		
24868.6	(26 ⁻)	3571 1		21297.5	(24 ⁻)		

[†] When a level is populated in two or more datasets, averages of values are taken where uncertainties are given. Additional comments are also provided for certain individual γ rays.

[‡] From $\gamma(\theta)$ and/or $\gamma(\text{lin pol})$ in ($\alpha, n\gamma$), (¹¹B, 2n γ). (1980Si02, 1978Si02). RUL (for E2 and M2) also used when $T_{1/2}$ known. Mult=Q (most likely E2) from $\Delta J=2$, quadrupole transition; mult=D or D+Q for $\Delta J=1$ (or in rare cases $\Delta J=0$), dipole or dipole+quadrupole transition. For some of the transitions, especially, for high-spin ($J \geq 4$) levels, multipolarities are also established in ⁵¹V(¹⁶O, p2n γ), ⁵⁹Co(⁷Li, 2n γ) from $\gamma(\theta)$, $\gamma(\text{pol})$ and level lifetimes.

Ordering of 1314-1046-1583 cascade above the 4636, 7⁻ level is from (²⁸Si, 4p γ) (2004Ka18 and 1997Fu08). Others: 1046-1314-1583 cascade in 1998Ga11, 1047-1582 cascade in 1994Cr05 with no 1314 γ reported, only the 1046 γ (placed above 4636, 7⁻ level) reported in ($\alpha, n\gamma$), (¹¹B, 2n γ) (1980Si02, 1978Ne02).

@ Total theoretical internal conversion coefficients, calculated using the BrIcc code (2008Ki07) with Frozen orbital approximation based on γ -ray energies, assigned multipolarities, and mixing ratios, unless otherwise specified.

& Multiply placed with intensity suitably divided.


^a Placement of transition in the level scheme is uncertain.

Adopted Levels, Gammas

Legend


Level Scheme

Intensities: Relative photon branching from each level

-----► γ Decay (Uncertain)

Adopted Levels, GammasLevel Scheme (continued)

Intensities: Relative photon branching from each level




Adopted Levels, Gammas

Legend


Level Scheme (continued)

Intensities: Relative photon branching from each level

-----► γ Decay (Uncertain)


Adopted Levels, GammasLevel Scheme (continued)

Intensities: Relative photon branching from each level




Adopted Levels, GammasLevel Scheme (continued)

Intensities: Relative photon branching from each level



Adopted Levels, Gammas**Level Scheme (continued)**

Intensities: Relative photon branching from each level
 @ Multiplied: intensity suitably divided




Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided

-----► γ Decay (Uncertain)


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level
@ Multiply placed: intensity suitably divided

-----► γ Decay (Uncertain)




Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level
@ Multiply placed: intensity suitably divided

-----► γ Decay (Uncertain)




Adopted Levels, Gammas

Legend


Level Scheme (continued)

Intensities: Relative photon branching from each level
@ Multiply placed: intensity suitably divided

-----▶ γ Decay (Uncertain)



⁶⁴Zn₃₄

Adopted Levels, Gammas

Adopted Levels, Gammas (continued)

