

## Math 110, Fall 2021, Test 1B Sample Answers

Note: Students were required to show their steps in solving the problems. For brevity these solutions omit intermediate steps of row reductions.

### Instructions:

- You may use a calculator on this test, but the only permitted calculators are SHARP brand calculators with model numbers beginning EL-510R. No other electronic devices are permitted.
- No notes, textbooks, or other outside materials or aids are permitted.
- For questions with numerical answers, either give your answer in exact form or give it as a decimal to two decimal places.
- For **all** questions you must show your work to be given credit, even if your answer is correct.
- For questions 1–3, show your work and then enter your final answer in the box provided.
- This test is printed double-sided - be sure not to miss the questions on the back of the first page! For the long-answer questions the backs of the pages are additional space for your solution.

[Do not write on this page]

[Do not write on this page]

(1 point) 1. Let  $\vec{v} = \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}$  and  $\vec{w} = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$ . Calculate  $2\vec{v} + \vec{w} - 2(\vec{w} + \vec{v})$ .

**Solution:**

$$2\vec{v} + \vec{w} - 2(\vec{w} + \vec{v}) = 2 \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} + \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} - 2 \left( \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} \right) = \begin{bmatrix} -2 \\ 0 \\ -1 \end{bmatrix}$$

Answer:

$$\begin{bmatrix} -2 \\ 0 \\ -1 \end{bmatrix}$$

(1 point) 2. Let  $\vec{v} = \begin{bmatrix} 3 \\ 1 \\ 1 \\ 0 \end{bmatrix}$  and  $\vec{w} = \begin{bmatrix} 1 \\ -2 \\ 2 \\ 0 \end{bmatrix}$ . Let  $\theta$  be the angle between  $\vec{v}$  and  $\vec{w}$ . Find  $\cos(\theta)$ .

**Solution:**

$$\cos(\theta) = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} = \frac{3}{\sqrt{11}\sqrt{9}} = \frac{1}{\sqrt{11}}$$

(The last step of simplification was not required).

Answer:

$$\frac{1}{\sqrt{11}}$$

- (1 point) 3. Find all values of  $x$  such that  $\begin{bmatrix} 0 \\ x \\ x+1 \end{bmatrix}$  has length 1.

**Solution:** We have

$$\left\| \begin{bmatrix} 0 \\ x \\ x+1 \end{bmatrix} \right\| = \sqrt{0^2 + x^2 + (x+1)^2} = \sqrt{2x^2 + 2x + 1}.$$


To give this vector length 1 we therefore need  $2x^2 + 2x + 1 = 1$ , so  $2x(x+1) = 0$ . That is, the possible values of  $x$  are  $x = 0$  and  $x = -1$ .

Answer:

$$x = 0 \text{ and } x = -1$$

- (1 point) 4. Sketch the line in  $\mathbb{R}^2$  that has vector equation  $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} -2 \\ 1 \end{bmatrix}$ . Very briefly (one sentence is enough) tell us how you decided where to draw the line.

**Solution:** By choosing  $t = 0$  and  $t = 1$  we find that the line passes through  $(0, 1)$  and  $(-2, 2)$ , so we draw the only line through those two points.



- (4 points) 5. Determine whether the following system of linear equations in variables  $x_1, x_2, x_3, x_4$  has no solution, exactly one solution, or infinitely many solutions.

$$x_1 - 3x_2 + 2x_3 = 1$$

$$-2x_1 + 5x_2 + x_4 = 2$$

$$x_2 + x_3 + x_4 = 0$$

**Solution:** We set up an augmented matrix and row reduce.

$$\left[ \begin{array}{cccc|c} 1 & -3 & 2 & 0 & 1 \\ -2 & 5 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cccc|c} 1 & 0 & 0 & 1 & -3 \\ 0 & 1 & 0 & 3/5 & -4/5 \\ 0 & 0 & 1 & 2/5 & 4/5 \end{array} \right].$$

We see from the reduced row echelon form that  $x_4$  is a free variable, and therefore there are infinitely many solutions.

(4 points) 6. Find all values of  $k$  for which  $\begin{bmatrix} 1 \\ 2 \\ k \end{bmatrix}$  is a linear combination of  $\begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}$  and  $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$ .

**Solution:** We want to know for which  $k$  there are  $a$  and  $b$  such that

$$a \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix} + b \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ k \end{bmatrix}.$$

We treat this vector equation as a system of linear equations in variables  $a$  and  $b$ , and solve:

$$\left[ \begin{array}{cc|c} -1 & 1 & 1 \\ 3 & -1 & 2 \\ 2 & 1 & k \end{array} \right] \rightarrow \left[ \begin{array}{cc|c} 1 & 0 & 3/2 \\ 0 & 1 & 5/2 \\ 0 & 0 & 2k - 11 \end{array} \right].$$

This system has a solution if and only if  $2k - 11 = 0$ , that is, if and only if  $k = 11/2$ .

Therefore the only  $k$  for which  $\begin{bmatrix} 1 \\ 2 \\ k \end{bmatrix}$  is a linear combination of  $\begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}$  and  $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$  is  $k = 11/2$ .

- (4 points) 7. Let  $L$  be the line in  $\mathbb{R}^3$  that passes through the points  $(2, 1, 0)$  and  $(0, 1, 1)$ . Let  $P$  be the plane in  $\mathbb{R}^3$  that is orthogonal to  $L$  and passes through the point  $(1, 2, 1)$ . Find, with justification, a vector equation for  $P$ .

**Solution:** Since  $P$  is orthogonal to  $L$ , a direction vector for  $L$  will be a normal vector to  $P$ . Such a vector is  $\vec{n} = \begin{bmatrix} 2-0 \\ 1-1 \\ 0-1 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix}$ . We know that  $P$  passes through  $(1, 2, 1)$ , so in normal form the equation for  $P$  is

$$\begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}.$$

Expanding the dot products we obtain the general form

$$2x - z = 1.$$

We rearrange this equation to say

$$z = -1 + 2x,$$

and then by substituting we obtain

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x \\ y \\ -1 + 2x \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix} + x \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} + y \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

The equation above is a vector equation for  $P$  (it is not the only possible correct answer).

(4 points) 8. Suppose that  $\vec{v}$  and  $\vec{w}$  are vectors in  $\mathbb{R}^n$ , and that  $c$  is a scalar. Show that

$$\text{proj}_{\vec{w}}(c\vec{v}) = c \text{proj}_{\vec{w}}(\vec{v}).$$

*Note:* In this question we want you to write a general argument, so you should not choose specific numbers for any of the objects in the question.

**Solution:** We use properties of the dot product to calculate:

$$\begin{aligned}\text{proj}_{\vec{w}}(c\vec{v}) &= \left( \frac{\vec{w} \cdot (c\vec{v})}{\vec{w} \cdot \vec{w}} \right) \vec{w} \\ &= \left( \frac{c(\vec{w} \cdot \vec{v})}{\vec{w} \cdot \vec{w}} \right) \vec{w} \\ &= c \left( \frac{\vec{w} \cdot \vec{v}}{\vec{w} \cdot \vec{w}} \right) \vec{w} \\ &= c \text{proj}_{\vec{w}}(\vec{v})\end{aligned}$$