

System Failure Example: Tacoma Narrows Bridge

- The (original) Tacoma Narrows Bridge was a suspension bridge linking Tacoma and Gig Harbor (WA, USA).
- This mile-long bridge, with a 2,800-foot main span, was the third largest suspension bridge at the time of opening.
- Construction began in Nov. 1938 and took about 19 months to build at a cost of \$6,400,000.
- On July 1, 1940, the bridge opened to traffic.
- On Nov. 7, 1940 at approximately 11:00, the bridge collapsed during a moderate (42 miles/hour) wind storm.
- The bridge was supposed to withstand winds of up to 120 miles/hour.
- The collapse was due to wind-induced vibrations and an *unstable mechanical system*.
- Repair of the bridge was not possible.
- Fortunately, a dog trapped in an abandoned car was the only fatality.

Image of bridge collapse omitted for copyright reasons.

A video of the bridge collapse can be found at

<https://youtu.be/j-zczJXSxw>.

Part 2

Preliminaries

Section 2.1

Functions, Sequences, System Operators, and Transforms

- A **rational number** is a number of the form x/y , where x and y are integers and $y \neq 0$ (i.e., a ratio of integers).
- For example, $-\frac{5}{3}$, $\frac{17}{11}$, and $0 = \frac{0}{1}$ are rational numbers, whereas π and e are irrational numbers (i.e., not rational).
- The symbols employed to denote several commonly-used sets are as follows:

Symbol	Set
\mathbb{Z}	integers
\mathbb{R}	real numbers
\mathbb{C}	complex numbers
\mathbb{Q}	rational numbers

- For two integers a and b , we define the following notation for sets of consecutive integers:

$$[a..b] = \{x \in \mathbb{Z} : a \leq x \leq b\},$$

$$[a..b) = \{x \in \mathbb{Z} : a \leq x < b\},$$

$$(a..b] = \{x \in \mathbb{Z} : a < x \leq b\}, \quad \text{and}$$

$$(a..b) = \{x \in \mathbb{Z} : a < x < b\}.$$

- In this notation, a and b indicate the endpoints of the range for the set, and the type of brackets used (i.e., parenthesis versus square bracket) indicate whether each endpoint is included in the set.
- For example:
 - $[0..4]$ denotes the set of integers $\{0, 1, 2, 3, 4\}$;
 - $[0..4)$ denotes the set of integers $\{0, 1, 2, 3\}$; and
 - $[0..N-1]$ and $[0..N)$ both denote the set of integers $\{0, 1, 2, \dots, N-1\}$.

Notation for Intervals on the Real Line

- For two real numbers a and b , we define the following notation for intervals on the real line:

$$[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\},$$

$$(a, b) = \{x \in \mathbb{R} : a < x < b\},$$


$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\}, \quad \text{and}$$

$$(a, b] = \{x \in \mathbb{R} : a < x \leq b\}.$$

- In this notation, a and b indicate the endpoints of the interval for the set, and the type of brackets used (i.e., parenthesis versus square bracket) indicate whether each endpoint is included in the set.
- For example:
 - $[0, 100]$ denotes the set of all real numbers from 0 to 100, including both 0 and 100;
 - $(-\pi, \pi]$ denotes the set of all real numbers from $-\pi$ to π , excluding $-\pi$ but including π ; and
 - $[-\pi, \pi)$ denotes the set of all real numbers from $-\pi$ to π , including $-\pi$ but excluding π .

Mappings

- A **mapping** is a relationship involving two sets that associates each element in one set, called the **domain**, with an element from the other set, called the **codomain**.
- The notation $f : A \rightarrow B$ denotes a mapping f whose domain is the set A and whose codomain is the set B .
- Example:



$$f : A \rightarrow B$$
$$A = \{1, 2, 3, 4\}$$
$$B = \{0, 1, 2, 3\}$$
$$f(x) = \begin{cases} 0 & x \in \{1, 2\} \\ 1 & x = 4 \\ 2 & x = 3. \end{cases}$$

- Although many types of mappings exist, the types of most relevance to our study of signals and systems are: functions, sequences, system operators, and transforms.

Functions

- A **function** is a mapping where the domain is a set that is *continuous* in nature, such as the real numbers or complex numbers.
- In practice, the codomain is typically either the real numbers or complex numbers.
- Functions are also commonly referred to as **continuous-time (CT) signals**.
- Example:
 - Let $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(t) = t^2$ (i.e., f is the squaring function).
 - The function f maps each real number t to the real number $f(t) = t^2$.
 - The domain and codomain are the real numbers.
 - Note that f is a *function*, whereas $f(t)$ is a *number* (namely, the value of the function f evaluated at t).
- Herein, we will focus almost exclusively on functions of a single independent variable (i.e., one-dimensional functions).

Sequences

- A **sequence** is a mapping where the domain is a set that is **discrete** in nature, such as the integers, or a subset thereof.
- In practice, the codomain is typically either the real numbers or complex numbers.
- Sequences are also commonly referred to as **discrete-time (DT) signals**.
- Example:
 - Let $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ such that $f(n) = n^2$, where \mathbb{Z}^+ denotes the set of (strictly) positive integers (i.e., f is the sequence of perfect squares).
 - The sequence f maps each (strictly) positive integer n to the (strictly) positive integer $f(n) = n^2$.
 - The domain and codomain are \mathbb{Z}^+ (i.e., the positive integers).
 - Note that f is a **sequence**, whereas $f(n)$ is a **number** (namely, the value of the sequence f evaluated at n).
- As a matter of notation, the n th element of a sequence x is denoted as either $x(n)$ or x_n .
- Herein, we will focus almost exclusively on sequences with a single independent variable (i.e., one-dimensional sequences).

Remarks on Notation for Functions and Sequences

SKIP SLIDE

- For a real-valued function f of a real variable and an arbitrary real number t , the expression f denotes the function f itself and the expression $f(t)$ denotes the value of the function f evaluated at t .
- That is, f is a *function* and $f(t)$ is a *number*.
- Unfortunately, the practice of using $f(t)$ to denote the function f is quite common, although strictly speaking this is an abuse of notation.
- In contexts where imprecise notation may lead to problems, one should be careful to clearly distinguish between a function and its value.
- For the real-valued functions f and g of a real variable and an arbitrary real number t :
 - The expression $f + g$ denotes a *function*, namely, the function formed by adding the functions f and g .
 - The expression $f(t) + g(t)$ denotes a *number*, namely, the sum of: 1) the value of the function f evaluated at t ; and 2) the value of the function g evaluated at t .
- Similar comments as the ones made above for functions also hold in the case of sequences.

- To express that two functions f and g are equal, we can write either:
 - 1 $f = g$; or
 - 2 $f(t) = g(t)$ for all t .
- Of the preceding two expressions, the first (i.e., $f = g$) is usually preferable, as it is less verbose.
- For the functions f and g and an operation \circ that is defined pointwise for functions (such as addition, subtraction, multiplication, and division), the following relationship holds:

$$(f \circ g)(t) = f(t) \circ g(t).$$

- Some operations \circ involving functions (such as convolution, to be discussed later) cannot be defined in a pointwise manner, in which case $(f \circ g)(t)$ is a valid mathematical expression, while $f(t) \circ g(t)$ is not.
- Again, similar comments as the ones made above for functions also hold in the case of sequences.

System Operators

- A **system operator** is a mapping used to represent a system.
- We will focus exclusively on the case of single-input single-output systems.
- A (single-input single-output) **system operator** maps a function or sequence representing the input of a system to a function or sequence representing the output of the system.
- The domain and codomain of a system operator are sets of *functions or sequences*, not sets of numbers.
- Example:
 - Let $\mathcal{H} : F \rightarrow F$ such that $\mathcal{H}x(t) = 2x(t)$ (for all $t \in \mathbb{R}$) and F is the set of functions mapping \mathbb{R} to \mathbb{R} .
 - The system \mathcal{H} maps a function to a function.
 - In particular, the domain and codomain are each F , which is a set of functions.
 - The system \mathcal{H} multiplies its input function x by a factor of 2 in order to produce its output function $\mathcal{H}x$.
 - Note that $\mathcal{H}x$ is a function, not a number.

Remarks on Operator Notation for CT Systems

- For a system operator \mathcal{H} and a function x , $\mathcal{H}x$ is the function produced as the output of the system \mathcal{H} when the input is the function x .
- Brackets around the operand of an operator are *often omitted when not required* for grouping.
- For example, for an operator \mathcal{H} , a function x , and a real number t , we would normally prefer to write:
 - 1 $\mathcal{H}x$ instead of the equivalent expression $\mathcal{H}(x)$; and
 - 2 $\mathcal{H}x(t)$ instead of the equivalent expression $\mathcal{H}(x)(t)$.
- Also, note that $\mathcal{H}x$ is a *function* and $\mathcal{H}x(t)$ is a *number* (namely, the value of the function $\mathcal{H}x$ evaluated at t).
- In the expression $\mathcal{H}(x_1 + x_2)$, the brackets are needed for grouping, since $\mathcal{H}(x_1 + x_2) \neq \mathcal{H}x_1 + x_2$ (where “ \neq ” means “not equivalent”).
- When multiple operators are applied, they group from *right to left*.
- For example, for the operators \mathcal{H}_1 and \mathcal{H}_2 , and the function x , the expression $\mathcal{H}_2\mathcal{H}_1x$ means $\mathcal{H}_2[\mathcal{H}_1(x)]$.

Remarks on Operator Notation for DT Systems

SKIP SLIDE

- For a system operator \mathcal{H} and a sequence x , $\mathcal{H}x$ is the sequence produced as the output of the system \mathcal{H} when the input is the sequence x .
- Brackets around the operand of an operator are *often omitted when not required* for grouping.
- For example, for an operator \mathcal{H} , a sequence x , and an integer n , we would normally prefer to write:
 - 1 $\mathcal{H}x$ instead of the equivalent expression $\mathcal{H}(x)$; and
 - 2 $\mathcal{H}x(n)$ instead of the equivalent expression $\mathcal{H}(x)(n)$.
- Also, note that $\mathcal{H}x$ is a *sequence* and $\mathcal{H}x(n)$ is a *number* (namely, the value of the sequence $\mathcal{H}x$ evaluated at n).
- In the expression $\mathcal{H}(x_1 + x_2)$, the brackets are needed for grouping, since $\mathcal{H}(x_1 + x_2) \neq \mathcal{H}x_1 + x_2$ (where “ \neq ” means “not equivalent”).
- When multiple operators are applied, they group from *right to left*.
- For example, for the operators \mathcal{H}_1 and \mathcal{H}_2 , and the sequence x , the expression $\mathcal{H}_2\mathcal{H}_1x$ means $\mathcal{H}_2[\mathcal{H}_1(x)]$.

- Later, we will be introduced to several types of mappings known as transforms.
- Transforms have a mathematical structure similar to system operators.
- That is, transforms map functions/sequences to functions/sequences.
- Due to this similar structure, many of the earlier comments about system operators also apply to the case of transforms.
- For example, the Fourier transform (introduced later) is denoted as \mathcal{F} and the result of applying the Fourier transform operator to the function/sequence x is denoted as $\mathcal{F}x$.
- Some examples of transforms of interest in the study of signals and systems are listed on the next slide.

Examples of Transforms

Name	Domain	Codomain
CT Fourier Series	T -periodic functions (with domain \mathbb{R})	sequences (with domain \mathbb{Z})
CT Fourier Transform	functions (with domain \mathbb{R})	functions (with domain \mathbb{R})
Laplace Transform	functions (with domain \mathbb{R})	functions (with domain \mathbb{C})
DT Fourier Series	N -periodic sequences (with domain \mathbb{Z})	N -periodic sequences (with domain \mathbb{Z})
DT Fourier Transform	sequences (with domain \mathbb{Z})	2π -periodic functions (with domain \mathbb{R})
Z Transform	sequences (with domain \mathbb{Z})	functions (with domain \mathbb{C})

Section 2.2

Properties of Signals

Even Symmetry



- A function x is said to be **even** if it satisfies

$$x(t) = x(-t) \quad \text{for all } t \text{ (where } t \text{ is a real number).}$$

- A sequence x is said to be **even** if it satisfies

$$x(n) = x(-n) \quad \text{for all } n \text{ (where } n \text{ is an integer).}$$

- Geometrically, the graph of an even signal is **symmetric** with respect to the vertical axis.
- Some examples of even signals are shown below.



Odd Symmetry



- A function x is said to be **odd** if it satisfies

$$x(t) = -x(-t) \quad \text{for all } t \text{ (where } t \text{ is a real number).}$$

- A sequence x is said to be **odd** if it satisfies

$$x(n) = -x(-n) \quad \text{for all } n \text{ (where } n \text{ is an integer).}$$

- An odd signal x must be such that $x(0) = 0$.
- Geometrically, the graph of an odd signal is **symmetric** with respect to the origin.
- Some examples of odd signals are shown below.



Conjugate Symmetry

- A function x is said to be **conjugate symmetric** if it satisfies

$$x(t) = x^*(-t) \quad \text{for all } t \text{ (where } t \text{ is a real number).}$$

- A sequence x is said to be **conjugate symmetric** if it satisfies

$$x(n) = x^*(-n) \quad \text{for all } n \text{ (where } n \text{ is an integer).}$$

- The real part of a conjugate symmetric function or sequence is even.
- The imaginary part of a conjugate symmetric function or sequence is odd.
- An example of a conjugate symmetric function is a complex sinusoid $x(t) = \cos \omega t + j \sin \omega t$, where ω is a real constant.

Periodicity



- A function x is said to be **periodic** with **period** T (or **T -periodic**) if, for some strictly-positive real constant T , the following condition holds:

$$x(t) = x(t + T) \quad \text{for all } t \text{ (where } t \text{ is a real number).}$$

- A sequence x is said to be **periodic** with **period** N (or **N -periodic**) if, for some strictly-positive integer constant N , the following condition holds:

$$x(n) = x(n + N) \quad \text{for all } n \text{ (where } n \text{ is an integer).}$$


- Some examples of periodic signals are shown below.



- A function/sequence that is not periodic is said to be **aperiodic**.
- A T -periodic function x is said to have **frequency** $\frac{1}{T}$ and **angular frequency** $\frac{2\pi}{T}$.
- An N -periodic sequence x is said to have **frequency** $\frac{1}{N}$ and **angular frequency** $\frac{2\pi}{N}$.

Periodicity (Continued 2)

- The period of a periodic signal is *not unique*. That is, a signal that is periodic with period T is also periodic with period kT , for every (strictly) positive integer k .



- The smallest period with which a signal is periodic is called the **fundamental period** and its corresponding frequency is called the **fundamental frequency**.

Part 3

Continuous-Time (CT) Signals and Systems