

---

ECE 260 A4

5.1 For each case below, find the Fourier series representation (in complex exponential form) of the function  $x$ , explicitly identifying the fundamental period of  $x$  and the Fourier series coefficient sequence  $c$ .

(a)  $x(t) = 1 + \cos(\pi t) + \sin^2(\pi t)$ ; ✓

(b)  $x(t) = \cos(4t) \sin(t)$ ; and

(c)  $x(t) = |\sin(2\pi t)|$ . [Hint:  $\int e^{ax} \sin(bx) dx = \frac{e^{ax} [a \sin(bx) - b \cos(bx)]}{a^2 + b^2} + C$ , where  $a$  and  $b$  are arbitrary complex and nonzero real constants, respectively.] ✓

a

$$x(t) = 1 + \cos \pi t + \sin^2 \pi t$$

$$= 1 + \frac{1}{2} [e^{j\pi t} + e^{-j\pi t}] + \left( \frac{1}{2j} [e^{j\pi t} - e^{-j\pi t}] \right)^2$$

$$= 1 + \frac{1}{2} e^{j\pi t} + \frac{1}{2} e^{-j\pi t} - \frac{1}{4} e^{j(2\pi t)} + \frac{1}{4} - \frac{1}{4} e^{-j(2\pi t)}$$

$$= -\frac{1}{4} e^{-j2\pi t} + \frac{1}{2} e^{-j\pi t} + \frac{3}{2} + \frac{1}{2} e^{j\pi t} - \frac{1}{4} e^{j2\pi t}$$

$$x(t) = \sum_{k=-\infty}^{\infty} c_k e^{jk\omega_0 t} \quad \begin{cases} \omega_0 = \pi \\ c_k = \begin{cases} \frac{3}{2} & \text{for } k=0 \\ \frac{1}{2} & \text{for } k=\pm 1 \\ -\frac{1}{4} & \text{for } k=\pm 2 \\ 0 & \text{otherwise} \end{cases} \end{cases}$$

c

$x(t)$  is periodic w/ period  $T = \frac{1}{2}$  and frequency  $\omega_0 = \frac{2\pi}{T} = \frac{2\pi}{1/2} = 4\pi$

$$\begin{aligned} c_k &= \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt = 2 \int_0^{1/2} e^{-j4\pi k t} \sin \pi t dt \\ &= \frac{2(2\pi)}{16\pi^2 k^2 + 4\pi^2} \left[ e^{-4j\pi k t} [-j2k \sin 2\pi t - \cos 2\pi t] \right]_0^{1/2} \end{aligned}$$

$$= \frac{1}{\pi(1-4k^2)} \left[ e^{(-j4\pi k/2)} \left[ -j2k \sin \frac{2\pi}{2} - \cos \frac{2\pi}{2} \right] - (-\cos 0) \right]$$


$$= \frac{1}{\pi(1-4k^2)} (2) = \frac{2}{\pi(1-16k^2)}$$


---

$$x(t) = \sum_{k=-\infty}^{\infty} c_k e^{jk\omega_0 t}$$

$\omega_0 = 4\pi$   
 $c_k = \frac{2}{\pi(1-16k^2)}$

5.2 For each of the periodic functions shown in the figures below, find the corresponding Fourier series coefficient sequence.



(c)

c  $x(t)$  is periodic w/  $T=5$  & frequency  $\omega_0 = \frac{2\pi}{5}$

$$\begin{aligned}
 c_k &= \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt = \frac{1}{5} \int_{-5/2}^{5/2} x(t) e^{-j2\pi kt/5} dt \\
 &= \frac{1}{5} \left[ \int_{-2}^{-1} e^{-j2\pi kt/5} dt + \int_{-1}^1 2e^{-j2\pi kt/5} dt + \int_1^2 e^{-j2\pi kt/5} dt \right] \\
 &= \frac{1}{5} \left[ \int_{-2}^2 e^{-j2\pi kt/5} dt + \int_{-1}^1 e^{-j2\pi kt/5} dt \right] \\
 &= \frac{1}{-j2\pi k} \left[ e^{-j4\pi k/5} - e^{j4\pi k/5} \right]_2^{-2} + \left[ e^{-j4\pi k/5} - e^{j4\pi k/5} \right]_{-1}^1 \\
 &= \frac{1}{-j2\pi k} \left[ e^{-j4\pi k/5} - e^{j4\pi k/5} + e^{-j2\pi k/5} - e^{j2\pi k/5} \right]
 \end{aligned}$$

$$= \frac{1}{\pi k} \left( -2j \sin 4\pi k/5 - 2j \sin 2\pi k/5 \right) = \frac{\sin 4\pi k}{5\pi k} + \frac{\sin 2\pi k}{5\pi k}$$

$$= \frac{4}{5} \sin \frac{4\pi k}{5} + \frac{2}{5} \sin \frac{2\pi k}{5}$$

$$C_0 = \frac{1}{T} \int_T u(t) dt = \frac{1}{5} \int_{-5/2}^{5/2} u(t) dt = \frac{1}{5} \left[ \int_{-2}^{-1} 1 dt + \int_{-1}^1 2 dt + \int_1^2 1 dt \right]$$

$$= \frac{1}{5} (1 + 4 + 1) = 6/5$$

$$C_k = \begin{cases} 6/5 & \dots k=0 \\ 4/5 \sin\left(\frac{4\pi k}{5}\right) + 2/5 \sin\left(\frac{2\pi k}{5}\right) & \dots \text{others} \end{cases}$$

$$C_0 = 1.2$$

$$C_1 = C_{-1} \approx 0.489828 \quad \text{and} \quad C_2 = C_{-2} \approx -0.57846$$

5.7 A periodic function  $x$  with period  $T$  and Fourier series coefficient sequence  $c$  is said to be odd harmonic if  $c_k = 0$  for all even  $k$ .

(a) Show that if  $x$  is odd harmonic, then  $x(t) = -x(t - \frac{T}{2})$  for all  $t$ .

(b) Show that if  $x(t) = -x(t - \frac{T}{2})$  for all  $t$ , then  $x$  is odd harmonic. ✓

$$c_k = \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt = \frac{1}{T} \left[ \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt + \int_{T/2}^T x(t) e^{-jk\omega_0 t} dt \right]$$

$$= \frac{1}{T} \left[ \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt + \int_T^{3T/2} x(\lambda - \frac{T}{2}) e^{-jk\omega_0 (\lambda - \frac{T}{2})} d\lambda \right]$$


$$= \frac{1}{T} \left[ \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt + \int_T^{3T/2} -x(\lambda) e^{-jk\omega_0 (\lambda - \frac{T}{2})} d\lambda \right]$$

$$= \frac{1}{T} \left[ \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt - e^{jk\omega_0 \frac{T}{2}} \int_T^{3T/2} x(\lambda) e^{-jk\omega_0 \lambda} d\lambda \right]$$

$$= \frac{1}{T} \left[ \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt - (-1)^k \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt \right]$$

$$= \frac{1}{T} \left[ 1 - (-1)^k \right] \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt = \begin{cases} \frac{2}{T} \int_0^{T/2} x(t) e^{-jk\omega_0 t} dt & \text{for odd } k \\ 0 & \text{for even } k \end{cases}$$

5.9 Find the Fourier series coefficient sequence  $c$  of the periodic function  $x$  shown in the figure below. Plot the frequency spectrum of  $x$ , including the first five harmonics.



$x(t)$  is periodic w/ period  $T=2$  and frequency  $\omega_0 = \frac{2\pi}{T} = \frac{2\pi}{2} = \pi$

$$\begin{aligned} c_k &= \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt = \frac{1}{2} \int_0^2 x(t) e^{-j\pi k t} dt \\ &= \frac{1}{2} \left[ \frac{1}{-j\pi k} e^{-j\pi k t} \right] \Big|_0^1 \\ &= \frac{1}{j2\pi k} \left( 1 - (-1)^k \right) \\ &= \begin{cases} \left( -\frac{j}{\pi k} \right) & \text{for odd } k \\ 0 & \text{for even } k \end{cases} \end{aligned}$$



---


$$c_0 = \frac{1}{T} \int_T x(t) dt = \frac{1}{2} \int_0^2 x(t) dt$$

$$= \frac{1}{2} \int_0^1 dt = \frac{1}{2} (t) \Big|_0^1 = \frac{1}{2}$$

Therefore,  $c_k = \begin{cases} \frac{1}{2} & \dots k=0 \\ -j/\pi k & \dots \text{odd } k \\ 0 & \dots \text{even } k, k \neq 0 \end{cases}$

<u>k</u>	<u><math> c_k </math></u>	<u><math>\arg c_k</math></u>
0	$\frac{1}{2}$	0
1	$\frac{1}{\pi}$	$-\pi/2$
2	0	0
3	$\frac{3}{\pi}$	$-\pi/2$
4	0	0
5	$\frac{1}{5\pi}$	$-\pi/2$





5.10 Consider a LTI system with frequency response

$$H(\omega) = \begin{cases} 1 & |\omega| \geq 5 \\ 0 & \text{otherwise.} \end{cases}$$

Using frequency-domain methods, find the output  $y$  of the system if the input  $x$  is given by

$$x(t) = 1 + 2\cos(2t) + 2\cos(4t) + \frac{1}{2}\cos(6t).$$

$$\begin{aligned} x(t) &= 1 + 2\cos(2t) + 2\cos(4t) + \frac{1}{2}\cos(6t) = 1 + 2\left[\frac{1}{2}e^{j2t} + e^{-j2t}\right] \\ &\quad + 2\left[\frac{1}{2}(e^{j4t} + e^{-j4t})\right] + \frac{1}{2}\left(\frac{1}{2}(e^{j6t} + e^{-j6t})\right) \\ &= 1 + e^{j2t} + e^{-j2t} + e^{j4t} + e^{-j4t} + \frac{1}{4}e^{j6t} + \frac{1}{4}e^{-j6t} \end{aligned}$$

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t}$$

$\omega_0 = 2$

$a_k = \begin{cases} 1 & \dots k=0 \\ 1 & \dots k=\pm 1 \\ 1 & \dots k=\pm 2 \\ \frac{1}{4} & \dots k=\pm 3 \\ 0 & \dots \text{otherwise} \end{cases}$

$$y(t) = \sum_{k=-\infty}^{\infty} b_k e^{jk\omega_0 t}$$

$$\begin{aligned} \dots \quad b_0 &= a_0 H(j[0][2]) = 0 & \left\{ \begin{aligned} b_3 &= a_3 H(j[5][2]) = \frac{1}{4}(1) = \frac{1}{4} \\ b_{-3} &= a_{-3} H(j[-5][2]) \\ &= \frac{1}{4}(1) = \frac{1}{4} \end{aligned} \right. \\ b_1 &= a_1 H(j[1][2]) = 0 \\ b_{-1} &= a_{-1} H(j[-1][2]) = 0 \\ b_2 &= a_2 H(j[2][2]) = 0 \\ b_{-2} &= a_{-2} H(j[-2][2]) = 0 \end{aligned}$$

$$y(t) = \frac{1}{4} e^{-j6t} + \frac{1}{4} e^{j6t}$$

$$= \frac{1}{4} (e^{-j6t} + e^{j6t})$$

$$= \frac{1}{4} (2\cos 6t) = \frac{1}{2} \cos 6t$$

```

main.m x +
/Users/arfaz/Library/CloudStorage/OneDrive-UniversityofVictoria/0 UVIC/2 ENGR Y3/4 ECE 260 A01 T03/1 Assignments/0 Solutions/2 M

1 % Clear all variables and close all figures.
2 clear all;
3 close all;
4
5 % Define the time variable.
6 t = linspace(-1, 1, 1000);
7
8 % Values for n.
9 n_values = [1, 5, 10, 50, 100];
10
11 for n = n_values
12     % Symbolic expression for the square wave.
13     syms k w;
14     f = symsum(0.5 * mysinc(pi * k / 2) * exp(j * k * w * t), 'k', -n, n);
15     f = subs(f, w, 2 * pi);
16
17     % Plot the result.
18     figure;
19     plot(t, real(f), 'LineWidth', 1.5);
20     title(['x_{', num2str(n), '}(t)']);
21     xlabel('Time');
22     ylabel('Amplitude');
23     grid on;
24
25     % Save the plot to a file.
26     print(['data/sqrwav_', num2str(n)], '-depsc');
27
28 end
29
30 function y = mysinc(x)
31     y = ones(size(x));
32     i = find(x);
33     y(i) = sin(x(i)) ./ x(i);
34
35 end

```