

Lecture 31: Structure theorem for f.g. modules over PID

25 November 2020

19:04

Defⁿ: Let R be an int domain and M be an R -mod.

Then $\text{rank}(M) = \text{vdim}_{S^{-1}R}(S^{-1}M)$ where $S = R \setminus \{0\}$
 $= \dim_K(S^{-1}M)$ where $K = \text{frac}(R)$
 as vector space

① $\text{rank}(M) = \text{size of the largest l.i. subset of } M$.

Defⁿ: Let R be ring & M be an R -mod.

$$T(M) = \{m \in M : \exists r \in R, r \neq 0 \text{ and } rm = 0\} \leftrightarrow \{m \in M \mid \text{Ann}(m) \neq 0\}$$

$T(M)$ is a submod of M if R is an int domain.

M is called torsion free R -mod if $T(M) = 0$.

② Let R be an int domain M an R -mod then

$M/T(M)$ is torsion free.

such modules are called torsion modules

③ R an int dom M an R -mod s.t. $M = T(M)$ then $\text{rank}(M) = 0$.

Pf: Let $\alpha \in S^{-1}M$ then $\alpha = \frac{m}{s}$ for $m \in M$ & $s \in R \setminus \{0\}$

$$\exists r \in R \text{ s.t. } rm = 0 \text{ in } M \Rightarrow \alpha = \frac{m}{s} = \frac{rm}{rs} \quad (\because r \in R \setminus \{0\})$$

$$= \frac{0}{rs} = \frac{0}{1}$$

Hence $S^{-1}M = 0 \Rightarrow \text{rank}(M) = 0$

④ Even converse holds. Because $\text{rank}(M) = 0 \Rightarrow S^{-1}M = 0$

$$\Rightarrow \frac{m}{s} = \frac{0}{1} \text{ in } S^{-1}M$$

$$\Rightarrow \frac{m}{1} = \frac{0}{1} \text{ in } S^{-1}M$$

$$\Rightarrow \exists r \in S \text{ s.t. } r(m - 0) = 0 \Rightarrow rm = 0$$

$$\Rightarrow m \in T(M) \quad \forall m \in M.$$

Thm (Str thm): Let R be a PID and M be a f.g. R -mod. Then

$$M \cong R^k \oplus R/(a_1) \oplus R/(a_2) \oplus \dots \oplus R/(a_m)$$

where $k = \text{rank}(M)$ and $a_1, \dots, a_m \in R$ are nonzero nonunits s.t. $a_1 | a_2 | a_3 | \dots | a_m$. Here k and m could be 0.

Cor: R a PID and M a f.g. torsion free R -mod then M is free R -mod.

Pf: $M \cong R^k \oplus R/(a_1) \oplus \dots \oplus R/(a_m)$. But if $m \geq 1$

then $m \cdot (0, \bar{1}, 0, \dots, 0) \in \text{RHS}$ then $a_m = a(0, \bar{1}, 0, \dots, 0) = (0, 0, \dots, 0)$.

Contradicting M is torsion free.

Cor: Let G be a f.g. abelian group then

$$G \cong \mathbb{Z}^k \oplus \mathbb{Z}/a_1\mathbb{Z} \oplus \mathbb{Z}/a_2\mathbb{Z} \oplus \dots \oplus \mathbb{Z}/a_m\mathbb{Z} \text{ for}$$

some $n \geq 0$ and $a_1, \dots, a_m \in \mathbb{Z}$ are s.t.

$$a_i | a_{i+1} \quad 1 \leq i \leq m-1.$$

Pf: $R = \mathbb{Z}$ then G is f.g. \mathbb{Z} -mod and apply str thm.

Cor: R a PID and M a submod of a f.g. free mod
then M is free.

Pf: $M \subseteq R^n \Rightarrow M$ is also torsion free.
 R noeth $\Rightarrow R^n$ is a noeth R -mod $\Rightarrow M$ is noeth R -mod.
Hence M is f.g. R -mod. □

Example: 1) $R = \mathbb{Z}[X]$ or $\mathbb{Q}[X, Y]$.

$$M = (2, X) \quad \text{or} \quad (X, Y) = M$$

M is f.g. torsion free R -mod.

Is M free? Note $\text{rank}(M) = 1$
But M is not a principal ideal.

$2, X$ are not l.i. as

$$X \cdot 2 + (-2) \cdot X = 0 \text{ in } M.$$

2) $R \subset \text{PID}$. M torsion free $\stackrel{?}{\Rightarrow} M$ is free

Example: \mathbb{Q} as a \mathbb{Z} -module

Let $S \subseteq \mathbb{Q}$ s.t. $|S| \geq 2$
then S is lin dep.

$$\text{sp} \left(\frac{a}{s} \right) - \text{sp} \left(\frac{b}{s} \right) = 0 \quad \& \quad \mathbb{Q} \cong \mathbb{Z}.$$

Prop: Let R be a PID and F be a free R -module of rank n . Let N be a submodule of F . Then N is a free R -mod of rank $m \leq n$. Moreover there is a basis x_1, \dots, x_n of F and $\exists a_1, \dots, a_m \in R^*$ s.t. $a_1 | a_2 | \dots | a_m$ and $\{a_1 x_1, a_2 x_2, \dots, a_m x_m\}$ is a basis of N .

Prop \Rightarrow str thm: Let M be a f.g. R -mod.

Let $m_1, \dots, m_n \in M$ s.t. $M = Rm_1 + \dots + Rm_n$.

$$\Rightarrow \quad \begin{array}{ccc} \varphi: F & \longrightarrow & M \\ e_i & \longmapsto & m_i \end{array}$$

where F is a free mod of rank n and e_i 's are std. basis vectors.

then φ extends to R -lin surj map.

$$\varphi((b_1, \dots, b_n)) = \sum_{i=1}^n b_i m_i \quad \text{We know } \varphi \text{ is } R\text{-lin}$$

and since $\{m_i \mid 1 \leq i \leq n\}$ gen M , φ is surj.

$N = \ker(\varphi) \subseteq F$. By prop. 1, \exists

a basis $\{x_1, \dots, x_n\}$ of F and $a_1, \dots, a_m \in R^*$ s.t. $a_1 | a_2 | \dots | a_m$ and $\{a_1 x_1, a_2 x_2, \dots, a_m x_m\}$ is a basis of N .

$$N \subseteq F = Rx_1 \oplus Rx_2 \oplus \dots \oplus Rx_n$$

$$Ra_1x_1 \oplus Ra_2x_2 \oplus \dots \oplus Ra_mx_m$$

$$M \cong F/N \quad (\text{by 1st isom thm})$$

$$= \frac{Rx_1 \oplus Rx_2 \oplus \dots \oplus Rx_n}{Ra_1x_1 \oplus Ra_2x_2 \oplus \dots \oplus Ra_mx_m}$$

$$= \frac{Rx_1}{Ra_1x_1} \oplus \frac{Rx_2}{Ra_2x_2} \oplus \dots \oplus \frac{Rx_m}{Ra_mx_m} \oplus \underbrace{Rx_{m+1} \oplus \dots \oplus Rx_n}_{\text{red bracket}}$$

($\because M_1, M_2, \dots, M_n$ are R -mod &
 $N_i \subseteq M_i$ are R -submod then

$$M_1 \oplus M_2 \oplus \dots \oplus M_n / N_1 \oplus N_2 \oplus \dots \oplus N_n$$

$$\cong M_1/N_1 \oplus M_2/N_2 \oplus \dots \oplus M_n/N_n$$

$$\cong \frac{Rx_1}{Ra_1x_1} \oplus \frac{Rx_2}{Ra_2x_2} \oplus \dots \oplus \frac{Rx_m}{Ra_mx_m} \oplus R^{n-m}$$

$$\left(\begin{array}{l} Rx_1 \cong R \\ x_1 \mapsto 1 \\ x_1 \mapsto 1 \end{array} \right)$$

$$\alpha(Ra_1x_1) = (a_1)$$

$$a_1 | a_2 | \dots | a_m \quad a_i \in R^\times.$$

Let a_1, \dots, a_r be units $r \leq m$.

$$\text{then } M \cong R/(a_{r+1}) \oplus \dots \oplus R/(a_m) \oplus R^{n-m}$$

and we get the str from by replacing

m by $m-r$ and a_i by a_{r+i} .

Finally note that $\text{rank}(M) = \dim_{S^{-1}R} (\bar{S}^{-1}M)$

where $S = R \setminus \{0\}$

$$\Rightarrow \bar{S}^{-1}(\text{RHS}) = (\bar{S}^{-1}R)^{n-m}$$

$$\Rightarrow \text{rank}(M) = n-m = k.$$

