► Set theory 101

- > Set theory 101
- $ightharpoonup \mathbb{P}$ is a set-function.

- > Set theory 101
- ightharpoonup is a set-function.
- ightharpoonup All sets in $\mathcal{P}(\Omega)$ need not be measurable.

- ► Set theory 101
- $ightharpoonup \mathbb{P}$ is a set-function.
- ightharpoonup All sets in $\mathcal{P}(\Omega)$ need not be measurable.
- ightharpoonup We restrict domain of $\mathbb P$ to sigma-algebra of measurable sets.

- ► Set theory 101
- $ightharpoonup \mathbb{P}$ is a set-function.
- ightharpoonup All sets in $\mathcal{P}(\Omega)$ need not be measurable.
- ightharpoonup We restrict domain of $\mathbb P$ to sigma-algebra of measurable sets.
- $ightharpoonup (\Omega, \mathcal{F}, \mathbb{P})$ is known as probability space.

- ► Set theory 101
- $ightharpoonup \mathbb{P}$ is a set-function.
- ightharpoonup All sets in $\mathcal{P}(\Omega)$ need not be measurable.
- ightharpoonup We restrict domain of $\mathbb P$ to sigma-algebra of measurable sets.
- $ightharpoonup (\Omega, \mathcal{F}, \mathbb{P})$ is known as probability space.
- Formal definition of probability measure with its axioms.

- ► Set theory 101
- $ightharpoonup \mathbb{P}$ is a set-function.
- ightharpoonup All sets in $\mathcal{P}(\Omega)$ need not be measurable.
- ightharpoonup We restrict domain of $\mathbb P$ to sigma-algebra of measurable sets.
- $ightharpoonup (\Omega, \mathcal{F}, \mathbb{P})$ is known as probability space.
- Formal definition of probability measure with its axioms.
- ightharpoonup We looked at $\mathcal{B}([0,1])$ and $\mathcal{B}(\mathbb{R})$.

ightharpoonup Continuity of set-function \mathbb{P} .

ightharpoonup Continuity of set-function \mathbb{P} .

Conditional Probability.

- ightharpoonup Continuity of set-function \mathbb{P} .
- Conditional Probability.
- ► Law of Total Probability.

- ightharpoonup Continuity of set-function \mathbb{P} .
- Conditional Probability.
- ► Law of Total Probability.
- ► Independence and Correlation between sets.

- ightharpoonup Continuity of set-function \mathbb{P} .
- Conditional Probability.
- ► Law of Total Probability.
- ► Independence and Correlation between sets.
- Conditional independence.

► How do we define limit of a sequence $\{a_1, a_2, \ldots, \}$?

- ► How do we define limit of a sequence $\{a_1, a_2, \dots, \}$?
- Notation: $\lim_{n\to\infty} a_n = L$.

- ► How do we define limit of a sequence $\{a_1, a_2, \ldots, \}$?
- Notation: $\lim_{n\to\infty} a_n = L$.
- ► How do you define limit of a function at a point c?

- ► How do we define limit of a sequence $\{a_1, a_2, \ldots, \}$?
- Notation: $\lim_{n\to\infty} a_n = L$.
- ► How do you define limit of a function at a point c?
- Notation: $\lim_{x\to c} f(x) = L$

- ▶ How do we define limit of a sequence $\{a_1, a_2, \ldots, \}$?
- Notation: $\lim_{n\to\infty} a_n = L$.
- ► How do you define limit of a function at a point c?
- Notation: $\lim_{x\to c} f(x) = L$
- ▶ How do you define continuity of a function f(x) at c?

- ▶ How do we define limit of a sequence $\{a_1, a_2, \ldots, \}$?
- Notation: $\lim_{n\to\infty} a_n = L$.
- ► How do you define limit of a function at a point c?
- Notation: $\lim_{x\to c} f(x) = L$
- ▶ How do you define continuity of a function f(x) at c?
- ► When do you say a function is continuous?

- ▶ How do we define limit of a sequence $\{a_1, a_2, \ldots, \}$?
- Notation: $\lim_{n\to\infty} a_n = L$.
- ► How do you define limit of a function at a point c?
- Notation: $\lim_{x\to c} f(x) = L$
- ▶ How do you define continuity of a function f(x) at c?
- When do you say a function is continuous?
- $ightharpoonup (\epsilon, \delta)$ -definition of limits and continuity?

Definition in terms of limits of sequences.

For a continuous function $f(\cdot)$, as $x \to c$, we have $f(x) \to f(c)$

Definition in terms of limits of sequences.

For a continuous function $f(\cdot)$, as $x \to c$, we have $f(x) \to f(c)$

For a continuous set-function S, as $A_n \to A$, we have $S(A_n) \to S(A)$

Definition in terms of limits of sequences.

For a continuous function $f(\cdot)$, as $x \to c$, we have $f(x) \to f(c)$

For a continuous set-function S, as $A_n \to A$, we have $S(A_n) \to S(A)$

ightharpoonup Recall that \mathbb{P} is a set-function.

Definition in terms of limits of sequences.

For a continuous function $f(\cdot)$, as $x \to c$, we have $f(x) \to f(c)$

For a continuous set-function S, as $A_n \to A$, we have $S(A_n) \to S(A)$

 \triangleright Recall that \mathbb{P} is a set-function. Is it continuous?

Definition in terms of limits of sequences.

For a continuous function $f(\cdot)$, as $x \to c$, we have $f(x) \to f(c)$

For a continuous set-function S, as $A_n \to A$, we have $S(A_n) \to S(A)$

- \triangleright Recall that \mathbb{P} is a set-function. Is it continuous?
- We will see the proof shortly.

▶ Given $(Ω, \mathcal{F})$, If $A_1 \subset A_2 \ldots$ is an increasing sequence of events defined on \mathcal{F} and $\bigcup_{n=1}^{\infty} A_n = A \in \mathcal{F}$, then we say thats the sequence of sets A_n are increasing to A $(A_n \uparrow A)$.

- ▶ Given $(Ω, \mathcal{F})$, If $A_1 \subset A_2 \ldots$ is an increasing sequence of events defined on \mathcal{F} and $\bigcup_{n=1}^{\infty} A_n = A \in \mathcal{F}$, then we say thats the sequence of sets A_n are increasing to A $(A_n \uparrow A)$.
- Similarly when $A_1 \supset A_2 \dots$ is a decreasing sequence of events and $\bigcap_{n=1}^{\infty} A_n = A$, then we have $A_n \downarrow A$.

- ▶ Given $(Ω, \mathcal{F})$, If $A_1 \subset A_2 \ldots$ is an increasing sequence of events defined on \mathcal{F} and $\bigcup_{n=1}^{\infty} A_n = A \in \mathcal{F}$, then we say thats the sequence of sets A_n are increasing to A $(A_n \uparrow A)$.
- Similarly when $A_1 \supset A_2 \dots$ is a decreasing sequence of events and $\bigcap_{n=1}^{\infty} A_n = A$, then we have $A_n \downarrow A$.
- Alternative notation: For an increasing sequence of sets A_n we often write $\lim_{n\to\infty} A_n$ for $\bigcup_{n=1}^{\infty} A_n$ and for a decreasing sequence of sets A_n that $\lim_{n\to\infty} A_n = \bigcap_{n=1}^{\infty} A_n$.

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

Proof

Consider increasing sequence first. Similar arguments follow for decreasing seq.

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

- Consider increasing sequence first. Similar arguments follow for decreasing seq.

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

- Consider increasing sequence first. Similar arguments follow for decreasing seq.

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

- Consider increasing sequence first. Similar arguments follow for decreasing seq.
- ightharpoonup Define $F_n = A_n A_{n-1}$
- $\triangleright \cup_{n=1}^{\infty} A_n = \cup_{n=1}^{\infty} F_n.$

Continuity of set-function \mathbb{P}

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

Proof

- Consider increasing sequence first. Similar arguments follow for decreasing seq.
- $\triangleright \cup_{n=1}^{\infty} A_n = \cup_{n=1}^{\infty} F_n.$

Continuity of set-function \mathbb{P}

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

Proof

- Consider increasing sequence first. Similar arguments follow for decreasing seq.
- ightharpoonup Define $F_n = A_n A_{n-1}$
- $ightharpoonup \mathbb{P}(A) = \mathbb{P}(\bigcup_{n=1}^{\infty} A_n) = \mathbb{P}(\bigcup_{n=1}^{\infty} F_n)$

Continuity of set-function \mathbb{P}

Lemma

For sequence of events of the type $A_n \uparrow A$ or $A_n \downarrow A$, we have $\lim_{n\to\infty} \mathbb{P}(A_n) = \mathbb{P}(A)$.

Proof

- Consider increasing sequence first. Similar arguments follow for decreasing seq.
- ightharpoonup Define $F_n = A_n A_{n-1}$
- $\triangleright \cup_{n=1}^{\infty} A_n = \cup_{n=1}^{\infty} F_n.$

Equivalently if $An \to \emptyset$, then $\mathbb{P}(A_n) \to 0$.

► Given/If dice rolls odd, what is the probability that the outcome is 1?

- ► Given/If dice rolls odd, what is the probability that the outcome is 1?
- ▶ Given/If $\bar{\omega} \in [0, 0.5]$ what is the probability that $\bar{\omega} \in [0, 0.25]$?

- ► Given/If dice rolls odd, what is the probability that the outcome is 1?
- ▶ Given/If $\bar{\omega} \in [0, 0.5]$ what is the probability that $\bar{\omega} \in [0, 0.25]$?
- The conditional probability of event B given event A is defined as $\mathbb{P}(B/A) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)}$ when $\mathbb{P}(A) > 0$.

- ► Given/If dice rolls odd, what is the probability that the outcome is 1?
- ▶ Given/If $\bar{\omega} \in [0, 0.5]$ what is the probability that $\bar{\omega} \in [0, 0.25]$?
- The conditional probability of event B given event A is defined as $\mathbb{P}(B/A) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)}$ when $\mathbb{P}(A) > 0$.

- ► Given/If dice rolls odd, what is the probability that the outcome is 1?
- ▶ Given/If $\bar{\omega} \in [0, 0.5]$ what is the probability that $\bar{\omega} \in [0, 0.25]$?
- The conditional probability of event B given event A is defined as $\mathbb{P}(B/A) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)}$ when $\mathbb{P}(A) > 0$.

$$ightharpoonup \Omega_{/A} = A$$
,

- ► Given/If dice rolls odd, what is the probability that the outcome is 1?
- ▶ Given/If $\bar{\omega} \in [0, 0.5]$ what is the probability that $\bar{\omega} \in [0, 0.25]$?
- The conditional probability of event B given event A is defined as $\mathbb{P}(B/A) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)}$ when $\mathbb{P}(A) > 0$.

- $ightharpoonup \Omega_{/A} = A,$
- $ightharpoonup \mathcal{F}_{/A} = \{ C \cap A : C \in \mathcal{F} \}$ and

- ► Given/If dice rolls odd, what is the probability that the outcome is 1?
- ▶ Given/If $\bar{\omega} \in [0, 0.5]$ what is the probability that $\bar{\omega} \in [0, 0.25]$?
- The conditional probability of event B given event A is defined as $\mathbb{P}(B/A) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)}$ when $\mathbb{P}(A) > 0$.

- $ightharpoonup \Omega_{/A} = A$,
- $ightharpoonup \mathcal{F}_{/A} = \{\mathit{C} \cap \mathit{A} : \mathit{C} \in \mathcal{F}\}$ and

► Show that P(A/B)P(B) = P(B/A)P(A).

► Show that P(A/B)P(B) = P(B/A)P(A).

► Bayes rule: $P(B/A) = \frac{P(A/B)P(B)}{P(A)}$.

Show that P(A/B)P(B) = P(B/A)P(A).

- ► Bayes rule: $P(B/A) = \frac{P(A/B)P(B)}{P(A)}$.
- ▶ What is $P(A/(B \cap C))$?.

Show that P(A/B)P(B) = P(B/A)P(A).

- ► Bayes rule: $P(B/A) = \frac{P(A/B)P(B)}{P(A)}$.
- ▶ What is $P(A/(B \cap C))$? This is also denoted as P(A/BC)

- ► Show that P(A/B)P(B) = P(B/A)P(A).
- ► Bayes rule: $P(B/A) = \frac{P(A/B)P(B)}{P(A)}$.
- ▶ What is $P(A/(B \cap C))$?. This is also denoted as P(A/BC)
- Prove the chain rule $P(A \cap B \cap C) = P(A)P(B/A)P(C/(AB)).$

- ► Show that P(A/B)P(B) = P(B/A)P(A).
- ► Bayes rule: $P(B/A) = \frac{P(A/B)P(B)}{P(A)}$.
- ▶ What is $P(A/(B \cap C))$?. This is also denoted as P(A/BC)
- Prove the chain rule $P(A \cap B \cap C) = P(A)P(B/A)P(C/(AB)).$

HW: Prove the chain rule for conditional probability given by

$$P(A_1 \cap A_2 \dots A_n) = P(A_1)P(A_2/A_1)P(A_3/A_1A_2)\dots P(A_n/A_{n-1} \dots A_1).$$

➤ Suppose you draw 4 cards from a deck at random without replacement. What is the probability that (in order) these cards are 9 of club, 8 of diamond, king of spade and king of club?

- Suppose you draw 4 cards from a deck at random without replacement. What is the probability that (in order) these cards are 9 of club, 8 of diamond, king of spade and king of club?
- What if you do the above with replacement?

- Suppose you draw 4 cards from a deck at random without replacement. What is the probability that (in order) these cards are 9 of club, 8 of diamond, king of spade and king of club?
- What if you do the above with replacement?
- Consider a finite sample space Ω where each outcome is equally likely. Then what is P(B/A)?

- Suppose you draw 4 cards from a deck at random without replacement. What is the probability that (in order) these cards are 9 of club, 8 of diamond, king of spade and king of club?
- What if you do the above with replacement?
- Consider a finite sample space Ω where each outcome is equally likely. Then what is P(B/A)?
- $P(B/A) = \frac{|A \cap B|}{|A|}.$

 $ightharpoonup A = (A \cap B) \cup (A \cap B^c)$. What is P(A)?

 $ightharpoonup A = (A \cap B) \cup (A \cap B^c)$. What is P(A)?

$$P(A) = P(A \cap B) + P(A \cap B^c).$$

- $ightharpoonup A = (A \cap B) \cup (A \cap B^c)$. What is P(A)?
- $ightharpoonup P(A \cap B) + P(A \cap B^c).$
- ► This is same as $P(A) = P(A/B)P(B) + P(A/B^c)P(B^c)$.

- $ightharpoonup A = (A \cap B) \cup (A \cap B^c)$. What is P(A)?
- $P(A) = P(A \cap B) + P(A \cap B^c).$
- ► This is same as $P(A) = P(A/B)P(B) + P(A/B^c)P(B^c)$.
- This formula is useful when P(A) is not given or is difficult to find but P(B) or P(A/B) is readily available.

- $ightharpoonup A = (A \cap B) \cup (A \cap B^c)$. What is P(A)?
- $P(A) = P(A \cap B) + P(A \cap B^c).$
- ► This is same as $P(A) = P(A/B)P(B) + P(A/B^c)P(B^c)$.
- This formula is useful when P(A) is not given or is difficult to find but P(B) or P(A/B) is readily available.

Let $B_1, B_2, \dots B_n$ be the partition of the sample space Ω . Then for any event A we have

$$P(A) = \sum_{i=1}^{n} P(A \cap B_i) = \sum_{i=1}^{n} P(A/B_i)P(B_i).$$

- 1. If an item is defective, a robot can spot it with 98% accuracy.
- 2. If an item is not defective, a robot will declare it so with 99% accuracy.
- 3. A total of 0.1% items are defective.
- 4. If the robot says that the item you drew at random is defective, what is the probability that the robot is correct?

- 1. If an item is defective, a robot can spot it with 98% accuracy.
- 2. If an item is not defective, a robot will declare it so with 99% accuracy.
- 3. A total of 0.1% items are defective.
- 4. If the robot says that the item you drew at random is defective, what is the probability that the robot is correct?
- P(defective/robot says defective) =

- 1. If an item is defective, a robot can spot it with 98% accuracy.
- 2. If an item is not defective, a robot will declare it so with 99% accuracy.
- 3. A total of 0.1% items are defective.
- 4. If the robot says that the item you drew at random is defective, what is the probability that the robot is correct?
- P(defective/robot says defective) = P(robot says defective/defective)P(defective) P(robot says defective)

- 1. If an item is defective, a robot can spot it with 98% accuracy.
- 2. If an item is not defective, a robot will declare it so with 99% accuracy.
- 3. A total of 0.1% items are defective.
- 4. If the robot says that the item you drew at random is defective, what is the probability that the robot is correct?
- P(defective/robot says defective) = P(robot says defective/defective)P(defective) P(robot says defective)
- What is P(robot says defective)?