

Recap

Recap

- ▶ Continuity of set-function \mathbb{P} .

Recap

- ▶ Continuity of set-function \mathbb{P} .
- ▶ Conditional Probability.

Recap

- ▶ Continuity of set-function \mathbb{P} .
- ▶ Conditional Probability.
- ▶ Law of Total Probability.

This class

This class

- ▶ Independence and Correlation between sets.

This class

- ▶ Independence and Correlation between sets.
- ▶ Conditional independence.

This class

- ▶ Independence and Correlation between sets.
- ▶ Conditional independence.
- ▶ Principles of counting.

Bayes rule revisited

Let B_1, B_2, \dots, B_n be the partition of the sample space Ω .
Then for any event A with $P(A) > 0$ we have

$$P(B_j/A) = \frac{P(A/B_j)P(B_j)}{\sum_{i=1}^n P(A/B_i)P(B_i)}.$$

Independence

Independence

- ▶ Consider the experiment of tossing a coin and a dice simultaneously.

Independence

- ▶ Consider the experiment of tossing a coin and a dice simultaneously.
- ▶ Identify its underlying probability space.

Independence

- ▶ Consider the experiment of tossing a coin and a dice simultaneously.
- ▶ Identify its underlying probability space.
- ▶ What is $\mathbb{P}(\{H, 6\})$?

Independence

- ▶ Consider the experiment of tossing a coin and a dice simultaneously.
- ▶ Identify its underlying probability space.
- ▶ What is $\mathbb{P}(\{H, 6\})$?
- ▶ What is $\mathbb{P}(\{T, \text{odd}\}) = \mathbb{P}(\{\cup_{i=1,3,5} \{T, i\}\})$?

Independence

- ▶ Consider the experiment of tossing a coin and a dice simultaneously.
- ▶ Identify its underlying probability space.
- ▶ What is $\mathbb{P}(\{H, 6\})$?
- ▶ What is $\mathbb{P}(\{T, \text{odd}\}) = \mathbb{P}(\{\cup_{i=1,3,5} \{T, i\}\})$?
- ▶ In both cases above we have $\mathbb{P}(A \cap B) = P(A)P(B)$.

Independence

- ▶ Consider the experiment of tossing a coin and a dice simultaneously.
- ▶ Identify its underlying probability space.
- ▶ What is $\mathbb{P}(\{H, 6\})$?
- ▶ What is $\mathbb{P}(\{T, \text{odd}\}) = \mathbb{P}(\{\cup_{i=1,3,5} \{T, i\}\})$?
- ▶ In both cases above we have $\mathbb{P}(A \cap B) = P(A)P(B)$.
- ▶ This implies that $\mathbb{P}(A/B) = \mathbb{P}(A)$.

Independence

- ▶ Consider the experiment of tossing a coin and a dice simultaneously.
- ▶ Identify its underlying probability space.
- ▶ What is $\mathbb{P}(\{H, 6\})$?
- ▶ What is $\mathbb{P}(\{T, \text{odd}\}) = \mathbb{P}(\{\cup_{i=1,3,5} \{T, i\}\})$?
- ▶ In both cases above we have $\mathbb{P}(A \cap B) = P(A)P(B)$.
- ▶ This implies that $\mathbb{P}(A/B) = \mathbb{P}(A)$.

- ▶ Two events A, B are independent iff $P(A/B) = P(A)$ and $P(B/A) = P(B)$.
- ▶ Two events A, B are independent iff $P(A \cap B) = P(A)P(B)$.

Independence

- ▶ Two events A, B are independent if and only if $P(A \cap B) = P(A)P(B)$.

Independence

▶ Two events A, B are independent if and only if $P(A \cap B) = P(A)P(B)$.

▶ If A and B are independent, then so are A^c and B^c .

Independence

▶ Two events A, B are independent if and only if $P(A \cap B) = P(A)P(B)$.

- ▶ If A and B are independent, then so are A^c and B^c .
- ▶ What about A and B^c ? Are they independent?

Independence

▶ Two events A, B are independent if and only if $P(A \cap B) = P(A)P(B)$.

- ▶ If A and B are independent, then so are A^c and B^c .
- ▶ What about A and B^c ? Are they independent?
- ▶ If A_1, A_2, \dots, A_n are independent, then prove that

$$P(\cup_{i=1}^n A_i) = 1 - \prod_{i=1}^n [1 - P(A_i)]$$

Mutual and Pairwise Independence

Mutual and Pairwise Independence

- ▶ A collection of events $\{A_i, i \in I\}$ are said to be **mutually independent** if the $P(\cap_{j \in J} A_j) = \prod_{j \in J} P(A_j)$ for any subset J of I .

Mutual and Pairwise Independence

- ▶ A collection of events $\{A_i, i \in I\}$ are said to be **mutually independent** if the $P(\cap_{j \in J} A_j) = \prod_{j \in J} P(A_j)$ for any subset J of I .
- ▶ A collection of events $\{A_i, i \in I\}$ are said to be **pairwise independent** if any pair of events from the collection are independent.

Mutual and Pairwise Independence

- ▶ A collection of events $\{A_i, i \in I\}$ are said to be **mutually independent** if the $P(\cap_{j \in J} A_j) = \prod_{j \in J} P(A_j)$ for any subset J of I .
- ▶ A collection of events $\{A_i, i \in I\}$ are said to be **pairwise independent** if any pair of events from the collection are independent.
- ▶ Mutual independence implies pairwise independence but not the other way around.

Mutual and Pairwise Independence

- ▶ A collection of events $\{A_i, i \in I\}$ are said to be **mutually independent** if the $P(\cap_{j \in J} A_j) = \prod_{j \in J} P(A_j)$ for any subset J of I .
- ▶ A collection of events $\{A_i, i \in I\}$ are said to be **pairwise independent** if any pair of events from the collection are independent.
- ▶ Mutual independence implies pairwise independence but not the other way around.
- ▶ HW: Find an example where pairwise independence does not imply mutual independence.

Independence - Example

Independence - Example

- ▶ Pick a number randomly from the set $\{1, \dots, 10\}$.

Independence - Example

- ▶ Pick a number randomly from the set $\{1, \dots, 10\}$.
- ▶ Event A says that the number is less than 7.

Independence - Example

- ▶ Pick a number randomly from the set $\{1, \dots, 10\}$.
- ▶ Event A says that the number is less than 7.
- ▶ Event B says that the number is less than 8.

Independence - Example

- ▶ Pick a number randomly from the set $\{1, \dots, 10\}$.
- ▶ Event A says that the number is less than 7.
- ▶ Event B says that the number is less than 8.
- ▶ Event C says that the number is even.

Independence - Example

- ▶ Pick a number randomly from the set $\{1, \dots, 10\}$.
- ▶ Event A says that the number is less than 7.
- ▶ Event B says that the number is less than 8.
- ▶ Event C says that the number is even.
- ▶ Are the events mutually independent?

Independence - Example

- ▶ Pick a number randomly from the set $\{1, \dots, 10\}$.
- ▶ Event A says that the number is less than 7.
- ▶ Event B says that the number is less than 8.
- ▶ Event C says that the number is even.
- ▶ Are the events mutually independent?
- ▶ Which pair of event is independent?

Correlation between events

Correlation between events

- ▶ Two events A, B are independent iff $P(A \cap B) = P(A)P(B)$.

Correlation between events

- ▶ Two events A, B are independent iff $P(A \cap B) = P(A)P(B)$.
- ▶ Two events A and B are positively correlated iff $P(A/B) > P(A)$.

Correlation between events

- ▶ Two events A, B are independent iff $P(A \cap B) = P(A)P(B)$.
- ▶ Two events A and B are positively correlated iff $P(A/B) > P(A)$.
- ▶ Two events A and B are negatively correlated iff $P(A/B) < P(A)$.

Correlation between events

- ▶ Two events A, B are independent iff $P(A \cap B) = P(A)P(B)$.
- ▶ Two events A and B are positively correlated iff $P(A/B) > P(A)$.
- ▶ Two events A and B are negatively correlated iff $P(A/B) < P(A)$.
- ▶ A and B have the same correlation as A^c and B^c .

Correlation between events

- ▶ Two events A, B are independent iff $P(A \cap B) = P(A)P(B)$.
- ▶ Two events A and B are positively correlated iff $P(A/B) > P(A)$.
- ▶ Two events A and B are negatively correlated iff $P(A/B) < P(A)$.
- ▶ A and B have the same correlation as A^c and B^c .
- ▶ A and B have the opposite correlation as A and B^c .

Mutually exclusive and Independence

Mutually exclusive and Independence

- ▶ Two events A and B are mutually exclusive if occurrence of one implies that the other event cannot occur.

Mutually exclusive and Independence

- ▶ Two events A and B are mutually exclusive if occurrence of one implies that the other event cannot occur. Are they independent?

Mutually exclusive and Independence

- ▶ Two events A and B are mutually exclusive if occurrence of one implies that the other event cannot occur. Are they independent?
- ▶ If A and B are mutually exclusive, then they are not independent (and vice versa). This can be seen as follows.

Mutually exclusive and Independence

- ▶ Two events A and B are mutually exclusive if occurrence of one implies that the other event cannot occur. Are they independent?
- ▶ If A and B are mutually exclusive, then they are not independent (and vice versa). This can be seen as follows.

A and B are Mutually Exclusive

- ▶ $P(A \cap B) = 0$
- ▶ $P(A/B) = 0$
- ▶ $P(A/B^c) = \frac{P(A \cap B^c)}{P(B^c)} = \frac{P(A)}{P(B^c)}$

Mutually exclusive and Independence

- ▶ Two events A and B are mutually exclusive if occurrence of one implies that the other event cannot occur. Are they independent?
- ▶ If A and B are mutually exclusive, then they are not independent (and vice versa). This can be seen as follows.

A and B are Mutually Exclusive

- ▶ $P(A \cap B) = 0$
- ▶ $P(A/B) = 0$
- ▶ $P(A/B^c) = \frac{P(A \cap B^c)}{P(B^c)} = \frac{P(A)}{P(B^c)}$

A and B are Independent

- ▶ $P(A \cap B) = P(A)P(B)$
- ▶ $P(A/B) = P(A)$
- ▶ $P(A/B^c) = P(A)$

Mutually exclusive and Independence

- ▶ Two events A and B are mutually exclusive if occurrence of one implies that the other event cannot occur. Are they independent?
- ▶ If A and B are mutually exclusive, then they are not independent (and vice versa). This can be seen as follows.

A and B are Mutually Exclusive

- ▶ $P(A \cap B) = 0$
- ▶ $P(A/B) = 0$
- ▶ $P(A/B^c) = \frac{P(A \cap B^c)}{P(B^c)} = \frac{P(A)}{P(B^c)}$

A and B are Independent

- ▶ $P(A \cap B) = P(A)P(B)$
- ▶ $P(A/B) = P(A)$
- ▶ $P(A/B^c) = P(A)$

- ▶ If $A \subseteq B$, then two events are neither mutually exclusive nor independent.

Zero probability events and Independence

Zero probability events are always independent!

Zero probability events and Independence

Zero probability events are always independent!

- ▶ Let E be a zero probability event, i.e. $P(E) = 0$.

Zero probability events and Independence

Zero probability events are always independent!

- ▶ Let E be a zero probability event, i.e. $P(E) = 0$.
- ▶ Then for any set F , we want to show that $P(E \cap F) = 0$.

Zero probability events and Independence

Zero probability events are always independent!

- ▶ Let E be a zero probability event, i.e. $P(E) = 0$.
- ▶ Then for any set F , we want to show that $P(E \cap F) = 0$.
- ▶ Note that $E \cap F \subseteq E$.

Zero probability events and Independence

Zero probability events are always independent!

- ▶ Let E be a zero probability event, i.e. $P(E) = 0$.
- ▶ Then for any set F , we want to show that $P(E \cap F) = 0$.
- ▶ Note that $E \cap F \subseteq E$.
- ▶ This implies that $P(E \cap F) \leq P(E)$.

Conditional independence

Conditional independence

► Recall : $P(A/B) = \frac{P(AB)}{P(B)}$.

Conditional independence

► Recall : $P(A/B) = \frac{P(AB)}{P(B)}$.

► Also recall : $P(A/BC) = \frac{P(AB/C)}{P(B/C)}$

Conditional independence

- ▶ Recall : $P(A/B) = \frac{P(AB)}{P(B)}$.
- ▶ Also recall : $P(A/BC) = \frac{P(AB/C)}{P(B/C)}$
- ▶ This implies $P(AB/C) = P(A/BC)P(B/C)$.

Two events A and B are said to be conditionally independent of event C ($P(C) > 0$) if $P((AB)/C) = P(A/C).P(B/C)$

- ▶ As a consequence $P(A/BC) = P(A/C)$

Conditional independence

- ▶ Recall : $P(A/B) = \frac{P(AB)}{P(B)}$.
- ▶ Also recall : $P(A/BC) = \frac{P(AB/C)}{P(B/C)}$
- ▶ This implies $P(AB/C) = P(A/BC)P(B/C)$.

Two events A and B are said to be conditionally independent of event C ($P(C) > 0$) if $P((AB)/C) = P(A/C).P(B/C)$

- ▶ As a consequence $P(A/BC) = P(A/C)$

HW: Verify if events A and B are conditionally independent of event C (in the experiment of picking number randomly in $\{1, \dots, 10\}$)

Conditional independence – example

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H.

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability?

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability?
 $P(A) = 3/4$.

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability?
 $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H.

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability?
 $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability?

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads). The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability? $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability?
 $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.
- ▶ Event C: Coin 1 is chosen.

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability?
 $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.
- ▶ Event C: Coin 1 is chosen.
- ▶ What is $P(A/C)$ and $P(B/C)$?

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads). The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability? $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.
- ▶ Event C: Coin 1 is chosen.
- ▶ What is $P(A/C)$ and $P(B/C)$? $1/2$

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads). The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability? $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.
- ▶ Event C: Coin 1 is chosen.
- ▶ What is $P(A/C)$ and $P(B/C)$? $1/2$
- ▶ What is $P((A \cap B)/C)$?

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads).
The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability?
 $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.
- ▶ Event C: Coin 1 is chosen.
- ▶ What is $P(A/C)$ and $P(B/C)$? $1/2$
- ▶ What is $P((A \cap B)/C)$? $1/4$


Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads). The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability? $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.
- ▶ Event C: Coin 1 is chosen.
- ▶ What is $P(A/C)$ and $P(B/C)$? $1/2$
- ▶ What is $P((A \cap B)/C)$? $1/4$ Hence A and B are conditionally independent given C.

Conditional independence – example

- ▶ There are two coins, one fair and other fake (both heads). The experiment is to choose a coin uniformly and toss twice.
- ▶ Event A: First coin toss results in H. What is its probability? $P(A) = 3/4$.
- ▶ Event B: Second coin toss results in H. What is its probability? $P(B) = 3/4$.
- ▶ Event C: Coin 1 is chosen.
- ▶ What is $P(A/C)$ and $P(B/C)$? $1/2$
- ▶ What is $P((A \cap B)/C)$? $1/4$ Hence A and B are conditionally independent given C.
- ▶ Are A and B independent? HW

First principle of counting



Principle of counting

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them?

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them? $n!$

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them? $n!$
- ▶ Given n objects, how many distinct pairs can you form?

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them? $n!$
- ▶ Given n objects, how many distinct pairs can you form?

$${}^nC_2 = \binom{n}{2} = \frac{n!}{n-2!2!}.$$

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them? $n!$
- ▶ Given n objects, how many distinct pairs can you form?
 ${}^nC_2 = \binom{n}{2} = \frac{n!}{n-2!2!}.$
- ▶ In general, given n objects, we can make ${}^nC_k = \binom{n}{k} = \frac{n!}{n-k!k!}$ distinct combination of k objects.

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them? $n!$
- ▶ Given n objects, how many distinct pairs can you form?
 ${}^nC_2 = \binom{n}{2} = \frac{n!}{n-2!2!}.$
- ▶ In general, given n objects, we can make ${}^nC_k = \binom{n}{k} = \frac{n!}{n-k!k!}$ distinct combination of k objects.
- ▶ Note that in each combination or group of k objects, the ordering within each group is immaterial.

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them? $n!$
- ▶ Given n objects, how many distinct pairs can you form?
 ${}^nC_2 = \binom{n}{2} = \frac{n!}{n-2!2!}.$
- ▶ In general, given n objects, we can make ${}^nC_k = \binom{n}{k} = \frac{n!}{n-k!k!}$ distinct combination of k objects.
- ▶ Note that in each combination or group of k objects, the ordering within each group is immaterial. What if we also want to count this?

Principle of counting

- ▶ Given n objects, in how many ways can you arrange them? $n!$
- ▶ Given n objects, how many distinct pairs can you form?
 ${}^nC_2 = \binom{n}{2} = \frac{n!}{n-2!2!}.$
- ▶ In general, given n objects, we can make ${}^nC_k = \binom{n}{k} = \frac{n!}{n-k!k!}$ distinct combination of k objects.
- ▶ Note that in each combination or group of k objects, the ordering within each group is immaterial. What if we also want to count this?
- ▶ ${}^nP_k = {}^nC_k \times k!$

Experiments with Sampling

Experiments with Sampling

- ▶ Sampling: Sampling from a set means choosing an element from the set.

Experiments with Sampling

- ▶ Sampling: Sampling from a set means choosing an element from the set.
- ▶ Sampling uniformly at random: All items in the set have equal probability of being chosen.

Experiments with Sampling

- ▶ Sampling: Sampling from a set means choosing an element from the set.
- ▶ Sampling uniformly at random: All items in the set have equal probability of being chosen.
- ▶ Sampling can be with replacement or without replacement.

Experiments with Sampling

- ▶ Sampling: Sampling from a set means choosing an element from the set.
- ▶ Sampling uniformly at random: All items in the set have equal probability of being chosen.
- ▶ Sampling can be with replacement or without replacement.
- ▶ Sampling can be ordered or unordered.

Experiments with Sampling

- ▶ Sampling: Sampling from a set means choosing an element from the set.
- ▶ Sampling uniformly at random: All items in the set have equal probability of being chosen.
- ▶ Sampling can be with replacement or without replacement.
- ▶ Sampling can be ordered or unordered.
- ▶ In ordered sampling, $(a, b, c) \neq (c, b, a)$.

Experiments with Sampling

- ▶ Sampling: Sampling from a set means choosing an element from the set.
- ▶ Sampling uniformly at random: All items in the set have equal probability of being chosen.
- ▶ Sampling can be with replacement or without replacement.
- ▶ Sampling can be ordered or unordered.
- ▶ In ordered sampling, $(a, b, c) \neq (c, b, a)$.
- ▶ This leaves us with 4 combinations.
 1. Ordered sampling with replacement
 2. Ordered sampling without replacement
 3. Unordered sampling with replacement
 4. Unordered sampling without replacement

Ordered sampling with replacement

Ordered sampling with replacement

- ▶ Suppose you want to sample k out of n objects with replacement and where the ordering of the k objects matters.

Ordered sampling with replacement

- ▶ Suppose you want to sample k out of n objects with replacement and where the ordering of the k objects matters.
- ▶ Because we sample with replacement, repetition is allowed.

Ordered sampling with replacement

- ▶ Suppose you want to sample k out of n objects with replacement and where the ordering of the k objects matters.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?

Ordered sampling with replacement

- ▶ Suppose you want to sample k out of n objects with replacement and where the ordering of the k objects matters.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) n^k ?

Ordered sampling with replacement

- ▶ Suppose you want to sample k out of n objects with replacement and where the ordering of the k objects matters.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) n^k ?
- ▶ There are k positions and n choices for every position.

Ordered sampling with replacement

- ▶ Suppose you want to sample k out of n objects with replacement and where the ordering of the k objects matters.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) n^k ?
- ▶ There are k positions and n choices for every position.
- ▶ Total n^k .

Ordered sampling without replacement

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.
- ▶ Because we sample without replacement, repetition is not allowed.

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?
- ▶ There are k positions and $n - i + 1$ choices for every i^{th} position.

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?
- ▶ There are k positions and $n - i + 1$ choices for every i^{th} position.
- ▶ Total $n \times (n - 1) \times \dots (n - k + 1) =$

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?
- ▶ There are k positions and $n - i + 1$ choices for every i^{th} position.
- ▶ Total $n \times (n - 1) \times \dots (n - k + 1) = \frac{n!}{(n-k)!}$

Ordered sampling without replacement

- ▶ Suppose you want to sample k out of n objects now without replacement and where the ordering of the k objects matters.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?
- ▶ There are k positions and $n - i + 1$ choices for every i^{th} position.
- ▶ Total $n \times (n - 1) \times \dots (n - k + 1) = \frac{n!}{(n-k)!} = {}^n P_k$.

Unordered sampling without replacement

Unordered sampling without replacement

- ▶ Here you want to sample k out of n objects without replacement and the ordering of the k objects does not matter.

Unordered sampling without replacement

- ▶ Here you want to sample k out of n objects without replacement and the ordering of the k objects does not matter.
- ▶ Because we sample without replacement, repetition is not allowed.

Unordered sampling without replacement

- ▶ Here you want to sample k out of n objects without replacement and the ordering of the k objects does not matter.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?

Unordered sampling without replacement

- ▶ Here you want to sample k out of n objects without replacement and the ordering of the k objects does not matter.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?

Unordered sampling without replacement

- ▶ Here you want to sample k out of n objects without replacement and the ordering of the k objects does not matter.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?
- ▶ Essentially we want to count distinct k sized subsets from n objects without caring for ordering.

Unordered sampling without replacement

- ▶ Here you want to sample k out of n objects without replacement and the ordering of the k objects does not matter.
- ▶ Because we sample without replacement, repetition is not allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ a) nk ? b) $\binom{n}{k}$ c) k^n d) none ?
- ▶ Essentially we want to count distinct k sized subsets from n objects without caring for ordering.
- ▶ nC_k .

Unordered sampling with replacement

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matter.

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matter.
- ▶ Because we sample with replacement, repetition is allowed.

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matter.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matters.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ In any such sampling, any object i can appear atmost k times.

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matters.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ In any such sampling, any object i can appear atmost k times.
- ▶ Let x_i denote the number of times object i is chosen in k samples.

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matter.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ In any such sampling, any object i can appear at most k times.
- ▶ Let x_i denote the number of times object i is chosen in k samples.
- ▶ Then any sampling satisfies $\sum_{i=1}^n x_i = k$

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matter.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ In any such sampling, any object i can appear at most k times.
- ▶ Let x_i denote the number of times object i is chosen in k samples.
- ▶ Then any sampling satisfies $\sum_{i=1}^n x_i = k$
- ▶ How many solutions to the above equation tells you how many ways you can do the above sampling.

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matter.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ In any such sampling, any object i can appear at most k times.
- ▶ Let x_i denote the number of times object i is chosen in k samples.
- ▶ Then any sampling satisfies $\sum_{i=1}^n x_i = k$
- ▶ How many solutions to the above equation tells you how many ways you can do the above sampling.
- ▶ $\binom{n+k-1}{k}$

Unordered sampling with replacement

- ▶ Here you want to sample k out of n objects with replacement and the ordering of the k objects does not matter.
- ▶ Because we sample with replacement, repetition is allowed.
- ▶ How many ways can you choose k objects out of n this way?
- ▶ In any such sampling, any object i can appear at most k times.
- ▶ Let x_i denote the number of times object i is chosen in k samples.
- ▶ Then any sampling satisfies $\sum_{i=1}^n x_i = k$
- ▶ How many solutions to the above equation tells you how many ways you can do the above sampling.
- ▶ $\binom{n+k-1}{k}$ Think(HW).