


Quantum Fisher Information as a Thermal and Dynamical Probe in Frustrated spin system

Cooperators





Zi Yang Meng University of Hongkong Professor

Menghan Song University of Hongkong Ph.D. Candidate



Yong Baek Kim University of Toronto Professor

Zhengbang Zhou University of Toronto Ph.D. Candidate



Felix Desrochers Harvard University Postdoctoral Fellow



The k -producibility state $|\psi\rangle = \bigotimes_{i=0}^M |\psi_i\rangle$
 Entanglement depth m means $m \geq k_i, \forall i$



$|\psi\rangle = (|\uparrow\rangle_0 + |\downarrow\rangle_0) \otimes (|\uparrow\rangle_1 + |\downarrow\rangle_1) \otimes (|\uparrow\rangle_1 + |\downarrow\rangle_1) \otimes \dots$



$|\psi\rangle = (|\uparrow\downarrow\rangle_{01} - |\downarrow\uparrow\rangle_{01}) \otimes (|\uparrow\downarrow\rangle_{23} - |\downarrow\uparrow\rangle_{23}) \otimes \dots$



$|\psi\rangle = (|\uparrow\uparrow\uparrow\uparrow \dots\rangle + |\downarrow\downarrow\downarrow\downarrow \dots\rangle)$



$|\psi\rangle = (|\uparrow\downarrow\rangle_{01} - |\downarrow\uparrow\rangle_{01}) \otimes (|\uparrow\uparrow\uparrow\uparrow \dots\rangle + |\downarrow\downarrow\downarrow\downarrow \dots\rangle)$

Quantum Fisher information

$$F_Q[\rho; \hat{O}(\mathbf{q})] = 2 \sum_{i,j} \frac{(p_i - p_j)^2}{p_i + p_j} |\langle i | \hat{O}(\mathbf{q}) | j \rangle|^2,$$



with $\rho = \sum_i p_i |i\rangle\langle i|$ and $p_i = \exp(-E_i/T)/Z$.

And

$$f_{QFI} = F_Q/N < m (\hat{O}_{max} - \hat{O}_{min})^2$$

Also,




$$f_{QFI} = \int_0^\infty d\omega \, 4 \tanh\left(\frac{\omega\beta}{2}\right) \pi [1 - \exp(-\beta\omega)] A_{\hat{O}}(\beta, \omega)$$



Heisenberg spin chain

V. Menon, et al., [Phys. Rev. B 107, 054422 \(2023\)](#)


A. Scheie, et al., [Phys. Rev. B .103, 224434 \(2021\)](#)




Triangle Lattice

A. O. Scheie, et al., [Nat. Phys. 20, 74–81 \(2024\)](#)

1, Pyrochlore lattice arXiv:2510.14813 (2025)



2, Balents-Fisher-Girvin (BFG) model arXiv:2603.19951 (2026)



$$H = J_z \sum_{\langle i,j \rangle} S_i^z S_j^z - J_{\pm} \sum_{\langle i,j \rangle} (S_i^+ S_j^- + S_i^- S_j^+)$$

Ring Exchange in Quantum Spin Ice regime:

$$H_{ring} = J_{ring} \sum_{\langle i,j,k,l,m,n \rangle} (S_i^+ S_j^- S_k^+ S_l^- S_m^+ S_n^- + S_i^- S_j^+ S_k^- S_l^+ S_m^- S_n^+)$$





$$J_{ring} = \frac{12J_{\pm}^3}{J_z^2}, T \sim J_{ring}$$

If $J_z = 1, J_{\pm} = 0.045$, then $T \sim 0.0010935$

Sign problem:

Exact diagonalization


Gauge mean-field theory calculation



$$Z = \sum_{n=0}^{\infty} \frac{\beta^n}{n!} \sum_{\{S_n\}} \sum_{\{\alpha\}} \langle \alpha | \Pi(-H_b) | \alpha \rangle$$






Directed loop (DL) update:

$$C_1 \xrightarrow[1 \text{ pair of } S^+ S^-]{\text{evolution}} C_2$$




Multi-directed loop (MDL) update:

$$C_1 \xrightarrow[3 \text{ pairs of } S^+ S^-]{\text{evolution}} C_3$$



Thermal annealing process: $\beta = 1/T$ from 1 to 1500 with $\delta\beta = 1$



Exact diagonalization:


The microcanonical thermal pure quantum method (mTPQ)

Starting from a state $|\psi_0\rangle$ at $T \rightarrow 0$, the real time correlation is measured by

$$\langle \psi_0 | O(t) O(0) | \psi_0 \rangle = \langle \psi_0 | e^{iHt} O e^{-iHt} O | \psi_0 \rangle$$


Size limitation: $N = 16$


Geometry Preferences



QFI in $S_{\pm}$ channel, $L = 4$
 $\Gamma = (0,0,0)$ and $\Gamma' = (4\pi, 4\pi, 0)$

Understanding: The absence of
spontaneous symmetry breaking
 $[U(1)]$ in a finite system






$$\text{Ker} = \tanh\left(\frac{\omega\beta}{2}\right) [1 - \exp(-\beta\omega)]$$

$$f_{QFI} = \int_0^\infty d\omega \, 4\pi \text{Ker} A(\beta, \omega)$$

$$S(\beta) = \int_0^\infty d\omega \, A(\beta, \omega)$$








Neutron scattering cross section:
Cerium-based pyrochlore (global frame)

$$A^{DO}(\mathbf{q}, \omega) = \sum_{\mu, \nu} \left(\mathbf{z}_{\mu} \mathbf{z}_{\nu} - \frac{(\mathbf{z}_{\mu} - \mathbf{q})(\mathbf{z}_{\nu} - \mathbf{q})}{|\mathbf{q}|^2} \right) A_{\mu, \nu}^{\pm}(\mathbf{q}, \omega)$$

$$\mathbf{z}_0 = \frac{1}{\sqrt{3}}(1, 1, 1), \mathbf{z}_1 = \frac{1}{\sqrt{3}}(1, -1, -1)$$

$$\mathbf{z}_2 = \frac{1}{\sqrt{3}}(-1, 1, -1), \mathbf{z}_3 = \frac{1}{\sqrt{3}}(-1, -1, 1)$$




$$H = -J_{\pm} \sum_{\langle i,j \rangle} (S_i^+ S_j^- + S_i^- S_j^+) + \frac{J_z}{2} \sum_{hex} \left(\sum_{i \in hex} S_i^z \right)^2$$


Ring Exchange in BFG model regime:

$$H_{ring} = J_{ring} \sum_{\langle 0,1,2,3 \rangle} (S_0^+ S_1^- S_2^+ S_3^- + S_0^- S_1^+ S_2^- S_3^+)$$

$$J_{ring} = \frac{J_{\pm}^2}{J_z}, T \sim J_{ring}$$

If $J_z = 1, J_{\pm} = 0.07$, then $T \sim 0.0049$







$$f_Q = \lambda^{\Delta_Q} \phi \left(T \lambda^z, L^{-1} \lambda, h \lambda^{\frac{1}{\nu}} \right)$$


$$h = 0, z = 1, \Delta_Q = 2 - (1 + \eta)$$



$$\eta \approx 1.495 \text{ (XY*) and } \eta \approx 0.04 \text{ (XY)}$$

$$\Delta_Q = -0.495 \text{ (XY*) and } \Delta_Q = 0.96 \text{ (XY)}$$

$$T \rightarrow 0, \lambda = L, f_Q = \lambda^{\Delta_Q} \phi(T \lambda)$$







Ground state change

No Spin order

No dimer order

Entangle structure change





Thank you.

arXiv:2510.14813 (2025)

arXiv:2603.19951 (2026)



The Quantum Fisher Information is given by

$$F_Q[\rho; \hat{O}(\mathbf{q})] = 2 \sum_{i,j} \frac{(p_i - p_j)^2}{p_i + p_j} |\langle i | \hat{O}(\mathbf{q}) | j \rangle|^2,$$

with $\rho = \sum_i p_i |i\rangle\langle i|$ and $p_i = \exp(-E_i/T)/Z$ referring to a thermal state.

And the dynamic susceptibilities is given by

$$\chi(\omega, \hat{O}(\mathbf{q})) = \sum_{i,j} (p_i - p_j) |\langle i | \hat{O}(\mathbf{q}) | j \rangle|^2 \pi \delta(\omega - E_j + E_i).$$

Noting

$$\int_{-\infty}^{\infty} d\omega \tanh\left(\frac{\omega}{2T}\right) \pi \delta(\omega - E_i + E_j) = \tanh\left(\frac{E_i - E_j}{2T}\right) = \frac{p_j - p_i}{p_j + p_i}.$$

Thus

$$F_Q[\rho; \hat{O}(\mathbf{q})] = \frac{4}{\pi} \int_0^{\infty} d\omega \tanh\left(\frac{\omega}{2T}\right) \chi(\omega, \hat{O}(\mathbf{q})).$$

For given operator $\hat{O}$, such as $\hat{O} = \sum_r e^{iqr} \hat{S}^z(r)$, the corresponding QFI is defined as

$$f_{QFI} = F_Q/N = \int_0^\infty d\omega 4 \tanh\left(\frac{\omega\beta}{2}\right) \pi[1 - \exp(-\beta\omega)] A(\beta, \omega),$$

with A is the dynamic structure factor of operator $\hat{O}$,

If

$$f_{QFI} > m(O_{max} - O_{min})^2,$$

the corresponding state is at-least $(m + 1)$ -partite entangled.










$$\frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) \otimes \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle), m < 2$$



$$\frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) \otimes |\uparrow\rangle \otimes |\downarrow\rangle, m < 1$$

Philipp Hauke, et al., [Nature Phys 12, 778–782 \(2016\)](#)
 Luca Pezzé, et al., [Phys. Rev. Lett. 102, 100401 \(2009\)](#)



Heisenberg spin chain

V. Menon, et al., [Phys. Rev. B 107, 054422 \(2023\)](#)
 A. Scheie, et al., [Phys. Rev. B .103, 224434 \(2021\)](#)



Triangle Lattice

A. O. Scheie, et al., [Nat. Phys. 20, 74–81 \(2024\)](#)

What is the physical meaning of Quantum Fisher information's independence of the dynamic spectrum?

Quantum Cramér-Rao bound: Given a state ρ_0 , if a unitary transform is applied as $\rho_\theta = e^{-i\theta G} \rho_0 e^{i\theta G}$, with θ as the parameter to be estimated and G a Hermitian operator, and an observable O is measured to infer θ , then the estimation variance satisfies

$$\text{Var}(\theta) \geq \frac{1}{mF_Q}$$


F_Q can be observed from the dynamic spectra of G

m : independent measuring time

F_Q : Quantum Fisher Information:

$$F_Q = N f_{QFI}$$

$$f_{QFI} = \int_0^\infty d\omega \, 4 \tanh\left(\frac{\omega\beta}{2}\right) \pi [1 - \exp(-\beta\omega)] A_G(\beta, \omega)$$



For pure state $|\psi_0\rangle$, operation on two spin, such as $U(\theta) = \exp[-i(\sigma_1^z + \sigma_2^z)\theta]$, the Quantum Fisher Information is given by

$$F_Q = 4 \left[\left\langle \frac{\partial \psi_\theta}{\partial x} \middle| \frac{\partial \psi_\theta}{\partial x} \right\rangle - \left| \left\langle \psi_\theta \middle| \frac{\partial \psi_\theta}{\partial x} \right\rangle \right|^2 \right],$$


and the Cramér–Rao bound is $\Delta\theta^2 = \text{Var}(\theta) \geq 1/mF_Q$.

Heisenberg limit: Best entangled

$$\frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle),$$

$$F_Q = 4$$

$$\Delta\theta \propto \sqrt{\frac{1}{F_Q}} = \frac{1}{N}$$




Classical limit: Standard quantum

$$\frac{1}{2}(|\uparrow\rangle + |\downarrow\rangle) \otimes (|\uparrow\rangle + |\downarrow\rangle),$$

$$F_Q = 2$$

$$\Delta\theta \propto \sqrt{\frac{1}{F_Q}} = \frac{1}{\sqrt{N}}$$



For $A^z(\mathbf{q}, \omega)$, $(O_{max} - O_{min})^2 = \left(\frac{1}{2} - \left(-\frac{1}{2}\right)\right)^2 = 1$, therefore

$$f_{QFI, A^z} < m$$

For $A^\pm(\mathbf{q}, \omega) = A^x(\mathbf{q}, \omega) + A^y(\mathbf{q}, \omega)$, $(O_{max} - O_{min})^2 = 2 \left(\frac{1}{2} - \left(-\frac{1}{2}\right)\right)^2 = 2$, therefore

$$f_{QFI, A^\pm} < \frac{m}{2}$$

For $A^{DO}(\mathbf{q}, \omega) = A^{NSF}(\mathbf{q}, \omega) + A^{SF}(\mathbf{q}, \omega)$, where (SF/NSF) means spin-flip/non-spin-flip channel with respect to the neutron polarization vector $\mathbf{p}$. For $A^{NSF/SF}(\mathbf{q}, \omega)$,

$$(O_{max} - O_{min})^2 = \frac{1}{4} \sum_{\mu} \left| \Pi_{(N)SF}^{\mu} \right|^2,$$

where

$$\Pi_{NSF}^{\mu} = \mathbf{p} \mathbf{z}_{\mu}, \quad \Pi_{SF}^{\mu} = \mathbf{v} \mathbf{z}_{\mu}, \quad \mathbf{p} \mathbf{v} = 0$$



Thus

$$f_{QFI, A^{(N)SF}} < m \text{Tr} \left[\sum_{\mu} \mathbf{p}^T \mathbf{z}_{\mu}^T \mathbf{z}_{\mu} \mathbf{p} \right] = \frac{m}{3}$$

$$f_{QFI, A^{DO}} < \frac{2m}{3}$$

$$L = 4$$

$$\Gamma' = (4\pi, 4\pi, 0) \text{ and } X = (0, 0, 2\pi)$$



QFI in $S_{\pm}$ channel, $L = 4$
 $\Gamma = (0,0,0)$ and $\Gamma' = (4\pi, 4\pi, 0)$

Understanding (1): The absence of spontaneous symmetry breaking
 $[U(1)]$ in a finite system

Understanding (2): Operator chosen

