Extremal Problems in Combinatorics

Your Name

Abstract. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat voluptatem. Ut enim aeque doleamus animo, cum corpore dolemus, fieri tamen permagna accessio potest, si aliquod aeternum et infinitum impendere malum nobis opinemur. Quod idem licet transferre in voluptatem, ut postea variari voluptas distinguique possit, augeri amplificarique non possit. At etiam Athenis, ut e patre audiebam facete et urbane Stoicos irridente, statua est in quo a nobis philosophia defensa et collaudata est, cum id, quod maxime placeat, facere possimus, omnis voluptas assumenda est, omnis dolor repellendus. Temporibus autem quibusdam et aut officiis debitis aut rerum necessitatibus saepe eveniet, ut et voluptates repudiandae sint et molestiae non recusandae. Itaque earum rerum.

1. Preliminaries

1.1. Graphons

We follow the notation and terminology of Chatterjee's monograph [1], to which we refer the reader for proofs and further background. This section collects the definitions and facts from graphon theory that we will use throughout. Before discussing the theory, we fix our graph-theoretic terminology.

Definition 1.1 A **simple graph** is a pair $G = (V, E)$ where V is finite set, identified with $\{1, ..., |V|\}$, and E is a set of unordered pairs of distinct elements of V. We call elements of V vertices and elements of E edges. When we need to refer to the vertex and edge sets of a particular graph G, we can write $V(G)$ and $E(G)$. Let the set of simple graphs be $\mathcal{G}$.

Graphons were introduced by Lovász and Szegedy [2] as a limit object for sequences of dense simple graphs. Before giving the formal definition, we motivate the choice of object by describing how a graphon parametrizes a natural random graph model — a perspective that will make the equivalence relations in the formal definition feel inevitable rather than arbitrary.

Given a symmetric measurable function $W : [0, 1]^2 \to [0, 1]$ and an integer n, the ***W*-random graph** $G(n, W)$ on a vertex set $\{1, ..., n\}$ is constructed in two steps. First, independently sample n random variables $U_1, ..., U_n$ uniformly on $[0, 1]$ and assign the label U_i to vertex i. Secondly, independently on each pair $i < j$, include the edge $\{i, j\}$ with probability $W(U_i, U_j)$. The value $W(x, y)$ is then the connection probability between vertices labeled x and y and the

uniform distribution on $[0,1]$ is the distribution over the labels. This construction recovers the Erdös–Rényi model $G(n,p)$ when $W \equiv p$ is constant, and recovers the stochastic block model when W is a step function, which we will call finite podal.

With this perspective in mind, we define the space of graphons.

Definition 1.2 The space $\mathcal{W}$ of **graphons** is the set of Borel measurable functions $W : [0,1]^2 \to [0,1]$ satisfying $W(x,y) = W(y,x)$, modulo the equivalence relation identifying functions that agree almost everywhere.

The simplest types of graphons are finite podal graphons which are defined below.

Definition 1.3 W is a **finite podal graphon** if there exists a natural number N, a partition of $[0,1]$ by measurable sets of positive measure $I_1, ..., I_N$, and weights $w_{ij} \in [0,1]$ with $w_{ij} = w_{ji}$ for $i,j \in \{1,...,N\}$ such that

$$W = \sum_{i=1}^{N} \sum_{j=1}^{N} w_{ij} \mathbb{1}_{I_i \times I_j}. \tag{1}$$

We call I_i the **podes** of W and the values w_{ij} the block weights of W. When we wish to specify the number of podes, we call W an ***N*-podal graphon**.

The almost everywhere quotient deserves comment. If W and W' differ only on a set of Lebesgue measure zero in $[0,1]^2$, then almost surely no sampled pair (U_i, U_j) ever lands on the set of disagreement, and so $G(n,W)$ and $G(n,W')$ have identical distributions. Two graphons that agree almost everywhere are thus genuinely indistinguishable as random graph models, and the a.e. quotient reflects this equality rather than being a technical convenience (though this is true too).

The natural topology on $\mathcal{W}$ is induced by the **cut norm**.

Definition 1.4 *(Cut Metric)* For $W_1, W_2 \in \mathcal{W}$,

$$d_\square(W_1, W_2) = \sup_{S,T \subset [0,1]} \left| \int_{S \times T} (W_1(x,y) - W_2(x,y))\, \mathrm{d}x\, \mathrm{d}y \right| \tag{2}$$

where the supremum is taken over measurable sets.

The cut metric captures the idea that two graphons are close if, for every pair of label regions S and T, they assign the same edge mass to $S \times T$. However, $d_\square$ is too fine an equivalence for our purposes: it distinguishes graphons that ought to represent the same object. To see what additional equivalence is needed, consider the analogous issue for finite graphs.

To motivate the next quotient, recall the analogous situation for simple graphs. If G_1 and G_2 are simple graphs on the same vertex set and G_2 arises from G_1 by a permutation of vertices, that is there exists a bijection $\sigma : V(G_1) \to V(G_2)$ with $\{u,v\} \in E(G_1) \Leftrightarrow \{\sigma(u), \sigma(v)\} \in E(G_2)$, then we say G_1 is isomorphic to G_2. Any property of a graph that we care about is invariant under vertex relabeling.

The analogue for graphons is a **measure preserving bijection** $\sigma : [0,1] \to [0,1]$, which serves as a permutation of labeling space. Given such a σ and W, define $W^\sigma(x,y) = W(\sigma(x), \sigma(y))$. If $\sigma(U_1), ..., \sigma(U_n)$ are obtained from i.i.d. uniform samples $U_1, ..., U_n$, then they themselves are i.i.d. uniform on $[0,1]$. So W^σ generates the exact same distribution of random graphs as W. Just as isomorphic simple graphs differ only in how their vertices are labeled, W and W^σ differ only in how $[0,1]$ is labeled.

Definition 1.5 For $W_1, W_2 \in \mathcal{W}$,

$$\delta_\square(W_1, W_2) = \inf_\sigma d_\square(W_1, W_2) \tag{3}$$

where the infimum is taken over the all measure preserving bijections $\sigma : [0,1] \to [0,1]$.

The functional $\delta_\square$ is a pseudometric on $\mathcal{W}$. It descends to a genuine metric on a quotient space:

$$\widetilde{\mathcal{W}} = \mathcal{W}/(W_1 \sim W_2 \Leftrightarrow \delta_\square(W_1, W_2) = 0). \tag{4}$$

The central topological fact, due to Lovász and Szegedy, is the following.

Theorem 1.1 *(Graphon Compactness Theorem [2])* The metric space $\left(\widetilde{\mathcal{W}}, \delta_\square\right)$ is compact.

Throughout the remainder of the thesis, we will work implicitly in the quotient space $\widetilde{\mathcal{W}}$.

Definition 1.6 *(Graphon Blowup)* Let G be a simple graph on vertex set $\{1, ..., n\}$. Partition $[0,1]$ into intervals $I_1, ..., I_n$ of equal length, and define the graphon $W_G : [0,1]^2 \to \{0,1\}$ by

$$W_G(x,y) = \begin{cases} 1 \text{ if } (x,y) \in I_i \times I_j \text{ and } \{i,j\} \in E(G) \\ 0 \text{ otherwise} \end{cases}. \tag{5}$$

We call W_G the **graphon blowup** of G.

The graphon W_G is symmetric by construction and takes values in $\{0,1\}$. It encodes the adjacency matrix of G as a finite podal graphon with n equal-sized parts. The terminology "limit object" is justified by the fact that for any graphon W, the graphon blowup of $G(n, W)$ converges to W in cut metric almost surely as $n \to \infty$ [2]. Thus graphons simultaneously parametrize random graph models and arise as limits of dense graph sequences.

1.1.1. Homomorphism Densities

The primary numerical invariants of graphons are homomorphism densities, which generalize subgraph densities of finite graphs.

Definition 1.7 For a simple graph G with k vertices and a graphon W, the **homomorphism density** of G in W is

$$t(G, W) := \int_{[0,1]^k} \prod_{\{i,j\} \in E(G)} W\left(x_i, x_j\right) \mathrm{d}x_1 ... \mathrm{d}x_k \tag{6}$$

Two special cases are central to this thesis. The **edge density** of W is

$$e(W) = \|W\|_1 = \int_{[0,1]^2} W(x,y)\,\mathrm{d}x\,\mathrm{d}y \tag{7}$$

and the **triangle density** is

$$t(W) = t(K_3, W) = \int_{\{[0,1]^3\}} W(x,y)W(y,z)W(x,z)\,\mathrm{d}x\,\mathrm{d}y\,\mathrm{d}z \tag{8}$$

From the stochastic perspective, $t(G,W)$ has a clean interpretation. It is the limiting probability, as $n \to \infty$, that a fixed injection $V(G) \hookrightarrow \{1,...,n\}$ is a graph homomorphism from G into $G(n,W)$. In particular, $e(W)$ is the edge probability and $t(W)$ is the triangle probability of the associated random graph model.

A foundational property, which motivates the choice of cut topology, is that the homomorphism densities are continuous in $\widetilde{\mathcal{W}}$.

Theorem 1.2 *(Continuity of Homomorphism Density [2])* For every simple graph G, the map $t(G,\cdot)$ is continuous on $\left(\widetilde{\mathcal{W}}, \delta_\square\right)$.

This continuity is what makes graphons the correct limit objects for the problems we study: the quantities of interest, edge density, triangle density, and the homomorphism density $t(G,W)$ whose asymptotics we analyze, all vary continuously in the cut topology, so the extremal questions over $\widetilde{\mathcal{W}}$ are well posed.

1.1.2. Approximation by Finite Podal Graphons

We now return to the subclass of finite podal graphons, which will play a central role in the analysis to follow: they are often used as proxies for the full space of graphons. The following density result is what makes finite podal graphons important.

Theorem 1.3 *([2])* The set of finite podal graphons is dense in $\left(\widetilde{\mathcal{W}}, \delta_\square\right)$.

For finite podal graphons, the homomorphism density reduces to a finite sum of vertex to pode assignments.

Lemma 1.4 *(Reduction of Homomorphism Density)* Let W be a N-podal graphon with podes $\{S_i\}_{i=1}^N$ and edge weights w_{ij}. For any simple graph G,

$$t(G,W) = \sum_{\varphi: V(G)\to\{1,...,N\}} \left(\prod_{v\in V(G)} \mu\left(S_{\varphi(v)}\right)\right)\left(\prod_{\{i,j\}\in E(G)} w_{\varphi(i)\varphi(j)}\right) \tag{9}$$

Proof Let $V(G) = \{1,...,k\}$. Since $\{S_i\}_{i=1}^N$ partitions $[0,1]$, the product space $[0,1]^k$ is partitioned into cells $\prod_{i=1}^k S_{\varphi(i)}$ indexed by functions $\varphi : V(G) \to \{1,...,N\}$. Hence

$$\begin{aligned} t(G,W) &= \int_{[0,1]^k} \prod_{\{i,j\}\in E(G)} W\left(x_i, x_j\right)\mathrm{d}x_1 ... \mathrm{d}x_k \\ &= \sum_{\varphi: V(G)\to\{1,...,N\}} \int_{\prod_{a=1}^k S_{\varphi(a)}} \prod_{\{i,j\}\in E(G)} W\left(x_i, x_j\right)\mathrm{d}x_1 ... \mathrm{d}x_k. \end{aligned} \tag{10}$$

By the definition of W,

$$\begin{aligned} t(G,W) &= \sum_{\varphi:V(G)\to\{1,\dots,N\}} \int_{\prod_{a=1}^{k} S_{\varphi(a)}} \prod_{\{i,j\}\in E(G)} w_{\varphi(i)\varphi(j)}\,\mathrm{d}x_1\dots\mathrm{d}x_k \\ &= \sum_{\varphi:V(G)\to\{1,\dots,N\}} \left(\prod_{\{i,j\}\in E(G)} w_{\varphi(i)\varphi(j)}\right) \int_{\prod_{a=1}^{k} S_{\varphi(a)}} \mathrm{d}x_1\dots\mathrm{d}x_k \\ &= \sum_{\varphi:V(G)\to\{1,\dots,N\}} \left(\prod_{\{i,j\}\in E(G)} w_{\varphi(i)\varphi(j)}\right)\left(\prod_{v\in V(G)} \mu\big(S_{\varphi(v)}\big)\right) \end{aligned} \tag{11}$$

□

Specializing the reduction lemma to $G = K_2$ and $G = K_3$, for which $\varphi : V(G) \to \{1, ..., N\}$ are ordered pairs and triples respectively, gives the following expressions for the edge and triangle densities

$$\begin{aligned} e(W) &= \sum_{i,j=1}^{N} \mu(S_i)\mu\big(S_j\big)w_{ij} \\ t(W) &= \sum_{i,j,k=1}^{N} \mu(S_i)\mu\big(S_j\big)\mu(S_k)w_{ij}w_{ik}w_{jk} \end{aligned} \tag{12}$$

1.1.3. Razborov Triangle

The image of the function $(e,t) : \widetilde{\mathcal{W}} \to [0,1]^2$ where $W \to (e(W), t(W))$ is called the **Razborov Triangle $\boldsymbol{R}$**, after Razborov found the minimal triangle density for a given edge density in his seminal paper [3].

Theorem 1.5 *(Maximum Triangle Density)* For a graphon W with edge density ε

$$t(W) \le \varepsilon^{\frac{3}{2}}. \tag{13}$$

The bound is uniquely satisfied by $W = \mathbb{1}_{[0,\sqrt{\varepsilon}]^2}$.

Theorem 1.6 *(Minimum Triangle Density I (Mantel's Theorem))* For a graphon W with edge density $\varepsilon \in \left[0, \frac{1}{2}\right]$, $t(W) \ge 0$. An example of a graphon which achieves this lower bound is

$$W = 2\varepsilon\Big(\mathbb{1}_{[0,\frac{1}{2}]\times[\frac{1}{2},1]} + \mathbb{1}_{[\frac{1}{2},1]\times[0,\frac{1}{2}]}\Big) \tag{14}$$

Theorem 1.7 *(Minimum Triangle Density II [3])* For a graphon W with edge density $\varepsilon \in \left[1-\frac{1}{k}, 1-\frac{1}{k+1}\right]$ for $k \ge 2$

$$t(W) \ge \frac{k(k-1)}{(k+1)^2}\left(1+\sqrt{1-\varepsilon\frac{k+1}{k}}\right)^2\left(1-2\sqrt{1-\varepsilon\frac{k+1}{k}}\right) \tag{15}$$

Remark 1.8 Razborov proved this using Flag Algebras, which can be thought of as a dual object to graphons. The map $W \to (t(G,W))_{G\in\mathcal{G}}$ is a continuous injective embbedding into

$\prod_{G \in \mathcal{G}}[0,1]$, a foundational result of graph limit theory. Flag Algebras seek to understand how the collections of homomorphism densities behave. Furthermore, one can define an algebra over such objects to do interesting computations, which often reduce to semidefinite programming problems. See [3], [4], [5] for further reference on this topic.

Theorem 1.9 *(Minimum Triangle Density III [6])* Let $\varepsilon \in \left[1 - \frac{1}{k}, 1 - \frac{1}{k+1}\right]$. Let

$$c = \frac{1 + \sqrt{1 - \varepsilon \frac{k+1}{k}}}{k+1}. \tag{16}$$

Partition $[0,1]$ into k intervals $I_1, ..., I_k$ where $|I_i| = c$ for $i \in \{1, ..., k-1\}$ and $|I_k| = 1 - (k-1)c$. Define W a graphon as follows. We set $W = 1$ when $I_i \times I_j$ for all $i \neq j$. For $i < k$, $W = 0$ on $I_i \times I_i$. Finally, on $I_k \times I_k$ the graphon is not constrained, except to have the correct average value, and to not allow any triangles where all three vertices are in I_k. For such a graphon $t(W)$ achieves the lower bound in Theorem 1.7.

Remark 1.10 This strategy, partitioning the graphon such that one localized region exhausts the allowed triangle density while the remainder contributes to the edge density without forming triangles, will be directly reflected in the constructions later in this thesis.

Putting the results of Theorem 1.5, Theorem 1.6, Theorem 1.7, and Theorem 1.9 together, we see R looks like the region below in Figure 1.



Figure 1: The Razborov Triangle R: The shaded interior represents the achievable triangle densities t for a given edge density e.

Our region of interest is studying the behavior $T_{\max}$ infinitesimally close to $(0, 0.5] \times \{0\} \in R$.

1.2. Linear Programming

Lets switch gears a bit, and discuss Linear Programming. We will follow the notation and terminology of Vanderbei, R.J. [7]. A **Linear Program (LP)** is a optimization problem in

which a linear function is minimized or maximized subject to finitely many linear inequality or equality constraints. The general minimization form which we will use is

$$\min c^T x \text{ subject to } Ax \geq b, x \geq 0, \tag{17}$$

where $A \in \mathbb{R}^{m \times n}$, $c \in \mathbb{R}^n$, $b \in \mathbb{R}^m$ are the **input data** and $x \in \mathbb{R}^n$ is the vector of **variables**. The linear function is called the **objective function**, and the conditions $Ax \geq b$ and $x \geq 0$ (both interpreted component wise) are the **constraints**.

A vector $x \in \mathbb{R}^n$ which satisfies all the constraints is a **feasible point**; the set of all feasible points is the **feasible region**, which is a polyhedron. An LP is **feasible** if the feasible region is nonempty, and bounded if $c^T x$ is bounded below on the feasible region. The **optimal value** is the infimum $\boldsymbol{c^T x}$ over the feasible region, and a feasible point achieving this value is an **optimal solution**.

Other standard forms (maximization, $\leq$, equality constraints, or free variables) can be converted into the form above by standard transformations, and all results we state below apply equally in those forms.

Geometrically, an LP asks for the minimum of a linear function over a polyhedron. Since the objective is linear, whenever an optimal solution exists, one can be found at a vertex of the feasible region. This underlies much of the theory.

Example 1.11 Consider the LP $\min x + y$ subject to the constraints that

$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \geq \begin{pmatrix} 2 \\ 2 \end{pmatrix} \tag{18}$$

and $x, y \geq 0$. The feasible region is an unbounded polyhedron in the first quadrant lying above the lines $x + 2y = 2$ and $2x + y = 2$. The level sets of the objective function are lines of slope -1. Thus starting at $x + y = 0$ and then moving the line upward, the first point we hit of the feasible region is the intersection of the two boundary lines, namely $(x, y) = \left(\frac{2}{3}, \frac{2}{3}\right)$, with objective value $\frac{4}{3}$.



Figure 2: Graphical solution to the linear program.

Two features seen in the example will recur. First the optimum is obtained at a vertex of the feasible region. It will specifically occur at the intersection of two binding constraints. Second, although the input data are integers, the optimal solution has rational (non-integer) coordinates; this is typical of LPs arising from combinatorial problems and reflects the fact that an LP is often a relaxation of a discrete problem.

1.2.1. Existence of Solutions to an LP

Given an LP of the form above. We have 3 possibilities regarding its behavior:

1. The LP is **infeasible**: There exists no $x \in \mathbb{R}^n$ satisfying the constraints
2. The LP is **feasible** but **unbounded**: For all $M \in \mathbb{R}$, there exists $x \in \mathbb{R}^n_{\geq 0}$ such that $c^T x < M$.
3. The LP is **feasible and bounded**: $c^T x$ has a finite infimum over the feasible region

In the third case, it is not obvious a priori that the infimum is attained. The feasible region is closed (as an intersection of finitely many closed half-spaces) but it need not be compact (see Example 1.11), and a continuous function on a non-compact closed set need not attain its infimum. However a foundational result in the theory of LPs tells us that the infimum is always attained.

Theorem 1.12 *(Existence of Optimal Solutions)* If an LP is feasible and bounded, then it has an optimal solution; moreover, one can always be chosen at a vertex of the feasible region

The second part of the theorem gives the intuition on why the first part holds. Although the polyhedron is unbounded, its set of vertices form a finite set, and a minimum of a linear

function over the feasible region, if finite, must be attained at one of them. A full proof uses the structure of the polyhedra (every feasible point is a convex combination of the vertices plus a nonnegative combination of extreme rays) and the observation that the objective can only decrease by moving along the extreme ray if the LP is unbounded.

The LPs that occur in the thesis are of a particularly nice form. Their feasible regions are nonempty. Every LP we will consider has a feasible point obtained by setting all the variables sufficiently large. Furthermore, their objectives are bounded below by 0, since all the variables are nonnegative and the objective coefficients are nonnegative. Hence every LP we consider has an optimal solution.

1.2.2. Dual Linear Programs

Before defining the dual, we will motivate the definition a bit. Given an LP (which we will call the original LP)

$$\min c^T x \text{ subject to } Ax \geq b, x \geq 0, \tag{19}$$

we can ask for a bound on the optimal value? In other words, how do we prove that the optimal value is at least some number L.

Consider our earlier example. Can we prove the lower bound algebraically, rather than graphically? Here is a way: if we add the two constraints, we see that $3x + 3y \geq 4$ which gives $x + y \geq \frac{4}{3}$. Since $x + y$ is the objective function, we have shown that every feasible point has an objective value of at least 4/3, which matches our optimal value.

The move here was to take a **non-negative linear combination of the constraints** and then observe that the resulting inequality had the objective function on one side and a number on the other side. Whenever such a combination can be arranged, the right-hand side is a valid lower bound on the primal optimum. The dual LP formalizes and generalizes this idea.

Going back to our general LP, suppose $y \in \mathbb{R}^m_{\geq 0}$ is a vector. Multiplying the k-th constraint $\sum_k A_{ki} x_i \geq b_k$ by y_k and summing gives us

$$y^T A x \geq y^T b. \tag{20}$$

This inequality holds for every feasible x. Now, suppose further we choose y such that $A^T y \leq c$ componentwise. Then for all feasible $x \geq 0$:

$$c^T x \geq \left(A^T y\right)^T x = y^T A x \geq y^T b = b^T y \tag{21}$$

so $b^T y$ is a lower bound on the primal optimum. To get the best lower bound this way, we should maximize over all y where $A^T y \leq c$. So we get

$$\max b^T y \text{ subject to } A^T y \leq c, y \geq 0, \tag{22}$$

This LP is called the **dual** to the original, which we will now call **primal**.

Definition 1.8 Given the primal LP,

$$\min c^T x \text{ subject to } Ax \geq b, x \geq 0, \tag{23}$$

its **dual** is the LP

$$\max b^T y \text{ subject to } A^T y \leq c, y \geq 0, \tag{24}$$

Variables of the dual are called **dual variables** and constraints of the dual are called **dual constraints**. By construction there is one dual variable per primal constraint, and one dual constraint per primal variable.

The dual of the dual, formed by applying this transformation again (after converting the maximization to standard minimization form), is the primal. So **primal** and **dual** are symmetric. The argument above gives us the following theorem for free.

Theorem 1.13 *(Weak Duality)* Let x be a feasible point of the primal LP and y be a feasible point of the dual LP. Then

$$c^T x \geq b^T y. \tag{25}$$

In particular, the optimal value of the dual is at most the optimal value of the primal.

Weak duality tells us the dual gives a lower bound on the primal. The fundamental theorem of LP duality says that this bound is in fact tight.

Theorem 1.14 *(Strong Duality)* Suppose the primal and dual are both feasible. Then they have optimal solutions, and their optimal values are equal:

$$\min\{c^T x : Ax \geq b, x \geq 0\} = \max\{b^T y : A^T y \leq c, y \geq 0\} \tag{26}$$

Strong Duality is a deep result. Its proof relies on a classical theorem of linear inequalities called Farkas' lemma. We refer to Vanderbei, R.J. [7] for full treatment. For our purposes, Strong Duality will be the key structural tool in the thesis. It allows us to prove statements about the primal optimum by reasoning about the dual, and vice versa.

Linear programs of the form considered here, minimizing a linear functional over a polyhedron defined by combinatorially-indexed constraints, arise throughout combinatorics as the natural continuous analogues to discrete optimization problems over graphs. LP duality then provides a structural link between the problem and its dual that is often surprising and powerful. In Section 4 we will define a linear program $\alpha(G)$, whose constraints are indexed by the triangles of a graph G. Strong duality applied to $\alpha(G)$ will be essential to the upper bound arguments of later sections.

1.3. Finner's Generalized Holder Inequality

The following is a classical inequality in measure theory, together with a corollary that will be convenient for our application.

Theorem 1.15 *(Finner's Generalized Hölder Inequality [8])* Let $n \in \mathbb{N}$, and let $\{(X_i, \mathcal{M}_i, \mu_i)\}_{i=1}^n$ be a collection of measure spaces. Let $X = X_1 \times ... \times X_n$ be their full product, equipped with the product measure μ. For any nonempty subset of indices $S \subset [n]$, let $X_S =$

$\prod_S X_k$ denote the lower-dimensional product space with the corresponding product measure μ_S. Suppose we have m non-empty index subsets $S_1, ..., S_m \subset [n]$, and corresponding exponents $p_1, ..., p_m \in [1, \infty)$. Assume that for every coordinate $k \in \{1, ..., n\}$, the sum of $\frac{1}{p_j}$ over the subsets S_j that contain k is exactly 1:

$$\sum_{\{j:k \in S_j\}} \frac{1}{p_j} = 1 \tag{27}$$

For each $j \in [m]$, let $f_j : X_{S_j} \to \mathbb{R}$ be a measurable function such that $f_j \in L^{p_j}\left(X_{S_j}, \mu_{S_j}\right)$. Then, the product $\prod_{i=1}^m |f_i|$, evaluated as a function on the full space X, is integrable with respect to μ, and the following inequality holds:

$$\int_X \prod_{i=1}^m \left|f_i\left(x_{S_i}\right)\right| \mathrm{d}\mu \leq \prod_{i=1}^m \|f_i\|_{p_i} \tag{28}$$

Corollary 1.16 Let $n \in \mathbb{N}$, and let $\{(X_i, \mathcal{M}_i, \mu_i)\}_{i=1}^n$ be a collection of probability spaces. Let $X = X_1 \times ... \times X_n$ be their full product, equipped with the product measure μ. For any nonempty subset of indices $S \subset [n]$, let $X_S = \prod_S X_k$ denote the lower-dimensional product space with the corresponding product measure μ_S. Suppose we have m nonempty index subsets $S_1, ..., S_m \subset [n]$, and corresponding exponents $p_1, ..., p_m \in [1, \infty)$. Assume that for every coordinate $k \in \{1, ..., n\}$, the sum of $\frac{1}{p_j}$ over the subsets S_j that contain k is less than or equal to 1:

$$\sum_{\{j:k \in S_j\}} \frac{1}{p_j} \leq 1 \tag{29}$$

For each $j \in [m]$, let $f_j : X_{S_j} \to \mathbb{R}$ be a measurable function such that $f_j \in L^{p_j}\left(X_{S_j}, \mu_{S_j}\right)$. Then, the product $\prod_{i=1}^m |f_i|$, evaluated as a function on the full space X, is integrable with respect to μ, and the following inequality holds:

$$\int_X \prod_{i=1}^m \left|f_i\left(x_{S_i}\right)\right| \mathrm{d}\mu \leq \prod_{i=1}^m \|f_i\|_{p_i} \tag{30}$$

Proof Let $A = \{a_1, ..., a_k\}$ be the set of coordinates where

$$\sum_{\{j:a_i \in S_j\}} \frac{1}{p_j} < 1. \tag{31}$$

Let $S_{m+1} = \{a_1\}, ..., S_{m+k} = \{a_k\}$. Let

$$p_{m+i}^{-1} = 1 - \sum_{\{j:a_i \in S_j, j \leq m\}} p_j^{-1}, \tag{32}$$

and let $f_{m+i} = 1$ for all $i \in [k]$. Thus

$$\int_X \prod_{i=1}^{m} \left| f_i\left(x_{S_i}\right) \right| \mathrm{d}\mu = \int_X \prod_{i=1}^{m+k} \left| f_i\left(x_{S_i}\right) \right| \mathrm{d}\mu$$
$$\leq \prod_{i=1}^{m+k} \|f_i\|_{p_i} = \prod_{i=1}^{m} \|f_i\|_{p_i}, \tag{33}$$

where the first equality comes from the fact that $f_{S_{m+i}} \equiv 1$. The inequality comes from Theorem 1.15. The last inequality comes from the fact $\left\|f_{m+i}\right\|_{p_{m+i}} = \mu\left(X_{a_i}\right)^{\frac{1}{p_{a_i}}}$ (using the probability space assumption). □

1.4. Notation

Sets: For any natural number N, define $[N] := \{1, ..., N\}$. For a finite set S and a natural number $n < |S|$, we write $\binom{S}{n}$ for the set of all n-element subsets of S. Let $\mathbb{R}_{\geq 0} := [0, \infty)$, $\mathbb{R}_{>0} := (0, \infty)$. Let μ be the Lebesgue measure on $[0, 1]$ or on $[0, 1]^2$, where context will make it clear whether we are talking about $[0, 1]$ or $[0, 1]^2$.

Graphs: For a graph G, we will write $V(G)$ and $E(G)$ as the vertex and edge sets, respectively. An edge between vertices u and v is written as $\{u, v\}$ or sometimes uv when there is no ambiguity. For a subgraph $T \subset G$, we write $V(T)$ and $E(T)$ for its vertex and edge sets. We write $\mathcal{T}(G)$ for the set of triangles in G. A triangle in G is a 3-vertex subgraph of G isomorphic to K_3. For any natural number n, let K_n be a complete graph on n vertices, in other words $K_n := \left([n], \binom{[n]}{2}\right)$ as a vertex set and edge set pair. In particular, K_2 is an edge and K_3 is a triangle.

Graphons: For (ε, τ) in the Razborov Triangle, we will define the following spaces of graphons for convenience:

$$\begin{aligned} \widetilde{\mathcal{W}}(N) &:= \left\{ W \;\middle|\; W \in \widetilde{\mathcal{W}}, W \text{ is } N\text{-podal} \right\} \\ \widetilde{\mathcal{W}}(\varepsilon, \tau) &:= \left\{ W \;\middle|\; W \in \widetilde{\mathcal{W}}, e(W) = \varepsilon, t(W) \leq \tau \right\} \\ \widetilde{\mathcal{W}}(N; \varepsilon, \tau) &:= \widetilde{\mathcal{W}}(N) \cap \widetilde{\mathcal{W}}(\varepsilon, \tau). \end{aligned} \tag{34}$$

Note if (ε, τ) are not in the Razborov Triangle, $\widetilde{\mathcal{W}}(\varepsilon, \tau)$ will be empty, by definition.

Asymptotics: For functions $f, g : D \to \mathbb{R}_{>0}$ and a limit point $p \in D$, we write $f(x) = O(g(x))$ if there exists $C > 0$ such that $f(x) \leq Cg(x)$ in a punctured neighborhood of p. We write $f(x) = \Omega(g(x))$ if there exists $c > 0$ such that $f(x) \geq cg(x)$ in a punctured neighborhood of p. We write $f(x) = \Theta(g(x))$ if $f(x) = O(g(x))$ and $f(x) = \Omega(g(x))$.

2. The Extremal Problem

Having developed the relavent machinery, we will state the problem this thesis addresses. Fix a simple graph G, and a pair of real numbers $(\varepsilon, \tau) \in \mathrm{int}(R)$. We ask: among all graphons with

edge density ε and triangle density at most τ, what is the tightest upper bound we can get for the homomorphism density of G? That is, we study the quantity

$$T^G_{\max}(\varepsilon,\tau) \coloneqq \sup\Big\{t(G,W) \;\Big|\; W \in \widetilde{\mathcal{W}}(\varepsilon,\tau)\Big\} \tag{35}$$

and how it behaves near the boundary of the Razborov Triangle. In this thesis, we will seek to understand the asymptotics of $T_{\max}$ for $\tau \to 0$ and $\varepsilon \in \big(0, \frac{1}{2}\big]$.

Intuitively, this is an extremal question about how much subgraph structure G can survive in a graphon with a few triangles. The regime $\tau \to 0$ forces the graphon toward triangle-free behavior while holding the edge density, and the question is whether (and how quickly) G-density must also decay.

Before turning to the asymptotics, we note that $T^G_{\varepsilon,\tau}$ is a genuine maximum, and not just a supremum.

Proposition 2.1 For every simple graph G, every $(\varepsilon,\tau) \in R$, there exists a graphon $W \in \widetilde{\mathcal{W}}(\varepsilon,\tau)$ such that $t(G,W) = T^G_{\max}(\varepsilon,\tau)$.

Proof Let $F : \widetilde{\mathcal{W}} \to R \subset [0,1]^2$ where $F(W) = (e(W), t(W))$. By Theorem 1.2, e and t are continuous, meaning F is continuous. By definition, $\widetilde{\mathcal{W}}(\varepsilon,\tau) = F^{-1}(\varepsilon \times [0,\tau])$. $\varepsilon \times [0,\tau]$ is closed, and as the preimage of a closed set $\widetilde{\mathcal{W}}(\varepsilon,\tau)$ is closed. Since $\big(\widetilde{\mathcal{W}}, \delta_\square\big)$ is compact and all closed sets of a compact set are compact, $\widetilde{\mathcal{W}}(\varepsilon,\tau)$ is compact. Finally, the map $t(G,\cdot)$ is continuous by Theorem 1.2, and a continuous real-valued function on a nonempty compact set attains its supremum. Therefore, $T^G_{\max}(\varepsilon,\tau)$ is attained by some $W \in \widetilde{\mathcal{W}}(\varepsilon,\tau)$. $\square$

So we can redefine $T^G_{\max}(\varepsilon,\tau)$ as

$$T^G_{\max}(\varepsilon,\tau) \coloneqq \max\Big\{t(G,W) \;\Big|\; W \in \widetilde{\mathcal{W}}(\varepsilon,\tau)\Big\}. \tag{36}$$

To show $T^G_{\max}(\varepsilon,\tau) = \Theta(g(\tau))$ for some function g, we have to show both $O(g(\tau))$ and $\Omega(g(\tau))$. $\Omega(g(\tau))$ is a bit finicky but straight forward, for fixed $\varepsilon \in (0, 0.5]$ we just have to exhibit a continuous function $W : (0,\tau_*] \to \widetilde{\mathcal{W}}$ for some $\tau_* > 0$ where $\tau \to W_\tau \in \widetilde{\mathcal{W}}(\varepsilon,\tau)$. We show that $t(G, W_\tau) = g(\tau)$ as $\tau \to 0$. We do this in Proposition 7.2 and Proposition 7.3 for $\varepsilon \in (0, 0.5)$ and $\varepsilon \in 0.5$ respectively. $O(g(\tau))$ requires a bit more work, which is why we look for simplifications to give us intuition, and hopefully reductions of the problem.

3. Triangle Spanning Decomposition

To establish the upper bound,we first prove the result for a simpler family of graphs: those where every edge belongs to a triangle (see Proposition 7.1). In this section, we demonstrate that any arbitrary graph can be decomposed into a set of these triangle-spanning subgraphs, denoted $\mathcal{C}$, alongside a collection of isolated edges and vertices. Subsequently, in Section 7, we will show that these residual components do not asymptotically affect the homomorphism density, allowing us to cleanly reduce the general upper bound to the triangle-spanning case.

Definition 3.1 A simple graph G is triangle spanning if every edge of G lies in a triangle of G.

Definition 3.2 Let G be a simple graph. For triangles $S, T \in \mathcal{T}(G)$, $S \leftrightarrow T$ if theres a sequence $S = T_0, T_1, ..., T_k = T$ of triangles in G where $V(T_i) \cap V(T_{i+1}) \neq \emptyset$ for all i. Let $\mathcal{T}_1, ..., \mathcal{T}_n$ be the equivalence classes under $\leftrightarrow$. Define C_i a subgraph of G with vertex set

$$V(C_i) := \bigcup_{T \in \mathcal{T}_i} V(T) \tag{37}$$

and edge set

$$E(C_i) := \bigcup_{T \in \mathcal{T}_i} E(T). \tag{38}$$

Define the **bridge edges** $B := E(G) \setminus \bigcup_i E(C_i)$, and the **leftover vertices** $L := V(G) \setminus \bigcup_i V(C_i)$.

Lemma 3.1 The C_i are vertex disjoint and edge disjoint triangle spanning subgraphs of G, B consists of edges not lying in any triangle of G, and L be the vertices in no triangles, $\{E(C_i)\}_{i=1}^n \cup B$ is a partition of $E(G)$, and $\{V(C_i)\}_{i=1}^n \cup L$ is a partition of $V(G)$.

Proof

1. **B consists of edges not in any triangle of G**: Let $e \in E(G)$. If e lies in some triangle $T \in \mathcal{T}(G)$, then $T \in \mathcal{T}_i$ for some i, so $e \in E(T) \subset E(C_i)$. Hence $e \notin B$. Conversely, if $e \in C_i$, then by definition $e \in E(T)$ for some triangle $T \in \mathcal{T}_i \subset \mathcal{T}(G)$, so e lies in some triangle of G. Thus $e \in B$ if and only if e lies in no triangle of G. The same argument with vertices in place of edges shows $v \in L$ if and only if v lies in no triangle.
2. **$\{E(C_i)\}$ and $\{V(C_i)\}$ are pairwise disjoint**: Suppose there exists $e \in C_i \cap C_j$ for some $i, j \in [n]$. Then $e \in E(A)$ for some $A \in \mathcal{T}_i$ and $B \in \mathcal{T}_j$. Since A and B share an edge e, hence share two vertices, so $A \leftrightarrow B$ directly (chain of length 1). So $i = j$. There for $E(C_i)$ are pairwise disjoint. The same argument works for the vertices. Combining with the definitions for B and L, this gives us the partitions $E(G) = \sqcup_i E(C_i) \sqcup B$ and $V(G) = \sqcup_i V(C_i) \cup L$.
3. **Each C_i is triangle spanning**: By construction, $\forall e \in E(C_i)$ there exists $T \in \mathcal{T}_i$ such that $e \in E(T)$. We must show T is a triangle in C_i, but this is trivial. By construction, $V(T) \subset V(C_i)$ and $E(T) \subset E(C_i)$. So e is contained in a triangle T in C_i.

□

Definition 3.3 The C_i are the **maximal triangle-spanning components** of G, note by assumption $V(C_i)$. The triple $(\mathcal{C}, B, L)$, where $\mathcal{C} = \{C_i\}_{i=1}^n$, is the **triangle spanning decomposition** of G. Let $V(\mathcal{C}) := V(G) \setminus V$ and $E(\mathcal{C}) := E(G) \setminus B$.

Thus we will prove the desired upper bound for triangle spanning graphs first, then we will use the decomposition and reduce the general graph case to a finite number of triangle spanning subgraphs.

4. The Graph Parameter $\alpha(G)$

Let $\alpha : \mathcal{G} \to \mathbb{R}_{>0}$ where

$$\alpha(G) := \min\left\{ \sum_{v \in V(G)} x_v + \sum_{e \in E} y_e \right\} \tag{39}$$

where $x : V(G) \to [0, \infty)$, $y : E(G) \to [0, \infty)$ and for all triangles τ in G,

$$\sum_{v \in V(\tau)} x_v + \sum_{e \in E(\tau)} y_e \geq 1. \tag{40}$$

Lemma 4.1 Let G be a finite simple graph where $\mathcal{T}(G) \neq \emptyset$. Let x^* and y^* be the vertex and edge weight functions which are the solutions to $\alpha(G)$. There exists a $T \in \mathcal{T}(G)$ such that

$$\sum_{v \in V(T)} x_v^* + \sum_{e \in E(T)} y_e^* = 1 \tag{41}$$

Proof Assume the contrary that for all $T \in \mathcal{T}(G)$

$$\sum_{v \in V(T)} x_v^* + \sum_{e \in E(T)} y_e^* > 1. \tag{42}$$

Thus there exists a $\delta > 0$ such that for all $T \in \mathcal{T}(G)$

$$\sum_{v \in V(T)} x_v^* + \sum_{e \in E(T)} y_e^* \geq 1 + \delta. \tag{43}$$

Fix some $T_0 \in \mathcal{T}(G)$. Let $S = \{x_v^* \mid v \in V(T_0)\} \cup \{y_e^* \mid e \in E(T_0)\}$. Let $z = \max S$ and f be the object which gives z (ie if f is a vertex $z = x_f$ and if its an edge $z = y_f$).

$$(|V(G)| + |E(G)|)z \geq \sum_{v \in V(T_0)} x_v^* + \sum_{e \in E(T_0)} y_e^* \geq 1 + \delta. \tag{44}$$

Thus

$$z \geq \frac{1 + \delta}{|V(G)| + |E(G)|} > 0 \tag{45}$$

Let $\delta' = \min(z, \delta)$. Let $(\tilde{x}, \tilde{y})$ be identical to (x^*, y^*) except at f where $\tilde{z}_f = z_f - \delta'$. The only sums which change are the sums which f is a part of. For all $T \in \mathcal{T}(G)$ such that $f \in V(T)$ or $f \in E(T)$,

$$\begin{aligned} \sum_{i \in V(T)} \tilde{x_i} + \sum_{e \in E(T)} y_e^* &= \sum_{i \in V(T)} x_i^* + \sum_{e \in E(T)} y_e^* - \delta' \\ &\geq 1 + \delta - \delta' \geq 1 \end{aligned} \tag{46}$$

because $\delta \geq \delta'$ by definition. Finally,

$$\begin{aligned}\sum_{v\in V(G)} \tilde{x}_v + \sum_{e\in E(G)} y_e^* &= \sum_{v\in V(G)} x_v^* + \sum_{e\in E(G)} y_e^* - \delta' \\ &= \alpha(G) - \delta' < \alpha(G)\end{aligned} \tag{47}$$

contradicting optimality. □

Proposition 4.2 Let G be a simple graph. Let $(\mathcal{C}, B, L)$ be its triangle spanning decomposition. Then

$$\alpha(G) = \sum_{C\in\mathcal{C}} \alpha(C) \tag{48}$$

Proof Recall $\alpha(G)$ is the optimal value of

$$\sum_{v\in V(G)} x_v + \sum_{e\in E} y_e \tag{49}$$

subject to $x_v, y_e \geq 0$ and for all triangles $T \in \mathcal{T}(G)$,

$$\sum_{v\in V(T)} x_v + \sum_{e\in E(T)} y_e \geq 1. \tag{50}$$

By Lemma 3.1, no vertex of L lies in triangle, so for every $v \in L$, x_v appears in no constraint. Hence at the optimum we may take $x_v = 0$. By a symmetric argument, $y_e = 0$ for all $e \in B$ at the optimum. Thus $\alpha(G)$ equals the optimal value of

$$\sum_{v\in V(\mathcal{C})} x_v + \sum_{e\in E(\mathcal{C})} y_e \tag{51}$$

subject to the same constraints. By Lemma 3.1, because $V(\mathcal{C})$ and $E(\mathcal{C})$ can be partitioned into $\{V(C)\}_{C\in\mathcal{C}}$ and $\{E(C)\}_{C\in\mathcal{C}}$,

$$\sum_{v\in V(\mathcal{C})} x_v + \sum_{e\in E(\mathcal{C})} y_e = \sum_{C\in\mathcal{C}} \left(\sum_{v\in V(C)} x_v + \sum_{e\in E(C)} y_e \right). \tag{52}$$

Moreover, every triangle $T \in \mathcal{T}(G)$ lives entirely within a unique $C \in \mathcal{C}$. Conversely for all $C \in \mathcal{C}$, every triangle in C is a triangle in G. So the constraint set decouples: the constraints involving variables from a fixed $C \in \mathcal{C}$ are exactly the triangle constraints of $\mathcal{T}(C)$, and no constraint mixes variables from different components. Hence the LP seperates into a sum of n independent sub-LPs, one per component, and

$$\begin{aligned}\alpha(G) &= \sum_{C\in\mathcal{C}} \min\left\{ \sum_{v\in V(C)} x_v + \sum_{e\in E(C)} y_e \;\middle|\; x, y \geq 0, (x, y) \text{ feasible for } \mathcal{T}(C) \right\} \\ &= \sum_{C\in\mathcal{C}} \alpha(C)\end{aligned} \tag{53}$$

□

5. Finite Podal Case

Definition 5.1 For a simple graph G, let

$$T^G_{\max,N}(\varepsilon,\tau) := \max\{t(G,W) \mid W \text{ a N-podal graphon}, e(W)=\varepsilon, t(W)\le\tau\} \tag{54}$$

Lemma 5.1 *(Triangle Lemma)* For any N-podal graphon W with podes A_i with measure $\alpha_i \in (0,1)$ and edge weights $w_{ij} \in [0,1]$, if $t(W) \le \tau$, then for all $i,j,k \in [N]$ $w_{ij}w_{ik}w_{jk}\alpha_i\alpha_j\alpha_k \le \tau$

Proof Since

$$t(W) = \sum_{i,j,k\in[N]} w_{ij}w_{ik}w_{jk}\alpha_i\alpha_j\alpha_k \le \tau \tag{55}$$

and all the terms in the expansion are non-negative, thus

$$w_{ij}w_{ik}w_{jk}\alpha_i\alpha_j\alpha_k \le \tau \tag{56}$$

□

Proposition 5.2 Let G be a triangle spanning graph and let $V = |V(G)|$, then for all natural numbers $N \ge V$

$$T^G_{\max,N}(\varepsilon,\tau) = O\big(\tau^{\alpha(G)}\big) \tag{57}$$

for e and N fixed and $\tau \to 0$.

Proof Let $\{z^*_\tau\}_{\tau\in\mathcal{T}(G)}$ be optimal solution to the dual LP to α. By Strong Duality,

$$\alpha(G) = \sum_{T\in\mathcal{T}(G)} z^*_T. \tag{58}$$

Let W be any V-podal graphon where $e(W) = e$ and $t(W) \le \tau$. As a sum over all possible mappings $\varphi : V(G) \to [N]$

$$t(G,W) = \sum_{\varphi:V(G)\to[N]} \Pi_\varphi \tag{59}$$

where

$$\Pi_\varphi := \prod_{v\in V(G)} \alpha_{\varphi(v)} \prod_{e\in E(G)} w_{\varphi(e)}. \tag{60}$$

Fix an arbitary mapping φ. For each triangle $\tau \in \mathcal{T}$ with vertices (v_1, v_2, v_3) and edges (e_1, e_2, e_3) by Lemma 5.1:

$$w_{\varphi(e_1)}w_{\varphi(e_2)}w_{\varphi(e_3)}\alpha_{\varphi(v_1)}\alpha_{\varphi(v_2)}\alpha_{\varphi(v_3)} \le t \tag{61}$$

Thus,

$$\prod_{\tau\in\mathcal{T}}\left(\prod_{v\in V(\tau)}\alpha_{\varphi(v)}\prod_{e\in E(\tau)}w_{\varphi(e)}\right)^{z^*_\tau}\leq\prod_{\tau\in\mathcal{T}}t^{z^*_\tau}. \tag{62}$$

We can easily simplify the right hand side:

$$\prod_{\tau\in\mathcal{T}}\tau^{z^*_\tau}=\tau^{\sum_{\tau\in\mathcal{T}}z^*_\tau}=t^{\alpha(G)}. \tag{63}$$

For the left hand side:

$$\prod_{\tau\in\mathcal{T}}\left(\prod_{v\in V(\tau)}\alpha_{\varphi(v)}\prod_{e\in E(\tau)}w_{\varphi(e)}\right)^{z^*_\tau}=\left(\prod_{\tau\in\mathcal{T}}\prod_{v\in V(\tau)}\alpha^{z^*_\tau}_{\varphi(v)}\right)\left(\prod_{\tau\in\mathcal{T}}\prod_{e\in E(\tau)}w^*_{\varphi(e)}\right). \tag{64}$$

$\forall v\in V(G)$, when you collect their corresponding terms, they appear with an exponent of z^*_τ for every $\tau\in\mathcal{T}$ where $v\in\tau$. Similarly for the edges. Thus,

$$\prod_{\tau\in\mathcal{T}}\left(\prod_{v\in V(\tau)}\alpha_{\varphi(v)}\prod_{e\in E(\tau)}w_{\varphi(e)}\right)^{z^*_\tau}=\left(\prod_{v\in V(G)}\alpha^{\sum_{v\in\tau\in\mathcal{T}}z^*_\tau}_{\varphi(v)}\right)\left(\prod_{e\in E(G)}w^{\sum_{e\in\tau\in\mathcal{T}}z^*_\tau}_{\varphi(v)}\right) \tag{65}$$

By the constraints of the Dual LP, $\sum_{v\in\tau\in\mathcal{T}}z^*_\tau\leq 1$ and $\sum_{e\in\tau\in\mathcal{T}}z^*_\tau$. Since $\alpha_i\leq 1$ and $w_{ij}\leq 1$:

$$\begin{aligned}\Pi_\varphi=\prod_{v\in V(G)}\alpha_{\varphi(v)}\prod_{e\in E(G)}w_{\varphi(v)}&\leq\left(\prod_{v\in V(G)}\alpha^{\sum_{v\in\tau\in\mathcal{T}}z^*_\tau}_{\varphi(v)}\right)\left(\prod_{e\in E(G)}w^{\sum_{e\in\tau\in\mathcal{T}}z^*_\tau}_{\varphi(v)}\right)\\&\leq t^{\alpha(G)}\end{aligned} \tag{66}$$

Since this holds for all maps φ and there are $N^{|V(G)|}$ such maps (which is a constant with respect to τ):

$$T^G_{\max,N}(e,t)\leq N^{|V(G)|}t^{\alpha(G)}\Rightarrow T^G_{\max,N}(e,t)=O\big(t^{\alpha(G)}\big) \tag{67}$$

□

6. General Graphons with Edge Density in $\left(0,\frac{1}{2}\right]$

Proposition 6.1 Let G be a triangle spanning graph. $T^G_{\max}(e,\tau)=O\big(\tau^{\alpha(G)}\big)$

Proof Let $\{z^*_\tau\}_{\tau\in\mathcal{T}(G)}$ be optimal solution to the dual LP to α. By Strong Duality, $\alpha(G)=\sum_{\tau\in\mathcal{T}(G)}z^*_\tau$. For $x\in[0,1]^V$ and for all $T=\{i,j,k\}\in\mathcal{T}(G)$, let

$$x_T:=\begin{pmatrix}x_i\\x_j\\x_j\end{pmatrix}. \tag{68}$$

Let $W_\triangle : [0,1]^3 \to [0,1]$ where

$$W_\triangle(x_1, x_2, x_3) = W(x_1, x_2)W(x_2, x_3)W(x_1, x_3). \tag{69}$$

By the Dual LP,$\forall e \in E(G)$ $\sum_{T \ni e} z_e^* \leq 1$. Since $W \leq 1$, raising W to a power $n \in [0,1]$ can only increase the value. Thus

$$\prod_{ij \in E(G)} W(x_i, x_j) \leq \prod_{ij \in E(G)} W(x_i, x_j)^{\sum_{T \ni ij} z_T^*} = \prod_{T \in \mathcal{T}(G)} W_\triangle(x_T)^{z_T^*}, \tag{70}$$

where the last equality comes from grouping all the edges into their respective triangles. Thus,

$$t(G, W) \leq \int_{[0,1]^V} \prod_{T \in \mathcal{T}(G)} W_\triangle(x_T)^{z_T^*} \, \mathrm{d}\vec{x} \tag{71}$$

For all $T = \{i, j, k\} \in \mathcal{T}(G)$, let $S_T = T$, $p_T = (z_T^*)^{-1}$, and $f_T = W_\triangle(x_T)^{z_T}$. Thus by Corollary 1.16,

$$\begin{aligned}
t(G, W) &\leq \int_{[0,1]^V} \prod_{T \in \mathcal{T}(G)} f_T(x_T) \, \mathrm{d}\vec{x} \\
&\leq \prod_{T \in \mathcal{T}(G)} \|f_T\|_{p_T} \\
&= \prod_{T \in \mathcal{T}(G)} \left(\int_{[0,1]^3} \left(W_\triangle(x_1, x_2, x_3)^{z_T^*} \right)^{p_T} \mathrm{d}\vec{x} \right)^{p_T} \\
&= \prod_{T \in \mathcal{T}(G)} \left(\int_{[0,1]^3} W_\triangle(x_1, x_2, x_3) \, \mathrm{d}\vec{x} \right)^{z_T^*} \\
&= \prod_{T \in \mathcal{T}(G)} t(W)^{z_T^*} \\
&= t(W)^{\sum_{T \in \mathcal{T}(G)} z_T^*} \\
&= t(W)^{\alpha(G)} \leq \tau^{\alpha(G)}
\end{aligned} \tag{72}$$

Thus

$$t(G, W) = O\big(\tau^{\alpha(G)}\big) \tag{73}$$

□

7. General Graphs

Proposition 7.1 Let G be simple graph, where $V = |V(G)|$, and $(\mathcal{C}, B, L)$ is its triangle spanning decomposition. Then

$$T_{\max}^G(\varepsilon, \tau) = O\big(\tau^{\alpha(G)}\big) \tag{74}$$

Proof Let W be a graphon where $e(W) = e$ and $t(W) = \tau$. By definition $W \leq 1$, so we have the following for $\vec{x} \in [0,1]^V$

$$\begin{aligned}\prod_{\{i,j\}\in E(G)} W(x_i, x_j) &\leq \prod_{\{i,j\}\in E(\mathcal{C})} W(x_i, x_j) \\ &= \prod_{C\in\mathcal{C}} \prod_{\{i,j\}\in E(C)} W(x_i, x_j),\end{aligned} \tag{75}$$

where the last equality comes from Lemma 3.1. Thus

$$t(G, W) \leq \int_{[0,1]^V} \prod_{C\in\mathcal{C}} \prod_{\{i,j\}\in E(C)} W(x_i, x_j)\, \mathrm{d}\vec{x}. \tag{76}$$

Again note by Lemma 3.1, for all $v \in L$ their corresponding variable x_v occurs nowhere in the integrand. By the same lemma, since $\mathcal{C}$ forms a partition of $V(\mathcal{C})$ each vertex in $V(\mathcal{C})$ appears in one and only one maximal triangle spanning component. Thus,

$$\begin{aligned}t(G, W) &\leq \int_{[0,1]^{V-|L|}} \prod_{C\in\mathcal{C}} \prod_{\{i,j\}\in E(C)} W(x_i, x_j)\, \mathrm{d}\vec{x} \\ &= \prod_{C\in\mathcal{C}} \int_{[0,1]^{|V(C)|}} \prod_{\{i,j\}\in E(C)} W(x_i, x_j)\, \mathrm{d}\vec{x} \\ &= \prod_{C\in\mathcal{C}} t(C, W) \\ &\leq \prod_{C\in\mathcal{C}} T^C_{\max}(\varepsilon, \tau) \leq \prod_{C\in\mathcal{C}} \tau^{\alpha(C_i)}\end{aligned} \tag{77}$$

where the last inequality comes from Proposition 6.1. Thus

$$t(G, W) \leq \tau^{\sum_{C\in\mathcal{C}} \alpha(C)} = \tau^{\alpha(G)} \tag{78}$$

where the equality $\alpha(G) = \sum_{C\in\mathcal{C}} \alpha(C)$ comes from Proposition 4.2. Thus, if for all valid graphons W, $t(G, W) \leq \tau^{\alpha(G)}$ then

$$T^G_{\max}(\varepsilon, \tau) \leq \tau^{\alpha(G)} \tag{79}$$

and

$$T^G_{\max}(\varepsilon, \tau) = O\big(\tau^{\alpha(G)}\big) \tag{80}$$

for fixed $\varepsilon \in \big(0, \frac{1}{2}\big)$ and $\tau \to 0$. □

Proposition 7.2 Let G simple graph with $\mathcal{T}(G) \neq \emptyset$. Let $V = |V(G)|$ and $(\mathcal{C}, B, L)$ is its triangle spanning decomposition.

$$T^G_{\max,N}(\varepsilon, \tau) = \Omega\big(\tau^{\alpha(G)}\big) \tag{81}$$

for fixed $\varepsilon \in (0, 0.5)$ and fixed $N \geq V + 2$ and $\tau \to 0$.

Proof Let $\varepsilon \in (0, 0.5)$ be fixed and let $\tau > 0$ be the parameter in $T^G_{\max,N}(\varepsilon, \tau)$. Let $\gamma = \tau/6|\mathcal{T}(G)|$. Let $n = |\mathcal{C}|$, thus $\mathcal{C} = \{C_1, ..., C_n\}$ through a bijection with $[n]$. Let $V_C = |V(\mathcal{C})|$ and $V_L = |L|$.

Let B_0, B_1, B_2 be a partition of B where B_0 is the L-to-L edges, B_1 are the L-to-$V(\mathcal{C})$ edges, and B_2 are the $V(\mathcal{C})$-to-$V(\mathcal{C})$ edges.

Let (x^*, y^*) be optimal for $\alpha(G)$, with the convention that $x^*(L) = \{0\}$ and $y^*(B) = \{0\}$ (optimal, since these variables appear in no triangle constraint).

Fix any $a_0 \in \left(\sqrt{\varepsilon/2}, 0.5\right)$; this interval is nonempty because $\varepsilon < 0.5$.

Let $\nu \in (0, \nu_{\max}]$ where

$$\nu_{\max} := \min\left(\frac{1 - 2a_0}{V_L}, \sqrt{\frac{\varepsilon}{2|B_0| + 1}}\right). \tag{82}$$

Let

$$\sigma(\gamma) := \sum_{v \in V(\mathcal{C})} \gamma^{x_i^*}. \tag{83}$$

Note as $\gamma \to 0$, $\sigma(\gamma) \to 0$.

Let $\{S_i\}_{i=1}^N$ be pairwise disjoint measurable sets in $[0, 1]$ where:

$$\mu(S_i) = \begin{cases} \gamma^{x_i^*} \text{ if } i \in V(\mathcal{C}) \\ \nu \text{ if } i \in L \\ a(N) \text{ if } i \in \{V+1, V+2\} \\ b(\gamma) \text{ otherwise} \end{cases} \tag{84}$$

where

$$\begin{aligned} a(N) &:= \begin{cases} a_0 + \frac{1}{2}(1 - 2a_0 - \sigma(\gamma) - V_L\nu) \text{ if } N = V + 2 \\ a_0 \text{ otherwise} \end{cases} \\ b(\gamma) &:= \frac{1 - 2a_0 - \sigma(\gamma) - V_L\nu}{N - V - 2} \text{ when } (N \geq V + 2) \end{aligned} \tag{85}$$

Since $\gamma \to 0 \Rightarrow \sigma(\gamma) \to 0$ and $2a_0 + V_L\nu > 1$, there exists a $\gamma_0 > 0$ such that $b(\gamma) > 0$ for all $\gamma \in (0, \gamma_0]$, and thus a valid measure. Furthermore, by construction

$$\sum_{i=1}^N \mu(S_i) = 1 \tag{86}$$

So $\{S_i\}_{i=1}^N$ is a partition of $[0, 1]$.

Let

$$
\begin{aligned}
R(\gamma) &:= 2 \sum_{ij \in E(\mathcal{C})} \gamma^{x_i^* + x_j^* + y_{ij}^*}, \\
Q_0 &:= 2 \sum_{e \in B_0} \nu^2 = 2|B_0|\nu^2, \\
Q_1(\gamma) &:= 2 \sum_{\{i,j\} \in B_1} \nu \gamma^{x^*_{\text{non}-V_0}} \\
Q_2(\gamma) &:= 2 \sum_{\{i,j\} \in B_2} \gamma^{y_{ij}^*} \\
\beta(\gamma) &= \frac{\varepsilon - R(\gamma) - Q_0 - Q_1(\gamma) - Q_2(\gamma)}{2\mu(S_{V+1})\mu(S_{V+2})}
\end{aligned}
\tag{87}
$$

Let $w : [N]^2 \to [0,1]$ where

$$
w_{ij} = \begin{cases} \gamma^{y_{ij}^*} \text{ if } \{i,j\} \in E(\mathcal{C}) \\ 1 \text{ if } \{i,j\} \in B \\ \beta(\gamma) \text{ if } \{i,j\} = \{V+1, V+2\} \\ 0 \text{ otherwise} \end{cases}.
\tag{88}
$$

As $\gamma \to 0$,

$$
\beta(\gamma) \to \frac{\varepsilon - Q_0}{2a_0^2} = \frac{\varepsilon - 2|B_0|\nu^2}{2a_0^2} > 0.
\tag{89}
$$

Since $a_0^2 > \varepsilon/2$, the limit is less than 1. Hence there exists a $\gamma_1 \in (0, \gamma_0]$ such that $\beta(\gamma) \in (0,1)$ for all $\gamma \in (0, \gamma_1)$.

Let W be an N-podal graphon with parts $\{S_i\}_{i=1}^N$ and edge weights w_{ij}.

$$
\begin{aligned}
e(W) &= \sum_{i,j \in [N]} \mu(S_i)\mu(S_j) w_{ij} \\
&= 2\beta(\gamma)\mu(S_{V+1})\mu(S_{V+2}) + R(\gamma) + Q_0 + Q_1(\gamma) + Q_2(\gamma) \\
&= \varepsilon
\end{aligned}
\tag{90}
$$

Thus W has the correct edge density. For the triangle density,

$$
t(W_0) = \sum_{i,j,k \in [N]} \mu(S_i)\mu(S_j)\mu(S_k) w_{ij} w_{ik} w_{jk}.
\tag{91}
$$

The only way that w_{ij}, w_{ik}, w_{jk} are all nonzero is if $\{i,j,k\} \in \mathcal{T}(G)$. Triangles of G are contained in components $C_l \in \mathcal{C}$, so all three edges must be in $E(C_l) \subset E(\mathcal{C}) \subset$. No edge in B may participate. Thus

$$
\begin{aligned}
t(W_0) &= 6 \sum_{\{i,j,k\}\in\mathcal{T}(G)} \mu(S_i)\mu(S_j)\mu(S_k) w_{ij} w_{ik} w_{jk} \\
&= 6 \sum_{\{i,j,k\}\in\mathcal{T}(G)} \gamma^{x_i+x_j+x_k} \gamma^{y_{ij}+y_{jk}+y_{ik}} \\
&= 6 \sum_{T\in\mathcal{T}(G)} \gamma^{\sum_{i\in V(T)} x_i + \sum_{e\in E(T)} y_e} \\
&= 6 \sum_{T\in\mathcal{T}(G)} \gamma^{\sum_{i\in V(T)} x_i + \sum_{e\in E(T)} y_e}
\end{aligned} \tag{92}
$$

By the feasibility constraint on $\alpha(G)$, $\forall T \in \mathcal{T}$

$$
\sum_{i\in V(T)} x_i^* + \sum_{e\in E(T)} y_e^* \geq 1. \tag{93}
$$

Since $\gamma \to 0 \Rightarrow \gamma < 1$, that means for all $T \in \mathcal{T}$,

$$
\gamma^{\sum_{i\in V(T)} x_i + \sum_{e\in E(T)} y_e} \leq \gamma^1 \tag{94}
$$

Thus

$$
t(W_0) \leq 6|\mathcal{T}(G)|\gamma = \tau. \tag{95}
$$

Furthermore by Lemma 4.1, there exists a $T_0 \in \mathcal{T}(G)$ such that

$$
\sum_{v\in V(T_0)} x_v + \sum_{e\in E(T_0)} y_e = 1. \tag{96}
$$

Since all terms in the sum of $t(W)$ are positive, $t(W) \geq 6\tau = |\mathcal{T}(G)|^{-1}\tau$. Thus $t(W) = \Theta(\tau)$.

Now for $t(G, W)$, the proof follows identically to Proposition 5.2.

$$
t(G, W) \geq \gamma^{\alpha(G)} = \left(\frac{\tau}{6|\mathcal{T}(G)|}\right)^{\alpha(G)} = \frac{1}{(6|\mathcal{T}(G)|)^{\alpha(G)}} \tau^{\alpha(G)} \tag{97}
$$

Since W is a valid N-podal graphon with $e(W) = \varepsilon$ and $t(W) \leq \tau$, so W is feasible in the definition of $T^G_{\max,N}(\varepsilon, \tau)$. Therefore,

$$
T^G_{\max,N}(e, \tau) \geq t(G, W) \geq \frac{1}{(6|\mathcal{T}(G)|)^{\alpha(G)}} \tau^{\alpha(G)}, \tag{98}
$$

giving $T^G_{\max,N}(\varepsilon, \tau) = \Omega\big(\tau^{\alpha(G)}\big)$ as $\tau \to 0$. □

Proposition 7.3 Let G simple graph with $\mathcal{T}(G) \neq \emptyset$. Let $V = |V(G)|$ and $(\mathcal{C}, B, L)$ is its triangle spanning decomposition.

$$
T^G_{\max,V+2}\left(\frac{1}{2}, \tau\right) = \Omega\big(\tau^{\alpha(G)}\big) \tag{99}
$$

for $\tau \to 0$.

Proof Let $\tau > 0$ be the parameter in $T^G_{\max,N}\left(\frac{1}{2}, \tau\right)$. Let $\gamma = \tau/6|\mathcal{T}(G)|$. Let $n = |\mathcal{C}|$, thus $\mathcal{C} = \{C_1, ..., C_n\}$ through a bijection with $[n]$. Let $V_C = |V(\mathcal{C})|$ and $V_L = |L|$.

Let B_0, B_1, B_2 be a partition of B where B_0 is the L-to-L edges, B_1 are the L-to-$V(\mathcal{C})$ edges, and B_2 are the $V(\mathcal{C})$-to-$V(\mathcal{C})$ edges.

Let (x^*, y^*) be optimal for $\alpha(G)$, with the convention that $x^*(L) = \{0\}$ and $y^*(B) = \{0\}$ (optimal, since these variables appear in no triangle constraint).

Let $\nu \in (0, \nu_{\max}]$ where

$$\nu_{\max} := \min\left(\frac{1}{V_L + 1}, \sqrt{\frac{1}{4|B_0| + 1}}\right). \tag{100}$$

Let

$$\sigma(\gamma) := \sum_{v \in V(\mathcal{C})} \gamma^{x_i^*}. \tag{101}$$

Note as $\gamma \to 0$, $\sigma(\gamma) \to 0$.

Let $\{S_i\}_{i=1}^N$ be pairwise disjoint measurable sets in $[0, 1]$ where:

$$\mu(S_i) = \begin{cases} \gamma^{x_i^*} \text{ if } i \in V(\mathcal{C}) \\ \nu \text{ if } i \in L \\ a(\gamma) \text{ if } i \in \{V+1, V+2\} \end{cases} \tag{102}$$

where

$$a(\gamma) := \frac{1}{2}(1 - V_L\nu - \sigma(\gamma)) \tag{103}$$

Since $\gamma \to 0 \Rightarrow \sigma(\gamma) \to 0$ and $1 - V_L\nu > 0$, there exists a $\gamma_0 > 0$ such that $b(\gamma) > 0$ for all $\gamma \in (0, \gamma_0]$, and thus a valid measure. Furthermore, by construction

$$\sum_{i=1}^N \mu(S_i) = 1 \tag{104}$$

So $\{S_i\}_{i=1}^N$ is a partition of $[0, 1]$. Let

$$
\begin{aligned}
R(\gamma) &:= 2 \sum_{ij \in E(\mathcal{C})} \gamma^{x_i^* + x_j^* + y_{ij}^*}, \\
Q_0 &:= 2 \sum_{e \in B_0} \nu^2 = 2|B_0|\nu^2, \\
Q_1(\gamma) &:= 2 \sum_{\{i,j\} \in B_1} \nu \gamma^{x_{\text{non}-V_0}^*} \\
Q_2(\gamma) &:= 2 \sum_{\{i,j\} \in B_2} \gamma^{y_{ij}^*} \\
\beta(\gamma) &= \frac{0.5 - R(\gamma) - Q_0 - Q_1(\gamma) - Q_2(\gamma)}{2\mu(S_{V+1})\mu(S_{V+2})}
\end{aligned}
\tag{105}
$$

Let $w : [N]^2 \to [0,1]$ where

$$
w_{ij} = \begin{cases} \gamma^{y_{ij}^*} \text{ if } \{i,j\} \in E(\mathcal{C}) \\ 1 \text{ if } \{i,j\} \in B \\ \beta(\gamma) \text{ if } \{i,j\} = \{V+1, V+2\} \end{cases}. \tag{106}
$$

As $\gamma \to 0$,

$$
\beta(\gamma) \to \frac{0.5 - Q_0}{2a_0^2} = \frac{0.5 - 2|B_0|\nu^2}{2a_0^2} > 0. \tag{107}
$$

Since $a_0^2 > \varepsilon/2$, the limit is less than 1. Hence there exists a $\gamma_1 \in (0, \gamma_0]$ such that $\beta(\gamma) \in (0,1)$ for all $\gamma \in (0, \gamma_1)$.

Let W be an N-podal graphon with parts $\{S_i\}_{i=1}^N$ and edge weights w_{ij}.

$$
\begin{aligned}
e(W) &= \sum_{i,j \in [N]} \mu(S_i)\mu(S_j) w_{ij} \\
&= 2\beta(\gamma)\mu(S_{V+1})\mu(S_{V+2}) + R(\gamma) + Q_0 + Q_1(\gamma) + Q_2(\gamma) \\
&= 0.5
\end{aligned}
\tag{108}
$$

Thus W has the correct edge density. For $t(W) \leq \tau$ and

$$
t(G, W) \geq \frac{1}{(6|\mathcal{T}(G)|)^{\alpha(G)}} \tau^{\alpha(G)}, \tag{109}
$$

the proof follows identically to Proposition 7.2. Since W is a valid $(V+2)$-podal graphon with $e(W) = 0.5$ and $t(W) \leq \tau$, so W is feasible in the definition of $T^G_{\max,V+2}(\varepsilon, \tau)$. Therefore,

$$
T^G_{\max,V+2}\left(\frac{1}{2}, \tau\right) \geq t(G, W) \geq \frac{1}{(6|\mathcal{T}(G)|)^{\alpha(G)}} \tau^{\alpha(G)}, \tag{110}
$$

giving $T^G_{\max,V+2}\left(\frac{1}{2}, \tau\right) = \Omega\left(\tau^{\alpha(G)}\right)$ as $\tau \to 0$. □

Theorem 7.4 G be a simple graph where $\mathcal{T}(G) \neq 0$.

$$T^G_{\max}(e,\tau) = \Theta\big(\tau^{\alpha(G)}\big) \tag{111}$$

for $.e \in \big(0, \frac{1}{2}\big]$ and $\tau \to 0$.

Proof By Proposition 7.2 and Proposition 7.3, for $V = |V(G)|$

$$T^G_{\max,V+2}(\varepsilon,\tau) \geq c_1\tau^{\alpha(G)}. \tag{112}$$

Since

$$T^G_{\max,V+2}(\varepsilon,\tau) \leq T^G_{\max}(\varepsilon,\tau) \Rightarrow c_1\tau^{\alpha(G)} \leq T^G_{\max}(\varepsilon,\tau). \tag{113}$$

Combining this with Proposition 7.1, we have

$$T^G_{\max}(\varepsilon,\tau) = \Theta\big(\tau^{\alpha(G)}\big) \tag{114}$$

□

References

1. Chatterjee, S., others: Large deviations for random graphs. (2017)
2. Lovász, L., Szegedy, B.: Limits of dense graph sequences. Journal of Combinatorial Theory, Series B. 96, 933–957 (2006)
3. Razborov, A.A.: On the minimal density of triangles in graphs. Combinatorics, Probability and Computing. 17, 603–618 (2008)
4. Razborov, A.A.: Flag algebras: an interim report, (2013)
5. Silva, M.K., Sato, C.M., others: Flag algebras: A first glance. arXiv preprint arXiv:1607.04741. (2016)
6. Pikhurko, O., Razborov, A.: Asymptotic structure of graphs with the minimum number of triangles. Combinatorics, Probability and Computing. 26, 138–160 (2017)
7. Vanderbei, R.J.: Linear programming: foundations and extensions. Journal of the Operational Research Society. 49, 94 (1998)
8. Finner, H.: A generalization of Hölder's inequality and some probability inequalities. The Annals of probability. 1893–1901 (1992)